

UNIVERSITY OF CALIFORNIA
RIVERSIDE

THESIS TITLE

A Dissertation submitted in partial satisfaction
of the requirements for the degree of

Doctor of Philosophy

in

Mathematics

by

Alexander Pokorny

MONTH 2021

Dissertation Committee:

Dr. Peter Samuelson, Chairperson
Dr. Jacob Greenstein
Dr. Stephano Vidussi

The Dissertation of Alexander Pokorny is approved:

Committee Chairperson

University of California, Riverside

Acknowledgements

ACKNOWLEDGEMENTS

This section may be omitted.

DEDICATION

This section may be omitted.

ABSTRACT OF THE DISSERTATION

THESIS TITLE

by

Alexander Pokorny

Doctor of Philosophy, Graduate Program in Mathematics
University of California, Riverside, ***MONTH*** 2021
Dr. Peter Samuelson, Chairperson

The text of the abstract goes here. An abstract is required, and must not exceed 350 words.

Further details can be found in UCR's formatting guidelines:

<https://graduate.ucr.edu/document/format-guide-201920>

Contents

List of Figures	viii
List of Tables	ix
1 Introduction	1
2 Background	2
2.1 Skein Theory	2
Foundations and General Notions	2
HOMFLYPT and Kauffman Skein Modules	8
The Iwahori-Hecke Algebra and the BMW Algebra	8
Skein Algebras of the Annulus	8
Connections to Representation Theory	8
A Relative Skein Algebra of the Annulus	8
The HOMFLYPT Skein Algebra of the Torus	8
2.2 The Ring of Symmetric Functions	8
Character Rings of Classical Groups	8
Bases of Λ and Identities	8
3 The Kauffman Skein Algebra of the Torus	9
3.1 Power Sum Type Elements	9
3.2 All Relations	9
3.3 Perpendicular Relations	9
3.4 Main Theorem	9
3.5 Compatibility With the Kauffman Bracket Skein Algebra of the Torus	9
4 Closures of Minimal Idempotents in BMW_n	10
??	10
References	11

List of Figures

List of Tables

Chapter 1

Introduction

Summarize the main background and new results here.

Chapter 2

Background

2.1 Skein Theory

Foundations and General Notions

In this work, we will be forced to discuss a few different variants of skein modules. For this reason, it will be useful to first describe some general framework of skein theory so that each of these variants will be a special case. Unless otherwise stated, we will assume M is an oriented 3-manifold with boundary ∂M (possibly empty), Σ is an oriented surface, I is the real interval $[0, 1]$, R is a commutative and unital ring.

Definition 2.1.1. Let $T_1, T_2 : X \rightarrow M$ be smooth embeddings of a smooth manifold X into M . A **smooth ambient isotopy** $H : T_1 \Rightarrow T_2$ is a smooth homotopy of diffeomorphisms H_t such that $H_0 = \text{id}_M$ and $H_1 \circ T_1 = T_2$. Furthermore, we demand that the boundary ∂M is fixed by the homotopy.

The relation

$$T_1 \sim T_2 \text{ if and only if there exists a smooth ambient isotopy } H : T_1 \Rightarrow T_2$$

is an equivalence relation. The smoothness requirement is important when considering knots. Without it, all knots would fall into the same equivalence class.

Definition 2.1.2. Let N be a finite set of points contained in the boundary ∂M . An N -**tangle** in M (or just *tangle* for short) is the smooth ambient isotopy class of a smooth embedding

$$T : \underbrace{\bigsqcup_{j \in J} S^1}_{:=L} \sqcup \underbrace{\bigsqcup_{k \in K} I}_{:=B} \rightarrow M$$

for some finite sets J and K such that

1. the image of L lies in the interior of M ,
2. the image of the interior of B lies in the interior of M ,
3. the image of the boundary of B equals N .


If B is empty, then the result is called a **link** in M . Similarly, if L is empty, then it's called a **braid** in M .

One may also consider *oriented* or *framed* tangles by choosing an orientation or framing for each point in N and for each connected component of L and B such that the choices are compatible with each other with respect to the smooth embedding. If $M = \Sigma \times I$, then we will assume that the points in N are contained in $\Sigma \times \{\frac{1}{2}\}$ and that their framings are thought to be embedded orthogonally to $\Sigma \times \{*\}$.


We say a framed tangle in $\Sigma \times I$ has **blackboard framing** if the entire framing is embedded orthogonally to Σ . Every framed link in $\Sigma \times I$ is isotopic to one with a blackboard framing by turning each twist into a loop with a local blackboard framing:



This suggests that we may represent framed links in $\Sigma \times I$ as link diagrams on Σ . Indeed, equivalence under ambient isotopy is captured by the Reidemeister moves 2, 3, and a modified Reidemeister move 1:



* Add pictures of RII and RIII for good measure. Below are some examples of framed tangle diagrams.



Define the **writhe** of a tangle diagram is the number of positive crossings minus the number of negative crossings. It is easy to see that the Reidemeister moves above preserve the writhe of a diagram, so the concept is well defined. Writhe should be thought of as a grading on the free R -module on the set of tangles in a given space, which provides a good reason to work with framed links over ordinary links. Such a module is a main ingredient of this theory, so let's honor it with a proper discussion.

Definition 2.1.3. Let $\mathcal{T}(M, N)$ be the free R -module generated by the set of framed N -tangles in M . Analogously, we can define $\mathcal{T}^{or}(M, N)$ to be the free R -module generated by the set of oriented framed N -tangles in M . All definitions which are to follow in this subsection have an analogous definition using oriented tangles. Also, we will formally define $\mathcal{S}_R(\emptyset, \emptyset) := R$.

The construction $\mathcal{T}(-, -)$ is actually a symmetric monoidal functor $\mathcal{T} : \mathbf{C} \rightarrow R\text{-Mod}$ for a careful choice of category \mathbf{C} which we now describe. The objects of \mathbf{C} are pairs (M, N) of the same type as discussed previously. A morphism $(f, W) : (M', N') \rightarrow (M, N)$ is a pair of a smooth, orientation-preserving embedding $f : M' \rightarrow M$ such that $M - f(M')$, which is either a smooth 3-manifold or the empty set, and choice of $W \in \mathcal{T}(M - f(M'), N \sqcup f(N'))$

(unless $M - f(M')$ is empty, in which case W is a formal symbol for the "empty link" in the empty set). Composition is given by $(g, W') \circ (f, W) = (g \circ f, W' \cup W)$, which is associative since \circ and \cup are associative.

* Give a picture of composition in \mathcal{C} .

The induced map denoted $W : \mathcal{T}(M', N') \rightarrow \mathcal{T}(M, N)$ is a linear map defined by $W(T) = W \cup T$, and we will refer to such a linear map W as a **wiring**. We are abusing notation by denoting this linear map by W , but it should be clear from the context what f is since it is technically encoded in the data of the element $W \in \mathcal{T}(M - f(M'), N \sqcup f(N'))$. It is a stimulating exercise in verifying definitions to check that \mathcal{T} preserves composition and identity morphisms, making and so \mathcal{T} is functorial. \mathcal{C} can now be equipped with a symmetric monoidal structure via disjoint union. It is clear that

$$\mathcal{S}_R(M \sqcup M', N \sqcup N') \cong \mathcal{S}_R(M, N) \otimes_R \mathcal{S}_R(M', N')$$

for any sets of framed points $N \subset \partial M$ and $N' \subset \partial M'$. The unit is given by the object $(\emptyset, \emptyset) \in \mathcal{C}$ and define $\mathcal{T}(\emptyset, \emptyset) := R$, which makes \mathcal{T} a symmetric monoidal functor.

Definition 2.1.4. Let B be the smooth closed 3-ball, N_B be some boundary points of B , and let $X \subset \mathcal{T}(B, N_B)$ be some (typically finite) set, which we will call a set of **skein relations**. Given any tangle module $\mathcal{T}(M, N_M)$, there exists a submodule $\mathcal{I}(X)$ generated by the set

$$\{W(x) \mid x \in X \text{ and } W : \mathcal{T}(B, N_B) \rightarrow \mathcal{T}(M, N_M) \text{ is a wiring diagram}\}.$$

A quotient of the form $\mathcal{S}_X(M, N) := \mathcal{T}(M, N) / \mathcal{I}(X)$ is called a **skein module** of M relative to N . If $N = \emptyset$ is the empty set, we may use the notation $\mathcal{S}_X(M) := \mathcal{S}_X(M, \emptyset)$. Similar definitions may be given using oriented and/or unframed tangles instead.

The construction $\mathcal{S}_X(-, -)$ is a functor in the same way that $\mathcal{T}(-, -)$ is; a smooth embedding $f : M \rightarrow M'$ and an element $W \in \mathcal{S}_X(M - f(M'), N \sqcup f(N'))$ defines a linear map

$W : \mathcal{S}_X(M, N) \rightarrow \mathcal{S}_X(M', N')$. In fact, the quotient maps $\alpha_{(M, N)} : \mathcal{T}(M, N) \rightarrow \mathcal{S}_X(M, N)$ yield a natural transformation. In other words, given a morphism $(M, N) \rightarrow (M', N')$ in \mathbb{C} , the diagram

$$\begin{array}{ccc} \mathcal{T}(M, N) & \xrightarrow{W} & \mathcal{T}(M', N') \\ \downarrow \alpha_{(M, N)} & & \downarrow \alpha_{(M', N')} \\ \mathcal{S}_X(M, N) & \xrightarrow{W} & \mathcal{S}_X(M', N') \end{array}$$

commutes.

For any oriented surface Σ , we can define a category $\text{Skein}_X(\Sigma)$. The objects of this category are finite sets of framed points N in Σ , and the morphisms $N \rightarrow N'$ are elements of $\mathcal{S}_X(\Sigma \times I, (N \times \{0\}) \sqcup (N' \times \{1\}))$, so the category is R -linear. Write composition of morphisms by concatenation. If $y : N \rightarrow N'$ and $z : N' \rightarrow N''$ are morphisms, then their composite $yz : N \rightarrow N''$ is constructed by gluing z on y through N' and rescaling the interval coordinate appropriately.

* [Picture of composition in \$\text{Skein}_X\(\Sigma\)\$.](#)

The endomorphism algebras in this category are called **skein algebras** and are denoted by $\mathcal{S}_X(\Sigma, N) := \mathcal{S}_X(\Sigma \times I, (N \times \{0\}) \sqcup (N \times \{1\}))$. If N is the empty set, then we reduce the notation to simply $\mathcal{S}_X(\Sigma)$.

If $f : \Sigma \rightarrow \Sigma'$ is a smooth, orientation-preserving embedding of surfaces, then there is an induced functor

$$f_* : \text{Skein}_X(\Sigma') \rightarrow \text{Skein}_X(\Sigma)$$

defined on objects by $f_*(N) = f(N)$ and on morphisms in the following way. First, extend f trivially to $f \times \text{id}_I : \Sigma \times I \rightarrow \Sigma' \times I$. Then, in the skein algebra of the complement of the image of $f \times \text{id}_I$, choose the multiplicative identity element $e \in \mathcal{S}_X(\Sigma' - \text{Im}(f))$ which is the empty tangle. The pair $(f \times \text{id}_I, e)$ is an object in the category \mathbb{C} , which gives rise to a wiring

$$e : \mathcal{S}_X(\Sigma \times I, (N \times \{0\}) \sqcup (N' \times \{1\})) \rightarrow \mathcal{S}_X(\Sigma' \times I, (f(N) \times \{0\}) \sqcup (f(N') \times \{1\}))$$

via the functor \mathcal{S}_X . Now we may define what f_* does to morphisms: $f_*(y) = e(y)$ for any $y \in \mathcal{S}_X(\Sigma \times I, (N \times \{0\}) \sqcup (N' \times \{1\}))$.

* [Picture of how \$f_*\$ works on morphisms.](#)

It is clear that f_* preserves composition. In particular, f_* defines algebra homomorphisms on the skein algebras

$$e : \mathcal{S}_X(\Sigma, N) \rightarrow \mathcal{S}_X(\Sigma', f(N)).$$

* [We use this type of algebra homomorphism when we embed the annulus into the torus.](#)

The above homomorphisms are a special case of a more general type of map. If N is a set of framed points on Σ , then a smooth embedding $f : \Sigma \rightarrow \partial M$ induces a $\mathcal{S}_X(\Sigma, N)$ -module structure on $\mathcal{S}_X(M, N')$ for any N' with $f(N) \subseteq N'$. The action is given by “pushing tangles in through the boundary”. In other words, the pre-composition of a smooth embedding of a collar neighborhood $g : \partial M \times I \rightarrow M$ with $f \times \text{id}_I : \Sigma \times I \rightarrow \partial M \times I$ induces a bilinear map

$$\mathcal{S}_X(\Sigma, N) \times \mathcal{S}_X(M, N') \rightarrow \mathcal{S}_X(M, N')$$

because M minus a collar neighborhood is diffeomorphic to itself. Alternatively, a choice of element in $\mathcal{S}_X(\Sigma, N')$ produces a wiring $\mathcal{S}_X(M, N') \rightarrow \mathcal{S}_X(M, N')$.

* [Picture of action.](#)

HOMFLYPT and Kauffman Skein Modules

The Iwahori-Hecke Algebra and the BMW Algebra

Skein Algebras of the Annulus

Connections to Representation Theory

A Relative Skein Algebra of the Annulus

The HOMFLYPT Skein Algebra of the Torus

2.2 The Ring of Symmetric Functions

Character Rings of Classical Groups

Bases of Λ and Identities

Chapter 3

The Kauffman Skein Algebra of the Torus

3.1 Power Sum Type Elements

3.2 All Relations

3.3 Perpendicular Relations

3.4 Main Theorem

3.5 Compatibility With the Kauffman Bracket Skein Algebra of the Torus

Chapter 4

Closures of Minimal Idempotents in BMW_n

??

References