

than one way to represent a rational as such a ratio, e.g., $6/4 = 3/2$.

Since ten = 10, we will now use 10 instead of ten throughout, as is customary.

Exercise B.1.1. The purpose of this exercise is to demonstrate that the procedure of long addition taught to you in elementary school is actually valid. Let $A = a_n \dots a_0$ and $B = b_m \dots b_0$ be positive integer decimals. Let us adopt the convention that $a_i = 0$ when $i > n$, and $b_i = 0$ when $i > m$; for instance, if $A = 372$, then $a_0 = 2$, $a_1 = 7$, $a_2 = 3$, $a_3 = 0$, $a_4 = 0$, and so forth. Define the numbers c_0, c_1, \dots and $\varepsilon_0, \varepsilon_1, \dots$ recursively by the following *long addition algorithm*.

- We set $\varepsilon_0 := 0$.
- Now suppose that ε_i has already been defined for some $i \geq 0$. If $a_i + b_i + \varepsilon_i < 10$, we set $c_i := a_i + b_i + \varepsilon_i$ and $\varepsilon_{i+1} := 0$; otherwise, if $a_i + b_i + \varepsilon_i \geq 10$, we set $c_i := a_i + b_i + \varepsilon_i - 10$ and $\varepsilon_{i+1} = 1$. (The number ε_{i+1} is the “carry digit” from the i^{th} decimal place to the $(i+1)^{\text{th}}$ decimal place.)

Prove that the numbers c_0, c_1, \dots are all digits, and that there exists an l such that $c_l \neq 0$ and $c_i = 0$ for all $i > l$. Then show that $c_l c_{l-1} \dots c_1 c_0$ is the decimal representation of $A + B$.

Note that one could in fact use this algorithm to *define* addition, but it would look extremely complicated, and to prove even such simple facts as $(a+b)+c = a+(b+c)$ would be rather difficult. This is one of the reasons why we have avoided the decimal system in our construction of the natural numbers. The procedure for long multiplication (or long subtraction, or long division) is even worse to lay out rigorously; we will not do so here.

B.2 The decimal representation of real numbers

We need a new symbol: the *decimal point* “.”.

Definition B.2.1 (Real decimals). A *real decimal* is any sequence of digits, and a decimal point, arranged as

$$\pm a_n \dots a_0 . a_{-1} a_{-2} \dots$$

which is finite to the left of the decimal point (so n is a natural number), but infinite to the right of the decimal point, where \pm is either $+$ or $-$, and $a_n \dots a_0$ is a natural number decimal (i.e., either a positive integer decimal, or 0). This decimal is equated to the real number

$$\pm a_n \dots a_0.a_{-1}a_{-2} \dots \equiv \pm 1 \times \sum_{i=-\infty}^n a_i \times 10^i.$$

The series is always convergent (Exercise B.2.1). Next, we show that every real number has at least one decimal representation:

Theorem B.2.2 (Existence of decimal representations). *Every real number x has at least one decimal representation*

$$x = \pm a_n \dots a_0.a_{-1}a_{-2} \dots$$

Proof. We first note that $x = 0$ has the decimal representation $0.000 \dots$. Also, once we find a decimal representation for x , we automatically get a decimal representation for $-x$ by changing the sign \pm . Thus it suffices to prove the theorem for positive real numbers x (by Proposition 5.4.4).

Let $n \geq 0$ be any natural number. From the Archimedean property (Corollary 5.4.13) we know that there is a natural number M such that $M \times 10^{-n} > x$. Since $0 \times 10^{-n} \leq x$, we thus see that there must exist a natural number s_n such that $s_n \times 10^{-n} \leq x$ and $s_n++ \times 10^{-n} > x$. (If no such natural number existed, one could use induction to conclude that $s \times 10^{-n} \leq x$ for all natural numbers s , contradicting the Archimedean property.)

Now consider the sequence s_0, s_1, s_2, \dots . Since we have

$$s_n \times 10^{-n} \leq x < (s_n + 1) \times 10^{-n}$$

we thus have

$$(10 \times s_n) \times 10^{-(n++)} \leq x < (10 \times s_n + 10) \times 10^{-(n++)}.$$

On the other hand, we have

$$s_{n+1} \times 10^{-(n+1)} \leq x < (s_{n+1} + 1) \times 10^{-(n+1)}$$

and hence we have

$$10 \times s_n < s_{n+1} + 1 \text{ and } s_{n+1} < 10 \times s_n + 10.$$

From these two inequalities we see that we have

$$10 \times s_n \leq s_{n+1} \leq 10 \times s_n + 9$$

and hence we can find a digit a_{n+1} such that

$$s_{n+1} = 10 \times s_n + a_n$$

and hence

$$s_{n+1} \times 10^{-(n+1)} = s_n \times 10^{-n} + a_{n+1} \times 10^{-(n+1)}.$$

From this identity and induction, we can obtain the formula

$$s_n \times 10^{-n} = s_0 + \sum_{i=0}^n a_i \times 10^{-i}.$$

Now we take limits of both sides (using Exercise B.2.1) to obtain

$$\lim_{n \rightarrow \infty} s_n \times 10^{-n} = s_0 + \sum_{i=0}^{\infty} a_i \times 10^{-i}.$$

On the other hand, we have

$$x - 10^{-n} \leq s_n \times 10^{-n} \leq x$$

for all n , so by the squeeze test (Corollary 6.4.14) we have

$$\lim_{n \rightarrow \infty} s_n \times 10^{-n} = x.$$

Thus we have

$$x = s_0 + \sum_{i=0}^{\infty} a_i \times 10^{-i}.$$

Since s_0 already has a positive integer decimal representation by Theorem B.1.4, we thus see that x has a decimal representation as desired. \square

There is however one slight flaw with the decimal system: it is possible for one real number to have two decimal representations.

Proposition B.2.3 (Failure of uniqueness of decimal representations). *The number 1 has two different decimal representations: $1.000\dots$ and $0.999\dots$*

Proof. The representation $1 = 1.000\dots$ is clear. Now let's compute $0.999\dots$. By definition, this is the limit of the Cauchy sequence

$$0.9, 0.99, 0.999, 0.9999, \dots$$

But this sequence has 1 as a formal limit by Proposition 5.2.8. \square

It turns out that these are the only two decimal representations of 1 (Exercise B.2.2). In fact, as it turns out, all real numbers have either one or two decimal representations - two if the real is a terminating decimal, and one otherwise (Exercise B.2.3).

Exercise B.2.1. If $a_n \dots a_0.a_{-1}a_{-2}\dots$ is a real decimal, show that the series $\sum_{i=-\infty}^n a_i \times 10^i$ is absolutely convergent.

Exercise B.2.2. Show that the only decimal representations

$$1 = \pm a_n \dots a_0.a_{-1}a_{-2}\dots$$

of 1 are $1 = 1.000\dots$ and $1 = 0.999\dots$

Exercise B.2.3. A real number x is said to be a *terminating decimal* if we have $x = n/10^{-m}$ for some integers n, m . Show that if x is a terminating decimal, then x has exactly two decimal representations, while if x is not at terminating decimal, then x has exactly one decimal representation.

Exercise B.2.4. Rewrite the proof of Corollary 8.3.4 using the decimal system.

Index

- $++$ (increment), 18, 56
 - on integers, 87
- $+C$, 342
- α -length, 334
- ε -adherent, 161, 244
 - contually ε -adherent, 161
- ε -close
 - eventual, 115
 - functions, 253
 - local, 253
 - rational, 99
 - reals, 146
 - sequences, 115, 147
- ε -steady, 110, 146
 - eventually ε -steady, 111, 146
- π , 506, 508
- σ -algebra, 576, 595
- a posteriori*, 20
- a priori*, 20
- Abel's theorem, 484
- absolute convergence
 - for series, 192, 220
 - test, 192
- absolute value
 - for complex numbers, 498
 - for rational, 98
 - for reals, 129
- absolutely integrable, 617
- absorption laws, 52
- abstraction, 24-25, 390
- addition
 - long, 385
 - of cardinals, 82
 - of functions, 252
 - of complex numbers, 496
 - of integers, 86
 - of natural numbers, 27
 - of rational, 93
 - of reals, 119
- (countably) additive measure, 576, 592
- adherent point
 - infinite, 286
 - of sequences: *see* limit point
 - of sequences
 - of sets: 245, 402, 435
- alternating series test, 193
- ambient space, 406
- analysis, 1
- and: *see* conjunction
- antiderivative, 340
- approximation to the identity, 466, 469, 523
- Archimedian property, 132
- arctangent: *see* trigonometric functions

Aristotlean logic, 373–374

associativity

of addition in \mathbf{C} , 496

of addition in \mathbf{N} , 29

of composition, 59–60

of multiplication in \mathbf{N} , 34

of scalar multiplication, 538

of vector addition, 534

see also: ring, field, laws
of algebra

asymptotic discontinuity, 268

Axiom(s)

in mathematics, 24–25

of choice, 40, 73, 229

of comprehension: *see* Ax-
iom of universal spec-
ification

of countable choice, 230

of equality, 377

of foundation: *see* Axiom
of regularity

of induction: *see* princi-
ple of mathematical
induction

of infinity, 50

of natural numbers: *see*
Peano axioms

of pairwise union, 42

of power set, 66

of reflexivity, 377

of regularity, 54

of replacement, 49

of separation, 45

of set theory, 38, 40–42, 45, 49–
50, 54, 66

of singleton sets and pair

sets, 41

of specification, 45

of symmetry, 377

of substitution, 57, 377

of the empty set, 40

of transitivity, 377

of universal specification,
52

of union, 67

ball, 400

Banach-Tarski paradox, 575,
590

base of the natural logarithm:
see e

basis

standard basis of row vec-
tors, 535

bijection, 62

binomial formula, 189

Bolzano-Weierstrass theorem,
174

Boolean algebra, 47, 576, 591

Boolean logic, 367

Borel property, 575, 596

Borel-Cantelli lemma, 615

bound variable, 180, 369, 376

boundary (point), 401, 435

bounded

from above and below, 269

function, 269, 451

interval, 244

sequence, 113, 150

sequence away from zero,
123, 127

set, 248, 413

- C, C^0, C^1, C^2, C^k , 556
- cancellation law
 - of addition in \mathbf{N} , 29
 - of multiplication in \mathbf{N} , 35
 - of multiplication in \mathbf{Z} , 91
 - of multiplication in \mathbf{R} , 126
- Cantor's theorem, 224
- cardinality
 - arithmetic of, 81
 - of finite sets, 80
 - uniqueness of, 80
- Cartesian product, 70–71
 - infinite, 228–229
- Cauchy criterion, 197
- Cauchy sequence, 111, 146, 409
- Cauchy-Schwarz inequality, 399, 516
- chain: *see* totally ordered set
- chain rule, 293
 - in higher dimensions, 552, 555
- change of variables formula, 346–358
- character, 518
- characteristic function, 602
- choice
 - single, 40
 - finite, 73
 - countable, 230
 - arbitrary, 229
- closed
 - box, 580
 - interval, 243
 - set, 403, 435
- Clairaut's theorem: *see* interchanging derivatives
 - with derivatives
- closure, 245, 403, 435
- cluster point: *see* limit point
- cocountable topology, 438
- coefficient, 476
- cofinite topology, 437
- column vector, 534
- common refinement, 311
- commutativity
 - of addition in \mathbf{C} , 496
 - of addition in \mathbf{N} , 29
 - of addition in vector spaces, 534
 - of convolution, 467, 522
 - of multiplication in \mathbf{N} , 34
 - see also*: ring, field, laws of algebra
- compactness, 413, 436
 - compact support, 465
- comparison principle (or test)
 - for finite series, 181
 - for infinite series, 196
 - for sequences, 166
- completeness
 - of the space of continuous functions, 454
 - of metric spaces, 410
 - of the reals, 168
- completion of a metric space, 412
- complex numbers \mathbf{C} , 495
 - complex conjugation, 498
- composition of functions, 59
- conjunction (and), 354
- connectedness, 307, 430
 - connected component, 433

- constant
 - function, 58, 312
 - sequence, 170
- continuity, 261, 420, 436
 - and compactness, 427
 - and connectedness, 431
 - and convergence, 255, 421
- continuum, 242
 - hypothesis, 227
- contraction, 558
 - mapping theorem, 559
- contrapositive, 362
- convergence
 - in L^2 , 517
 - of a function at a point, 254, 441
 - of sequences, 148, 394, 434
 - of series, 190
 - pointwise: *see* pointwise convergence
 - uniform: *see* uniform convergence
- converse, 362
- convolution, 466, 487, 522
- corollary, 28
- coset, 587
- cosine: *see* trigonometric functions
- cotangent: *see* trigonometric functions
- countability, 208
 - of the integers, 212
 - of the rationals, 214
- cover, 578
 - see also*: open cover
- critical point, 572
- de Moivre identities, 507
- de Morgan laws, 47
- decimal
 - negative integer, 384
 - non-uniqueness of representation, 388
 - point, 385
 - positive integer, 384
 - real, 385–386
- degree, 464
- denumerable: *see* countable
- dense, 465
- derivative, 288
 - directional, 544
 - in higher dimensions, 542, 544, 546, 551
 - partial, 546
 - matrix, 551
 - total, 542, 544
 - uniqueness of, 542
- difference rule, 293
- difference set, 47
- differential matrix: *see* derivative matrix
- differentiability
 - at a point, 288
 - continuous, 556
 - directional, 544
 - in higher dimensions, 542
 - infinite, 478
 - k -fold, 478, 556
- digit, 381
- dilation, 536
- diophantine, 616
- Dirac delta function, 466
- direct sum

- of functions, 75, 425
- discontinuity: *see* singularity
- discrete metric, 393
- disjoint sets, 47
- disjunction (or), 354
 - inclusive vs. exclusive, 354
- distance
 - in \mathbf{C} , 499
 - in \mathbf{Q} , 98
 - in \mathbf{R} , 145, 391
- distributive law
 - for natural numbers, 34
 - for complex numbers, 497
 - see also*: laws of algebra
- divergence
 - of series, 3, 190
 - of sequences, 4
 - see also*: convergence
- divisibility, 237
- division
 - by zero, 3
 - formal ($//$), 93
 - of functions, 252
 - of rationals, 96
- domain, 55
- dominate: *see* majorize
 - dominated convergence: *see*
 - Lebesgue dominated convergence theorem
- doubly infinite, 244
- dummy variable: *see* bound variable
- e , 491
- Egoroff's theorem, 617
- empty
 - Cartesian product, 73
 - function, 58
 - sequence, 73
 - series, 185
 - set, 40, 576, 579
- equality, 377
 - for functions, 58
 - for sets, 39
 - of cardinality, 77
- equivalence
 - of sequences, 116, 281
 - relation, 378
- error-correcting codes, 392
- Euclidean algorithm, 35
- Euclidean metric, 391
- Euclidean space, 391
- Euler's formula, 503, 506
- Euler's number: *see* e
- exponential function, 490, 501
- exponentiation
 - of cardinals, 81
 - with base and exponent in \mathbf{N} , 36
 - with base in \mathbf{Q} and exponent in \mathbf{Z} , 101, 102
 - with base in \mathbf{R} and exponent in \mathbf{Z} , 140
 - with base in \mathbf{R}^+ and exponent in \mathbf{Q} , 142
 - with base in \mathbf{R}^+ and exponent in \mathbf{R} , 177
- expression, 353
- extended real number system
 - \mathbf{R}^* , 137, 153
- extremum: *see* maximum, minimum
- exterior (point), 401, 435

- factorial, 189
- family, 67
- Fatou's lemma, 614
- Fejér kernel, 524
- field, 95
 - ordered, 97
- finite intersection property, 419
- finite set, 80
- fixed point theorem, 276, 559
- forward image: *see* image
- Fourier
 - coefficients, 520
 - inversion formula, 520
 - series, 520
 - series for arbitrary periods, 531
 - theorem, 536
 - transform, 520
- fractional part, 512
- free variable, 368
- frequency, 518
- Fubini's theorem, 624
 - for finite series, 188
 - for infinite series, 217
 - see also*: interchanging integrals/sums with integrals/sums
- function, 55
 - implicit definition, 57
- fundamental theorems of calculus, 338, 341
- geometric series, 190, 196
 - formula, 197, 200, 460
- geodesic, 394
- gradient, 550
- graph, 58, 75, 251, 568
- greatest lower bound: *see* least upper bound
- hairy ball theorem, 559
- half-infinite, 244
- half-open, 243
- half-space, 591
- harmonic series, 199
- Hausdorff space, 437, 438
- Hausdorff maximality principle, 240
- Heine-Borel theorem, 414
 - for the real line, 248
- Hermitian form, 515
- homogeneity, 516, 535
- hypersurface, 568
- identity map (or operator), 63, 536
- if: *see* implication
- iff (if and only if), 30
- ill-defined, 351, 353
- image
 - of sets, 64
 - inverse image, 65
- imaginary, 498
- implication (if), 357
- implicit differentiation, 568
- implicit function theorem, 568
- improper integral, 318
- inclusion map, 63
- inconsistent, 227, 228, 502
- index of summation: *see* dummy variable
- index set, 67
- indicator function: *see* characteristic function

- induced
 - metric, 391, 407
 - topology, 407, 435
- induction: *see* Principle of mathematical induction
- infinite
 - interval, 244
 - set, 80
- infimum: *see* supremum
- injection: *see* one-to-one function
- inner product, 514
- integer part, 103, 133, 512
- integers \mathbf{Z}
 - definition, 85
 - identification with rationals, 94
 - interspersing with rationals, 103
- integral test, 332
- integration
 - by parts, 343-345, 484
 - laws, 315, 321
 - piecewise constant, 313, 315
 - Riemann: *see* Riemann integral
- interchanging
 - derivatives with derivatives, 10, 556
 - finite sums with finite sums, 187, 188
 - integrals with integrals, 7, 614, 624
 - limits with derivatives, 9, 463
 - limits with integrals, 9, 462, 610, 619
 - limits with length, 12
 - limits with limits, 8, 9, 450
 - limits with sums, 617
 - sums with derivatives, 463, 476
 - sums with integrals, 459, 476, 613, 614, 616
 - sums with sums, 6, 217
- interior (point), 401, 435
- intermediate value theorem, 274, 432
- intersection
 - pairwise, 46
- interval, 243
- intrinsic, 413
- inverse
 - function theorem, 301, 562
 - image, 65
 - in logic, 362
 - of functions, 63
- invertible function: *see* bijection
 - local, 562
- involution, 498
- irrationality, 108
 - of $\sqrt{2}$, 104, 137
- isolated point, 247
- isometry, 406
- jump discontinuity, 268
- l^1 , l^2 , l^∞ , L^1 , L^2 , L^∞ , 391-393, 516, 617

- equivalence of in finite dimensions, 396
- see also*: absolutely integrable
- see also*: supremum as norm
- L'Hôpital's rule, 11, 303
- label, 67
- laws of algebra
 - for complex numbers, 496, 497
 - for integers, 89
 - for rationals, 95
 - for reals, 122
- laws of arithmetic: *see* laws of algebra
- laws of exponentiation, 101, 102, 141, 143, 177, 490
- least upper bound, 134
 - least upper bound property, 135, 158
 - see also* supremum
- Lebesgue dominated convergence theorem, 619
- Lebesgue integral
 - of absolutely integrable functions, 618
 - of nonnegative functions, 608
 - of simple functions, 604
 - upper and lower, 620
 - vs. the Riemann integral, 622
- Lebesgue measurable, 590
- Lebesgue measure, 577
 - motivation of, 575-577
- Lebesgue monotone convergence theorem, 610
- Leibnitz rule, 293, 554
- lemma, 28
- length of interval, 308
- limit
 - at infinity, 286
 - formal (LIM), 118, 150, 412
 - laws, 150, 256, 500
 - left and right, 265
 - limiting values of functions, 5, 254, 441
 - of sequences, 148
 - pointwise, 444
 - uniform, *see* uniform limit
 - uniqueness of, 148, 256, 397, 442
- limit inferior, *see* limit superior
- limit point
 - of sequences, 160, 409
 - of sets, 247
- limit superior, 162
- linear combination, 535
- linearity
 - approximate, 541
 - of convolution, 471, 522
 - of finite series, 186
 - of limits, 151
 - of infinite series, 194
 - of inner product, 515
 - of integration, 315, 321, 606, 612
 - of transformations, 535
- Lipschitz constant, 298
- Lipschitz continuous, 298

- logarithm (natural), 492
 - power series of, 460, 492
- logical connective, 354
- lower bound: *see* upper bound
- majorize, 317, 608
- manifold, 572
- map: *see* function
- matrix, 536
 - identification with linear transformations, 537-540
- maximum, 233, 296
 - local, 296
 - of functions, 252, 271
 - principle, 271, 427
- mean value theorem, 297
- measurability
 - for functions, 597, 598
 - for sets, 590
 - motivation of, 574
 - see also*: Lebesgue measure, outer measure
- meta-proof, 140
- metric, 390
 - ball: *see* ball
 - on \mathbf{C} , 499
 - on \mathbf{R} , 391
 - space, 390
 - see also*: distance
- minimum, 233, 296
 - local, 296
 - of a set of natural numbers, 210
 - of functions, 252, 271
- minorize: *see* majorize
- monomial, 518
- monotone (increasing or decreasing)
 - convergence: *see* Lebesgue monotone convergence theorem
 - function, 276, 336
 - measure, 576, 580
 - sequence, 159
- morphism: *see* function
- moving bump example, 446, 614
- multiplication
 - of cardinals, 81
 - of complex numbers, 497
 - of functions, 252
 - of integers, 86
 - of matrices, 536, 540
 - of natural numbers, 33
 - of rationals, 93, 94
 - of reals, 120
- Natural numbers \mathbf{N}
 - are infinite, 80
 - axioms: *see* Peano axioms
 - identification with integers, 87
 - informal definition, 17
 - in set theory: *see* Axiom of infinity
 - uniqueness of, 76
- negation
 - in logic, 355
 - of extended reals, 154
 - of complex numbers, 497
 - of integers, 88
 - of rationals, 93
 - of reals, 121

- negative: *see* negation, positive
- neighbourhood, 434
- Newton's approximation, 291, 544
- non-constructive, 229
- non-degenerate, 516
- nowhere differentiable function, 464, 508
- objects, 38
 - primitive, 53
- one-to-one function, 61
- one-to-one correspondence: *see* bijection
- onto, 61
- open
 - box, 578
 - cover, 414
 - interval, 243
 - set, 403
- or: *see* disjunction
- order ideal, 238
- order topology, 437
- ordered pair, 70
 - construction of, 74
- ordered n -tuple, 71
- ordering
 - lexicographical, 239
 - of cardinals, 227
 - of orderings, 240
 - of partitions, 310
 - of sets, 233
 - of the extended reals, 154
 - of the integers, 91
 - of the natural numbers, 31
 - of the rationals, 97
 - of the reals, 129
- orthogonality, 516
- orthonormal, 519
- oscillatory discontinuity, 268
- outer measure, 579
 - non-additivity of, 587, 589
- pair set, 41
- partial function, 69
- partially ordered set, 45, 232
- partial sum, 190
- Parseval identity, 531
 - see also*: Plancherel formula
- partition, 308
- path-connected, 432
- Peano axioms, 18-21, 23
- perfect matching: *see* bijection
- periodic, 511
 - extension, 512
- piecewise
 - constant, 312
 - constant Riemann-Stieltjes integral, 335
 - continuous, 330
- pigeonhole principle, 83
- Plancherel formula (or theorem), 520, 528
- pointwise convergence, 444
 - of series, 456
 - topology of, 455
- polar representation, 507
- polynomial, 265, 464
 - and convolution, 467
 - approximation by, 465, 470

- positive
 - complex number, 498, 502
 - integer, 88
 - inner product, 515
 - measure, 576, 580
 - natural number, 30
 - rational, 96
 - real, 128
- power series, 476
 - formal, 474
 - multiplication of, 487
 - uniqueness of, 481
- power set, 66
- pre-image: *see* inverse image
- principle of infinite descent, 106
- principle of mathematical induction, 21
 - backwards induction, 33
 - strong induction, 32, 234
 - transfinite, 237
- product rule, *see* Leibnitz rule
- product topology, 455
- projection, 536
- proof
 - by contradiction, 352, 363
 - abstract examples, 364-367, 375-377
- proper subset, 44
- property, 354
- proposition, 28
- propositional logic, 367
- Pythagoras' theorem, 516
- quantifier, 369
 - existential (for some), 371
 - negation of, 372
 - nested, 372
 - universal (for all), 370
- Quotient: *see* division
- Quotient rule, 293, 555
- radius of convergence, 475
- range, 55
- ratio test, 206
- rational numbers **Q**
 - definition, 93
 - identification with reals, 121
 - interspersing with rationals, 103
 - interspersing with reals, 132
- real analytic, 478
- real numbers **R**
 - are uncountable: *see* uncountability of the reals
 - definition, 117
- real part, 498
- real-valued, 455
- rearrangement
 - of absolutely convergent series, 202
 - of divergent series, 203, 222
 - of finite series, 185
 - of non-negative series, 200
- reciprocal
 - of complex numbers, 499
 - of rationals, 95
 - of reals, 125
- recursive definitions, 26, 76

- reductio ad absurdum*: *see* proof
 by contradiction
 relative topology: *see* induced
 topology
 removable discontinuity: *see*
 removable singularity
 removable singularity, 259, 268
 restriction of functions, 250
 Riemann hypothesis, 200
 Riemann integrability, 318
 closure properties, 321-326
 failure of, 332
 of bounded continuous func-
 tions, 328
 of continuous functions on
 compacta, 328
 of monotone functions, 330
 of piecewise continuous bounded
 functions, 329
 of uniformly continuous func-
 tions, 326
 Riemann integral, 318
 upper and lower, 317
 Riemann sums (upper and lower),
 321
 Riemann zeta function, 199
 Riemann-Stieltjes integral, 336
 ring, 89
 commutative, 89, 497
 Rolle's theorem, 297
 root, 140
 mean square: *see* L^2
 test, 204
 row vector, 533
 Russell's paradox, 52
 scalar multiplication, 533
 of functions, 252
 Schröder-Bernstein theorem,
 227
 sequence, 109
 finite, 74
 series
 finite, 179, 182
 formal infinite, 189
 laws, 194, 220
 of functions, 459
 on arbitrary sets, 220
 on countable sets, 216
 vs. sum, 180
 set
 axioms: *see* axioms of set
 theory
 informal definition, 38
 signum function, 258
 simple function, 602
 sine: *see* trigonometric func-
 tions
 singleton set, 41
 singularity, 268
 space, 390
 statement, 350
 sub-additive measure, 576, 580
 subset, 44
 subsequence, 172, 408
 substitution: *see* rearrange-
 ment
 subtraction
 formal ($--$), 86
 of functions, 252
 of integers, 91
 sum rule, 292
 summation by parts, 484

- sup norm: *see* supremum as norm
- support, 465
- supremum (and infimum)
 - as metric, 393
 - as norm, 393, 457
 - of a set of extended reals, 156, 157
 - of a set of reals, 137, 139
 - of sequences of reals, 158
- square root, 56
- square wave, 512, 518
- Squeeze test
 - for sequences, 167
- Stone-Weierstrass theorem, 472, 522
- strict upper bound, 235
- surjection: *see* onto
- taxi-cab metric, 392
- tangent: *see* trigonometric function
- Taylor series, 480
- Taylor's formula: *see* Taylor series
- telescoping series, 195
- ten, 381
- theorem, 28
- topological space, 433
- totally bounded, 418
- totally ordered set, 45, 233
- transformation: *see* function
- translation invariance, 577, 580, 591
- transpose, 534
- triangle inequality
 - in Euclidean space, 399
 - in inner product spaces, 516
- in metric spaces, 390
- in \mathbf{C} , 499
- in \mathbf{R} , 99
- for finite series, 181, 186
- for integrals, 618
- trichotomy of order
 - of extended reals, 155
 - for natural numbers, 31
 - for integers, 91
 - for rationals, 97
 - for reals, 129
- trigonometric functions, 503, 509
 - and Fourier series, 530
 - trigonometric polynomials, 518
 - power series, 504, 508
- trivial topology, 437
- two-to-one function, 61
- uncountability, 208
 - of the reals, 225
- undecidable, 228
- uniform continuity, 280, 428
- uniform convergence, 447
 - and anti-derivatives, 462
 - and derivatives, 451
 - and integrals, 459
 - and limits, 450
 - and radius of convergence, 476
 - as a metric, 453, 514
 - of series, 457
- uniform limit, 447
 - of bounded functions, 451

- of continuous functions, 450
- and Riemann integration, 458
- union, 67
 - pairwise, 42
- universal set, 53
- upper bound,
 - of a set of reals, 133
 - of a partially ordered set, 234
 - see also*: least upper bound
- variable, 368
- vector space, 534
- vertical line test, 55, 76, 567
- volume, 578
- Weierstrass approximation theorem, 465, 470-471, 521
- Weierstrass example: *see* nowhere differentiable function
- Weierstrass M -test, 457
- well-defined, 351
- well-ordered sets, 234
- well ordering principle
 - for natural numbers, 210
 - for arbitrary sets, 241
- Zermelo-Fraenkel(-Choice) axioms, 69
 - see also* axioms of set theory
- zero test
 - for sequences, 167
 - for series, 191
- Zorn's lemma, 237