


i-Tree Canopy Report


i-Tree Benefits and Cover Assessment

Estimated using random sampling statistics on 9/15/2026



Google

Land Cover



Abbr.	Cover Class	Description	Points	% Cover ± SE	Area (km²) ± SE
H	Grass/Steppe/Herbaceous		4185	20.67 ± 0.28	0.39 ± 0.01
IB	Impervious Buildings		76	0.38 ± 0.04	0.01 ± 0.00
IO	Impervious Other		10	0.05 ± 0.02	0.00 ± 0.00
IR	Impervious Road		588	2.90 ± 0.12	0.06 ± 0.00
RDS	Reeds/Aquatic Plants		31	0.15 ± 0.03	0.00 ± 0.00
S	Soil/Bare Ground		878	4.34 ± 0.14	0.08 ± 0.00
T	Tree/Shrub		14475	71.48 ± 0.32	1.36 ± 0.01
W	Water		7	0.03 ± 0.01	0.00 ± 0.00
Total			20250	100.00	1.90

Tree Benefit Estimates: Carbon (Metric units)

Description	Carbon (t)	±SE	CO ₂ Equiv. (t)	±SE	Value (USD)	±SE
Sequestered annually in trees	257.69	±1.14	944.87	±4.19	\$122,919	±546
Stored in trees (Note: this benefit is not an annual rate)	10,422.69	±46.26	38,216.52	±169.63	\$4,971,622	±22,067

Currency is in USD and rounded. Standard errors of removal and benefit amounts are based on standard errors of sampled and classified points. Amount sequestered is based on 190.000 t of Carbon, or 696.667 t of CO₂, per km²/yr and rounded. Amount stored is based on 7684.808 t of Carbon, or 28177.630 t of CO₂, per km² and rounded. Value (USD) is based on \$477.00/t of Carbon, or \$130.09/t of CO₂ and rounded. (Metric units: t = tonnes, metric tons, km² = square kilometers)

Tree Benefit Estimates: Air Pollution (Metric units)

Abbr.	Description	Amount (kg)	±SE	Value (USD)	±SE
CO	Carbon Monoxide removed annually	114.89	±0.51	\$169	±1
NO2	Nitrogen Dioxide removed annually	1,306.24	±5.80	\$839	±4
O3	Ozone removed annually	6,035.30	±26.79	\$15,826	±70
SO2	Sulfur Dioxide removed annually	87.63	±0.39	\$16	±0
PM2.5	Particulate Matter less than 2.5 microns removed annually	597.84	±2.65	\$80,424	±357
PM10*	Particulate Matter greater than 2.5 microns and less than 10 microns removed annually	1,951.26	±8.66	\$13,483	±60
Total		10,093.16	±44.80	\$110,756	±492

Currency is in USD and rounded. Standard errors of removal and benefit amounts are based on standard errors of sampled and classified points. Air Pollution Estimates are based on these values in kg/km²/yr @ \$/kg/yr and rounded:
CO 84.711 @ \$1.47 | NO2 963.109 @ \$0.64 | O3 4,449.923 @ \$2.62 | SO2 64.614 @ \$0.18 | PM2.5 440.794 @ \$134.53 | PM10* 1,438.693 @ \$6.91 (Metric units: kg = kilograms, km² = square kilometers)

Tree Benefit Estimates: Hydrological (Metric units)

Abbr.	Benefit	Amount (MI)	±SE	Value (USD)	±SE
AVRO	Avoided Runoff	16.58	±0.07	\$39,144	±174
E	Evaporation	100.79	±0.45	N/A	N/A
I	Interception	101.03	±0.45	N/A	N/A
T	Transpiration	188.11	±0.83	N/A	N/A
PE	Potential Evaporation	950.99	±4.22	N/A	N/A
PET	Potential Evapotranspiration	796.97	±3.54	N/A	N/A

Currency is in USD and rounded. Standard errors of removal and benefit amounts are based on standard errors of sampled and classified points. Hydrological Estimates are based on these values in MI/km²/yr @ \$/MI/yr and rounded:
AVRO 12.226 @ \$2,360.64 | E 74.311 @ N/A | I 74.493 @ N/A | T 138.695 @ N/A | PE 701.183 @ N/A | PET 587.615 @ N/A (Metric units: MI = megaliters, km² = square kilometers)

About i-Tree Canopy

The concept and prototype of this program were developed by David J. Nowak, Jeffery T. Walton, and Eric J. Greenfield (USDA Forest Service). The current version of this program was developed and adapted to i-Tree by David Ellingsworth, Mike Binkley, and Scott Maco (The Davey Tree Expert Company)

Limitations of i-Tree Canopy

The accuracy of the analysis depends upon the ability of the user to correctly classify each point into its correct class. As the number of points increase, the precision of the estimate will increase as the standard error of the estimate will decrease. If too few points are classified, the standard error will be too high to have any real certainty of the estimate.