

The pair distribution function describes how atomic positions are correlated as a function of real space distance rather than reciprocal space. Total scattering methods avoid the restrictions of Bragg averaging and therefore retain information contained in diffuse background intensity. Before refining a structural model the instrument parameters must be calibrated using a standard material measured under identical conditions. Peak broadening at high momentum transfer is dominated by the finite Q resolution function while low angle peaks reflect correlated microstrain in the lattice. A careful background subtraction removes the contribution of the capillary air scattering and leaves the sample signal alone for further processing. Compressed files load faster on slow connections which matters when a reviewer opens the supplementary material from a conference laptop. PDF documents that embed vector figures tend to shrink dramatically when the font subsets and image streams are reencoded losslessly. The refinement converged after twelve cycles with a low weighted residual factor and sensible anisotropic displacement parameters for every atom site. Care should be taken when truncating the data because sharp termination produces ripples in real space that mimic genuine short range order. Synchronization of the detector counter with the motor encoder is essential to keep the absorption correction geometrically consistent. The pair distribution function describes how atomic positions are correlated as a function of real space distance rather than reciprocal space. Total scattering methods avoid the restrictions of Bragg averaging and therefore retain information contained in diffuse background intensity. Before refining a structural model the instrument parameters must be calibrated using a standard material measured under identical conditions. Peak broadening at high momentum transfer is dominated by the finite Q resolution function while low angle peaks reflect correlated microstrain in the lattice. A careful background subtraction removes the contribution of the capillary air scattering and leaves the sample signal alone for further processing. Compressed files load faster on slow connections which matters when a reviewer opens the supplementary material from a conference laptop. PDF documents that embed vector figures tend to shrink dramatically when the font subsets and image streams are reencoded losslessly. The refinement converged after twelve cycles with a low weighted residual factor and sensible anisotropic displacement parameters for every atom site. Care should be taken when truncating the data because sharp termination produces ripples in real space that mimic genuine short range order. Synchronization of the detector counter with the motor encoder is essential to keep the absorption correction geometrically consistent. The pair distribution function describes how atomic positions are correlated as a function of real space distance rather than

The pair distribution function describes how atomic positions are correlated as a function of real space distance rather than reciprocal space. Total scattering methods avoid the restrictions of Bragg averaging and therefore retain information contained in diffuse background intensity. Before refining a structural model the instrument parameters must be calibrated using a standard material measured under identical conditions. Peak broadening at high momentum transfer is dominated by the finite Q resolution function while low angle peaks reflect correlated microstrain in the lattice. A careful background subtraction removes the contribution of the capillary air scattering and leaves the sample signal alone for further processing. Compressed files load faster on slow connections which matters when a reviewer opens the supplementary material from a conference laptop. PDF documents that embed vector figures tend to shrink dramatically when the font subsets and image streams are reencoded losslessly. The refinement converged after twelve cycles with a low weighted residual factor and sensible anisotropic displacement parameters for every atom site. Care should be taken when truncating the data because sharp termination produces ripples in real space that mimic genuine short range order. Synchronization of the detector counter with the motor encoder is essential to keep the absorption correction geometrically consistent. The pair distribution function describes how atomic positions are correlated as a function of real space distance rather than reciprocal space. Total scattering methods avoid the restrictions of Bragg averaging and therefore retain information contained in diffuse background intensity. Before refining a structural model the instrument parameters must be calibrated using a standard material measured under identical conditions. Peak broadening at high momentum transfer is dominated by the finite Q resolution function while low angle peaks reflect correlated microstrain in the lattice. A careful background subtraction removes the contribution of the capillary air scattering and leaves the sample signal alone for further processing. Compressed files load faster on slow connections which matters when a reviewer opens the supplementary material from a conference laptop. PDF documents that embed vector figures tend to shrink dramatically when the font subsets and image streams are reencoded losslessly. The refinement converged after twelve cycles with a low weighted residual factor and sensible anisotropic displacement parameters for every atom site. Care should be taken when truncating the data because sharp termination produces ripples in real space that mimic genuine short range order. Synchronization of the detector counter with the motor encoder is essential to keep the absorption correction geometrically consistent. The pair distribution function describes how atomic positions are correlated as a function of real space distance rather than

The pair distribution function describes how atomic positions are correlated as a function of real space distance rather than reciprocal space. Total scattering methods avoid the restrictions of Bragg averaging and therefore retain information contained in diffuse background intensity. Before refining a structural model the instrument parameters must be calibrated using a standard material measured under identical conditions. Peak broadening at high momentum transfer is dominated by the finite Q resolution function while low angle peaks reflect correlated microstrain in the lattice. A careful background subtraction removes the contribution of the capillary air scattering and leaves the sample signal alone for further processing. Compressed files load faster on slow connections which matters when a reviewer opens the supplementary material from a conference laptop. PDF documents that embed vector figures tend to shrink dramatically when the font subsets and image streams are reencoded losslessly. The refinement converged after twelve cycles with a low weighted residual factor and sensible anisotropic displacement parameters for every atom site. Care should be taken when truncating the data because sharp termination produces ripples in real space that mimic genuine short range order. Synchronization of the detector counter with the motor encoder is essential to keep the absorption correction geometrically consistent. The pair distribution function describes how atomic positions are correlated as a function of real space distance rather than reciprocal space. Total scattering methods avoid the restrictions of Bragg averaging and therefore retain information contained in diffuse background intensity. Before refining a structural model the instrument parameters must be calibrated using a standard material measured under identical conditions. Peak broadening at high momentum transfer is dominated by the finite Q resolution function while low angle peaks reflect correlated microstrain in the lattice. A careful background subtraction removes the contribution of the capillary air scattering and leaves the sample signal alone for further processing. Compressed files load faster on slow connections which matters when a reviewer opens the supplementary material from a conference laptop. PDF documents that embed vector figures tend to shrink dramatically when the font subsets and image streams are reencoded losslessly. The refinement converged after twelve cycles with a low weighted residual factor and sensible anisotropic displacement parameters for every atom site. Care should be taken when truncating the data because sharp termination produces ripples in real space that mimic genuine short range order. Synchronization of the detector counter with the motor encoder is essential to keep the absorption correction geometrically consistent. The pair distribution function describes how atomic positions are correlated as a function of real space distance rather than