

Illustrating Hilbert's Nullstellensatz

Timothy Cho

UC Irvine, MATH 195

30 November 2023

Motivation: Why do Algebraic Geometry?

Motivation: Why do Algebraic Geometry?

- It's cool

Motivation: Why do Algebraic Geometry?

- It's cool
- Historical: shapes/patterns \implies equations

Motivation: Why do Algebraic Geometry?

- It's cool
- Historical: shapes/patterns \implies equations
- Algebraic geometry: shapes/patterns \iff equations

Motivation: Why do Algebraic Geometry?

- It's cool
- Historical: shapes/patterns \implies equations
- Algebraic geometry: shapes/patterns \iff equations
- Connections: analysis, topology, commutative algebra, etc.

Roots of Polynomials

Find the real roots of $(x^2 + 4y^2 - 4)^2 \in \mathbb{R}[x, y]$.

Roots of Polynomials


Find the real roots of $(x^2 + 4y^2 - 4)^2 \in \mathbb{R}[x, y]$.

$$(x^2 + 4y^2 - 4)^2 = 0 \implies x^2 + 4y^2 - 4 = 0 \implies x^2 + 4y^2 = 4$$

Roots of Polynomials

Find the real roots of $(x^2 + 4y^2 - 4)^2 \in \mathbb{R}[x, y]$.

$$(x^2 + 4y^2 - 4)^2 = 0 \implies x^2 + 4y^2 - 4 = 0 \implies x^2 + 4y^2 = 4$$



Algebraic Varieties

Definition

Let k be a field, and let $S \subseteq k[x_1, \dots, x_n]$. The *algebraic variety* of S is the set

$$\mathbb{V}(S) := \{x \in k^n : f(x) = 0 \text{ for all } f \in S\}.$$

Similarly, $V \subseteq k^n$ is an *algebraic set* if $V = \mathbb{V}(S)$ for some $S \subseteq k[x_1, \dots, x_n]$.

Algebraic Varieties

Definition

Let k be a field, and let $S \subseteq k[x_1, \dots, x_n]$. The *algebraic variety* of S is the set

$$\mathbb{V}(S) := \{x \in k^n : f(x) = 0 \text{ for all } f \in S\}.$$

Similarly, $V \subseteq k^n$ is an *algebraic set* if $V = \mathbb{V}(S)$ for some $S \subseteq k[x_1, \dots, x_n]$.

Advantage: points \iff pictures \iff shapes

Disadvantages: hard to work with

Another Example

Let $S = \{(x^2 + y^2 - 4)(x - 1), (x^2 + y^2 - 4)(y - 1)\} \subseteq \mathbb{R}[x, y]$.
What is $\mathbb{V}(S) \subseteq \mathbb{R}^2$?

Another Example

Let $S = \{(x^2 + y^2 - 4)(x - 1), (x^2 + y^2 - 4)(y - 1)\} \subseteq \mathbb{R}[x, y]$.
What is $\mathbb{V}(S) \subseteq \mathbb{R}^2$?

$$\begin{cases} (x^2 + y^2 - 4)(x - 1) = 0 \\ (x^2 + y^2 - 4)(y - 1) = 0 \end{cases}$$

Another Example

Let $S = \{(x^2 + y^2 - 4)(x - 1), (x^2 + y^2 - 4)(y - 1)\} \subseteq \mathbb{R}[x, y]$.
What is $V(S) \subseteq \mathbb{R}^2$?

$$\begin{cases} (x^2 + y^2 - 4)(x - 1) = 0 \\ (x^2 + y^2 - 4)(y - 1) = 0 \end{cases}$$

$$\implies 0 = (x^2 + y^2 - 4)(x - 1) = (x^2 + y^2 - 4)(y - 1)$$

$$\implies x^2 + y^2 = 4 \text{ or } (x, y) = (1, 1)$$

Another Example




Figure: $\mathbb{V}\left((x^2 + y^2 - 4)(x - 1), (x^2 + y^2 - 4)(y - 1)\right)$

Ideals

Motivation: “the other way around”

Ideals

Motivation: “the other way around”

Definition

Let k be a field, and let $V \subseteq k^n$. The *ideal* of V to be the set

$$\mathbb{I}(V) := \{f \in k[x_1, \dots, x_n] : f(x) = 0 \text{ for all } x \in V\}.$$

Ideals

Motivation: “the other way around”

Definition

Let k be a field, and let $V \subseteq k^n$. The *ideal* of V to be the set

$$\mathbb{I}(V) := \{f \in k[x_1, \dots, x_n] : f(x) = 0 \text{ for all } x \in V\}.$$

Proposition

Let k be a field, and let $V \subseteq k^n$. Then $\mathbb{I}(V)$ is an ideal in $k[x_1, \dots, x_n]$.

Ideals

Motivation: “the other way around”

Definition

Let k be a field, and let $V \subseteq k^n$. The *ideal* of V to be the set


$$\mathbb{I}(V) := \{f \in k[x_1, \dots, x_n] : f(x) = 0 \text{ for all } x \in V\}.$$

Proposition

Let k be a field, and let $V \subseteq k^n$. Then $\mathbb{I}(V)$ is an ideal in $k[x_1, \dots, x_n]$.

Advantage: we understand ideals!

Computing Ideals



What is $\mathbb{I}(V)$?

Figure: $V \subseteq \mathbb{R}^2$

Computing Ideals




Figure: $V \subseteq \mathbb{R}^2$

What is $\mathbb{I}(V)$?

$$V = \{(x, y) \in \mathbb{R}^2 : x^2 + y - 4 = 0\}.$$

Computing Ideals




Figure: $V \subseteq \mathbb{R}^2$

What is $\mathbb{I}(V)$?

$$V = \{(x, y) \in \mathbb{R}^2 : x^2 + y - 4 = 0\}.$$

$f(x, y) = g(x, y)(x^2 + y - 4)$
vanishes for all $g(x, y) \in \mathbb{R}[x, y]$

Computing Ideals




Figure: $V \subseteq \mathbb{R}^2$

What is $\mathbb{I}(V)$?

$$V = \{(x, y) \in \mathbb{R}^2 : x^2 + y - 4 = 0\}.$$

$f(x, y) = g(x, y)(x^2 + y - 4)$
vanishes for all $g(x, y) \in \mathbb{R}[x, y]$

$$\mathbb{I}(V) = \langle x^2 + y - 4 \rangle.$$

\mathbb{V} versus \mathbb{I}

Let k be a field.

| Attribute | \mathbb{V} | \mathbb{I} |
|-----------|--------------|--------------|
|-----------|--------------|--------------|

\mathbb{V} versus \mathbb{I}

Let k be a field.

| Attribute | \mathbb{V} | \mathbb{I} |
|-----------|----------------------------------|-------------------|
| Input | $S \subseteq k[x_1, \dots, x_n]$ | $V \subseteq k^n$ |

\mathbb{V} versus \mathbb{I}

Let k be a field.

| Attribute | \mathbb{V} | \mathbb{I} |
|-----------|----------------------------------|--|
| Input | $S \subseteq k[x_1, \dots, x_n]$ | $V \subseteq k^n$ |
| Output | $\mathbb{V}(S) \subseteq k^n$ | $\mathbb{I}(V) \subseteq k[x_1, \dots, x_n]$ |

\mathbb{V} versus \mathbb{I}

Let k be a field.

| Attribute | \mathbb{V} | \mathbb{I} |
|------------|----------------------------------|--|
| Input | $S \subseteq k[x_1, \dots, x_n]$ | $V \subseteq k^n$ |
| Output | $\mathbb{V}(S) \subseteq k^n$ | $\mathbb{I}(V) \subseteq k[x_1, \dots, x_n]$ |
| Advantages | Very visual | Easy to manipulate |

\mathbb{V} versus \mathbb{I}

Let k be a field.

| Attribute | \mathbb{V} | \mathbb{I} |
|---------------|----------------------------------|--|
| Input | $S \subseteq k[x_1, \dots, x_n]$ | $V \subseteq k^n$ |
| Output | $\mathbb{V}(S) \subseteq k^n$ | $\mathbb{I}(V) \subseteq k[x_1, \dots, x_n]$ |
| Advantages | Very visual | Easy to manipulate |
| Disadvantages | Hard to manipulate | Not visual |

\mathbb{V} versus \mathbb{I}

Let k be a field.

| Attribute | \mathbb{V} | \mathbb{I} |
|---------------|----------------------------------|--|
| Input | $S \subseteq k[x_1, \dots, x_n]$ | $V \subseteq k^n$ |
| Output | $\mathbb{V}(S) \subseteq k^n$ | $\mathbb{I}(V) \subseteq k[x_1, \dots, x_n]$ |
| Advantages | Very visual | Easy to manipulate |
| Disadvantages | Hard to manipulate | Not visual |

In an ideal world: \mathbb{I} and \mathbb{V} are mutually inverse

$$\mathbb{V}(\mathbb{I}(V)) = V$$

Proposition

Let $V \subseteq k^n$ be an algebraic set. Then $\mathbb{V}(\mathbb{I}(V)) = V$. That is, \mathbb{V} is a left-inverse to \mathbb{I} .

Proof.

Definition-chase.



$\mathbb{V}(\mathbb{I}(V)) = \mathbb{V}$: Example




Figure: $V \subseteq \mathbb{R}^2$

Know:

$$I := \mathbb{I}(V) = \langle x^2 + y - 4 \rangle.$$

$\mathbb{V}(I)$: points in \mathbb{R}^2 that vanish
on all of I

$\mathbb{V}(\mathbb{I}(V)) = \mathbb{V}$: Example




Figure: $V \subseteq \mathbb{R}^2$

Know:

$$I := \mathbb{I}(V) = \langle x^2 + y - 4 \rangle.$$

$\mathbb{V}(I)$: points in \mathbb{R}^2 that vanish on all of I — but this is just V .

$$\mathbb{I}(\mathbb{V}(I)) = I?$$

Let $I := \langle (x^2 + 4y^2 - 4)^2 \rangle$.




Figure: $\mathbb{V}(I)$

$$\mathbb{I}(\mathbb{V}(I)) = I?$$




Figure: $\mathbb{V}(I)$

Let $I := \langle (x^2 + 4y^2 - 4)^2 \rangle$.

$\mathbb{I}(\mathbb{V}(I))$: polynomials in $\mathbb{R}[x, y]$
that vanish on $\mathbb{V}(I)$

$$\mathbb{I}(\mathbb{V}(I)) = \langle x^2 + 4y^2 - 4 \rangle \neq I.$$

$$\mathbb{I}(\mathbb{V}(I)) = I?$$




Figure: $\mathbb{V}(I)$

Let $I := \langle (x^2 + 4y^2 - 4)^2 \rangle$.

$\mathbb{I}(\mathbb{V}(I))$: polynomials in $\mathbb{R}[x, y]$ that vanish on $\mathbb{V}(I)$

$$\mathbb{I}(\mathbb{V}(I)) = \langle x^2 + 4y^2 - 4 \rangle \neq I.$$

But how are $\mathbb{I}(\mathbb{V}(I))$ and I related?

Radical of an Ideal

“Set of n th roots”

Radical of an Ideal

“Set of n th roots”

Definition

Let $I \subseteq R$ be an ideal. Then the *radical* of I is the set

$$\sqrt{I} := \{r \in R : r^n \in I \text{ for some } n \in \mathbb{Z}^+\}.$$

Similarly, we say that an ideal I is *radical* if $I = \sqrt{I}$.

Radical of an Ideal

“Set of n th roots”

Definition

Let $I \subseteq R$ be an ideal. Then the *radical* of I is the set

$$\sqrt{I} := \{r \in R : r^n \in I \text{ for some } n \in \mathbb{Z}^+\}.$$

Similarly, we say that an ideal I is *radical* if $I = \sqrt{I}$.

Proposition

The radical of an ideal is an ideal.

Proposition

Prime ideals are radical.

A Rad(ical) Example




Figure: $\mathbb{V}(I)$

Let $I := \langle (x^2 + 4y^2 - 4)^2 \rangle$.

$$\mathbb{I}(\mathbb{V}(I)) = \langle x^2 + 4y^2 - 4 \rangle.$$

A Rad(ical) Example




Figure: $\mathbb{V}(I)$

Let $I := \langle (x^2 + 4y^2 - 4)^2 \rangle$.

$$\mathbb{I}(\mathbb{V}(I)) = \langle x^2 + 4y^2 - 4 \rangle.$$

But how are $\mathbb{I}(\mathbb{V}(I))$ and I related?

A Rad(ical) Example




Figure: $\mathbb{V}(I)$

Let $I := \langle (x^2 + 4y^2 - 4)^2 \rangle$.

$$\mathbb{I}(\mathbb{V}(I)) = \langle x^2 + 4y^2 - 4 \rangle.$$

But how are $\mathbb{I}(\mathbb{V}(I))$ and I related?

$$\boxed{\mathbb{I}(\mathbb{V}(I)) = \sqrt{I}}$$

The Nullstellensatz

Theorem (Hilbert's Nullstellensatz)

*Let k be an algebraically closed field, and let $I \trianglelefteq k[x_1, \dots, x_n]$.
Then $\mathbb{I}(\mathbb{V}(I)) = \sqrt{I}$. In particular, if I is radical, then $\mathbb{I}(\mathbb{V}(I)) = I$.*

The Nullstellensatz

Theorem (Hilbert's Nullstellensatz)

*Let k be an algebraically closed field, and let $I \trianglelefteq k[x_1, \dots, x_n]$.
Then $\mathbb{I}(\mathbb{V}(I)) = \sqrt{I}$. In particular, if I is radical, then $\mathbb{I}(\mathbb{V}(I)) = I$.*

Restricting to the set of radical ideals:

$$\left\{ \text{algebraic sets in } k^n \right\} \longleftrightarrow \left\{ \text{radical ideals of } k[x_1, \dots, x_n] \right\}$$

The Nullstellensatz

Theorem (Hilbert's Nullstellensatz)

*Let k be an algebraically closed field, and let $I \trianglelefteq k[x_1, \dots, x_n]$.
Then $\mathbb{I}(\mathbb{V}(I)) = \sqrt{I}$. In particular, if I is radical, then $\mathbb{I}(\mathbb{V}(I)) = I$.*

Restricting to the set of radical ideals:

$$\left\{ \text{algebraic sets in } k^n \right\} \longleftrightarrow \left\{ \text{radical ideals of } k[x_1, \dots, x_n] \right\}$$

The field k **must** be algebraically closed!

The Nullstellensatz, Applications

- “Fundamental theorem of algebraic geometry”

The Nullstellensatz, Applications

- “Fundamental theorem of algebraic geometry”
- Nullstellensatz: Shapes \implies abstraction
- Abstraction good: equations \iff shapes

The Nullstellensatz, Applications

- “Fundamental theorem of algebraic geometry”
- Nullstellensatz: Shapes \implies abstraction
- Abstraction good: equations \iff shapes
- Curves in weird fields: $\overline{\mathbb{F}_2}$, etc.
- Elliptic curves, number theory, FLT, etc.