

# Practice with Spinors

## I Algebra with Dirac $\gamma$ matrices

Notation:

1.  $S = \bar{u}u$
2.  $P = \bar{u}\gamma^5 u$
3.  $V^\mu = \bar{u}\gamma^\mu u$
4.  $A^\mu = \bar{u}\gamma^\mu\gamma^5 u$
5.  $T^{\mu\nu} = \bar{u}\sigma^{\mu\nu} u$

1.  $(\bar{u}_1\gamma^\mu u_2)^* = ?$

Note:  $(\gamma^0)^\dagger = \gamma^0$  and  $(\gamma^\mu)^\dagger = \gamma^0\gamma^\mu\gamma^0$

$(\bar{u}_1\gamma^\mu u_2)$  is a  $1 \times 1$  matrix. Therefore, its complex conjugate is the same as its Hermitian conjugate, i.e. if we call  $L^\mu = (\bar{u}_1\gamma^\mu u_2)$ , then  $(L^\mu)^* = (L^\mu)^\dagger$ . We can then express this quantity as follows:

$$\begin{aligned}
 L^\mu &= \bar{u}_1\gamma^\mu u_2, \\
 \Rightarrow (L^\mu)^* &= (L^\mu)^\dagger, \\
 &= (\bar{u}_1\gamma^\mu u_2)^\dagger, \\
 &= ((u_1)^\dagger\gamma^0\gamma^\mu u_2)^\dagger \quad \text{using } (A \dots Z)^\dagger = Z^\dagger \dots A^\dagger, \\
 &= (u_2^\dagger)(\gamma^\mu)^\dagger(\gamma^0)^\dagger(u_1) \\
 &= (u_2^\dagger)\gamma^0\gamma^\mu\gamma^0\gamma^0(u_1) \\
 &= (u_2^\dagger)\gamma^0\gamma^\mu(u_1) \\
 &= \bar{u}_2\gamma^\mu(u_1)
 \end{aligned} \tag{1}$$

Therefore  $(\bar{u}_1\gamma^\mu u_2)^* = \bar{u}_2\gamma^\mu u_1$ . To solve for  $|L^\mu|^2$  we simply use  $(L^{\mu\nu})^2 = \text{Tr}[\bar{u}_1\gamma^\mu u_2\bar{u}_2\gamma^\nu u_1]$  (Why does  $L^\mu$  have only one index but  $L^{\mu\nu} = |L^\mu|^2$ ?)

Note:  $\text{Tr}[\gamma^\mu\gamma^\nu] = 4g^{\mu\nu}$ ,  $\text{Tr}[\gamma^\mu\gamma^\nu\gamma^\lambda\gamma^\sigma] = 4(g^{\mu\nu}g^{\lambda\sigma} - g^{\mu\lambda}g^{\nu\sigma} + g^{\mu\sigma}g^{\nu\lambda})$ , The trace of an odd number gamma matrix product is 0

$$\begin{aligned}
 (L^{\mu\nu})^2 &= \text{Tr}[\bar{u}_1\gamma^\mu u_2\bar{u}_2\gamma^\nu u_1] \\
 &= \text{Tr}[\bar{u}_1\gamma^\mu(\not{p}_2 + m)\gamma^\nu u_1] \\
 &= \text{Tr}[u_1\bar{u}_1\gamma^\mu(\not{p}_2 + m)\gamma^\nu] \\
 &= \text{Tr}[(\not{p}_1 + m)\gamma^\mu(\not{p}_2 + m)\gamma^\nu] \\
 &= \text{Tr}[\not{p}_1\gamma^\mu\not{p}_2\gamma^\nu] + m[\text{Tr}(\gamma^\mu\not{p}_1\gamma^\nu) + \text{Tr}(\gamma^\mu\gamma^\nu\not{p}_2)] + m^2\text{Tr}[\gamma^\mu\gamma^\nu] \\
 &= \text{Tr}[\not{p}_1\gamma^\mu\not{p}_2\gamma^\nu] + m^2\text{Tr}[\gamma^\mu\gamma^\nu] \\
 &= \text{Tr}[(p_1)_\lambda\gamma^\lambda\gamma^\mu(p_2)_\sigma\gamma^\sigma\gamma^\nu] + 4m^2g^{\mu\nu}
 \end{aligned} \tag{2}$$

$$\begin{aligned}
&= (p_1)_\lambda (p_2)_\sigma \text{Tr}[\gamma^\lambda \gamma^\mu \gamma^\sigma \gamma^\nu] + 4m^2 g^{\mu\nu} \\
&= (p_1)_\lambda (p_2)_\sigma 4(g^{\mu\nu} g^{\lambda\sigma} - g^{\mu\lambda} g^{\nu\sigma} + g^{\mu\sigma} g^{\nu\lambda}) + 4m^2 g^{\mu\nu} \\
&= 4[p_1^\mu p_2^\nu - g^{\mu\nu} (p_1 \cdot p_2) + p_2^\mu p_1^\nu] + 4m^2 g^{\mu\nu} \\
&= 4[p_1^\mu p_2^\nu - 4g^{\mu\nu} (p_1 \cdot p_2 + m^2) + p_2^\mu p_1^\nu]
\end{aligned} \tag{3}$$

2.  $(\bar{u}_1 \gamma^\mu \gamma^5 u_2)^*$  is also a  $1 \times 1$  Matrix so the same reasoning applies as above in 1. Note:  
 $(\gamma^5)^\dagger = \gamma^5$

We define:  $R^\mu$  as  $\bar{u}_1 \gamma^\mu \gamma^5 u_2$  thus:

$$\begin{aligned}
\Rightarrow (R^\mu)^* &= (R^\mu)^\dagger \\
&= (\bar{u}_1 \gamma^\mu \gamma^5 u_2)^\dagger \\
&= ((u_1)^\dagger \gamma^0 \gamma^\mu \gamma^5 u_2)^\dagger \\
&= (u_2^\dagger) (\gamma^5)^\dagger (\gamma^\mu)^\dagger (\gamma^0)^\dagger (u_1) \\
&= (u_2^\dagger) \gamma^5 \gamma^0 \gamma^\mu \gamma^0 u_1 \\
&= (u_2^\dagger) \gamma^5 \gamma^0 \gamma^\mu (1) u_1 \\
&= -(u_2^\dagger) \gamma^0 \gamma^5 \gamma^\mu u_1 \\
&= -\bar{u}_2 \gamma^5 \gamma^\mu u_1 \\
&= \bar{u}_2 \gamma^\mu \gamma^5 u_1
\end{aligned} \tag{4}$$

Therefore  $(\bar{u}_1 \gamma^\mu \gamma^5 u_2)^* = \bar{u}_2 \gamma^\mu \gamma^5 u_1$

We also are able to calculate  $(R^\mu)^2$

$$\begin{aligned}
(R^{\mu\nu})^2 &= \text{Tr}[\bar{u}_1 \gamma^\mu \gamma^5 u_2 \bar{u}_2 \gamma^\nu \gamma^5 u_1] \\
&= \text{Tr}[\bar{u}_1 \gamma^\mu \gamma^5 (\not{p}_2 + m) \gamma^\nu \gamma^5 u_1] \\
&= \text{Tr}[u_1 \bar{u}_1 \gamma^\mu \gamma^5 (\not{p}_2 + m) \gamma^\nu \gamma^5] \\
&= \text{Tr}[(\not{p}_1 + m) \gamma^\mu \gamma^5 (\not{p}_2 + m) \gamma^\nu \gamma^5] \\
&= \text{Tr}[\not{p}_1 \gamma^\mu \gamma^5 \not{p}_2 \gamma^\nu \gamma^5 + m(\not{p}_1 \gamma^\mu \gamma^5 \gamma^\nu \gamma^5 + \gamma^\mu \gamma^5 \not{p}_2 \gamma^\nu \gamma^5) + m^2(\gamma^\mu \gamma^5 \gamma^\nu \gamma^5)] \\
&= \text{Tr}[\not{p}_1 \gamma^\mu \gamma^5 \not{p}_2 \gamma^\nu \gamma^5 + m^2(\gamma^\mu \gamma^5 \gamma^\nu \gamma^5)] \\
&= \text{Tr}[(p_1)_\lambda \gamma^\lambda \gamma^\mu \gamma^5 (p_2)_\sigma \gamma^\sigma \gamma^\nu \gamma^5] + m^2 \text{Tr}[\gamma^\mu \gamma^5 \gamma^\nu \gamma^5] \\
&= (p_1)_\lambda (p_2)_\sigma \text{Tr}[\gamma^\lambda \gamma^\mu \gamma^5 \gamma^\sigma \gamma^\nu \gamma^5] - m^2 \text{Tr}[\gamma^\mu \gamma^5 \gamma^5 \gamma^\nu] \\
&= (p_1)_\lambda (p_2)_\sigma \text{Tr}[\gamma^\lambda \gamma^\mu \gamma^5 \gamma^5 \gamma^\sigma \gamma^\nu] - m^2 \text{Tr}[\gamma^\mu \gamma^\nu] \\
&= (p_1)_\lambda (p_2)_\sigma \text{Tr}[\gamma^\lambda \gamma^\mu \gamma^\sigma \gamma^\nu] - m^2(g^{\mu\nu}) \\
&= (p_1)_\lambda (p_2)_\sigma 4(g^{\mu\nu} g^{\lambda\sigma} - g^{\mu\lambda} g^{\nu\sigma} + g^{\mu\sigma} g^{\nu\lambda}) - 4m^2 g^{\mu\nu} \\
&= 4[p_1^\mu p_2^\nu - g^{\mu\nu} (p_1 \cdot p_2) + p_2^\mu p_1^\nu] - 4m^2 g^{\mu\nu} \\
&= 4[p_1^\mu p_2^\nu - 4g^{\mu\nu} (p_1 \cdot p_2 - m^2) + p_2^\mu p_1^\nu]
\end{aligned} \tag{-11}$$

3.  $(\bar{u}_1 u_2)^* = ?$

We let:  $P = \bar{u}_1 u_2$

$$\Rightarrow (P)^* = (P)^\dagger, \tag{-11}$$

$$\begin{aligned}
&= (\bar{u}_1 u_2)^\dagger, \\
&= ((u_1)^\dagger \gamma^0 u_2)^\dagger \\
&= (u_2)^\dagger (\gamma^0)^\dagger (u_1) \\
&= (u_2)^\dagger \gamma^0 (u_1) \\
&= \bar{u}_2 (u_1)
\end{aligned} \tag{-14}$$

Therefore  $(\bar{u}_1 u_2)^* = \bar{u}_2 u_1$

In order to find  $(P)^2$  we simply do the following:

$$\begin{aligned}
(P)^2 &= \text{Tr}[\bar{u}_1 u_2 \bar{u}_2 u_1] \\
&= \text{Tr}[(\not{p}_1 + m)(\not{p}_2 + m)] \\
&= \text{Tr}[\not{p}_1 \not{p}_2 + m(\not{p}_1 + \not{p}_2) + m^2] \\
&= \text{Tr}[\not{p}_1 \not{p}_2] + \text{Tr}[m(\not{p}_1 + \not{p}_2)] + \text{Tr}[m^2] \\
&= \text{Tr}[\not{p}_1 \not{p}_2] + m(\text{Tr}[\not{p}_1] + \text{Tr}[\not{p}_2]) + m^2 \text{Tr}[1] \\
&= \text{Tr}[\not{p}_1 \not{p}_2] + 4m^2 \\
&= 4(p_1 \cdot p_2) + 4m^2
\end{aligned} \tag{-18}$$

4. By the same reasoning as shown above it can be shown that  $(\bar{u}_1 \gamma^5 u_2)^* = \bar{u}_2 \gamma^5 u_1$   
 If we let  $T = \bar{u}_1 \gamma^5 u_2$  then:

$$\begin{aligned}
\Rightarrow (T)^* &= (T)^\dagger, \\
&= (\bar{u}_1 \gamma^5 u_2)^\dagger, \\
&= ((u_1)^\dagger \gamma^0 \gamma^5 u_2)^\dagger \\
&= (u_2)^\dagger (\gamma^5)^\dagger (\gamma^0)^\dagger (u_1) \\
&= (u_2)^\dagger (-\gamma^5) \gamma^0 (u_1) \\
&= (u_2)^\dagger \gamma^0 \gamma^5 (u_1) \\
&= \bar{u}_2 \gamma^5 (u_1)
\end{aligned} \tag{-22}$$

Therefore  $(\bar{u}_1 \gamma^5 u_2)^* = \bar{u}_2 \gamma^5 u_1$

In order to square  $T$  we do the following:

$$\begin{aligned}
T^2 &= \text{Tr}[\bar{u}_1 \gamma^5 u_2 \bar{u}_2 \gamma^5 u_1] \\
&= \text{Tr}[u_1 \bar{u}_1 \gamma^5 \not{p}_2 \gamma^5] \\
&= \text{Tr}[\not{p}_1 \gamma^5 \not{p}_2 \gamma^5] \\
&= \text{Tr}[(p_1)_\mu \gamma^\mu \gamma^5 (p_2)_\nu \gamma^\nu \gamma^5] \\
&= (p_1)_\mu (p_2)_\nu \text{Tr}[\gamma^\mu \gamma^5 \gamma^\nu \gamma^5] \\
&= -(p_1)_\mu (p_2)_\nu \text{Tr}[\gamma^\mu \gamma^5 \gamma^5 \gamma^\nu] \\
&= -(p_1)_\mu (p_2)_\nu \text{Tr}[\gamma^\mu \gamma^\nu] \\
&= -(p_1)_\mu (p_2)_\nu (4g^{\mu\nu}) \\
&= -4(p_1)(p_2)
\end{aligned} \tag{-21}$$

(-28)

5. While the above identities could be shown to be trivial, the identity:  $(\bar{u}_1 \sigma^{\mu\nu} u_2)^* = \bar{u}_2 \sigma^{\mu\nu} u_1$  is more difficult to solve

The identity:  $(\sigma^{\mu\nu})^\dagger = \sigma^{\mu\nu}$  is needed

$$\begin{aligned}
(\sigma^{\mu\nu})^\dagger &= \left(\frac{i}{2}[\gamma^\mu, \gamma^\nu]\right)^\dagger \\
&= (u_2)^\dagger (\sigma^{\mu\nu})^\dagger (\gamma^0)^\dagger u_1 \\
&= \frac{i}{2}([\gamma^\mu, \gamma^\nu])^\dagger \\
&= \frac{i}{2}(\gamma^\mu \gamma^\nu - \gamma^\nu \gamma^\mu)^\dagger \\
&= \frac{i}{2}((\gamma^\mu)^\dagger (\gamma^\nu)^\dagger - (\gamma^\nu)^\dagger (\gamma^\mu)^\dagger) \\
&= \frac{i}{2}(\gamma^0 \gamma^\mu \gamma^0 \gamma^0 \gamma^\nu \gamma^0 - \gamma^0 \gamma^\nu \gamma^0 \gamma^0 \gamma^\mu \gamma^0) \\
&= \frac{i}{2}(\gamma^0 \gamma^\mu \gamma^\nu \gamma^0 - \gamma^0 \gamma^\nu \gamma^\mu \gamma^0) \\
&= \frac{i}{2}((-1)^2 \gamma^\mu \gamma^\nu - (-1)^2 \gamma^\nu \gamma^\mu) \\
&= \frac{i}{2}(\gamma^\mu \gamma^\nu - \gamma^\nu \gamma^\mu) \\
&= \sigma^{\mu\nu}
\end{aligned} \tag{-28}$$

After showing  $(\sigma^{\mu\nu})^\dagger = \sigma^{\mu\nu}$  is true it is trivial to show  $(\bar{u}_1 \sigma^{\mu\nu} u_2)^* = \bar{u}_2 \sigma^{\mu\nu} u_1$   
We let  $B^{\mu\nu} = \bar{u}_1 \sigma^{\mu\nu} u_2$

$$\begin{aligned}
\Rightarrow (B^{\mu\nu})^* &= (B^{\mu\nu})^\dagger, \\
&= (\bar{u}_1 \sigma^{\mu\nu} u_2)^\dagger, \\
&= ((u_1)^\dagger \gamma^0 \sigma^{\mu\nu} u_2)^\dagger \\
&= (u_2^\dagger) (\sigma^{\mu\nu})^\dagger (\gamma^0)^\dagger (u_1) \\
&= (u_2^\dagger) \sigma^{\mu\nu} \gamma^0 (u_1) \\
&= (u_2^\dagger) (\gamma^0) \sigma^{\mu\nu} (u_1) \\
&= \bar{u}_2 \sigma^{\mu\nu} (u_1)
\end{aligned} \tag{-35}$$

An interesting thing to note is that the expression  $\bar{u} \sigma^{\mu\nu} \gamma^5 u$  is not an independent quantity. Since  $\gamma^5 = i\gamma^0 \gamma^1 \gamma^2 \gamma^3$  it follows that the product of  $\sigma^{\mu\nu}$  and  $\gamma^5$  can be simplified to an expression with only 2  $\gamma$  matrices which has been defined as a pseudoscalar.

For example, let  $\mu = 0$  and  $\nu = 1$ :

$$\begin{aligned}
\bar{u} \sigma^{01} \gamma^5 u &= \bar{u} \sigma^{01} (i\gamma^0 \gamma^1 \gamma^2 \gamma^3) u \\
&= \bar{u} \left(\frac{i}{2}(\gamma^0 \gamma^1 - \gamma^1 \gamma^0)\right) (i\gamma^0 \gamma^1 \gamma^2 \gamma^3) u \\
&= \bar{u} \left(\frac{i}{2}\right) [\gamma^0 \gamma^1 (i\gamma^0 \gamma^1 \gamma^2 \gamma^3) - \gamma^1 \gamma^0 (i\gamma^0 \gamma^1 \gamma^2 \gamma^3)] u \\
&= \bar{u} \left(\frac{-1}{2}\right) [\gamma^0 \gamma^1 \gamma^0 \gamma^1 \gamma^2 \gamma^3 - \gamma^1 \gamma^0 \gamma^0 \gamma^1 \gamma^2 \gamma^3] u
\end{aligned} \tag{-39}$$

$$\begin{aligned}
&= \bar{u}\left(\frac{-1}{2}\right)[- \gamma^2 \gamma^3 - \gamma^2 \gamma^3]u \\
&= \bar{u}\left(\frac{-1}{2}\right)[-2\gamma^2 \gamma^3]u \\
&= 2\bar{u}\gamma^2 \gamma^3 u
\end{aligned} \tag{-43}$$

This expression (because it contains two gamma matrices) is a pseudoscalar. Any values of  $\mu$  and  $\nu$  can be shown to be similar to this because of the commutative and identity properties of the gamma matrices.

In order to find the value of  $(B^{\mu\nu})^2$  one needs to do the following:

$$\begin{aligned}
(B^{\mu\nu\sigma\lambda})^2 &= \text{Tr}[\bar{u}_1 \sigma^{\mu\nu} u_2 \bar{u}_2 \sigma^{\sigma\lambda} u_1] \\
&= \text{Tr}[\bar{u}_1 \sigma^{\mu\nu} \not{p}_2 \sigma^{\sigma\lambda} u_1] \\
&= \text{Tr}[\not{p}_1 \sigma^{\mu\nu} \not{p}_2 \sigma^{\sigma\lambda}] \\
&= \text{Tr}[p_\kappa \gamma^\kappa \sigma^{\mu\nu} p_\gamma \gamma^\gamma \sigma^{\sigma\lambda}] \\
&= p_\kappa p_\gamma \text{Tr}[\gamma^\kappa \sigma^{\mu\nu} \gamma^\gamma \sigma^{\sigma\lambda}] \\
&= -\frac{1}{2} p_\kappa p_\gamma \text{Tr}[\gamma^\kappa (\gamma^\mu \gamma^\nu - \gamma^\nu \gamma^\mu) \gamma^\gamma (\gamma^\sigma \gamma^\lambda - \gamma^\lambda \gamma^\sigma)] \\
&= -\frac{1}{2} p_\kappa p_\gamma \text{Tr}[2\gamma^\kappa \gamma^\mu \gamma^\nu \gamma^\gamma \gamma^\sigma \gamma^\lambda - 2\gamma^\kappa \gamma^\nu \gamma^\mu \gamma^\gamma \gamma^\sigma \gamma^\lambda] \\
&= -\frac{1}{2} p_\kappa p_\gamma \text{Tr}[4\gamma^\kappa \gamma^\mu \gamma^\nu \gamma^\gamma \gamma^\sigma \gamma^\lambda] \\
&= -2p_\kappa p_\gamma \text{Tr}[\gamma^\kappa \gamma^\gamma] \text{ Using a similar identity as shown in eq. 23} \\
&= -8p_\kappa p_\gamma g^{\kappa\gamma}
\end{aligned} \tag{-51}$$