

# Faster mixing of the Jerrum-Sinclair chain

Weiming Feng

The University of Hong Kong

Joint work with

Xiaoyu Chen, Zhe Ju, Tianshun Miao, Yitong Yin, Xinyuan Zhang


Nanjing University

Conference on Mixing Times between Probability, Computer Science and Statistical Physics  
ICTP, Italy, 9 May 2025

# Monomer-dimer model

Simple graph  $G = (V, E)$  and edge weight  $\lambda > 0$


$\forall$  matching  $M \subseteq E$ ,  $\mu(M) \propto \lambda^{|M|}$



# Jerrum-Sinclair chain (Metropolis chain)

Jerrum-Sinclair chain updates matching  $X_t \rightarrow X_{t+1}$  by


- select an edge  $e = \{u, v\} \in E$  u.a.r.
- propose a candidate matching  $M$  from  $X_t$  by
  - 1) **down transition**: if  $e \in X_t$ , set  $M \leftarrow X_t - e$



# Jerrum-Sinclair chain (Metropolis chain)

Jerrum-Sinclair chain updates matching  $X_t \rightarrow X_{t+1}$  by


- select an edge  $e = \{u, v\} \in E$  u.a.r.
- propose a candidate matching  $M$  for  $X_{t+1}$  by
  - 1) **down transition**: if  $e \in X_t$ , set  $M \leftarrow X_t - e$
  - 2) **up transition**: if both  $u, v$  are **not saturated** in  $X_t$ , set  $M \leftarrow X_t + e$



# Jerrum-Sinclair chain (Metropolis chain)

Jerrum-Sinclair chain updates matching  $X_t \rightarrow X_{t+1}$  by


- select an edge  $e = \{u, v\} \in E$  u.a.r.
- propose a candidate matching  $M$  for  $X_{t+1}$  by
  - 1) **down transition**: if  $e \in X_t$ , set  $M \leftarrow X_t - e$
  - 2) **up transition**: if both  $u, v$  are **not saturated** in  $X_t$ , set  $M \leftarrow X_t + e$
  - 3) **exchange transition**: if one endpoint is **saturated** and the other is **not**, say  $u$  is saturated by edge  $f$  and  $v$  is not, set  $M \leftarrow X_t + e - f$



# Jerrum-Sinclair chain (Metropolis chain)

Jerrum-Sinclair chain updates matching  $X_t \rightarrow X_{t+1}$  by

- select an edge  $e = \{u, v\} \in E$  u.a.r.
- propose a candidate matching  $M$  for  $X_{t+1}$  by
  - 1) **down transition**: if  $e \in X_t$ , set  $M \leftarrow X_t - e$
  - 2) **up transition**: if both  $u, v$  are **not saturated** in  $X_t$ , set  $M \leftarrow X_t + e$
  - 3) **exchange transition**: if one endpoint is **saturated** and the other is **not**, say  $u$  is saturated by edge  $f$  and  $v$  is not, set  $M \leftarrow X_t + e - f$
  - 4) **otherwise** (both  $u, v$  are saturated and  $e \notin X_t$ ): set  $M \leftarrow X_t$



# Jerrum-Sinclair chain (Metropolis chain)

Jerrum-Sinclair chain  $P_{JS}$  updates matching  $X_t \rightarrow X_{t+1}$  by

- select an edge  $e = \{u, v\} \in E$  u.a.r.
- propose a candidate matching  $M$  for  $X_{t+1}$  by
  - 1) **down transition**: if  $e \in X_t$ , set  $M \leftarrow X_t - e$
  - 2) **up transition**: if both  $u, v$  are **not saturated** in  $X_t$ , set  $M \leftarrow X_t + e$
  - 3) **exchange transition**: if one endpoint is **saturated** and the other is **not**, say  $u$  is saturated by edge  $f$  and  $v$  is not, set  $M \leftarrow X_t + e - f$
  - 4) **otherwise** (both  $u, v$  are saturated and  $e \notin X_t$ ): set  $M \leftarrow X_t$
- with prob.  $\min \left\{ 1, \frac{\mu(M)}{\mu(X_t)} \right\}$ , accept  $M$  and set  $X_{t+1} \leftarrow M$ ; otherwise,  $X_{t+1} \leftarrow X_t$  (**Metropolis filter**)


1/2-Lazy Jerrum-Sinclair chain:  $P_{JS\_zz} = \frac{1}{2}(P_{JS} + I)$

$$\textbf{Mixing time: } T_{\text{mix}}(P_{JS\_zz}) = \max_{X_0} \min \left\{ t > 0 \mid \|X_t - \mu\|_{TV} \leq \frac{1}{4e} \right\}$$

# Mixing time results

**Jerrum-Sinclair (1989):** Graph with  $n$  vertices and  $m$  edges; constant  $\lambda > 0$ ,

$$T_{\text{mix}}(P_{JS\_zz}) = \tilde{O}(mn^2)$$

**Canonical Path:**  $O(nm)$ -congestion with  $O(n)$  path length  **spectral gap**  $\gamma = \Omega\left(\frac{1}{nm}\right)$

**Mixing time:**  $T_{\text{mix}}(P_{JS\_zz}) = O\left(\frac{1}{\gamma} \log \frac{1}{\mu_{\min}}\right)$ ,  $\mu_{\min} = \min_M \mu(M)$  and  $\log \frac{1}{\mu_{\min}} = \tilde{O}(n)$

**Chen-Liu-Vigoda (2021):** Graph with  $n$  vertices,  $m$  edges, and max degree  $\Delta$ ; constant  $\lambda > 0$ ,

$$T_{\text{mix}}(\text{Glauber dynamics}) = O(\Delta^{\Delta^2} \cdot m \log n)$$

**Spectral independence (local-to-global in HDX)**  **modified log-Sobolev const.**  $\alpha = \Omega_{\Delta}\left(\frac{1}{m}\right)$

**Our Result:**  $T_{\text{mix}}(P_{JS\_zz}) = O(\Delta m \cdot \min\{n, \Delta \log \Delta \log n\}) = \tilde{O}(m\Delta^2)$

- **Spectral gap:**  $\Omega\left(\frac{1}{m\Delta}\right)$
- **Log-Sobolev const.:**  $\Omega\left(\frac{1}{m\Delta^2}\right)$

**Corollary:**  $T_{\text{mix}}(\text{Glauber dynamics}) = \tilde{O}(\Delta^3 \cdot m)$



# General results

- **Distribution**  $\mu$  over  $\Omega \subseteq [q]^E$  for finite domain  $[q] = \{1, 2, \dots, q\}$  and variable set  $E$
- **Random variable**  $F = f(X)$  for a function  $f: \Omega \rightarrow \mathbb{R}$  and  $X \sim \mu$

**Variance:**  $\text{Var}[F] = \text{Var}_\mu[f] = \mathbb{E}[F^2] - \mathbb{E}[F]^2$

**Entropy:**  $\text{Ent}[F] = \text{Ent}_\mu[f] = \mathbb{E}[F \log F] - \mathbb{E}[F] \log \mathbb{E}[F]$

- **Reversible Markov chain**  $Q$  for  $\mu: \forall x, y, \mu(x)Q(x, y) = \mu(y)Q(y, x)$

**Dirichlet form:**  $\mathcal{E}_Q(f, f) = \frac{1}{2} \sum_{x, y \in \Omega} \mu(x)Q(x, y)(f(x) - f(y))^2$  for all  $f: \Omega \rightarrow \mathbb{R}$

## Poincaré Inequality (Spectral Gap)

$$\gamma(Q) \cdot \text{Var}_\mu[f] \leq \mathcal{E}_Q(f, f)$$

➡  $T_{\text{mix}}\left(\frac{Q + I}{2}\right) = O\left(\frac{1}{\gamma(Q)} \log \frac{1}{\mu_{\min}}\right)$


## Log-Sobolev Inequality

$$\rho(Q) \cdot \text{Ent}_\mu[F^2] \leq \mathcal{E}_Q(f, f)$$

➡  $T_{\text{mix}}(Q) = O\left(\frac{1}{\rho(Q)} \log \log \frac{1}{\mu_{\min}}\right)$

## Family of Markov chains

- For a subset  $\Lambda \subseteq E$ , a pinning  $\tau \in [q]^{E \setminus \Lambda}$  outside  $\Lambda$ , define **conditional distribution**  
 $\mu^\tau = (\text{distribution of } X \sim \mu \text{ conditional on } X_{E \setminus \Lambda} = \tau)$



## Family of Markov chains


- For a subset  $\Lambda \subseteq E$ , a pinning  $\tau \in [q]^{E \setminus \Lambda}$  outside  $\Lambda$ , define **conditional distribution**  
 $\mu^\tau = (\text{distribution of } X \sim \mu \text{ conditional on } X_{E \setminus \Lambda} = \tau)$
- Markov chain  $Q^\tau$  is a reversible chain for  $\mu^\tau$

a family of chains  $\mathcal{Q} = \{Q^\tau \mid \tau \text{ is a pinning}\}$

Example:  $\mathcal{Q}$  is a family of *Glauber dynamics* or a family of *Metropolis chains*.

**Concave Dirichlet forms:**  $\forall \Lambda \subseteq E, \forall \tau \in [q]^{E \setminus \Lambda}$ , the Markov chain  $Q^\tau$  for  $\mu^\tau$  satisfies

$$\frac{1}{|\Lambda|} \sum_{e \in \Lambda} \mathbb{E}_{c \sim \mu_e^\tau} \left[ \mathcal{E}_{Q^{\tau \wedge (e \leftarrow c)}}(f, f) \right] \leq \mathcal{E}_{Q^\tau}(f, f).$$



## Family of Markov chains

- For a subset  $\Lambda \subseteq E$ , a pinning  $\tau \in [q]^{E \setminus \Lambda}$  outside  $\Lambda$ , define **conditional distribution**  
 $\mu^\tau = (\text{distribution of } X \sim \mu \text{ conditional on } X_{E \setminus \Lambda} = \tau)$
- Markov chain  $Q^\tau$  is a reversible chain for  $\mu^\tau$

a family of chains  $\mathcal{Q} = \{Q^\tau \mid \tau \text{ is a pinning}\}$

Example:  $\mathcal{Q}$  is a family of *Glauber dynamics* or a family of *Metropolis chains*.

**Concave Dirichlet forms:**  $\forall \Lambda \subseteq E, \forall \tau \in [q]^{E \setminus \Lambda}$ , the Markov chain  $Q^\tau$  for  $\mu^\tau$  satisfies

$$\underbrace{\frac{1}{|\Lambda|} \sum_{e \in \Lambda} \mathbb{E}_{c \sim \mu_e^\tau} \left[ \mathcal{E}_{Q^{\tau \wedge (e \leftarrow c)}}(f, f) \right]}_{\text{average of Dirichlet forms}} \leq \underbrace{\mathcal{E}_{Q^\tau}(f, f)}_{\text{Dirichlet forms of } Q^\tau \text{ for } \mu^\tau}.$$

Markov chain for conditional distributions  
with **one more pinned variable**

## Local functional inequalities

Distribution  $\mu$  over  $\Omega \subseteq [q]^E$ , random variables  $F = f(X)$ , where  $f: \Omega \rightarrow \mathbb{R}$  and  $X \sim \mu$

### $\alpha$ -local Poincaré Inequality

$$\alpha \cdot \sum_{e \in E} \text{Var} \left[ \mathbb{E}[F \mid X_e] \right] \leq \mathcal{E}_Q(f, f)$$

sum over all  
variables in  $E$

a random variable in  $\mathbb{R}$   
support size at most  $q$

### $\alpha$ -local log-Sobolev Inequality

$$\alpha \cdot \sum_{e \in E} \text{Ent} \left[ \mathbb{E}[F^2 \mid X_e] \right] \leq \mathcal{E}_Q(f, f)$$

## Local functional inequalities for a family of Markov chains $\mathcal{Q}$

$(\alpha_1, \alpha_2, \dots, \alpha_{|E|})$ -local Poincaré Inequality

$\forall \Lambda \subseteq E, \forall \tau \in [q]^{E \setminus \Lambda}$ , the Markov chain  $Q^\tau$  for  $\mu^\tau$   
satisfies the  $\alpha_{|\Lambda|}$ -local Poincaré Inequality

set of  $k = |\Lambda|$   
**free** variables  
 $\Lambda$

**pinning**  $\tau$   
on  $E \setminus \Lambda$

## Local functional inequalities for a family of Markov chains $\mathcal{Q}$

$(\alpha_1, \alpha_2, \dots, \alpha_{|E|})$ -local Poincaré Inequality

$\forall \Lambda \subseteq E, \forall \tau \in [q]^{E \setminus \Lambda}$ , the Markov chain  $Q^\tau$  for  $\mu^\tau$  satisfies the  $\alpha_{|\Lambda|}$ -local Poincaré Inequality

$$\alpha_{|\Lambda|} \cdot \sum_{e \in E} \text{Var} [\mathbb{E}[F \mid X_e]] \leq \varepsilon_{Q^\tau}(f, f),$$

where  $F = f(X)$  and  $X \sim \mu^\tau$

## Local functional inequalities for a family of Markov chains $\mathcal{Q}$

$(\alpha_1, \alpha_2, \dots, \alpha_{|E|})$ -local Poincaré Inequality

$\forall \Lambda \subseteq E, \forall \tau \in [q]^{E \setminus \Lambda}$ , the Markov chain  $Q^\tau$  for  $\mu^\tau$  satisfies the  $\alpha_{|\Lambda|}$ -local Poincaré inequality

$(\alpha_1, \alpha_2, \dots, \alpha_{|E|})$ -local log-Sobolev Inequality

$\forall \Lambda \subseteq E, \forall \tau \in [q]^{E \setminus \Lambda}$ , the Markov chain  $Q^\tau$  for  $\mu^\tau$  satisfies the  $\alpha_{|\Lambda|}$ -local log-Sobolev inequality



## Local functional inequalities for a family of Markov chains $\mathcal{Q}$

$(\alpha_1, \alpha_2, \dots, \alpha_{|E|})$ -local Poincaré Inequality

$\forall \Lambda \subseteq E, \forall \tau \in [q]^{E \setminus \Lambda}$ , the Markov chain  $Q^\tau$  for  $\mu^\tau$  satisfies the  $\alpha_{|\Lambda|}$ -local Poincaré inequality

$(\alpha_1, \alpha_2, \dots, \alpha_{|E|})$ -local log-Sobolev Inequality

$\forall \Lambda \subseteq E, \forall \tau \in [q]^{E \setminus \Lambda}$ , the Markov chain  $Q^\tau$  for  $\mu^\tau$  satisfies the  $\alpha_{|\Lambda|}$ -local log-Sobolev inequality

### Local-to-global theorem for functional inequalities

For a family of Markov chains  $\mathcal{Q}$  with the concave Dirichlet forms

$(\alpha_1, \alpha_2, \dots, \alpha_{|E|})$ -local Poincaré Inequality



Poincaré constant  $\gamma(Q) \geq \left( \sum_{k=1}^{|E|} \frac{1}{k\alpha_k} \right)^{-1}$

$(\alpha_1, \alpha_2, \dots, \alpha_{|E|})$ -local log-Sobolev Inequality



Log-Sobolev constant  $\rho(Q) \geq \left( \sum_{k=1}^{|E|} \frac{1}{k\alpha_k} \right)^{-1}$

**$Q$ : Markov chain for  $\mu = \mu^\emptyset$  without pinning**

## How to establish local functional inequalities?

### $\alpha$ -local Poincaré Inequality

$$\alpha \cdot \sum_{e \in E} \text{Var} [\mathbb{E}[F \mid X_e]] \leq \mathcal{E}_Q(f, f)$$


### $\alpha$ -local log-Sobolev Inequality

$$\alpha \cdot \sum_{e \in E} \text{Ent} [\mathbb{E}[F^2 \mid X_e]] \leq \mathcal{E}_Q(f, f)$$

## Transport Flow



Given a Markov chain  $Q$ , a transport flow  $\Gamma$  from a **distribution  $\nu$**  to a **distribution  $\pi$**  is **a distribution of paths** such that  $\gamma = (x_0, x_1, \dots, x_\ell) \sim \Gamma$  satisfies

- The starting point  $s(\gamma) = x_0 \sim \nu$
  - The endpoint  $t(\gamma) = x_\ell \sim \pi$
  - Every pair of adjacent points  $(x_i, x_{i+1})$  is a **transition** in  $Q$
- }  $(s(\gamma), t(\gamma))$  forms a **coupling** of  $\nu$  and  $\pi$



For any variable  $e \in E$


- **partition** the space by **the value of  $e$**
- **routing** between different parts




$$\Omega(ea) = \{X \in \Omega \mid X_e = a\}$$



sending  $\mu_e(a)\mu_e(b)$   
units of flow in total




$$\Omega(eb) = \{X \in \Omega \mid X_e = b\}$$



For any variable  $e \in E$

- **partition** the space by **the value of  $e$**
- **routing** between different parts



$\forall \sigma \in \Omega(ea)$  needs to **send**  
 $\mu_e(a)\mu_e(b)\mu^{e \leftarrow a}(\sigma)$  unit of flow

$$\Omega(ea) = \{X \in \Omega \mid X_e = a\}$$

routing via  
transitions in  $Q$



$\forall \tau \in \Omega(eb)$  needs to **receive**  
 $\mu_e(a)\mu_e(b)\mu^{e \leftarrow b}(\tau)$  unit of flow


sending  $\mu_e(a)\mu_e(b)$   
units of flow in total

$$\Omega(eb) = \{X \in \Omega \mid X_e = a\}$$





Sending  $\mu_e(a)\mu_e(b)$  units of flow via a **random path**

from the **transport flow**  $\Gamma_e^{a \rightarrow b}$  from  $\mu^{e \leftarrow a}$  to  $\mu^{e \leftarrow b}$



For any variable  $e \in E$


- **partition** the space by **the value of  $e$**
- **routing** between different parts



$$\Omega(ea) = \{X \in \Omega \mid X_e = a\}$$



sending  $\mu_e(a)\mu_e(b)$   
units of flow in total



$$\Omega(eb) = \{X \in \Omega \mid X_e = a\}$$



Every path  $\gamma = (x_0, x_1, \dots, x_\ell)$

send  $\mu_e(a)\mu_e(b) \Pr_{X \sim \Gamma_e^{a \rightarrow b}}[X = \gamma]$  units of flow from  $x_0$  to  $x_\ell$

## Local Poincaré inequality via transport flow

If there exists a *family of transport flow*

$$\{\Gamma_e^{a \rightarrow b} \text{ from } \mu^{e \leftarrow a} \text{ to } \mu^{e \leftarrow b} \mid e \in E, a, b \in [q]\}$$


## Local Poincaré inequality via transport flow

If there exists a *family of transport flow*

$$\{\Gamma_e^{a \rightarrow b} \text{ from } \mu^{e \leftarrow a} \text{ to } \mu^{e \leftarrow b} \mid e \in E, a, b \in [q]\}$$

- ( $\kappa$ -*expected congestion*) For *any transition*  $(x \rightarrow y)$  in  $Q$ , and any  $a, b \in [q]$ ,

$$\sum_{e \in E} \mu_e(a) \mu_e(b) \cdot \Pr_{\gamma \sim \Gamma_e^{a \rightarrow b}} [(x \rightarrow y) \in \gamma] \leq \kappa \cdot \mu(x) Q(x, y).$$



## Local Poincaré inequality via transport flow

If there exists a *family of transport flow*

$$\{\Gamma_e^{a \rightarrow b} \text{ from } \mu^{e \leftarrow a} \text{ to } \mu^{e \leftarrow b} \mid e \in E, a, b \in [q]\}$$

- ( $\kappa$ -**expected congestion**) For **any transition**  $(x \rightarrow y)$  in  $Q$ , and any  $a, b \in [q]$ ,

$$\sum_{e \in E} \mu_e(a) \mu_e(b) \cdot \Pr_{\gamma \sim \Gamma_e^{a \rightarrow b}} [(x \rightarrow y) \in \gamma] \leq \kappa \cdot \mu(x) Q(x, y).$$

- ( $L$ -**expected length**) For any  $e \in E$ , any  $a, b \in [q]$ ,

$$\mathbb{E}_{\gamma \sim \Gamma_e^{a \rightarrow b}} [\ell(\gamma)] \leq L$$

*Slightly different definitions  
are used in the paper to  
improve the application*



$\alpha$ -local Poincaré Inequality

$$\alpha \cdot \sum_{e \in E} \text{Var} [\mathbb{E}[F \mid X_e]] \leq \mathcal{E}_Q(f, f) \text{ with } \alpha = \Omega\left(\frac{1}{q^2 \kappa L}\right)$$



## Local Poincaré inequality via transport flow

Family of transport flow

- low expected congestion
- low expected length



local Poincaré Inequality

Construct transport flow  $\Gamma_e^{a \rightarrow b}$  from  $\mu^{e \leftarrow a}$  to  $\mu^{e \leftarrow b}$  such that for  $\gamma = (x_0, x_1, \dots, x_\ell) \sim \Gamma$


- The starting point  $s(\gamma) = x_0 \sim \mu^{e \leftarrow a}$
  - The endpoint  $t(\gamma) = x_\ell \sim \mu^{e \leftarrow b}$
  - Every pair of adjacent points  $(x_i, x_{i+1})$  is a **transition** in  $Q$
- $(s(\gamma), t(\gamma))$  forms a **coupling** of  $\mu^{e \leftarrow a}$  and  $\mu^{e \leftarrow b}$

Find a **good coupling** with small **expected discrepancy** between  $\mu^{e \leftarrow a}$  and  $\mu^{e \leftarrow b}$

# Canonical path and multicommodity flow

The technique [Diaconis and Stroock 91] [Sinclair 92] is to bound **global variance**


$$\alpha \cdot \text{Var}[F] \leq \varepsilon_Q(f, f)$$



- **Canonical path**: sending flow through one path
- **Multicommodity flow**: sending flow through a distribution of paths

## Proof outline: Local Poincaré inequality via transport flow

$$\text{Var}[\mathbb{E}[F \mid X_e]] = \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a) \mu_e(b) (\mathbb{E}[F \mid X_e = a] - \mathbb{E}[F \mid X_e = b])^2$$

  
*local variance*

  
*amount of flow*

## Proof outline: Local Poincaré inequality via transport flow

$$\begin{aligned}\text{Var}[\mathbb{E}[F \mid X_e]] &= \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a) \mu_e(b) (\mathbb{E}[F \mid X_e = a] - \mathbb{E}[F \mid X_e = b])^2 \\ &= \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a) \mu_e(b) \left( \mathbb{E}_{X \sim \mu^{e \leftarrow a}}[f(X)] - \mathbb{E}_{Y \sim \mu^{e \leftarrow b}}[f(Y)] \right)^2\end{aligned}$$

By definition  $F = f(x)$

*beginning of the  
transport flow*

*ending of the  
transport flow*

## Proof outline: Local Poincaré inequality via transport flow

$$\text{Var}[\mathbb{E}[F \mid X_e]] = \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a)\mu_e(b) (\mathbb{E}[F \mid X_e = a] - \mathbb{E}[F \mid X_e = b])^2$$

By definition  $F = f(x)$

$$= \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a)\mu_e(b) \left( \mathbb{E}_{X \sim \mu^{e \leftarrow a}}[f(X)] - \mathbb{E}_{Y \sim \mu^{e \leftarrow b}}[f(Y)] \right)^2$$

Coupling :  $(x_0, x_\ell) \sim (\mu^{e \leftarrow a}, \mu^{e \leftarrow b})$

$$= \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a)\mu_e(b) \left( \mathbb{E}_{\gamma=(x_0, x_1, \dots, x_\ell) \sim \Gamma_e^{a \rightarrow b}} [f(x_0) - f(x_\ell)] \right)^2$$



*sample a random path from the flow*

## Proof outline: Local Poincaré inequality via transport flow

$$\text{Var}[\mathbb{E}[F \mid X_e]] = \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a) \mu_e(b) (\mathbb{E}[F \mid X_e = a] - \mathbb{E}[F \mid X_e = b])^2$$

By definition  $F = f(x)$

$$= \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a) \mu_e(b) \left( \mathbb{E}_{X \sim \mu^{e \leftarrow a}}[f(X)] - \mathbb{E}_{Y \sim \mu^{e \leftarrow b}}[f(Y)] \right)^2$$

Coupling :  $(x_0, x_\ell) \sim (\mu^{e \leftarrow a}, \mu^{e \leftarrow b})$

$$= \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a) \mu_e(b) \left( \mathbb{E}_{\gamma=(x_0, x_1, \dots, x_\ell) \sim \Gamma_e^{a \rightarrow b}}[f(x_0) - f(x_\ell)] \right)^2$$

Telescoping sum along the path

$$= \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a) \mu_e(b) \left( \mathbb{E}_{\gamma=(x_0, x_1, \dots, x_\ell) \sim \Gamma_e^{a \rightarrow b}} \left[ \sum_{1 \leq i \leq \ell} (f(x_i) - f(x_{i-1})) \right] \right)^2$$

## Proof outline: Local Poincaré inequality via transport flow

$$\text{Var}[\mathbb{E}[F \mid X_e]] = \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a) \mu_e(b) (\mathbb{E}[F \mid X_e = a] - \mathbb{E}[F \mid X_e = b])^2$$

By definition  $F = f(x)$

$$= \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a) \mu_e(b) \left( \mathbb{E}_{X \sim \mu^{e \leftarrow a}}[f(X)] - \mathbb{E}_{Y \sim \mu^{e \leftarrow b}}[f(Y)] \right)^2$$

Coupling :  $(x_0, x_\ell) \sim (\mu^{e \leftarrow a}, \mu^{e \leftarrow b})$

$$= \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a) \mu_e(b) \left( \mathbb{E}_{\gamma=(x_0, x_1, \dots, x_\ell) \sim \Gamma_e^{a \rightarrow b}}[f(x_0) - f(x_\ell)] \right)^2$$

Telescoping sum along the path

$$= \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a) \mu_e(b) \left( \mathbb{E}_{\gamma=(x_0, x_1, \dots, x_\ell) \sim \Gamma_e^{a \rightarrow b}} \left[ \sum_{1 \leq i \leq \ell} (f(x_i) - f(x_{i-1})) \right] \right)^2$$

Summing by  
enumerating transitions

$$= \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a) \mu_e(b) \left( \mathbb{E}_{\gamma \sim \Gamma_e^{a \rightarrow b}} \left[ \sum_{(x \rightarrow y) \in Q} (f(x) - f(y)) \mathbf{1}[(x \rightarrow y) \in \gamma] \right] \right)^2$$

## Proof outline: Local Poincaré inequality via transport flow

$$\text{Var}[\mathbb{E}[F \mid X_e]] = \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a)\mu_e(b) (\mathbb{E}[F \mid X_e = a] - \mathbb{E}[F \mid X_e = b])^2$$

By definition  $F = f(x)$

$$= \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a)\mu_e(b) \left( \mathbb{E}_{X \sim \mu^{e \leftarrow a}}[f(X)] - \mathbb{E}_{Y \sim \mu^{e \leftarrow b}}[f(Y)] \right)^2$$

Coupling :  $(x_0, x_\ell) \sim (\mu^{e \leftarrow a}, \mu^{e \leftarrow b})$

$$= \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a)\mu_e(b) \left( \mathbb{E}_{\gamma=(x_0, x_1, \dots, x_\ell) \sim \Gamma_e^{a \rightarrow b}} [f(x_0) - f(x_\ell)] \right)^2$$

Telescoping sum along the path

$$= \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a)\mu_e(b) \left( \mathbb{E}_{\gamma=(x_0, x_1, \dots, x_\ell) \sim \Gamma_e^{a \rightarrow b}} \left[ \sum_{1 \leq i \leq \ell} (f(x_i) - f(x_{i-1})) \right] \right)^2$$

Summing by  
enumerating transitions

$$= \frac{1}{2} \sum_{a,b \in [q]} \mu_e(a)\mu_e(b) \left( \mathbb{E}_{\gamma \sim \Gamma_e^{a \rightarrow b}} \left[ \sum_{(x \rightarrow y) \in Q} (f(x) - f(y)) \mathbf{1}[(x \rightarrow y) \in \gamma] \right] \right)^2$$

- using **Cauchy–Schwarz inequality** on the term
- the rest of the proof follows from the standard analysis in [\[Sinclair 92\]](#)



## Local log-Sobolev inequality via transport flow

If there exists a *family of transport flow*

$$\{\Gamma_e^{a \rightarrow b} \text{ from } \mu^{e \leftarrow a} \text{ to } \mu^{e \leftarrow b} \mid e \in E, a, b \in [q]\}$$

- ( $\kappa$ -**strong expected congestion**) For **any transition**  $(x \rightarrow y)$  in  $Q$ , and any  $a, b \in [q]$ ,

$$\sum_{e \in E} \mu_e(a) \mu_e(b) \cdot \mathbb{E}_{\gamma \sim \Gamma_e^{a \rightarrow b}} [\ell(\gamma) \cdot \mathbf{1}[(x \rightarrow y) \in \gamma]] \leq \kappa \cdot \mu(x) Q(x, y).$$



add the length of the path into the expectation

## Local log-Sobolev inequality via transport flow

If there exists a *family of transport flow*

$$\{\Gamma_e^{a \rightarrow b} \text{ from } \mu^{e \leftarrow a} \text{ to } \mu^{e \leftarrow b} \mid e \in E, a, b \in [q]\}$$

- ( $\kappa$ - **strong expected congestion**) For **any transition**  $(x \rightarrow y)$  in  $Q$ , and any  $a, b \in [q]$ ,

$$\sum_{e \in E} \mu_e(a) \mu_e(b) \cdot \mathbb{E}_{\gamma \sim \Gamma_e^{a \rightarrow b}} [\ell(\gamma) \cdot \mathbf{1}[(x \rightarrow y) \in \gamma]] \leq \kappa \cdot \mu(x) Q(x, y).$$




$\alpha$ -local log-Sobolev Inequality

$$\alpha \cdot \sum_{e \in E} \text{Ent} [\mathbb{E}[F^2 \mid X_e]] \leq \mathcal{E}_Q(f, f) \text{ with } \alpha = \Omega\left(\frac{1}{q^2 \kappa \log \frac{1}{\phi}}\right)$$

$\phi = \min\{\mu_e(c) \mid e \in E, c \in [q]\}$  is the **marginal lower bound**

## Proof outline: log-Sobolev inequality via transport flow




$$\text{Var}\left[\sqrt{\mathbb{E}[F^2 \mid X_e]}\right] \leq \sum_{a,b \in [q]} \mu_e(a)\mu_e(b) \cdot \mathbb{E}_{\gamma=(x_0, \dots, x_\ell) \sim \Gamma_e^{a \rightarrow b}} \left[ \left( \sum_{1 \leq i \leq \ell} f(x_i) - f(x_{i-1}) \right)^2 \right]$$

by convexity of  $h(x, y) = (\sqrt{x} - \sqrt{y})^2$

Using **Cauchy–Schwarz inequality** on the term



# Application to Jerrum-Sinclair chain



For  $\mu^\tau$  with pinnings  $\tau \in \{0,1\}^{E-\Lambda}$ , free variables in  $\Lambda$ , the Jerrum-Sinclair chain  $Q^\tau: X_t \rightarrow X_{t+1}$

- Pick an edge  $e \in \Lambda$  uniformly at random
- Construct a candidate matching  $M$  from  $X_t$
- Accept or reject  $M$  via Metropolis filter w.r.t.  $\mu^\tau$

The family of Jerrum-Sinclair chains  $\mathcal{Q} = \{ Q^\tau \mid \tau \}$  satisfies


- $(\alpha_1, \alpha_2, \dots, \alpha_{|E|})$ -**local Poincaré inequality** with  $\alpha_k = \Omega_\lambda \left( \frac{1}{k\Delta} \right)$
- $(\alpha_1, \alpha_2, \dots, \alpha_{|E|})$ -**log Sobolev inequality** with  $\alpha_k = \Omega_\lambda \left( \frac{1}{k\Delta^2 \log \Delta} \right)$

Proved by **transport flow**


Fix an edge  $e \in E$ , construct transport flow from  $\mu^{e \leftarrow \text{unmatched}}$  to  $\mu^{e \leftarrow \text{matched}}$

- Sample  $(X, Y)$  from the **local-flipping coupling** of  $\mu^{e \leftarrow \text{unmatched}}$  to  $\mu^{e \leftarrow \text{matched}}$
- Construct canonical path from  $X$  to  $Y$  using Jerrum and Sinclair's construction

**Local flipping  
coupling**





**Low discrepancy**  
between  $X$  and  $Y$



**Low congestion**  
and  
**low path length**



## Local Flipping Coupling

- Sample  $X \sim \mu^{e \leftarrow \text{unmatched}}$  and  $Z \sim \mu^{e \leftarrow \text{matched}}$  **independently**
- The difference between  $X$  and  $Y$  are paths and cycles, find the unique one  $B$  containing  $e$
- Let  $Y = Z_B \cup X_{E-B}$  (flipping  $B$  in  $X$  to obtain  $Y$ )





## Local Flipping Coupling

- Sample  $X \sim \mu^{e \leftarrow \text{unmatched}}$  and  $Z \sim \mu^{e \leftarrow \text{matched}}$  **independently**
- The **difference** between  $X$  and  $Z$  are paths and cycles, find the unique one  $B$  **containing  $e$**
- Let  $Y = Z_B \cup X_{E-B}$  (flipping  $B$  in  $X$  to obtain  $Y$ )



## Local Flipping Coupling


- Sample  $X \sim \mu^{e \leftarrow \text{unmatched}}$  and  $Z \sim \mu^{e \leftarrow \text{matched}}$  **independently**
- The **difference** between  $X$  and  $Z$  are paths and cycles, find the unique one  $B$  **containing  $e$**
- Let  $Y = Z_B \cup X_{E-B}$  (flipping  $B$  in  $X$  to obtain  $Y$ )






## Local Flipping Coupling

- Sample  $X \sim \mu^{e \leftarrow \text{unmatched}}$  and  $Z \sim \mu^{e \leftarrow \text{matched}}$  **independently**
- The **difference** between  $X$  and  $Z$  are paths and cycles, find the unique one  $B$  **containing  $e$**
- Let  $Y = Z_B \cup X_{E-B}$  (flipping  $B$  in  $X$  to obtain  $Y$ )




Copy




## Local Flipping Coupling

- Sample  $X \sim \mu^{e \leftarrow \text{unmatched}}$  and  $Z \sim \mu^{e \leftarrow \text{matched}}$  **independently**
- The **difference** between  $X$  and  $Z$  are paths and cycles, find the unique one  $B$  **containing  $e$**
- Let  $Y = Z_B \cup X_{E-B}$  (flipping  $B$  in  $X$  to obtain  $Y$ )




Flip




## Transport Flow

- Sample  $(X, Y)$  from the **local-flipping coupling** of  $\mu^{e \leftarrow \text{unmatched}}$  to  $\mu^{e \leftarrow \text{matched}}$
- Construct canonical path from  $X$  to  $Y$  using Jerrum and Sinclair's construction



# Proof overview of expected length and congestion

- Analyze coupling via **local reviewing process**
- Disagreement percolation



For  $(X, Y) \sim \mathcal{C}_e$  from local flipping coupling

- Length bound  $\mathbb{E}[|X \oplus Y|] \leq O_\lambda(\sqrt{\Delta})$
- One sided bound  $\mathbb{E}[|X \oplus Y| \mid X = x] \leq O_\lambda(\Delta)$

## Expected congestion and strong congestion analysis

$$\sum_{e \in E} \mu_e(a) \mu_e(b) \cdot \mathbb{E}_{\gamma \sim \Gamma_e} [\mathbf{1}[(x \rightarrow y) \in \gamma]] \leq \kappa \cdot \mu(x) Q(x, y).$$
$$\sum_{e \in E} \mu_e(a) \mu_e(b) \cdot \mathbb{E}_{\gamma \sim \Gamma_e} [\ell(\gamma) \cdot \mathbf{1}[(x \rightarrow y) \in \gamma]] \leq \kappa \cdot \mu(x) Q(x, y).$$

sum of  $|E|$  different couplings

$$\gamma \sim \Gamma_e$$




- Sample starting and ending points from coupling
- Construct the path deterministically

**The randomness is only from locally flipping coupling**

# Proof overview of expected length and congestion

- Analyze coupling via **local reviewing process**
- Disagreement percolation



For  $(X, Y) \sim \mathcal{C}_e$  from local flipping coupling

- Length bound  $\mathbb{E}[|X \oplus Y|] \leq O_\lambda(\sqrt{\Delta})$
- One sided bound  $\mathbb{E}[|X \oplus Y| \mid X = x] \leq O_\lambda(\Delta)$

## Expected congestion and strong congestion analysis

$$\sum_{e \in E} \mu_e(a) \mu_e(b) \cdot \mathbb{E}_{\gamma \sim \Gamma_e} [\mathbf{1}[(x \rightarrow y) \in \gamma]] \leq \kappa \cdot \mu(x) Q(x, y).$$
$$\sum_{e \in E} \mu_e(a) \mu_e(b) \cdot \mathbb{E}_{\gamma \sim \Gamma_e} [\ell(\gamma) \cdot \mathbf{1}[(x \rightarrow y) \in \gamma]] \leq \kappa \cdot \mu(x) Q(x, y).$$

sum of  $|E|$  different couplings



**decoupling lemma**



bound congestion by constructing **injection**  
(guided by Jerrum-Sinclair's analysis)

analyze **one coupling** with a different  
function inside the expectation

# Open problems

- Lower discrepancy coupling of  $\mu^{e \leftarrow a}$  and  $\mu^{e \leftarrow b}$
- Construction of canonical paths



- Poincaré inequality
- log-Sobolev inequality

- Sharp bound for Jerrum-Sinclair chain:  $\tilde{O}(m\sqrt{\Delta})$  mixing?
- More applications?

Improving the mixing bound for e.g. the permanent, the Ising model,  
the switch/flip chain for sampling regular graphs...