

Modern Difference-in-Difference Designs

Ryan T. Moore

American University

The Lab @ DC

2024-07-17

Table of contents I

Canonical Diff-in-Diff Designs, Calloway-Sant'Anna

Canonical Diff-in-Diff Designs,
Calloway-Sant'Anna

Canonical Difference-in-Difference Designs

- ▶ Goal: observational causal statements by adjusting for confounding

Canonical Difference-in-Difference Designs

- ▶ Goal: observational causal statements by adjusting for confounding
- ▶ (Similar to matching, regression, etc.)

Canonical Difference-in-Difference Designs

- ▶ Goal: observational causal statements by adjusting for confounding
- ▶ (Similar to matching, regression, etc.)
- ▶ Confounding must be “equi-additive”

Canonical Difference-in-Difference Designs

- ▶ Goal: observational causal statements by adjusting for confounding
- ▶ (Similar to matching, regression, etc.)
- ▶ Confounding must be “equi-additive”
 - ▶ same amt added to outcome over time in both Tr/Co

Canonical Difference-in-Difference Designs

- ▶ Goal: observational causal statements by adjusting for confounding
- ▶ (Similar to matching, regression, etc.)
- ▶ Confounding must be “equi-additive”
 - ▶ same amt added to outcome over time in both Tr/Co
- ▶ Adjustment using a “negative outcome control”

Canonical Difference-in-Difference Designs

- ▶ Goal: observational causal statements by adjusting for confounding
- ▶ (Similar to matching, regression, etc.)
- ▶ Confounding must be “equi-additive”
 - ▶ same amt added to outcome over time in both Tr/Co
- ▶ Adjustment using a “negative outcome control”
 - ▶ pre-tr outcome cannot be affected by treatment

Canonical Difference-in-Difference Designs

- ▶ Goal: observational causal statements by adjusting for confounding
- ▶ (Similar to matching, regression, etc.)
- ▶ Confounding must be “equi-additive”
 - ▶ same amt added to outcome over time in both Tr/Co
- ▶ Adjustment using a “negative outcome control”
 - ▶ pre-tr outcome cannot be affected by treatment
 - ▶ pre-tr outcome affected by (obs/unobs) predictors of post-tr outcome

Canonical Difference-in-Difference Designs

Before-After Designs:

- ▶ Before-After designs ask “Did the outcome go up or down?”

Canonical Difference-in-Difference Designs

Before-After Designs:

- ▶ Before-After designs ask “Did the outcome go up or down?”
- ▶ In NJ minwage study (Card and Krueger (1994)), Before-After: “After the new minimum wage, did employment rise or fall in NJ?”

Canonical Difference-in-Difference Designs

Before-After Designs:

- ▶ Before-After designs ask “Did the outcome go up or down?”
- ▶ In NJ minwage study (Card and Krueger (1994)), Before-After: “After the new minimum wage, did employment rise or fall in NJ?”
- ▶ The Before-After estimate for NJ:

Canonical Difference-in-Difference Designs

Before-After Designs:

- ▶ Before-After designs ask “Did the outcome go up or down?”
- ▶ In NJ minwage study (Card and Krueger (1994)), Before-After: “After the new minimum wage, did employment rise or fall in NJ?”
- ▶ The Before-After estimate for NJ:

Canonical Difference-in-Difference Designs

Before-After Designs:

- ▶ Before-After designs ask “Did the outcome go up or down?”
- ▶ In NJ minwage study (Card and Krueger (1994)), Before-After: “After the new minimum wage, did employment rise or fall in NJ?”
- ▶ The Before-After estimate for NJ:

$$\text{Before-After estimate} = \overline{\text{Employment}}_T^{\text{After}} - \overline{\text{Employment}}_T^{\text{Before}}$$

Canonical Difference-in-Difference Designs

Before-After Designs:

- ▶ Before-After designs ask “Did the outcome go up or down?”
- ▶ In NJ minwage study (Card and Krueger (1994)), Before-After: “After the new minimum wage, did employment rise or fall in NJ?”
- ▶ The Before-After estimate for NJ:

$$\text{Before-After estimate} = \overline{\text{Employment}}_T^{\text{After}} - \overline{\text{Employment}}_T^{\text{Before}}$$

What's weak about this question? What's being ignored?

Canonical Difference-in-Difference Designs

Before-After Designs:

- ▶ Before-After designs ask “Did the outcome go up or down?”
- ▶ In NJ minwage study (Card and Krueger (1994)), Before-After: “After the new minimum wage, did employment rise or fall in NJ?”
- ▶ The Before-After estimate for NJ:

$$\text{Before-After estimate} = \overline{\text{Employment}_T^{\text{After}}} - \overline{\text{Employment}_T^{\text{Before}}}$$

What's weak about this question? What's being ignored?

This is **not**

$$ATT = \overline{[Y(1)|T = 1]} - \overline{[Y(0)|T = 1]}$$

Why not?

Canonical Difference-in-Difference

- ▶ DiD estimate: difference between two Before-After estimates

Canonical Difference-in-Difference

- ▶ DiD estimate: difference between two Before-After estimates
- ▶ DiD: Before-After estimate for NJ, subtract off change that occurred in control state, PA.

Canonical Difference-in-Difference

- ▶ DiD estimate: difference between two Before-After estimates
- ▶ DiD: Before-After estimate for NJ, subtract off change that occurred in control state, PA.

Canonical Difference-in-Difference

- ▶ DiD estimate: difference between two Before-After estimates
- ▶ DiD: Before-After estimate for NJ, subtract off change that occurred in control state, PA.

$$\text{DiD estimate} = (\bar{Y}_T^{\text{After}} - \bar{Y}_T^{\text{Before}}) - (\bar{Y}_C^{\text{After}} - \bar{Y}_C^{\text{Before}})$$

Canonical Difference-in-Difference

- ▶ DiD estimate: difference between two Before-After estimates
- ▶ DiD: Before-After estimate for NJ, subtract off change that occurred in control state, PA.

$$\begin{aligned}\text{DiD estimate} &= (\bar{Y}_T^{\text{After}} - \bar{Y}_T^{\text{Before}}) - (\bar{Y}_C^{\text{After}} - \bar{Y}_C^{\text{Before}}) \\ &= \underbrace{(\bar{Y}_{NJ}^{\text{After}} - \bar{Y}_{NJ}^{\text{Before}})}_{\text{Change in Treated}} - \underbrace{(\bar{Y}_{PA}^{\text{After}} - \bar{Y}_{PA}^{\text{Before}})}_{\text{Change in Control}}\end{aligned}$$

Canonical Difference-in-Difference

- ▶ DiD estimate: difference between two Before-After estimates
- ▶ DiD: Before-After estimate for NJ, subtract off change that occurred in control state, PA.

$$\begin{aligned}\text{DiD estimate} &= (\bar{Y}_T^{\text{After}} - \bar{Y}_T^{\text{Before}}) - (\bar{Y}_C^{\text{After}} - \bar{Y}_C^{\text{Before}}) \\ &= \underbrace{(\bar{Y}_{NJ}^{\text{After}} - \bar{Y}_{NJ}^{\text{Before}})}_{\text{Change in Treated}} - \underbrace{(\bar{Y}_{PA}^{\text{After}} - \bar{Y}_{PA}^{\text{Before}})}_{\text{Change in Control}}\end{aligned}$$

- ▶ Change in Control: removes part of change in emp that would have occurred anyway, in absence of law change.

Canonical Difference-in-Difference

- ▶ DiD estimate: difference between two Before-After estimates
- ▶ DiD: Before-After estimate for NJ, subtract off change that occurred in control state, PA.

$$\begin{aligned}\text{DiD estimate} &= (\bar{Y}_T^{\text{After}} - \bar{Y}_T^{\text{Before}}) - (\bar{Y}_C^{\text{After}} - \bar{Y}_C^{\text{Before}}) \\ &= \underbrace{(\bar{Y}_{NJ}^{\text{After}} - \bar{Y}_{NJ}^{\text{Before}})}_{\text{Change in Treated}} - \underbrace{(\bar{Y}_{PA}^{\text{After}} - \bar{Y}_{PA}^{\text{Before}})}_{\text{Change in Control}}\end{aligned}$$

- ▶ Change in Control: removes part of change in emp that would have occurred anyway, in absence of law change.
- ▶ (Economy growing, or fast food restaurants hit by health scares and lay off workers, ...)

Canonical Difference-in-Difference

- ▶ Before-After uses 2 pieces of info; DiD design uses 4

Canonical Difference-in-Difference

- ▶ Before-After uses 2 pieces of info; DiD design uses 4
- ▶ Change in Control is our estimate of what *would have happened* in treated unit (NJ), in absence of treatment

Canonical Difference-in-Difference


- ▶ Before-After uses 2 pieces of info; DiD design uses 4
- ▶ Change in Control is our estimate of what *would have happened* in treated unit (NJ), in absence of treatment
- ▶ Key assumption: “parallel trends” – change in PA is what change in NJ *would have been* without new minimum wage.

Canonical Difference-in-Difference

- ▶ Before-After uses 2 pieces of info; DiD design uses 4
- ▶ Change in Control is our estimate of what *would have happened* in treated unit (NJ), in absence of treatment
- ▶ Key assumption: “parallel trends” – change in PA is what change in NJ *would have been* without new minimum wage.


Canonical Difference-in-Difference

- ▶ Before-After uses 2 pieces of info; DiD design uses 4
- ▶ Change in Control is our estimate of what *would have happened* in treated unit (NJ), in absence of treatment
- ▶ Key assumption: “parallel trends” – change in PA is what change in NJ *would have been* without new minimum wage.




Self-Reported Health in *Seguro Popular* Evaluation


King, et al. (2007)



Simplified Example



Simplified Example




The Difference-in-Differences Estimator

Units	Outcome Before	Outcome After	Difference
Treated	\bar{Y}^{T_B}	\bar{Y}^{T_A}	$\bar{Y}^{T_A} - \bar{Y}^{T_B}$
Control	\bar{Y}^{C_B}	\bar{Y}^{C_A}	$\bar{Y}^{C_A} - \bar{Y}^{C_B}$
	$\bar{Y}^{T_B} - \bar{Y}^{C_B}$	$\bar{Y}^{T_A} - \bar{Y}^{C_A}$	$(\bar{Y}^{T_A} - \bar{Y}^{T_B}) - (\bar{Y}^{C_A} - \bar{Y}^{C_B})$


Example: Effect of Minimum Wage on Employment

The Baseline



Example: Effect of Minimum Wage on Employment

The Policy Change: Significant Shock



Example: Effect of Minimum Wage on Employment

Variable	Difference, NJ – PA		
	PA (i)	NJ (ii)	NJ – PA (iii)
1. FTE employment before, all available observations	23.33 (1.35)	20.44 (0.51)	-2.89 (1.44)
2. FTE employment after, all available observations	21.17 (0.94)	21.03 (0.52)	-0.14 (1.07)
3. Change in mean FTE employment	-2.16 (1.25)	0.59 (0.54)	2.76 (1.36)

Extension 1: Regression Difference-in-Differences

Let

- ▶ $T_i \in \{0, 1\}$ be treatment status (“ever-treated”)
- ▶ $A_i \in \{0, 1\}$ be “after” period status

Extension 1: Regression Difference-in-Differences

Let

- ▶ $T_i \in \{0, 1\}$ be treatment status (“ever-treated”)
- ▶ $A_i \in \{0, 1\}$ be “after” period status

Model

$$y_i = \beta_0 + \beta_1 T_i + \beta_2 A_i + \beta_3 T_i A_i + \epsilon_i$$

Extension 1: Regression Difference-in-Differences

Let

- ▶ $T_i \in \{0, 1\}$ be treatment status (“ever-treated”)
- ▶ $A_i \in \{0, 1\}$ be “after” period status

Model

$$y_i = \beta_0 + \beta_1 T_i + \beta_2 A_i + \beta_3 T_i A_i + \epsilon_i$$

Then

$$\begin{aligned}\bar{Y}^{T_A} &= \beta_0 + \beta_1 + \beta_2 + \beta_3 \\ \bar{Y}^{T_B} &= \beta_0 + \beta_1 \\ \bar{Y}^{C_A} &= \beta_0 + \beta_2 \\ \bar{Y}^{C_B} &= \beta_0\end{aligned}$$

Extension 1: Regression Difference-in-Differences

Let

- ▶ $T_i \in \{0, 1\}$ be treatment status (“ever-treated”)
- ▶ $A_i \in \{0, 1\}$ be “after” period status

Model

$$y_i = \beta_0 + \beta_1 T_i + \beta_2 A_i + \beta_3 T_i A_i + \epsilon_i$$

Then

$$\begin{aligned}\bar{Y}^{T_A} &= \beta_0 + \beta_1 + \beta_2 + \beta_3 \\ \bar{Y}^{T_B} &= \beta_0 + \beta_1 \\ \bar{Y}^{C_A} &= \beta_0 + \beta_2 \\ \bar{Y}^{C_B} &= \beta_0\end{aligned}$$

And

$$\begin{aligned}(\bar{Y}^{T_A} - \bar{Y}^{T_B}) - (\bar{Y}^{C_A} - \bar{Y}^{C_B}) &= ([\beta_0 + \beta_1 + \beta_2 + \beta_3] - [\beta_0 + \beta_1]) \\ &\quad - ([\beta_0 + \beta_2] - \beta_0) \\ &= (\beta_2 + \beta_3) - (\beta_2) \\ &= \beta_3\end{aligned}$$

Extension 1: Regression Difference-in-Differences

Note: $T_i \in \{0, 1\}$ needs to be “ever-treated”
(not “currently treated”)

Simulation:

- ▶ 40 units (20 Tr, 20 Co)
- ▶ 2 time periods (2018, 2022)
- ▶ Baseline outcome difference $\text{Tr} - \text{Co} = 1$
- ▶ True TE = 2
- ▶ (TE occurs in 2022, only for Tr)

Extension 1: Regression Difference-in-Differences




Figure 2: Simulated Data DiD

Extension 1: Regression Difference-in-Differences

Two possible regressions:

$$Y = \beta_0 + \beta_1(2022) + \beta_2(\text{TrEver}) + \beta_3(2022 \times \text{TrEver})$$

Extension 1: Regression Difference-in-Differences

Two possible regressions:

$$Y = \beta_0 + \beta_1(2022) + \beta_2(\text{TrEver}) + \beta_3(2022 \times \text{TrEver})$$

versus

$$Y = \beta_0 + \beta_1(2022) + \beta_2(\text{TrNow}) + \beta_3(2022 \times \text{TrNow})$$

Extension 1: Regression Difference-in-Differences

Table 1: Ever-treated vs. Currently-treated DiD Regressions

	Outcome	
	y	
	(1)	(2)
2022	0.500 (0.313)	0.150 (0.278)
Tr (Ever)	0.700** (0.313)	
2022 x Tr (Ever)	2.000*** (0.442)	
Tr (Now)		2.700*** (0.321)
2022 x Tr (Now)		
(Intercept)	0.462** (0.221)	0.812*** (0.160)
Observations	80	80
R ²	0.623	0.598
Adjusted R ²	0.608	0.588
Residual Std. Error	0.989 (df = 76)	1.015 (df = 77)
F Statistic	41.827*** (df = 3; 76)	57.261*** (df = 2; 77)

Note:

*p<0.1; **p<0.05; ***p<0.01

Extension 1: Regression Difference-in-Differences

$$Y = \underbrace{0.46}_{\text{Co Mean 2018}} + \underbrace{0.5}_{\text{Time Trend}} (2022) + \underbrace{0.7}_{\text{Baseline Diff}} (\text{TrEver}) + \underbrace{2}_{\widehat{ATT}} (2022 \times \text{TrEver})$$

Extension 1: Regression Difference-in-Differences

$$Y = \underbrace{0.46}_{\text{Co Mean 2018}} + \underbrace{0.5}_{\text{Time Trend}} (2022) + \underbrace{0.7}_{\text{Baseline Diff}} (\text{TrEver}) + \underbrace{2}_{\widehat{ATT}} (2022 \times \text{TrEver})$$

versus

$$Y = \underbrace{0.81}_{\text{Part of Co Mean 2018 + Time Trend??}} + \underbrace{0.15}_{\text{Rest of Time Trend??}} (2022) + \underbrace{2.7}_{\text{Baseline Diff} + \widehat{ATT}??} (\text{TrNow}) + \underbrace{NA}_{\text{Unest Interaction for } \widehat{ATT}??} (2022 \times \text{TrNow})$$

Extension 2: Regression Difference-in-Differences

Related: Use change scores or regressor variable method?

(Allison 90)

Extension 2: Regression Difference-in-Differences

Related: Use change scores or regressor variable method?

(Allison 90)

$$y_{i2} = \alpha + \beta_1 y_{i1} + \beta_2 T_i + \epsilon_i \quad (1)$$

Extension 2: Regression Difference-in-Differences

Related: Use change scores or regressor variable method?

(Allison 90)

$$y_{i2} = \alpha + \beta_1 y_{i1} + \beta_2 T_i + \epsilon_i \quad (1)$$

$$y_{i1} = \alpha + \gamma G_i + \epsilon_{i1} \quad (2)$$

$$y_{i2} = \alpha + \underbrace{\tau}_{\text{secular trend}} + \gamma G_i + \delta T_i + \epsilon_{i2} \quad (3)$$

$$y_{i2} - y_{i1} = \tau + \delta T_i + (\epsilon_{i2} - \epsilon_{i1}) \quad (4)$$

Extension 2: Regression Difference-in-Differences

Related: Use change scores or regressor variable method?
(Allison 90)

$$y_{i2} = \alpha + \beta_1 y_{i1} + \beta_2 T_i + \epsilon_i \quad (1)$$

$$y_{i1} = \alpha + \gamma G_i + \epsilon_{i1} \quad (2)$$

$$y_{i2} = \alpha + \underbrace{\tau}_{\text{secular trend}} + \gamma G_i + \delta T_i + \epsilon_{i2} \quad (3)$$

$$y_{i2} - y_{i1} = \tau + \delta T_i + (\epsilon_{i2} - \epsilon_{i1}) \quad (4)$$

- ▶ Change score is a DiD estimate
- ▶ Worse if pretest not very strong, and restriction of $\beta_1 = 1$ creates bias.
- ▶ Change score is more restrictive on parameter, but makes different assumption about $Cor(y_1, \epsilon)$.

Extension 3: Regression Difference-in-Differences

Add other covariates:

$$y_i = \beta_0 + \beta_1 T_i + \beta_2 A_i + \beta_3 T_i A_i + \gamma' \mathbf{x}_i + \epsilon_i$$

Extension 3: Regression Difference-in-Differences

Add other covariates:

$$y_i = \beta_0 + \beta_1 T_i + \beta_2 A_i + \beta_3 T_i A_i + \gamma' \mathbf{x}_i + \varepsilon_i$$

But, careful about inference from SE's with many time periods (Bertrand, Duflo, and Mullainathan (2004)).

Extension 3: Regression Difference-in-Differences

Add other covariates:

$$y_i = \beta_0 + \beta_1 T_i + \beta_2 A_i + \beta_3 T_i A_i + \gamma' \mathbf{x}_i + \varepsilon_i$$

But, careful about inference from SE's with many time periods (Bertrand, Duflo, and Mullainathan (2004)).

(At $\alpha = 0.05$, false findings actually 0.45!)

Extension 3: Regression Difference-in-Differences

Add other covariates:

$$y_i = \beta_0 + \beta_1 T_i + \beta_2 A_i + \beta_3 T_i A_i + \gamma' \mathbf{x}_i + \varepsilon_i$$

But, careful about inference from SE's with many time periods (Bertrand, Duflo, and Mullainathan (2004)).

(At $\alpha = 0.05$, false findings actually 0.45!)

Actually, be careful about bias with many time periods, too!

(See here.)

Extension 4: Difference in (Differences-in-Differences)

- ▶ This is crazy! (?)

Extension 4: Difference in (Differences-in-Differences)

- ▶ This is crazy! (?)

Extension 4: Difference in (Differences-in-Differences)

- ▶ This is crazy! (?)
- ▶ Useful for estimating effect across two subgroups

Extension 4: Difference in (Differences-in-Differences)

- ▶ This is crazy! (?)
- ▶ Useful for estimating effect across two subgroups
- ▶ (E.g., Empl vs. unempl'd; or placebo robustness check if one subgroup should *not* be affected by treatment)

Extension 4: Difference in (Differences-in-Differences)

Group 1:

Units	Outcome Before	Outcome After	Difference
Treated	T_{1B}	T_{1A}	$T_{1A} - T_{1B}$
Control	C_{1B}	C_{1A}	$C_{1A} - C_{1B}$
	$T_{1B} - C_{1B}$	$T_{1A} - C_{1A}$	$(T_{1A} - T_{1B}) - (C_{1A} - C_{1B})$

Group 2:

Units	Outcome Before	Outcome After	Difference
Treated	T_{2B}	T_{2A}	$T_{2A} - T_{2B}$
Control	C_{2B}	C_{2A}	$C_{2A} - C_{2B}$
	$T_{2B} - C_{2B}$	$T_{2A} - C_{2A}$	$(T_{2A} - T_{2B}) - (C_{2A} - C_{2B})$

DiDiD Estimate:

$$[(T_{2A} - T_{2B}) - (C_{2A} - C_{2B})] - [(T_{1A} - T_{1B}) - (C_{1A} - C_{1B})]$$

Extension 5: Multiple Time Periods, Staggered Designs

- ▶ Callaway and Sant'Anna (2021b) on multiple time periods

Extension 5: Multiple Time Periods, Staggered Designs

- ▶ Callaway and Sant'Anna (2021b) on multiple time periods
- ▶ Goodman-Bacon (2021) on staggered treatment assignment

Extension 5: Multiple Time Periods, Staggered Designs

- ▶ Callaway and Sant'Anna (2021b) on multiple time periods
- ▶ Goodman-Bacon (2021) on staggered treatment assignment
- ▶ Package `did`

The Stepped Wedge Design

- ▶ NMAAHC experiment

The Stepped Wedge Design

- ▶ NMAAHC experiment
- ▶ All 2000 members of DC MPD sworn force

The Stepped Wedge Design

- ▶ NMAAHC experiment
- ▶ All 2000 members of DC MPD sworn force
- ▶ Gradually, month by month, discuss history of race and policing in DC and US, guided tour of exhibits, lunch, free time in museum

The Stepped Wedge Design

- ▶ NMAAHC experiment
- ▶ All 2000 members of DC MPD sworn force
- ▶ Gradually, month by month, discuss history of race and policing in DC and US, guided tour of exhibits, lunch, free time in museum
- ▶ Start with all untreated

The Stepped Wedge Design

- ▶ NMAAHC experiment
- ▶ All 2000 members of DC MPD sworn force
- ▶ Gradually, month by month, discuss history of race and policing in DC and US, guided tour of exhibits, lunch, free time in museum
- ▶ Start with all untreated
- ▶ Over ≈ 1.5 -2 years, treat all officers

The Stepped Wedge Design

- ▶ NMAAHC experiment
- ▶ All 2000 members of DC MPD sworn force
- ▶ Gradually, month by month, discuss history of race and policing in DC and US, guided tour of exhibits, lunch, free time in museum
- ▶ Start with all untreated
- ▶ Over ≈ 1.5 -2 years, treat all officers
- ▶ Measure admin data (use of force, etc.) and three waves of survey

Calloway-Sant'anna Approach: Multiple Time Periods

With *many* time periods, challenging to defining **the** estimand (like ATT).

Callaway-Sant'anna Approach: Multiple Time Periods

With *many* time periods, challenging to defining **the** estimand (like ATT).

The Callaway and Sant'Anna (2021b) approach:

- ▶ ID disaggregated parameters of interest

Callaway-Sant'anna Approach: Multiple Time Periods

With *many* time periods, challenging to defining **the** estimand (like ATT).

The Callaway and Sant'Anna (2021b) approach:

- ▶ ID disaggregated parameters of interest
- ▶ Decide how to aggregate them

Callaway-Sant'anna Approach: Multiple Time Periods

With *many* time periods, challenging to defining **the** estimand (like ATT).

The Callaway and Sant'Anna (2021b) approach:

- ▶ ID disaggregated parameters of interest
- ▶ Decide how to aggregate them
- ▶ Estimate them

Calloway-Sant'anna Approach: Multiple Time Periods

Central disaggregated component:

the *group-time average treatment effect*.

Calloway-Sant'anna Approach: Multiple Time Periods

Central disaggregated component:
the *group-time average treatment effect*.

► “Group”: treatment cohort.

Calloway-Sant'anna Approach: Multiple Time Periods

Central disaggregated component:
the *group-time average treatment effect*.

► “Group”: treatment cohort.

Calloway-Sant'anna Approach: Multiple Time Periods

Central disaggregated component:

the *group-time average treatment effect*.

- ▶ “Group”: treatment cohort. Set of units first-treated at the same time moment

Calloway-Sant'anna Approach: Multiple Time Periods

Central disaggregated component:

the *group-time average treatment effect*.

- ▶ “Group”: treatment cohort. Set of units first-treated at the same time moment
- ▶ $ATT(g, t)$

Calloway-Sant'anna Approach: Multiple Time Periods

Then, how to aggregate?

Calloway-Sant'anna Approach: Multiple Time Periods

Then, how to aggregate? Many ways!

Calloway-Sant'anna Approach: Multiple Time Periods

Then, how to aggregate? Many ways!

- ▶ Exposure/Dosage (“effect for those w/ 2 years treatment”)

Calloway-Sant'anna Approach: Multiple Time Periods

Then, how to aggregate? Many ways!

- ▶ Exposure/Dosage (“effect for those w/ 2 years treatment”)
- ▶ Group (“effect for those first treated in 2020”)

Calloway-Sant'anna Approach: Multiple Time Periods

Then, how to aggregate? Many ways!

- ▶ Exposure/Dosage (“effect for those w/ 2 years treatment”)
- ▶ Group (“effect for those first treated in 2020”)
- ▶ Cumulative (“total effect up to 2020”)

Calloway-Sant'anna Approach: Multiple Time Periods

Then, how to aggregate? Many ways!

- ▶ Exposure/Dosage (“effect for those w/ 2 years treatment”)
- ▶ Group (“effect for those first treated in 2020”)
- ▶ Cumulative (“total effect up to 2020”)
- ▶ Overall (“total effect up to 2024”)

Calloway-Sant'anna Approach: Multiple Time Periods

Then, how to aggregate? Many ways!

- ▶ Exposure/Dosage (“effect for those w/ 2 years treatment”)
- ▶ Group (“effect for those first treated in 2020”)
- ▶ Cumulative (“total effect up to 2020”)
- ▶ Overall (“total effect up to 2024”)
- ▶ Overall, group as unit (“total effect up to 2024: get state avgs, avg those”)

Calloway-Sant'Anna Approach: Multiple Time Periods

Key assumptions:

1. No treated at $t = 1$. If treated at t^* , then treated at $t^* + 1$. Once treated, stay “treated”.

Calloway-Sant'Anna Approach: Multiple Time Periods

Key assumptions:

1. No treated at $t = 1$. If treated at t^* , then treated at $t^* + 1$. Once treated, stay “treated”.
2. Random selection of units from big population

Calloway-Sant'Anna Approach: Multiple Time Periods

Key assumptions:

1. No treated at $t = 1$. If treated at t^* , then treated at $t^* + 1$. Once treated, stay “treated”.
2. Random selection of units from big population
3. Limited, known amt of anticipation

Calloway-Sant'Anna Approach: Multiple Time Periods

Key assumptions:

1. No treated at $t = 1$. If treated at t^* , then treated at $t^* + 1$. Once treated, stay “treated”.
2. Random selection of units from big population
3. Limited, known amt of anticipation
4. Parallel trends with a never-treated group (conditional on covariates)

Calloway-Sant'Anna Approach: Multiple Time Periods

Key assumptions:

1. No treated at $t = 1$. If treated at t^* , then treated at $t^* + 1$. Once treated, stay “treated”.
2. Random selection of units from big population
3. Limited, known amt of anticipation
4. Parallel trends with a never-treated group (conditional on covariates)
5. Parallel trends with not-yet-treated (conditional on covariates)

Calloway-Sant'Anna Approach: Multiple Time Periods

Key assumptions:

1. No treated at $t = 1$. If treated at t^* , then treated at $t^* + 1$. Once treated, stay “treated”.
2. Random selection of units from big population
3. Limited, known amt of anticipation
4. Parallel trends with a never-treated group (conditional on covariates)
5. Parallel trends with not-yet-treated (conditional on covariates)
6. Overlap: $\epsilon < \Pr(\text{Tr}) < 1 - \epsilon$ for each $t > 2, g$

Calloway-Sant'Anna Approach: Multiple Time Periods

Key assumptions:

1. No treated at $t = 1$. If treated at t^* , then treated at $t^* + 1$. Once treated, stay “treated”.
2. Random selection of units from big population
3. Limited, known amt of anticipation
4. Parallel trends with a never-treated group (conditional on covariates)
5. Parallel trends with not-yet-treated (conditional on covariates)
6. Overlap: $\epsilon < \Pr(\text{Tr}) < 1 - \epsilon$ for each $t > 2, g$

3/4/5 define valid comparison group, thus analysis sample.

C-S Example

Effect of US state min wages on *teen* employment.

C-S Example

Effect of US state min wages on *teen* employment.

- ▶ Annual data, 2003-2007

C-S Example

Effect of US state min wages on *teen* employment.

- ▶ Annual data, 2003-2007
- ▶ 500 counties

C-S Example

Effect of US state min wages on *teen* employment.

- ▶ Annual data, 2003-2007
- ▶ 500 counties
 - ▶ 309 never-treated

C-S Example

Effect of US state min wages on *teen* employment.

- ▶ Annual data, 2003-2007
- ▶ 500 counties
 - ▶ 309 never-treated
 - ▶ 20 first treated in 2004

C-S Example

Effect of US state min wages on *teen* employment.

- ▶ Annual data, 2003-2007
- ▶ 500 counties
 - ▶ 309 never-treated
 - ▶ 20 first treated in 2004
 - ▶ 40 first treated in 2006

C-S Example

Effect of US state min wages on *teen* employment.

- ▶ Annual data, 2003-2007
- ▶ 500 counties
 - ▶ 309 never-treated
 - ▶ 20 first treated in 2004
 - ▶ 40 first treated in 2006
 - ▶ 131 first treated in 2007

C-S Example

Effect of US state min wages on *teen* employment.

- ▶ Annual data, 2003-2007
- ▶ 500 counties
 - ▶ 309 never-treated
 - ▶ 20 first treated in 2004
 - ▶ 40 first treated in 2006
 - ▶ 131 first treated in 2007
- ▶ Panel data structure: $(5 \text{ yrs}) \times (500 \text{ cntys}) = (2500 \text{ cnty-yrs})$

C-S Example

Effect of US state min wages on *teen* employment.

- ▶ Annual data, 2003-2007
- ▶ 500 counties
 - ▶ 309 never-treated
 - ▶ 20 first treated in 2004
 - ▶ 40 first treated in 2006
 - ▶ 131 first treated in 2007
- ▶ Panel data structure: $(5 \text{ yrs}) \times (500 \text{ cntys}) = (2500 \text{ cnty-yrs})$

C-S Example

Effect of US state min wages on *teen* employment.

- ▶ Annual data, 2003-2007
- ▶ 500 counties
 - ▶ 309 never-treated
 - ▶ 20 first treated in 2004
 - ▶ 40 first treated in 2006
 - ▶ 131 first treated in 2007
- ▶ Panel data structure: $(5 \text{ yrs}) \times (500 \text{ cntys}) = (2500 \text{ cnty-yrs})$

Callaway and Sant'Anna (2021a): `did`

C-S Example

Calculate all the $ATT(g, t)$ effects, for each (state \times year)

C-S Example

Calculate all the $ATT(g, t)$ effects, for each (state \times year)

```
cs_out <- att_gt(ymame = "lemp",  
                 gname = "first.treat",  
                 idname = "countyreal",  
                 tname = "year",  
                 xformula = ~ 1,  
                 data = mpdta,  
                 est_method = "reg"  
)
```


C-S Example

Calculate all the $ATT(g, t)$ effects, for each (state \times year)

```
cs_out <- att_gt(yname = "lemp",  
                 gname = "first.treat",  
                 idname = "countyreal",  
                 tname = "year",  
                 xformula = ~ 1,  
                 data = mpdta,  
                 est_method = "reg"  
)
```

► `est_method`: regression, IPW, doubly-robust, etc.

C-S Example

Calculate all the $ATT(g, t)$ effects, for each (state \times year)

```
cs_out <- att_gt(yname = "lemp",  
                 gname = "first.treat",  
                 idname = "countyreal",  
                 tname = "year",  
                 xformla = ~ 1,  
                 data = mpdta,  
                 est_method = "reg"  
)
```

- ▶ `est_method`: regression, IPW, doubly-robust, etc.
- ▶ `xformla`: formula for covariate adjustment

C-S Example


Calculate all the $ATT(g, t)$ effects, for each (state \times year)

```
cs_out <- att_gt(ymame = "lemp",  
                 gname = "first.treat",  
                 idname = "countyreal",  
                 tname = "year",  
                 xformula = ~ 1,  
                 data = mpdta,  
                 est_method = "reg"  
)
```

- ▶ `est_method`: regression, IPW, doubly-robust, etc.
- ▶ `xformula`: formula for covariate adjustment
- ▶ `control_group`: "nevertreated", "notyettreated"

C-S Example

```
ggdid(cs_out)
```



C-S Example

- ▶ “Pre” C-S $ATT(g, t)$ values: *pre*-test of parallel trends assumption

C-S Example

- ▶ “Pre” C-S $ATT(g, t)$ values: *pre*-test of parallel trends assumption
- ▶ “Post” C-S $ATT(g, t)$ values: effects of min wage policy

C-S Example: Aggregate by g, t

Instead of fully-disaggregated $ATT(g, t)$'s, summarise into various effects.

C-S Example: Aggregate by g, t

Instead of fully-disaggregated $ATT(g, t)$'s, summarise into various effects.

First, avg $ATT(g, t)$'s (weight by group size):

```
aggte(cs_out, type = "simple")
```

Call:

```
aggte(MP = cs_out, type = "simple")
```

Reference: Callaway, Brantly and Pedro H.C. Sant'Anna.

ATT	Std. Error	[95% Conf. Int.]
-0.04	0.0124	-0.0643 -0.0156 *

C-S Example: Aggregate by group g

Second, get ATE for each group, avg across groups.

C-S Example: Aggregate by group g

Second, get ATE for each group, avg across groups.

```
group_effects <- aggte(cs_out, type = "group")
```

C-S Example: Aggregate by group g

Second, get ATE for each group, avg across groups.

```
group_effects <- aggte(cs_out, type = "group")
```

Overall summary of ATT's based on group/cohort aggregation:

ATT	Std. Error	[95% Conf. Int.]
-0.031	0.0127	-0.056 -0.006 *

Group Effects:

Group	Estimate	Std. Error	[95% Simult. Conf. Band]
2004	-0.0797	0.0301	-0.1455 -0.0140 *
2006	-0.0229	0.0172	-0.0606 0.0148
2007	-0.0261	0.0169	-0.0631 0.0109

C-S Example: Aggregate by Dosage

Third, by amt of treatment/exposure length/dosage
 (“event study”):


C-S Example: Aggregate by Dosage

Third, by amt of treatment/exposure length/dosage (“event study”):

```
cs_es_out <- aggte(cs_out, type = "dynamic")
```

C-S Example: Aggregate by Dosage

```
ggdid(cs_es_out)
```



C-S Example: Aggregate by Dosage

```
summary(cs_es_out)
```

C-S Example: Aggregate by Dosage

```
summary(cs_es_out)
```

Overall summary of ATT's based on event-study/dynamic aggregation:

ATT	Std. Error	[95% Conf. Int.]
-0.0772	0.0222	-0.1207 -0.0338 *

C-S Example: Aggregate by Dosage

```
summary(cs_es_out)
```

Overall summary of ATT's based on event-study/dynamic aggregation:

ATT	Std. Error	[95% Conf. Int.]
-0.0772	0.0222	-0.1207 -0.0338 *

“Overall” effect of participating: get avg effect at each exposure length, avg those across different exposure lengths.

C-S Example: Aggregate by Time t

Fourth, get avg effect for each time period, avg across periods.

```
cs_cal_out <- aggte(cs_out, type = "calendar")
```

Overall summary of ATT's based on calendar time aggregation:


ATT	Std. Error	[95% Conf. Int.]
-0.0417	0.018	-0.0769 -0.0065 *

Time Effects:

Time	Estimate	Std. Error	[95% Simult. Conf. Band]
2004	-0.0105	0.0240	-0.0656 0.0446
2005	-0.0704	0.0300	-0.1392 -0.0016 *
2006	-0.0488	0.0215	-0.0980 0.0003
2007	-0.0371	0.0144	-0.0700 -0.0041 *

C-S Example: Aggregate by Time t

```
ggdid(cs_cal_out)
```



Linear Models and Two-Way Fixed-Effects

Linear Models and Two-Way Fixed-Effects

Consider

$$Y_{i,t} = \alpha_i + \gamma_t + \beta \cdot T_{i,t} + \epsilon_{i,t}$$

Linear Models and Two-Way Fixed-Effects

Consider

$$Y_{i,t} = \alpha_i + \gamma_t + \beta \cdot T_{i,t} + \epsilon_{i,t}$$

► β : DiD estimate (if 2 units, 2 periods!)

Linear Models and Two-Way Fixed-Effects

Consider

$$Y_{i,t} = \alpha_i + \gamma_t + \beta \cdot T_{i,t} + \epsilon_{i,t}$$

- ▶ β : DiD estimate (if 2 units, 2 periods!)
- ▶ More generally, β is weighted avg of *all possible* 2×2 DiD estimators comparing C-S “groups”

Linear Models and Two-Way Fixed-Effects

Consider

$$Y_{i,t} = \alpha_i + \gamma_t + \beta \cdot T_{i,t} + \epsilon_{i,t}$$

- ▶ β : DiD estimate (if 2 units, 2 periods!)
- ▶ More generally, β is weighted avg of *all possible* 2×2 DiD estimators comparing C-S “groups”
- ▶ (Among these, some use “never treated”, others use “not yet treated”)

Linear Models and Two-Way Fixed-Effects

- ▶ Weights: prop to

Linear Models and Two-Way Fixed-Effects

- ▶ Weights: prop to
 - ▶ group size

Linear Models and Two-Way Fixed-Effects

- ▶ Weights: prop to
 - ▶ group size
 - ▶ var in treatment indicator

Linear Models and Two-Way Fixed-Effects

- ▶ Weights: prop to
 - ▶ group size
 - ▶ var in treatment indicator

Linear Models and Two-Way Fixed-Effects

- ▶ Weights: prop to
 - ▶ group size
 - ▶ var in treatment indicator

$$\text{var}(p) = \sqrt{\frac{p(1-p)}{n}}$$

Linear Models and Two-Way Fixed-Effects

- ▶ Weights: prop to
 - ▶ group size
 - ▶ var in treatment indicator

$$\text{var}(p) = \sqrt{\frac{p(1-p)}{n}}$$

- ▶ highest at $p = 0.5$

Linear Models and Two-Way Fixed-Effects

- ▶ Weights: prop to
 - ▶ group size
 - ▶ var in treatment indicator

$$\text{var}(p) = \sqrt{\frac{p(1-p)}{n}}$$

- ▶ highest at $p = 0.5$
- ▶ \leadsto (treated at midpoint) gets most weight

Linear Models and Two-Way Fixed-Effects

- ▶ Weights: prop to
 - ▶ group size
 - ▶ var in treatment indicator

$$\text{var}(p) = \sqrt{\frac{p(1-p)}{n}}$$

- ▶ highest at $p = 0.5$
- ▶ \rightsquigarrow (treated at midpoint) gets most weight
- ▶ Also, can be *negative* if ATE's vary over time!

Linear Models and Two-Way Fixed-Effects

- ▶ Weights: prop to
 - ▶ group size
 - ▶ var in treatment indicator

$$\text{var}(p) = \sqrt{\frac{p(1-p)}{n}}$$

- ▶ highest at $p = 0.5$
- ▶ \rightsquigarrow (treated at midpoint) gets most weight
- ▶ Also, can be *negative* if ATE's vary over time!

Linear Models and Two-Way Fixed-Effects

- ▶ Weights: prop to
 - ▶ group size
 - ▶ var in treatment indicator

$$\text{var}(p) = \sqrt{\frac{p(1-p)}{n}}$$

- ▶ highest at $p = 0.5$
- ▶ \rightsquigarrow (treated at midpoint) gets most weight
- ▶ Also, can be *negative* if ATE's vary over time!

Goodman-Bacon (2021)

Linear Models and Two-Way Fixed-Effects

- ▶ Weights: prop to
 - ▶ group size
 - ▶ var in treatment indicator

$$\text{var}(p) = \sqrt{\frac{p(1-p)}{n}}$$

- ▶ highest at $p = 0.5$
- ▶ \rightsquigarrow (treated at midpoint) gets most weight
- ▶ Also, can be *negative* if ATE's vary over time!

Goodman-Bacon (2021)

So, use C-S aggregation!

Linear Models and Two-Way Fixed-Effects

What if effect constant over time?

Linear Models and Two-Way Fixed-Effects

What if effect constant over time?

(Still, use C-S aggregation!)


Linear Models and Two-Way Fixed-Effects

What if effect constant over time?

(Still, use C-S aggregation!)

Callaway and Sant'Anna (2023)

Linear Models and Two-Way Fixed-Effects



The above plot shows the 1,000 individual values of $Y_{i,t}$, as well as the average by treatment-group (the thicker colorful lines), and the vertical lines show the period when treatment begins for each group. Because treatment effects here grows linearly with time elapsed since treatment started, we can see that the earliest treated units end up with the highest value of Y , and that the difference grows by the gap between treatment years.

Linear Models and Two-Way Fixed-Effects Event Study

Model

$$Y_{i,t} = \alpha_i + \beta_t + \gamma(Pre) + \delta_j(Lag_j) + \zeta_k(Lead_k) + \epsilon_{i,t}$$

- ▶ Indicators for units
- ▶ Indicators for time periods
- ▶ Indicators for lags (“tr how long ago?”)
- ▶ Indicators for leads (“tr how far into future?”)
- ▶ (Cluster SE’s on “states” of assignment)


Linear Models and Two-Way Fixed-Effects Event Study

Model


$$Y_{i,t} = \alpha_i + \beta_t + \gamma(Pre) + \delta_j(Lag_j) + \zeta_k(Lead_k) + \epsilon_{i,t}$$

- ▶ δ_j often interpreted as “ATE of j periods of treatment”
- ▶ ζ_j often interpreted as “pre-tr trends”

Linear Models and Two-Way Fixed-Effects Event Study




Linear Models and Two-Way Fixed-Effects Event Study



Severe bias for “effect of 3 years of Tr”, e.g.!

Linear Models and Two-Way Fixed-Effects Event Study

Instead, using C-S approach correctly ID's identical lead/pre-tr trends and cumulative effects of Tr!



Linear Models and Two-Way Fixed-Effects Event Study

- ▶ If you include a never-treated group, TWFE-ES can recover truth

Linear Models and Two-Way Fixed-Effects Event Study

- ▶ If you include a never-treated group, TWFE-ES can recover truth

Linear Models and Two-Way Fixed-Effects Event Study

- ▶ If you include a never-treated group, TWFE-ES can recover truth (but so does Calloway-Sant'Anna)

Linear Models and Two-Way Fixed-Effects Event Study

- ▶ If you include a never-treated group, TWFE-ES can recover truth (but so does Calloway-Sant'Anna)
- ▶ But, if there are heterogeneous treatment effects, TWFE-ES won't recover truth (even with never-treated group)

Linear Models and Two-Way Fixed-Effects Event Study

- ▶ If you include a never-treated group, TWFE-ES can recover truth (but so does Calloway-Sant'Anna)
- ▶ But, if there are heterogeneous treatment effects, TWFE-ES won't recover truth (even with never-treated group)
- ▶ Calloway-Sant'Anna does

Linear Models and Two-Way Fixed-Effects Event Study

- ▶ If you include a never-treated group, TWFE-ES can recover truth (but so does Calloway-Sant'Anna)
- ▶ But, if there are heterogeneous treatment effects, TWFE-ES won't recover truth (even with never-treated group)
- ▶ Calloway-Sant'Anna does
- ▶ (So, use C-S aggregation!)

Thanks!

rtm@american.edu
www.ryantmoore.org

References I

- Bertrand, Marianne, Esther Duflo, and Sendhil Mullainathan. 2004. “How Much Should We Trust Differences-in-Differences Estimates?” *The Quarterly Journal of Economics* 119 (1): 249–75.
- Callaway, Brantly, and Pedro H. C. Sant’Anna. 2021a. “Did: Difference in Differences.” <https://bcallaway11.github.io/did/>.
- . 2021b. “Difference-in-Differences with Multiple Time Periods.” *Journal of Econometrics* 225 (2): 200–230.
- . 2023. “Problems with Two-Way Fixed-Effects Event-Study Regressions.” <https://bcallaway11.github.io/did/articles/TWFE.html>.
- Card, David, and Alan B. Krueger. 1994. “Minimum Wages and Employment: A Case Study of the Fast-Food Industry in New Jersey and Pennsylvania.” *The American Economic Review* 90 (5): 1397–1420.
- Goodman-Bacon, Andrew. 2021. “Difference-in-Differences with Variation in Treatment Timing.” *Journal of Econometrics* 225: 254–77. <https://doi.org/10.1016/j.jeconom.2021.03.014>.