

Table 1: Goodness of fit of the weighted-cumulative-exposure log-hazard-ratio estimators across the six simulation scenarios of Figure ?? . Entries are the mean integrated squared error, $MISE = \frac{1}{300} \sum_{\text{sims}} \sum_{l=0}^{15} (f(l) - \hat{f}(l))^2$, of $\hat{f}(l)$ relative to the true curve over lags $l = 0, \dots, 15$, averaged over 300 simulations; the smallest value in each row is in bold. All entries are multiplied by 10^5 .

Scenario	Knot-search RCspline	Term-search RCspline	Polynomial (df=5)
Unimodal	1.54	12.47	1.19
Damped exponential	1.32	7.60	0.62
Bimodal 1	7.72	3.82	6.52
Bimodal 2	9.04	72.73	23.90
Null effects	0.21	2.94	0.28
Constant positive effects	12.54	143.69	14.53

Table 2: Goodness of fit of the term-search RCspline estimator under different candidate knot grids and model-selection criteria (unimodal scenario, 300 simulations). Columns compare the AIC and BIC selection criteria used inside the term search; the smaller of the two is in bold. MISE is multiplied by 10^5 .

Candidate knot grid	AIC	BIC
5 knots	1.63	2.66
8 knots	2.25	2.19
16 knots	13.12	3.81