

Documentation DELCODE Inflammation Markers

Concentrations of neurodegeneration and inflammation markers were determined in 309 matched baseline CSF and serum samples from the DELCODE study in conduct of project 095, DELCODE inflammation. Measurements were conducted using immunoassays optimized for detection range of the respective markers using technical duplicates, coefficient of variance <20%, and inter-run-normalization by internal control samples. Variable names are composed as follows: BIOMATERIAL_MARKER_UNIT, e.g. for the marker ferritin in CSF in ng/ml reads CSF_ferritin_ng_ml.

Contact for support / questions:

Dr. Frederic Brosseon (Frederic.brosseon@dzne.de)

Prof. Dr. Michael T. Heneka (Michael.heneka@ukbonn.de)

Overview of determined markers:

Neurodegeneration	FABP-3
	Neurogranin
Multifactorial	Apolipoprotein E
	Ferritin
Inflammation	CRP
	YKL-40 (CHI3L1)
	MIF
	MCP-1 (CCL2)
	IP-10 (CXCL10)
	IL-6
	IL-18
	sTREM2
	sAXL
	sTyro3
	Complement C1q
	Complement C3
	Complement C3b
	Complement C4
	Complement factor H
	Complement factor B

Graph of concentration range of the markers:

