

Introduction to Dynamical Systems

King's College London

2023-2024

4CCM131A

Homework 1

Instructions

This homework has been set to be similar to a potential exam question.

As it is a participation homework, so long as you attempt the homework, the actual mark will not matter for your module mark - a serious attempt at the homework will earn the participation mark.

You do not need to complete all parts of the question to earn the participation mark, but I hope you will try so long as you do not spend too long on it.

This is a typical question: it starts with an easy part and then gets progressively harder. The marks are also not distributed evenly - the last few marks are much harder to get than the first few marks.

1. (9 marks) A first order ODE is an equation of the form

$$\frac{dx}{dt} = f(x, t). \quad (1)$$

- (a) Find $f(x, t)$ such that the ODE is homogeneous but not separable.
- (b) Find $f(x, t)$ such that the ODE is separable but not homogeneous.
- (c) Find $f(x, t)$ such that the ODE both separable and homogeneous.

2. (9 marks) Consider the first order ODE

$$t^2 \frac{dx}{dt} = (t + x)^2 + tx. \quad (2)$$

- (a) Show that the ODE can be written in the form

$$\frac{dx}{dt} = G(v), \quad (3)$$

where $v = x/t$.

- (b) Find the general solution $x(t)$ of the ODE.

3. (7 marks) Consider the initial value problem that consists of the ODE of Eq. (2) and the initial condition

$$x(1) = 0. \quad (4)$$

- (a) Find the solution of the initial value problem.
- (b) Apply Picard's theorem and show that the solution to the initial value problem exists in $[t_0 - \delta_1, t_0 + \delta_1]$ and it is unique. What is the range of possible values of δ_1 ?