

# ARIMA - integrated ARMA

ARIMA MODELS IN R





**David Stoffer**

Professor of Statistics at the University  
of Pittsburgh

# Identifying ARIMA


- A time series exhibits ARIMA behavior if the differenced data has ARMA behavior

```
# Simulation  ARIMA(p = 1, d = 1, q = 0)
x <- arima.sim(list(order = c(1, 1, 0),  ar = .9), n = 200)
plot(x, main = "ARIMA(p = 1, d = 1, q = 0)")
plot(diff(x), main = "ARMA(p = 1, d = 0, q = 0)")
```




# ACF and PCF of an Integrated ARMA

```
x <- arima.sim(list(order = c(1, 1, 0), ar = .9), n = 200)
acf2(x)
```




# ACF and PCF of a Differenced ARIMA

```
x <- arima.sim(list(order = c(1, 1, 0), ar = .9), n = 200)
acf2(diff(x))
```




# ACF and PCF of a Differenced ARIMA

```
x <- arima.sim(list(order = c(1, 1, 0), ar = .9), n = 200)
acf2(diff(x))
```




Indicates an  
AR(1) for the  
differenced  
data

# Weekly Oil Prices



# Weekly Oil Prices



- Looks like `ARIMA(1, 1, 1)`

**Let's practice!**  
ARIMA MODELS IN R



# ARIMA diagnostics


ARIMA MODELS IN R




**David Stoffer**

Professor of Statistics at the University  
of Pittsburgh

# Weekly Oil Prices ARIMA(1, 1, 1)?



# Weekly Oil Prices ARIMA(1, 1, 1)?




# Weekly Oil Prices ARIMA(1, 1, 1)?

```
oil <- window(oil, end = 2006)
x <- sarima(oil, p = 1, d = 1, q = 1)
x$tttable
```

	Estimate	SE	t.value	p.value
ar1	-0.4987	0.0995	-5.0131	0.0000
ma1	0.7316	0.0734	9.9732	0.0000
constant	0.1091	0.0936	1.1664	0.2443

# Weekly Oil Prices ARIMA(1, 1, 1)!



# Overfit: ARIMA(2, 1, 1) and ARIMA(1, 1, 2)

```
oil_fit1 <- sarima(oil, p = 2, d = 1, q = 1)
oil_fit1$tttable
```

	Estimate	SE	t.value	p.value
ar1	-0.4704	0.1117	-4.2121	0.0000
ar2	-0.0738	0.0652	-1.1319	0.2586
ma1	0.6771	0.0986	6.8696	0.0000
constant	0.1088	0.0878	1.2391	0.2163

```
oil_fit2 <- sarima(oil, p = 1, d = 1, q = 2)
oil_fit2$tttable
```

	Estimate	SE	t.value	p.value
ar1	-0.3664	0.1816	-2.0178	0.0445
ma1	0.5777	0.1818	3.1777	0.0016
ma2	-0.0836	0.0837	-0.9989	0.3186
constant	0.1088	0.0884	1.2306	0.2194

# Let's practice!

ARIMA MODELS IN R

# Forecasting ARIMA

ARIMA MODELS IN R



**David Stoffer**

Professor of Statistics at the University  
of Pittsburgh




# Forecasting ARIMA Processes

- The model describes how the dynamics of the time series behave over time
- Forecasting simply continues the model dynamics into the future
- Use `sarima.for()` to forecast in the `astsa` package

# Forecasting ARIMA Processes

```
oil <- window(astsa::oil, end = 2006)
oilf <- window(astsa::oil, end = 2007)
sarima.for(oil, n.ahead = 52, 1, 1, 1)
lines(oilf)
```



# Let's practice!

ARIMA MODELS IN R