

AR and MA models

ARIMA MODELS IN R




David Stoffer

Professor of Statistics at the University
of Pittsburgh

AR and MA Models

```
x <- arima.sim(list(order = c(1, 0, 0), ar = -.7), n = 200)
y <- arima.sim(list(order = c(0, 0, 1), ma = -.7), n = 200)
par(mfrow = c(1, 2))
plot(x, main = "AR(1)")
plot(y, main = "MA(1)")
```




ACF and PACF

	AR(p)	MA(q)	ARMA(p, q)
ACF	Tails off	Cuts off lag q	Tails off
PACF	Cuts off lag p	Tails off	Tails off

ACF and PACF

	AR(p)	MA(q)	ARMA(p, q)
ACF	Tails off	Cuts off lag q	Tails off
PACF	Cuts off lag p	Tails off	Tails off




ACF and PACF

	AR(p)	MA(q)	ARMA(p, q)
ACF	Tails off	Cuts off lag q	Tails off
PACF	Cuts off lag p	Tails off	Tails off


ACF and PACF

	AR(p)	MA(q)	ARMA(p, q)
ACF	Tails off	Cuts off lag q	Tails off
PACF	Cuts off lag p	Tails off	Tails off



Estimation

- Estimation for time series is similar to using least squares for regression
- Estimates are obtained numerically using ideas of Gauss and Newton



Estimation with astsa

- AR(2) with mean 50:

$$W_t = 50 + 1.5(X_{t-1} - 50) - .75(X_{t-2} - 50) + W_t$$

```
x <- arima.sim(list(order = c(2, 0, 0),  
                    ar = c(1.5, -.75)),  
               n = 200) + 50  
x_fit <- sarima(x, p = 2, d = 0, q = 0)  
x_fit$tttable
```

	Estimate	SE	t.value	p.value
ar1	1.5429	0.0435	35.4417	0
ar2	-0.7752	0.0434	-17.8650	0
xmean	49.6984	0.3057	162.5788	0

Estimation with astsa

- MA(1) with mean 0:

$$X_t = W_t - .7W_{t-1}$$

```
y <- arima.sim(list(order = c(0, 0, 1), ma = -.7), n = 200)
y_fit <- sarima(y, p = 0, d = 0, q = 1)
y_fit$tttable
```

	Estimate	SE	t.value	p.value
ma1	-0.7459	0.0513	-14.5470	0.00000
xmean	0.0324	0.0191	1.6946	0.0917

Let's practice!
ARIMA MODELS IN R

AR and MA together

ARIMA MODELS IN R



David Stoffer

Professor of Statistics at the University
of Pittsburgh

AR and MA Together: ARMA

$$X_t = \phi X_{t-1} + W_t + \theta W_{t-1}$$

AR and MA Together: ARMA


$$X_t = \phi X_{t-1} + W_t + \theta W_{t-1}$$

auto-regression with correlated errors

AR and MA Together: ARMA

$$X_t = \phi X_{t-1} + W_t + \theta W_{t-1}$$

auto-regression with correlated errors



```
x <- arima.sim(list(order = c(1, 0, 1),  
                    ar = .9,  
                    ma = -.4),  
               n = 200)  
plot(x, main = "ARMA(1, 1)")
```



ACF and PACF of ARMA Models

	AR(p)	MA(q)	ARMA(p, q)
ACF	Tails off	Cuts off lag q	Tails off
PACF	Cuts off lag p	Tails off	Tails off

ACF and PACF of ARMA Models

	AR(p)	MA(q)	ARMA(p, q)
ACF	Tails off	Cuts off lag q	Tails off
PACF	Cuts off lag p	Tails off	Tails off

$$X_t = .9X_{t-1} + W_t - .4W_{t-1}$$



Estimation

$$X_t = .9X_{t-1} + W_t - .4W_{t-1}$$

```
x <- arima.sim(list(order = c(1, 0, 1),
                    ar = .9,
                    ma = -.4),
               n = 200)
x_fit <- sarima(x, p = 1, d = 0, q = 1)
x_fit$tttable
```

	Estimate	SE	t.value	p.value
ar1	0.9083	0.0424	21.4036	0
ma1	-0.4458	0.0879	-5.0716	0
xmean	49.5647	0.4079	121.5026	0

Let's practice!

ARIMA MODELS IN R

Model choice and residual analysis

ARIMA MODELS IN R



David Stoffer

Professor of Statistics at the University
of Pittsburgh

AIC and BIC

$$\begin{array}{ccc} \text{Error} & & \text{Number of Parameters} \\ \text{average}(\text{observed} - \text{predicted})^2 & + & k(p + q) \\ \downarrow & & \uparrow \end{array}$$

- AIC and BIC measure the error and penalize (differently) for adding parameters
- For example, AIC has $k = 2$ and BIC has $k = \log(n)$
- Goal: find the model with the smallest AIC or BIC



Model Choice: AR(1) vs. MA(2)

```
gnpgr <- diff(log(gnp))  
sarima(gnpgr, p = 1, d = 0, q = 0)
```

\$AIC	\$BIC
-8.294403	-9.263748

```
sarima(gnpgr, p = 0, d = 0, q = 2)
```


\$AIC	\$BIC
-8.297695	-9.251712



Residual Analysis


`sarima()` includes residual analysis graphic showing:

1. Standardized residuals
2. Sample ACF of residuals
3. Normal Q-Q plot
4. Q-statistic p-values




Bad Residuals

- Pattern in the residuals



Bad Residuals


✗ Pattern in the residuals



Bad Residuals

✗ Pattern in the residuals


- ACF has large values



Bad Residuals

✗ Pattern in the residuals

✗ ACF has large values




Bad Residuals

✗ Pattern in the residuals

✗ ACF has large values

- Q-Q plot suggests normality




Bad Residuals

✗ Pattern in the residuals

✗ ACF has large values

- Q-Q plot suggests normality
- Q-statistic - all points below line




Bad Residuals

✗ Pattern in the residuals

✗ ACF has large values

- Q-Q plot suggests normality

✗ Q-statistic - all points below line



Let's practice!

ARIMA MODELS IN R