



# X-RAY RESONANT RAMAN SCATTERING

SPECTRA SIMULATION FROM FIRST PRINCIPLES  
FOR COPPER BELOW IONIZATION THRESHOLD  
USING HIGH-PERFORMANCE COMPUTING

GONÇALO GARCÊS SOBREIRA RODRIGUES BAPTISTA

BSc in Physics Engineering

DOCTORATE IN PHYSICS ENGINEERING

NOVA University Lisbon  
September, 2023

# X-RAY RESONANT RAMAN SCATTERING

SPECTRA SIMULATION FROM FIRST PRINCIPLES  
FOR COPPER BELOW IONIZATION THRESHOLD  
USING HIGH-PERFORMANCE COMPUTING

GONÇALO GARCÊS SOBREIRA RODRIGUES BAPTISTA

BSc in Physics Engineering

**Advisers:** Jorge Felizardo Machado  
*Auxiliary Professor, NOVA University Lisbon*  
Mauro António Guerra  
*Auxiliary Professor, NOVA University Lisbon*

## Examination Committee

**Chair:** Name of the committee chairperson  
*Full Professor, FCT-NOVA*

**Rapporteur:** Name of a rapporteur  
*Associate Professor, Another University*

**Members:** Another member of the committee  
*Full Professor, Another University*  
Yet another member of the committee  
*Assistant Professor, Another University*

DOCTORATE IN PHYSICS ENGINEERING

NOVA University Lisbon

September, 2023

## **X-ray resonant Raman scattering**

### **Spectra simulation from first principles for Copper bellow ionization threshold using high-performance computing**

Copyright © Gonalo Garcês Sobreira Rodrigues Baptista, NOVA School of Science and Technology, NOVA University Lisbon.

The NOVA School of Science and Technology and the NOVA University Lisbon have the right, perpetual and without geographical boundaries, to file and publish this dissertation through printed copies reproduced on paper or on digital form, or by any other means known or that may be invented, and to disseminate through scientific repositories and admit its copying and distribution for non-commercial, educational or research purposes, as long as credit is given to the author and editor.

---

# Acknowledgements

Acknowledgments are personal text and should be a free expression of the author.

However, without any intention of conditioning the form or content of this text, I would like to add that it usually starts with academic thanks (instructors, etc.); then institutional thanks (Research Center, Department, Faculty, University, FCT / MEC scholarships, etc.) and, finally, the personal ones (friends, family, etc.).

But I insist that there are no fixed rules for this text, and it must, above all, express what the author feels.

---

”

*“Sometimes I’ll start a sentence, and I don’t even know where it’s going.  
I just hope I find it along the way.  
Like an improv conversation.  
An improversation.”*

— **Michael Scott**, *The Office*  
(Regional Manager of Dunder Mifflin Scranton)

---

”

*all work and no play makes Jack a dull boy*

[illegible]

— Jack Torrance, *The Shinning*  
(Caretaker of the Overlook)

---

## Abstract

The work performed on this thesis comes as part of the effort to further understand the highly convoluted structure present on Copper's X-ray emission spectra, where, as with many other transition metals, a skewness can be observed on the  $K_{\alpha_{1,2}}$ ,  $K_{\beta}$  and L transition lines. These lines originate due to the radiative relaxation of the atom's electronic structure post-ionization of inner shell electrons. However, it is very likely that the observed skewness is due to copper's satellite states' transitions.

Throughout this thesis, a study will be performed for the satellite states formed by the excitation of the inner-shell electrons, where, as opposed to the ionization process, usually considered in X-ray calculations, a photoexcitation process occurs.

Multiple atomic structure calculations will be performed using the *ab initio* state of the art [Multiconfiguration Dirac-Fock General Matrix Elements \(MCDFGME\)](#) code for different excited states configurations.

The obtained results will then be used in the analysis of experimental data obtained from a High-Precision [Double Crystal Spectrometer \(DCS\)](#), using a synchrotron X-ray source.

Due to the complexity of the calculations, the process can become substantial in terms of computational power and time. Therefore, further similar and more complex studies will be performed by implementing and running a script in the *Oblivion* supercomputer located at the University of Évora.

**Keywords:** Atomic Excitation, X-ray lines, [MCDFGME](#), [DCS](#), High Performance Computing



---

# Resumo

asdasdasdasd

---

# Contents

|  |           |
|--|-----------|
| List of Figures  | x         |
| List of Tables   | xi        |
| Listings   | xii       |
| Glossary   | xiii      |
| Acronyms   | xiv       |
| <b>1 Introduction</b>  | <b>2</b>  |
| 1.1 Theoretical Introduction . . . . .                                   | 2         |
| 1.1.1 Characteristic X-rays . . . . .                                    | 2         |
| 1.1.2 Radiative transitions . . . . .                                    | 5         |
| 1.2 Atomic Structure Calculations . . . . .                              | 6         |
| 1.2.1 The non-relativistic Hamiltonian . . . . .                         | 6         |
| 1.2.2 The Dirac Equation . . . . .                                       | 7         |
| 1.2.3 The Dirac-Breit Equation . . . . .                                 | 10        |
| 1.2.4 The Multiconfiguration Dirac-Fock ( <i>MCDF</i> ) Method . . . . . | 10        |
| 1.3 State of the Art . . . . .   | 11        |
| 1.3.1 Copper’s characteristic X-rays . . . . .                           | 11        |
| 1.3.2 MCDFGME capabilities . . . . .                                     | 11        |
| 1.4 Methodology . . . . .  | 12        |
| <b>2 Theoretical Introduction</b>  | <b>15</b> |
| 2.1 Characteristic X-rays . . . . .                                      | 15        |
| 2.1.1 Ionization as a vacancy generator . . . . .                        | 15        |
| 2.1.2 Transition notation . . . . .                                      | 17        |
| 2.1.3 Excitation as a vacancy generator . . . . .                        | 17        |
| 2.2 Radiative transitions . . . . .                                      | 18        |

|          |  |           |
|----------|--|-----------|
| 2.3      | Solving the atomic many-body problem . . . . . | 19        |
| 2.3.1    | The non-relativistic Hamiltonian . . . . .     | 19        |
| 2.3.2    | The Hartree-Fock Method . . . . .              | 20        |
| 2.3.3    | The Dirac Equation . . . . .                   | 21        |
| 2.3.4    | The <i>MCDF</i> Method . . . . .               | 24        |
| 2.4      | State of the Art . . . . .                     | 24        |
| 2.4.1    | Copper's characteristic X-rays . . . . .       | 24        |
| 2.4.2    | MCDFGME capabilities . . . . .                 | 25        |
| 2.5      | Methodology . . . . .                          | 25        |
| <b>3</b> | <b>Atomic Structure Calculations</b>           | <b>27</b> |
| 3.1      | The non-relativistic Hamiltonian . . . . .     | 27        |
| 3.1.1    | The Hartree-Fock Method . . . . .              | 27        |
| 3.2      | The Dirac Equation . . . . .                   | 28        |
| 3.3      | The Dirac-Breit Equation . . . . .             | 31        |
| 3.4      | The <i>MCDF</i> Method . . . . .               | 31        |
|          | <b>Bibliography</b>                            | <b>33</b> |
|          | <b>Appendices</b>                              |           |
| <b>A</b> | <b>The Breit Hamiltonian Operators</b>         | <b>37</b> |
| <b>B</b> | <b>Transition Diagram</b>                      | <b>39</b> |
| <b>C</b> | <b>QED considerations</b>                      | <b>40</b> |
| C.1      | Self-Energy . . . . .                          | 40        |
| C.2      | Vacuum Polarization . . . . .                  | 40        |
|          | <b>Annexes</b>                                 |           |
| <b>I</b> | <b>Input File .f05 Example</b>                 | <b>41</b> |

---

## List of Figures

|     |  |    |
|-----|--|----|
| 1.1 | Photoionization. . . . .                               | 3  |
| 1.2 | Principal atomic relaxation processes. . . . .         | 3  |
| 1.3 | Resonant Photoexcitation . . . . .                     | 5  |
| 1.4 | HF method's block diagram. . . . .                     | 8  |
| 2.1 | Photoionization. . . . .                               | 16 |
| 2.2 | Principal atomic relaxation processes. . . . .         | 16 |
| 2.3 | Resonant Photoexcitation . . . . .                     | 18 |
| 2.4 | HF method's block diagram. . . . .                     | 21 |
| 3.1 | HF method's block diagram. . . . .                     | 29 |
| B.1 | Transition notation scheme. Adapted from [1] . . . . . | 39 |
| C.1 | QED Feynman Diagrams . . . . .                         | 40 |

---

## List of Tables

|     |   |    |
|-----|---|----|
| 1.1 | Siegbahn VS IUPAC notation. Adapted from [1]. . . . . | 4  |
| 2.1 | Siegbahn VS IUPAC notation. Adapted from [1]. . . . . | 18 |

---

## Listings

---

## Glossary

|                                |   |
|--------------------------------|---|
| <b>Feynman Diagram</b>         | 2D diagram depicting various physical interactions between elementary particles. Fermions are depicted as straight lines and bosons as wavy lines. (p. 40)  |
| <b>four-vector</b>             | Vector used in special relativity composed of 4 components, one scalar time-like, and three vectorial space-like. These vectors behave in special way, such as their norm being <a href="#">Lorentz invariant</a> . Can be written in covariant, $X_\mu$ , and contravariant form, $X^\mu$ , with the difference being the sign of the time-like components. Example of a contravariant four vector: $X^\mu = (X^0, X^1, X^2, X^3) = (X^0, \mathbf{X})$ (pp. <a href="#">xiii</a> , <a href="#">7</a> , <a href="#">22</a> , <a href="#">28</a> ) |
| <b>Lorentz invariant</b>       | A Lorentz invariant scalar, obtained, for example, from a <a href="#">Minkowski norm</a> , does not change when operated by a Lorentz Transformation. (pp. <a href="#">xiii</a> , <a href="#">7</a> , <a href="#">22</a> , <a href="#">28</a> )   |
| <b>Minkowski norm</b>          | Yields the Lorentz Invariant norm for a <a href="#">four-vector</a> : $p_\mu p^\mu$ . Equivalent to the dot product of a classical vector. (pp. <a href="#">xiii</a> , <a href="#">7</a> , <a href="#">22</a> , <a href="#">28</a> )  |
| <b>Self Consistency Method</b> | The electron wavefunctions go through a variational process in order to reach convergence for an energetic minimum. (p. <a href="#">20</a> )  |
| <b>virtual photons</b>         | While in reality, during a Coulomb interaction, 'real' particles are not exchanged, the electromagnetic field is still mediated by photons. This way virtual photons are tools used in order to better represent electromagnetic interactions. (pp. <a href="#">10</a> , <a href="#">24</a> , <a href="#">31</a> )  |

---

## Acronyms

|                       |  |
|-----------------------|--|
| <b><i>Grasp2k</i></b> | General-purpose Relativistic Atomic Structure Package 2k ( <i>pp.</i> <a href="#">11</a> , <a href="#">25</a> )  |
| <b><i>MCDFGME</i></b> | Multiconfiguration Dirac-Fock General Matrix Elements ( <i>pp.</i> <a href="#">vi</a> , <a href="#">10–12</a> , <a href="#">24–26</a> , <a href="#">31</a> ) |
| <b><i>MCDF</i></b>    | Multiconfiguration Dirac-Fock ( <i>pp.</i> <a href="#">viii</a> , <a href="#">ix</a> , <a href="#">10</a> , <a href="#">24</a> , <a href="#">31</a> )        |
| <b>DCS</b>            | Double Crystal Spectrometer ( <i>p.</i> <a href="#">vi</a> )   |
| <b>EMF</b>            | Electromagnetic Field ( <i>p.</i> <a href="#">19</a> )   |
| <b>FAC</b>            | Flexible Atomic Code ( <i>pp.</i> <a href="#">11</a> , <a href="#">25</a> )  |
| <b>QED</b>            | Quantum Electrodynamics ( <i>pp.</i> <a href="#">2</a> , <a href="#">15</a> )  |



---

## Todo list

|   |    |
|---|----|
| Figure: Maybe put a Thomas model plot? . . . . .  | 17 |
| Explain antisymmetry and put it in the glossary . . . . .   | 20 |
| Explain in the glossary . . . . .   | 20 |
| Finish the explanation . . . . .  | 21 |
| eheheheh . . . . .  | 22 |
| spin? . . . . .   | 22 |
| Ler bem o manual e explicar bem o método e todas as coisas diferentes que se podem<br>fazer . . . . . | 24 |
| This really needs a re-do . . . . .   | 24 |
| Take this out . . . . .   | 25 |

---

# Introduction

## 1.1 Theoretical Introduction

In this section, the theoretical topics that served as base for the work will be discussed. They will include themes from the much used characteristic X-rays of elements, to the relativistic Dirac equation, with [Quantum Electrodynamics \(QED\)](#) effects corrections and methods for solving the many-body problem.

### 1.1.1 Characteristic X-rays

The X-rays emitted by an element when exposed to a high energy particle beam (usually made of electrons or photons), can be used to determine what the analyzed element is. Therefore, the study of this characteristic radiation is of high importance in many scientific areas.

#### 1.1.1.1 Ionization as a vacancy generator

When an atomic system is in a bound state, the electrons orbiting the nucleus will be occupying fixed quantum states, defined by their principal atomic number,  $n$ , angular momentum,  $l$ , and spin,  $s$ . Electrons are also fermions, thus must respect Pauli's exclusion principle, each occupying a single state, only occupied by that single electron. These quantum numbers will provide information of their specific electron's wavefunction, which, when evaluated by operation of the Hamiltonian, will yield that state's energy. Furthermore, beside the occupied states, more energetic unoccupied states will still be part of the system's eigenfunctions basis.

The work performed in this thesis will explore what would happen should the atomic system experience a perturbation that changes what the occupied and unoccupied states are, while still preserving the number of electrons within.

X-ray fluorescence spectroscopy has many uses and applications in a wide range of scientific areas. In this form of spectroscopy, the element under study, composed of the nucleus and  $N$  orbiting electrons, is bombarded with X-rays leading to the photoionization

of inner-shell electrons and leaving a vacancy where the ejected electron used to be (Figure 2.1). The atomic structure, now composed of  $N - 1$  electrons, will be left in an energetically unstable state, due to there being other possible lower energy configurations. This will lead to various processes of atomic relaxation, where the system will rearrange itself in order to find a lower energy configuration.




Figure 1.1: Photoionization.

The main processes for this rearrangement are two different competing processes: radiative relaxation, where an electron from a higher orbital changes state in order to now occupy the orbital where the vacancy was, emitting an X-ray photon of characteristic energy, allowing for the easy identification of the element at study; The other competing process, the non-radiative relaxation, more commonly known as the Auger process, will still have the electron transit between states, but where a photon was previously emitted, a new ionization will now occur.




Figure 1.2: Principal atomic relaxation processes.

In reality, when the initial vacancy is generated due to an ionization, two more processes can occur: shake-off and shake-up. When this processes occur, due to a sudden change in the potential felt by the remaining electrons, during the ionization, one other

electron may become unbound, occurring a second ionization, in the shake-off process, or may be excited to an upper state, during shake-up. These processes occur due to the fact that the system, while it may have had a set of eigenstates before ionization, was changed when an electron was removed, leading to the previous wavefunctions no longer being part of the new system's eigenfunction basis, and the new system not being able to be accurately described by the previous system's states. These processes, will not be studied throughout this thesis, as the work in question focuses on excitation, while disregarding the possibility of initial ionization, later explained in Section 2.1.3.

### 1.1.1.2 Transition notation

The characteristic radiation measured from the relaxation of a post-ionization unstable atomic system is one of the main ways of identifying an atomic element. This is due to the photons emitted from different elements possessing distinct, quantized values of energy, forming well-defined energy lines when observed in a spectrometer. These lines have specific notations denoting them by vacancy (hole) and the electron's orbital. A very illustrative diagram, exemplifying some transitions can be found in Appendix B.

Throughout this thesis, Siegbahn notation will be used, for the most part, but should the reader prefer IUPAC's, Table 2.1 explains the conversion between notations.

Table 1.1: Siegbahn VS IUPAC notation. Adapted from [1].

| Siegbahn           | IUPAC     | Siegbahn       | IUPAC           | Siegbahn        | IUPAC           |
|--------------------|-----------|----------------|-----------------|-----------------|-----------------|
| $K_{\alpha_1}$     | $K - L_3$ | $L_{\alpha_1}$ | $L_3 - M_5$     | $L_{\gamma_1}$  | $L_2 - N_4$     |
| $K_{\alpha_2}$     | $K - L_2$ | $L_{\alpha_2}$ | $L_3 - M_4$     | $L_{\gamma_2}$  | $L_1 - N_1$     |
| $K_{\beta_1}$      | $K - M_3$ | $L_{\beta_1}$  | $L_2 - M_4$     | $L_{\gamma_3}$  | $L_1 - N_2$     |
| $K_{\beta_2}^I$    | $K - N_3$ | $L_{\beta_2}$  | $L_3 - N_5$     | $L_{\gamma_4}$  | $L_1 - O_3$     |
| $K_{\beta_2}^{II}$ | $K - N_2$ | $L_{\beta_3}$  | $L_1 - M_3$     | $L_{\gamma_4}'$ | $L_1 - O_2$     |
| $K_{\beta_3}$      | $K - M_2$ | $L_{\beta_4}$  | $L_1 - M_2$     | $L_{\gamma_5}$  | $L_2 - N_1$     |
| $K_{\beta_4}^I$    | $K - N_5$ | $L_{\beta_5}$  | $L_3 - O_{4,5}$ | $L_{\gamma_5}$  | $L_2 - O_4$     |
| $K_{\beta_4}^{II}$ | $K - N_4$ | $L_{\beta_6}$  | $L_3 - N_1$     | $L_{\gamma_8}$  | $L_2 - O_1$     |
| $K_{\beta_4}^x$    | $K - N_4$ | $L_{\beta_7}$  | $L_3 - O_1$     | $L_{\gamma_8}'$ | $L_2 - N_{5,6}$ |
| $K_{\beta_5}^I$    | $K - M_5$ | $L_{\beta_8}$  | $L_3 - N_{6,7}$ | $L_{\eta}$      | $L_2 - M_1$     |
| $K_{\beta_4}^{II}$ | $K - M_4$ | $L_{\beta_9}$  | $L_1 - M_5$     | $L_l$           | $L_3 - M_1$     |
|                    |           | $L_{\beta_9}$  | $L_1 - M_4$     | $L_s$           | $L_3 - M_3$     |
|                    |           | $L_{\beta_9}$  | $L_3 - N_4$     | $L_t$           | $L_3 - M_2$     |
|                    |           | $L_{\beta_9}$  | $L_2 - M_3$     | $L_u$           | $L_3 - N_{6,7}$ |
|                    |           |                |                 | $L_v$           | $L_2 - N_{6,7}$ |

### 1.1.1.3 Excitation as a vacancy generator

As previously mentioned, throughout this thesis, while the study is focused on the characteristic radiation emitted during the atomic relaxation process, the vacancy generation method will be due to an excitation (Figure 2.3), instead of an ionization. This is one possibility of the many so-called satellite states, where the electronic configuration present during the relaxation process contains additional electrons or holes. The characteristic radiation from these states may be one of the keys needed to fully comprehend and deconvolute an element's emission spectra.




Figure 1.3: Resonant Photoexcitation

### 1.1.2 Radiative transitions

In terms of quantum mechanics, the properties of a system's change of state can be derived by perturbation theory, where the change of energy can be treated as perturbation, and takes into account angular momentum conservation laws as basis for the selection rules.

The transition's radiation type can be of two main types, which can be either Electric,  $E_k$ , or Magnetic,  $M_k$ , with  $k$  representing its multiplicity. In both these transitions the system's total angular momentum is allowed a change of  $\Delta J = 0, \pm k$ , and so does its projection,  $M_J$ . There are, however, different changes of parity in the system, depending on the occurred transition, and the selection rules have many other exceptions.

It is also of note that, usually, for the same multiplicity, an  $E$  transition is more intense than an  $M$ , and that it is also possible for rare 2-photon transitions to take place, where a combination of different transition types can occur.

The intensity of a transition is proportional to the squared norm of the perturbation's matrix element involving the initial and final state  $\left(\Gamma_{if} \propto |\langle \psi_i | H' | \psi_f \rangle|^2\right)$ . It should also be noted that the electron's initial state population will serve as a scaling factor for the transitions rate. For example, while the transition rate for a  $2p_{1/2} \rightarrow 1s$  and  $2p_{3/2} \rightarrow 1s$  should be about the same, the orbital  $2p_{3/2}$  has double the population  $2p_{1/2}$  has, hence why the  $K_{\alpha_2}$  line has about half the  $K_{\alpha_1}$  line's intensity.

One should also mention that monochromatic transitions do not exist. While the transition might have a well-defined energy, calculated by the difference in energy between the initial and final states, due to Heisenberg's uncertainty principle,  $\Delta E \Delta t > \frac{\hbar}{2}$ , there will exist a natural energy broadening, the natural width. The shapes representing these transitions are given by a Lorentzian distribution.

## 1.2 Atomic Structure Calculations

When studying a system composed of multiple charged bodies, one must consider all the existing interactions. Whilst there are known analytical solutions for the 2-bodies Hydrogenoid systems, with the presence of more non spatially-bound particles, these become impossible to find for the Schrödinger equation, even while using the Born-Oppenheimer approximation. This approximation considers the nucleus as static, at a fixed position, making the electrons the only non spatially-bound particles in the system. That way, there was a need for the development of numerical solutions to solve this problem.

### 1.2.1 The non-relativistic Hamiltonian

The first approach used in order to solve the many-bodies problem used a non-relativistic consideration. This way, the Hamiltonian consisted on the sum of the system's non-relativistic momentum-related energies and the Coulomb interactions between bodies, while still considering the Born-Oppenheimer approximation.

Essentially, and in atomic units:

$$\underbrace{\sum_i^N \left( \overbrace{\frac{1}{2} \nabla_i^2}^{E_1} - \overbrace{\frac{Z}{r_i}}^{E_2} \right)}_{\text{Individual Hamiltonian}} + \underbrace{\sum_{i < j}^j \overbrace{\frac{1}{r_{ij}}}^{E_3}}_{\text{Pair repulsion}}$$

$$E_1 \rightarrow \text{Momentum} \quad E_2 \rightarrow e^- \text{ nuc. Coulomb attraction} \quad E_3 \rightarrow e^- e^- \text{ Coulomb repulsion} \quad (1.1)$$

#### 1.2.1.1 The Hartree-Fock Method

*Note: This section used the works in [2–5] as reference.*

Hartree developed an iterative method, further enhanced by Fock and Slater, based on the field self-consistency method.

In this method, while studying a multi-electronic system, such as an atom, each electron's wavefunction is composed as a product of a spacial part,  $\psi$ , and one indicating the electron's spin,  $\chi$ , as to be able to account for relativistic effects.

$$u = \psi\chi \quad (1.2)$$

The wavefunction capable of describing the whole system,  $\Psi$ , should be somewhat of a product of all the wavefunctions describing each individual electron. However, one must not forget the need for this wavefunction to respect the antisymmetry principle, due to the electron's fermionic nature. In order to respect this,  $\Psi$  is to be composed of a Slater determinant:

$$\Psi = \frac{1}{\sqrt{N!}} \begin{vmatrix} u_1(x_1) & u_2(x_1) & \cdots & u_N(x_1) \\ u_1(x_2) & u_2(x_2) & \cdots & u_N(x_2) \\ \vdots & \vdots & \ddots & \vdots \\ u_1(x_N) & u_2(x_N) & \cdots & u_N(x_N) \end{vmatrix} \quad (1.3)$$

It is of high importance that the wavefunctions must form an orthonormal basis. These are to be initialized as trial wavefunctions.

The main goal of this method is to follow the variational principle and to minimize a functional, with the purpose of minimizing the system's energy. This optimal, yet unknown energy,  $E_0$  (calculated by Operating the Hamiltonian on the optimal wavefunctions), will try to be reached by varying the trial functions that provide a non-minimized solution, with  $\langle \Psi | H | \Psi \rangle \geq E_0$ .

The computational method consists on starting with the previously mentioned trial wavefunctions, using them to calculate Hartree-Fock's potential through the HF equations.

In a very simplified way, the self-consistent Hartree-Fock computational method can be represented by the block diagram in Figure 3.1.

### 1.2.2 The Dirac Equation

*Note: This section used the works in [6–9] as reference.*

It is no secret that the Schrödinger equation has some very considerable limitations. The fact that it does not account for the existence of the electron's spin and the lack of consideration of relativistic effects are some of the most impactful setbacks.

Many scientists, such as Klein, Gordon and later Fock, had already conceived a relativistic correction to Schrödinger's equation, where the free-particle energy makes use of the relativistic momentum-energy relation (3.4).

$$E = \sqrt{c^2 p^2 + m^2 c^4} \quad (1.4)$$

Which can be derived from the [Lorentz invariant Minkowski norm](#) of the momentum four-vector (3.5).

$$p^\mu p_\mu = m^2 c^2 \Leftrightarrow \frac{E^2}{c^2} - \mathbf{p}^2 = m^2 c^2 \Leftrightarrow \frac{E^2}{c^2} = \mathbf{p}^2 + m^2 c^2 \quad (1.5)$$




Figure 1.4: HF method's block diagram.

Now, inputting this new energy operator into Schrödinger's equation, yields the Klein-Gordon equation (3.6), allowing for Schrödinger's equation now to be Lorentz-invariant.

$$-\hbar^2 \frac{\partial^2}{\partial t^2} \psi = \left( -c^2 \hbar^2 \nabla^2 + m^2 c^4 \right) \psi \quad (1.6)$$

This new approach was, however, still faulty, due to only describing spin 0 particles (e.g., some mesons), and making use of a second order derivative in the time-like component.

That way, a new equation was developed by Paul Dirac, in 1928 [10], one taking into account not the classical 3 dimensional space components, but the relativistic four components.

Dirac took a spin at rewriting the energy-momentum relation, ending up with an equivalent equation (3.7), involving  $4 \times 4$  matrices, due to the 4 relativistic dimensions at play, and incorporating spins into the equation by making use of the now famous Pauli matrices (3.9).

$$E = c \boldsymbol{\alpha} \cdot \mathbf{p} + \beta m c^2, \quad \boldsymbol{\alpha} = (\alpha_1, \alpha_2, \alpha_3) \quad (1.7)$$



$$\alpha_i = \begin{pmatrix} 0 & \sigma_i \\ \sigma_i & 0 \end{pmatrix} \quad \beta = \begin{pmatrix} I_2 & 0 \\ 0 & -I_2 \end{pmatrix} \quad I_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (1.8)$$

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (1.9)$$

In order to fully comprehend this shift of notation, one should equate the square of the two equations, (3.4) and (3.7), and confirm if logic still stands.

$$c^2 \mathbf{p}^2 + m^2 c^4 = c^2 \boldsymbol{\alpha}^2 \mathbf{p}^2 + 2mc^3 \boldsymbol{\alpha} \cdot \mathbf{p} + \beta^2 m^2 c^4 \quad (1.10)$$

In order for this equation to make sense, the following conditions must be true (which in fact, they are):

$$\begin{cases} c^2 \mathbf{p}^2 = c^2 \boldsymbol{\alpha}^2 \mathbf{p}^2 & \Leftrightarrow \boldsymbol{\alpha}^2 = 1 \\ 0 = 2mc^3 \boldsymbol{\alpha} \cdot \mathbf{p} & \Leftrightarrow \boldsymbol{\alpha} \cdot \mathbf{p} = 0 \\ m^2 c^4 = \beta^2 m^2 c^4 & \Leftrightarrow \beta^2 = 1 \end{cases} \quad (1.11)$$

With all the previous considerations taken into account, one can now construct Dirac's free-particle equation(3.12):

$$i\hbar \frac{\partial}{\partial t} \boldsymbol{\psi} = (c \boldsymbol{\alpha} \cdot \mathbf{p} + \beta mc^2) \boldsymbol{\psi} = \begin{pmatrix} mc^2 I_2 & -i\hbar c \boldsymbol{\sigma} \cdot \nabla \\ -i\hbar c \boldsymbol{\sigma} \cdot \nabla & -mc^2 I_2 \end{pmatrix} \cdot \begin{pmatrix} \psi_1 \\ \vdots \\ \psi_4 \end{pmatrix} \quad (1.12)$$

This equation, however, as mentioned above, can only describe a single particle present in a field-free region. In order to account for the existence of a field, such as the electromagnetic field, derived from the four-potential  $A^\mu$ , composed by the electric scalar potential field,  $A^0 = \phi$ , and the vector potential,  $(A^1, A^2, A^3) = \mathbf{A}$ , the following change to the momentum four-vector must be made:

$$p^\mu \rightarrow p^\mu - eA^\mu, \quad A^\mu = (\phi, \mathbf{A}) \quad (1.13)$$

The Hamiltonian can now be rewritten to account for the presence of the electromagnetic field (3.14). This way it is possible to include, for example, the electron-nucleus Coulomb attraction.

$$H_D = -e\phi + \beta mc^2 + \boldsymbol{\alpha}(c\mathbf{p} + e\mathbf{A}) \quad (1.14)$$

For a central potential, as is the one generated by the nuclear charge (assuming Born-Oppenheimer's approximation), the 3 space-like components from the four-potential are null, and the time-like component,  $\phi = \frac{Ze}{r}$ . The Hamiltonian gains now a more recognizable form:

$$H_D = -\frac{e^2 Z}{r} + \beta mc^2 + \boldsymbol{\alpha} \cdot \mathbf{p} c \quad (1.15)$$

Something very interesting about Dirac's equation is the fact that it does not yield a single solution, but in fact, two: the large component (positive energy values), for particles, and the small component (negative energy values), for antiparticles.

### 1.2.3 The Dirac-Breit Equation

Once again, when considering a system composed of many bodies, one must consider all the present interactions, namely, the electron-electron repulsion in an atom. Breit, in 1929, had created a relativistic approach to treat the electron-electron interactions, consisting on a set of equations building upon the classical non-relativistic Hamiltonian from equation (3.1), which can be consulted in Appendix A. Breit's equations are able to account for angular momenta couplings and estimate level energy splittings, the change of a particle's apparent mass as a function of velocity, and even include the interaction of an applied external magnetic field [9].

It is quite obvious Breit's equations introduce a great complexity in the search of the new Hamiltonian's eigenfunctions. However, when trying to include an approximation of Breit's considerations into Dirac's equation, one must add the following operator to the one present in equation (3.15):

$$H_B = \sum_{i>j} \frac{e^2}{r_{ij}} - e^2 \left( \frac{\boldsymbol{\alpha}_i \boldsymbol{\alpha}_j}{r_{ij}} + \frac{(\boldsymbol{\alpha}_i \nabla_i)(\boldsymbol{\alpha}_j \nabla_j) r_{ij}}{2} \right) \quad (1.16)$$

This set of terms will account for the fact that Coulomb interactions, mediated by the electric field, and therefore, **virtual photons**, cannot occur at instantaneous velocities, but at the speed of light.

### 1.2.4 The **MCDF** Method

As previously mentioned in section 3.1.1, there is a need for a numerical method in order to compute and find the eigenfunctions for a many-body Hamiltonian. While the Hartree-Fock method was able to reasonably solve the non-relativistic problem, now, while considering the Dirac-Breit Hamiltonian from equations (3.15) and (3.16), there is a need for a new method.

Hence, the state of the art **MCDFGME** arises. This self-consistent iterative method, based on the same method present in section 3.1.1, is able to solve and find eigenfunctions for a multielectronic system, now taking into account the Dirac-Breit Hamiltonian. Moreover, it is also capable of incorporating electron correlation and many QED effects not yet considered in the relativistic equation, such as the Lamb-shift, vacuum polarization, and the electron's self energy. A brief description of these contributions can be found in appendix C

## 1.3 State of the Art

### 1.3.1 Copper's characteristic X-rays

Copper is a dominant element in today's society. While most of its uses are day-to-day related, it also has a high prevalence in many physical areas, namely, copper's  $K_\alpha$  transitions [11]. While these radiative transitions have been measured countless times, with very well recorded energy values [12–16]. It is also common knowledge Copper's  $K_\alpha$  lines do not have a symmetric distribution, since both  $K_{\alpha_1}$  and  $K_{\alpha_2}$  line display a negative (left-tailed) skewness. Due to this fact, most of the fitting models used in order to analyze both Copper's  $K_{\alpha_1}$  and  $K_{\alpha_2}$  transitions involve the usage of a Lorentzian doublet, for each line [17, 18]. While many associate these asymmetries due to satellite states formed by shake processes [11, 19], with some theoretical studies having been performed [20], some authors note it could be due to X-ray resonant Raman scattering [19]. This effect occurs when a sample is exposed to energies under to near the ionization threshold, when a bound electron is excited to an upper state [21]. While there are some studies exploring this topic, most are focused on the cascade of low energy transitions that follow the post-scattering excitation [22]. Nonetheless, a previous experimental study has been able to show that for Copper exposed to synchrotron radiation tuned to energies near K-shell's ionization energy, some  $K_{\alpha_1}$  transitions demonstrated to be narrower than expected [23].

### 1.3.2 MCDFGME capabilities

It has been noted multiple times that *MCDFGME* code excels in atomic structure calculations of super-heavy elements and highly-charged ions, where relativistic and QED effects are in prevalence [24–26]. However, it has also been proven to be an excellent tool for the calculation of less ionized and lighter atomic systems [27].

In addition, the *MCDFGME* code is able to calculate radiative and auger transition rates for the calculated configurations, which can be used in the simulation of theoretical spectra, due to being able to compute the transition's intensity and natural width. Since it is able to perform calculations, even for exotic atoms, it can be used to further understand many QED phenomena, further exploring the limits of our theory, and it's comparison to experimental data [28].

It should also be of note that there are many other code alternatives. While *MCDFGME*, which is a close-source project, provides a very high precision in the performed calculations at a high computational cost, *Flexible Atomic Code (FAC)*, is an open source code which requires less computational time for the calculations, however, it lacks *MCDFGME*'s precision, since it only is able to consider all the spin-orb couplings, but does not mix the possible configurations originating them. It can, however, calculate other collisional processes, such as electronic impact excitation cross-sections [29]. *General-purpose Relativistic Atomic Structure Package 2k (Grasp2k)* [30], and *AUTOSTRUCTURE* [31] are other codes with some of the same capabilities.

## 1.4 Methodology

As mentioned before, in this thesis, atomic relaxation transitions where the ejected inner shell electron, responsible for the generated hole, was not sent to the continuum, but excited to an upper state.

In order to use *MCDFGME*, since it is written in FORTRAN, there will be a multitude of input/output files, with the extension *.f0X*. The most important file extensions are *.f05*, the input file, where the calculation settings are indicated; *.f06*, where the energy results will be displayed, among others, such as some matrix element overlaps, both important to evaluate the convergence. Other file extensions, such as *.f09*, also store the final electrons' wavefunctions and can be quite useful for debugging processes, and for calculating other atomic parameters, such as shake probabilities, which will not be of relevance as of this thesis.

An example of a normal atomic structure calculation for an excited Copper atom (which has a ground state configuration of  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^1$ ) is as follows:

One starts by choosing the excitation upper orbital, and by writing in a text file all the possible 1 hole configurations from where the electron may have originated from. An example can be found in Annex I. This represents all the possible excited states where the electron is in the chosen orbital.

The same process is now performed for a 2 hole configuration, where an additional ionization occurred. This represents the possible states after the occurrence of the Auger process, which is in competition with the radiative decay.

Now, a *bash* script, *runMCDFEC.sh* is to be executed. Now, it will be possible to choose the atomic number of the element in study, the isotope mass, and, if one wishes so, to perform the calculations for an exotic atom, where, instead of electrons, other particles such as muons, negative hadrons and antiparticles can orbit the nucleus. The script will calculate all the possible spin-orbit couplings and eigenvalues, and then finally apply the *MCDFGME* method.

Depending on the excitation orbital, there will be thousands to tens of thousands of possibilities for the final configuration, with this number increasing with the quantum number of angular momentum,  $l$ , of the orbital.

However, not all the calculations performed will automatically converge. Each time, a few hundreds of computed states will fail convergence. In a file, generated for the compilation of the calculation results, these states will be missing the total system's energy, or will display a significant energy difference for the results obtained by using two different methods, summing and integrating.

The configurations that were not able to converge will now have to be converged by hand. In order to do this, their *.f05* will have to be edited, where different methods will be chosen for the calculation for each specific orbital. As there are multiple possible methods and combinations, this step can take up to a few weeks to complete.

After eliminating all the convergence problems, the radiative transition rates will

be calculated, by operating the transition operator on all the possible 1 hole - 1 hole configuration pair combinations. The same will follow for all the pairs of 1 hole - 2 holes configurations, in order to compute the auger transition rates. Finally, the process is repeated for the 2 holes - 2 holes configurations, for the radiative transitions of the auger-generated satellite states.

Lastly, all the calculated transitions' widths will be summed in order to calculate each subshell's fluorescence yield (calculated by the sum of all radiative transition widths divided by the sum of all the radiative and non-radiative transition widths).

---

## Todo list

|   |    |
|---|----|
| Figure: Maybe put a Thomas model plot? . . . . .  | 17 |
| Explain antisymmetry and put it in the glossary . . . . .   | 20 |
| Explain in the glossary . . . . .   | 20 |
| Finish the explanation . . . . .  | 21 |
| eheheheh . . . . .  | 22 |
| spin? . . . . .   | 22 |
| Ler bem o manual e explicar bem o método e todas as coisas diferentes que se podem<br>fazer . . . . . | 24 |
| This really needs a re-do . . . . .   | 24 |
| Take this out . . . . .   | 25 |

---

# Theoretical Introduction

Throughout this thesis, different topics regarding Quantum Mechanics, Atomic Physics and spectroscopy will be approached, hence why a theoretical introduction is needed. In it, themes such as the characteristic X-rays of elements, the relativistic Dirac equation and QED corrections and methods for solving the many-body problem will be discussed.

## 2.1 Characteristic X-rays

When subjected to a high energy beam of particles (usually photons or electrons), an element may go through the process of the ionization of inner-core electrons, which is followed by the emission of radiation of its own, in the form of X-rays and/or Auger electrons. The radiation emitted throughout this process presents well-defined energy values and thus provides a way to ascertain if an element is present in a sample under study. Due to this reason, these characteristic emissions are of an extremely high importance in a wide range of scientific areas.

When an atomic system is in a bound state, the electrons orbiting the nucleus are occupying fixed quantum states, defined by their principal atomic number,  $n$ , angular momentum,  $l$ , and spin,  $s$ . Electrons are also fermions, and thus must respect Pauli's exclusion principle, each occupying a single state, only occupied by that single electron. These quantum numbers provide information about each electron's wavefunction, and the collection of all the electrons' wavefunctions can be used in order to describe the whole system. When operating the Hamiltonian on this collection of wavefunctions, the atomic system's state's energy is obtained. Besides all the occupied states, an infinite more number of possible eigenfunctions will compose that system's base.

### 2.1.1 Ionization as a vacancy generator

As previously mentioned, X-ray fluorescence spectroscopy has many uses and applications in a wide range of scientific areas. In this form of spectroscopy, the element at study, composed of a nucleus and  $N$  orbiting electrons, is bombarded with radiation leading to the ionization of inner-shell electrons while leaving a vacancy in their place (Figure 2.1).




Figure 2.1: Photoionization.

The atomic structure, now composed of  $N - 1$  electrons, will be left in an energetically unstable state, due to there being other possible lower energy states. This will lead to various processes of atomic relaxation, where the system will rearrange itself in order to find a lower energy configuration.

The main processes for this rearrangement are two competing decay paths: radiative relaxation, and Auger electron emission.




Figure 2.2: Principal atomic relaxation processes.


In the case of radiative relaxation, an upper shell electron will move and occupy the hole left in the inner shell. During this process, energy is released through the emission of photons with energy in the range of X-rays. The detection of these photons is what allows for an analysis and detection of the element at study.

On the other hand, an Auger emission occurs when in the process of an upper electron shifting to a shell hole, energy is released not in the form of photons, but by the atomic system going through the process of the ionization of a lesser bound electron.

In reality, when the initial vacancy is generated, two more processes can occur: shake-off and shake-up. Due to a sudden change in the potential felt by the remaining electrons,



whilst going through the ionization, one other electron may become unbound, leading to a second ionization, the shake-off process, or may be excited to an upper state, during shake-up. These processes' occurrence probabilities are related to the rate at which the electron leaves the system during the first ionization. For lower electron emission energies, where the ionization process is rather "slow", and there is not much energy for shake up/off to occur, the adiabatic regime reigns. In this regime, the shake probabilities are small, and increase with the beam's energy. For higher energies, the transition occurs in the sudden regime, where the electrons' exit can be considered instantaneous. Now, while the shake probabilities still increase with the energy of the beam, they quickly saturate for a maximum value of probability, which can be calculated by computing overlaps between the electrons' initial and final states. These processes, however, are out of scope for this thesis, but should the reader be interested, the model conceived by Thomas [32] predicts the shake probability as a function of energy.



### 2.1.2 Transition notation

The characteristic radiation measured from the radiative relaxation of a post-ionization unstable atomic system is one of the main ways of identifying an atomic element. This is due to the photons emitted possessing quantized values of energy, forming well-defined energy lines when observed in a spectrometer. In order to understand the change that occurred in the atomic system which lead to a specific emission, the spectral lines get labels based on a notation which usually takes into account the initial and final orbitals where the vacancy was present. A very illustrative diagram, exemplifying some transitions can be found in Appendix B.

Throughout this thesis, Siegbahn notation will be used, for the most part, but should the reader prefer IUPAC's, Table 2.1 has the conversion between notations.

### 2.1.3 Excitation as a vacancy generator

As previously mentioned, throughout this thesis, while the study is focused on the characteristic radiation emitted during an atomic relaxation process, the main vacancy generation method at study shall be the photoexcitation process (Figure 2.3), instead of ionization. The levels obtained after a photoexcitation of core shell electrons has occurred could be some of the many so-called satellite states, where the electronic configuration present during the relaxation process contains additional electrons or holes, or is simply not

Table 2.1: Siegbahn VS IUPAC notation. Adapted from [1].

| Siegbahn           | IUPAC     | Siegbahn       | IUPAC           | Siegbahn        | IUPAC           |
|--------------------|-----------|----------------|-----------------|-----------------|-----------------|
| $K_{\alpha_1}$     | $K - L_3$ | $L_{\alpha_1}$ | $L_3 - M_5$     | $L_{\gamma_1}$  | $L_2 - N_4$     |
| $K_{\alpha_2}$     | $K - L_2$ | $L_{\alpha_2}$ | $L_3 - M_4$     | $L_{\gamma_2}$  | $L_1 - N_1$     |
| $K_{\beta_1}$      | $K - M_3$ | $L_{\beta_1}$  | $L_2 - M_4$     | $L_{\gamma_3}$  | $L_1 - N_2$     |
| $K_{\beta_2}^I$    | $K - N_3$ | $L_{\beta_2}$  | $L_3 - N_5$     | $L_{\gamma_4}$  | $L_1 - O_3$     |
| $K_{\beta_2}^{II}$ | $K - N_2$ | $L_{\beta_3}$  | $L_1 - M_3$     | $L_{\gamma_4}'$ | $L_1 - O_2$     |
| $K_{\beta_3}$      | $K - M_2$ | $L_{\beta_4}$  | $L_1 - M_2$     | $L_{\gamma_5}$  | $L_2 - N_1$     |
| $K_{\beta_4}^I$    | $K - N_5$ | $L_{\beta_5}$  | $L_3 - O_{4,5}$ | $L_{\gamma_5}$  | $L_2 - O_4$     |
| $K_{\beta_4}^{II}$ | $K - N_4$ | $L_{\beta_6}$  | $L_3 - N_1$     | $L_{\gamma_8}$  | $L_2 - O_1$     |
| $K_{\beta_4}^x$    | $K - N_4$ | $L_{\beta_7}$  | $L_3 - O_1$     | $L_{\gamma_8}'$ | $L_2 - N_{5,6}$ |
| $K_{\beta_5}^I$    | $K - M_5$ | $L_{\beta_8}$  | $L_3 - N_{6,7}$ | $L_{\eta}$      | $L_2 - M_1$     |
| $K_{\beta_4}^{II}$ | $K - M_4$ | $L_{\beta_9}$  | $L_1 - M_5$     | $L_l$           | $L_3 - M_1$     |
|                    |           | $L_{\beta_9}$  | $L_1 - M_4$     | $L_s$           | $L_3 - M_3$     |
|                    |           | $L_{\beta_9}$  | $L_3 - N_4$     | $L_t$           | $L_3 - M_2$     |
|                    |           | $L_{\beta_9}$  | $L_2 - M_3$     | $L_u$           | $L_3 - N_{6,7}$ |
|                    |           |                |                 | $L_v$           | $L_2 - N_{6,7}$ |

the standard configuration when talking about characteristic emissions. The characteristic radiation from transitions that originated from these states are one of the keys needed to fully comprehend and deconvolute an element's emission spectra.




Figure 2.3: Resonant Photoexcitation

## 2.2 Radiative transitions

In terms of quantum mechanics, the properties of a system's change of state can be derived using perturbation theory, where a change in energy can be treated as perturbation, and takes into physical properties ~~angular momentum~~ conservation laws as basis for the selection rules.

The transition's radiation type can be of two main flavors, Electric,  $E_k$ , or Magnetic,  $M_k$ ,

with  $k$  representing its multiplicity. These labels indicate which of the [Electromagnetic Field \(EMF\)](#) component had the strongest influence on the transition's occurrence. For both these types of transitions, the system's total angular momentum is allowed a change of  $\Delta J = 0, \pm k$ , and so does its projection,  $M_J$ . However, different transitions lead to different changes of parity in the system and to different selection rules.

It should also be of note that, usually, for the same multiplicity, an  $E$  transition is more intense than an  $M$ , and that it is also possible for rare 2-photon transitions to take place, where a combination of different transition types can occur.

The intensity of a transition is proportional to the squared norm of the perturbation's matrix element involving the initial and final state ( $\Gamma_{if} \propto |\langle \psi_i | H' | \psi_f \rangle|^2$ ). It should also be noted that the electron's initial state population will serve as a scaling factor for the transitions' rate. For example, while the transition rate for a  $2p_{1/2} \rightarrow 1s$  and  $2p_{3/2} \rightarrow 1s$  should be about the same, the orbital  $2p_{3/2}$  has double the population  $2p_{1/2}$  has, hence why the  $K_{\alpha_2}$  line has about half the  $K_{\alpha_1}$  line's intensity.

One should also mention that monochromatic transitions do not exist. While the transition might have a well-defined energy, calculated by the difference in energy between the initial and final levels, due to Heisenberg's uncertainty principle,  $\Delta E \Delta t > \frac{\hbar}{2}$ , there will exist a natural energy broadening, the transition's natural width. The shapes representing these transitions are given by a Lorentzian distribution.

## 2.3 Solving the atomic many-body problem

When studying a system composed of multiple charged bodies, one must consider all the existing interactions. Whilst there are known analytical solutions for 2-bodies Hydrogenoid systems, with the presence of more non spatially-bound particles, the Coulomb interaction pairs lead to the impossibility of finding a set of analytical wavefunctions which are a part of the Hamiltonian's eigenset. Consecutively, the need for a numerical method that is able to compute solutions for these complex systems arose.

### 2.3.1 The non-relativistic Hamiltonian

The first approach used in order to solve the many-bodies problem used a non-relativistic consideration. This way, the Hamiltonian consists on the sum of the system's non-relativistic momentum-related energies and the energy of Coulomb interactions between all the bodies in the system, while considering the nucleus as fixed in space, due to it being thousands of times more massive than the orbiting electrons.

Essentially, and in atomic units:

$$\underbrace{\sum_i^N \left( \overbrace{\frac{1}{2} \nabla_i^2}^{E_1} - \overbrace{\frac{Z}{r_i}}^{E_2} \right)}_{\text{Individual Hamiltonian}} + \underbrace{\sum_{i < j}^j \frac{1}{r_{ij}}}_{\text{Pair repulsion}} \overbrace{\quad}^{E_3}$$

$$E_1 \rightarrow \text{Momentum} \quad E_2 \rightarrow e^- \text{ nuc. Coulomb attraction} \quad E_3 \rightarrow e^- e^- \text{ Coulomb repulsion} \quad (2.1)$$

### 2.3.2 The Hartree-Fock Method

This numerical method is one of the staple and most enduring procedures for solving the problem associated with a many-body system. As a side note, an in-depth explanation on all the intricacies of this method can be found in many of the literature, but as for the writing of this thesis, the works [2–5] were the ones consulted.

Hartree developed an iterative method, further enhanced by Fock and Slater, based on the field's [Self Consistency Method](#). In this approach, when studying a multi-electronic system, such as an atom, each electron's wavefunction is composed as a product of a spacial part,  $\psi$ , and one related to the electron's spin,  $\chi$ , as to be able to account for Pauli's exclusion principle and, if so desired, relativistic effects.

$$u = \psi \chi \quad (2.2)$$

The wavefunction capable of describing the whole system,  $\Psi$ , should be somewhat of a product of all the wavefunctions describing each individual electron. However, one must not forget the need for this wavefunction to respect the antisymmetry principle, due to the electron's fermionic nature. In order to achieve this,  $\Psi$  is to be composed of a Slater determinant:

$$\Psi = \frac{1}{\sqrt{N!}} \begin{vmatrix} u_1(x_1) & u_2(x_1) & \cdots & u_N(x_1) \\ u_1(x_2) & u_2(x_2) & \cdots & u_N(x_2) \\ \vdots & \vdots & \ddots & \vdots \\ u_1(x_N) & u_2(x_N) & \cdots & u_N(x_N) \end{vmatrix} \quad (2.3)$$

It is of high importance that the wavefunction basis set respects orthonormality. These are to be initialized as trial wavefunctions for the numerical method.

The main goal for this algorithm is to, as per the [Self Consistency Method](#), follow the variational principle and to minimize a functional, as a way of minimizing the system's energy. This optimal, yet unknown energy,  $E_0$  (calculated by operating the Hamiltonian on the system's optimal wavefunctions), will attempt to be reached by the variation of trial wavefunctions that provide a non-minimized solution. Whilst the optimal wavefunctions will never be reached, the purpose of this method is to reach a solution that yields an

Explain anti-symmetry and put it in the glossary

Explain in the glossary

energetic value as close to  $E_0$  as possible, while always yielding  $\langle \Psi | H | \Psi \rangle \geq E_0$ , should the calculation be correct.

The computational method consists on starting with the previously mentioned trial wavefunctions, using them to calculate Hartree-Fock's potential through the HF equations.

In a very simplified way, the self-consistent Hartree-Fock computational method can be represented by the block diagram in Figure 3.1.

Finish the explanation




Figure 2.4: HF method's block diagram.

### 2.3.3 The Dirac Equation

*Note: This section used the works in [6–9] as reference.*

It is no secret that Schrödinger's equation has some very considerable limitations. The fact that it does not account for the existence of the electron's spin and the lack of consideration of relativistic effects are some of the most impactful setbacks.

Many scientists, such as Klein, Gordon and later Fock, had already conceived a relativistic correction to Schrödinger's equation, where the free-particle energy makes use of the relativistic momentum-energy relation (3.4).

$$E = \sqrt{c^2 \mathbf{p}^2 + m^2 c^4} \quad (2.4)$$

Which can be derived from the [Lorentz invariant Minkowski norm](#) of the momentum four-vector (3.5).

$$p^\mu p_\mu = m^2 c^2 \Leftrightarrow \frac{E^2}{c^2} - \mathbf{p}^2 = m^2 c^2 \Leftrightarrow \frac{E^2}{c^2} = \mathbf{p}^2 + m^2 c^2 \quad (2.5)$$

Now, inputting this new energy operator into Schrödinger's equation, yields the Klein-Gordon equation (3.6), allowing for Schrödinger's equation to now be Lorentz-invariant.

$$-\hbar^2 \frac{\partial^2}{\partial t^2} \psi = (-c^2 \hbar^2 \nabla^2 + m^2 c^4) \psi \quad (2.6)$$

This new approach was, however, still faulty, due to only describing spin 0 particles (e.g., some mesons), and making use of a second order derivative in the time-like component.

That way, a new equation was developed by Paul Dirac, in 1928 [10], one taking now into account not the classical 3 dimensional space components, but the relativistic four components.

eheheheh — Dirac took a spin at rewriting the energy-momentum relation, ending up with an equivalent equation (3.7), involving  $4 \times 4$  matrices, due to the 4 relativistic dimensions at play, and incorporating spins into the equation by making use of the now famous Pauli matrices (3.9).

$$E = c \boldsymbol{\alpha} \cdot \mathbf{p} + \beta m c^2, \quad \boldsymbol{\alpha} = (\alpha_1, \alpha_2, \alpha_3) \quad (2.7)$$

$$\alpha_i = \begin{pmatrix} 0 & \sigma_i \\ \sigma_i & 0 \end{pmatrix} \quad \beta = \begin{pmatrix} I_2 & 0 \\ 0 & -I_2 \end{pmatrix} \quad I_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (2.8)$$

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (2.9)$$

In order to fully comprehend this shift of notation, one should equate the square of the two equations, (3.4) and (3.7), and confirm if logic still stands.

$$c^2 \mathbf{p}^2 + m^2 c^4 = c^2 \boldsymbol{\alpha}^2 \mathbf{p}^2 + 2 m c^3 \boldsymbol{\alpha} \cdot \mathbf{p} + \beta^2 m^2 c^4 \quad (2.10)$$

In order for this equation to make sense, the following conditions must be true (which in fact, they are):

$$\begin{cases} c^2 \mathbf{p}^2 = c^2 \boldsymbol{\alpha}^2 \mathbf{p}^2 & \Leftrightarrow \boldsymbol{\alpha}^2 = 1 \\ 0 = 2 m c^3 \boldsymbol{\alpha} \cdot \mathbf{p} & \Leftrightarrow \boldsymbol{\alpha} \beta = 0 \\ m^2 c^4 = \beta^2 m^2 c^4 & \Leftrightarrow \beta^2 = 1 \end{cases} \quad (2.11)$$

With all the previous considerations taken into account, one can now construct Dirac's free-particle equation(3.12):

$$i\hbar \frac{\partial}{\partial t} \psi = (c \boldsymbol{\alpha} \cdot \mathbf{p} + \beta mc^2) \psi = \begin{pmatrix} mc^2 I_2 & -i\hbar c \boldsymbol{\sigma} \cdot \nabla \\ -i\hbar c \boldsymbol{\sigma} \cdot \nabla & -mc^2 I_2 \end{pmatrix} \cdot \begin{pmatrix} \psi_1 \\ \vdots \\ \psi_4 \end{pmatrix} \quad (2.12)$$

This equation, however, as mentioned above, can only describe a single particle present in a field-free region. In order to account for the existence of a field, such as the electromagnetic field, derived from the four-potential  $A^\mu$ , composed by the electric scalar potential field,  $A^0 = \phi$ , and the vector potential,  $(A^1, A^2, A^3) = \mathbf{A}$ , the following change to the momentum four-vector must be made:

$$p^\mu \rightarrow p^\mu - eA^\mu, \quad A^\mu = (\phi, \mathbf{A}) \quad (2.13)$$

The Hamiltonian can now be rewritten as to account for the presence of the electromagnetic field (3.14). This way it is possible to include, for example, the electron-nucleus Coulomb attraction.

$$H_D = -e\phi + \beta mc^2 + \boldsymbol{\alpha}(c\mathbf{p} + e\mathbf{A}) \quad (2.14)$$

For a central fixed potential, as is the one generated by the nuclear charge (assuming Born-Oppenheimer's approximation), the 3 space-like components from the four-potential are null, and the time-like component,  $\phi = \frac{Ze}{r}$ . The Hamiltonian gains now a more recognizable form:

$$H_D = -\frac{e^2 Z}{r} + \beta mc^2 + \boldsymbol{\alpha} \cdot \mathbf{p} c \quad (2.15)$$

Something very interesting about Dirac's equation is the fact that it does not yield a single solution, but in fact, two: the large component (positive energy values), for particles, and the small component (negative energy values), for antiparticles.

### 2.3.3.1 The Dirac-Breit Equation

Once again, when considering a system composed of many bodies, one must consider all the present interactions, namely, the electron-electron repulsion in an atom. Breit, in 1929, had created a relativistic approach to treat the electron-electron interactions, consisting on a set of equations building upon the classical non-relativistic Hamiltonian from equation (3.1), which can be consulted in Appendix A. Breit's equations are able to account for angular momenta couplings and estimate level energy splittings, the change of a particle's apparent mass as a function of velocity, and even include the interaction of an applied external magnetic field [9].

It is quite obvious Breit's equations introduce a great complexity in the search of the new Hamiltonian's eigenfunctions. Nonetheless, when trying to include an approximation

of Breit's considerations into Dirac's equation, one must add the following operator to the one present in equation (3.15):

$$H_B = \sum_{i>j} \frac{e^2}{r_{ij}} - e^2 \left( \frac{\boldsymbol{\alpha}_i \boldsymbol{\alpha}_j}{r_{ij}} + \frac{(\boldsymbol{\alpha}_i \nabla_i)(\boldsymbol{\alpha}_j \nabla_j) r_{ij}}{2} \right) \quad (2.16)$$

This set of terms will account for the fact that Coulomb interactions, mediated by the electric field, and therefore, **virtual photons**, cannot occur at instantaneous velocities, but at the speed of light.

### 2.3.4 The *MCDF* Method

As previously mentioned in section 3.1.1, there is a need for a numerical method in order to compute and find the eigenfunctions for a many-body Hamiltonian. While the Hartree-Fock method was able to reasonably solve the non-relativistic problem, now, while considering the Dirac-Breit Hamiltonian from equations (3.15) and (3.16), there is a need for a new method.

Hence, the state of the art *MCDFGME* arises. This self-consistent iterative method, based on the same method present in section 3.1.1, is able to solve and find eigenfunctions for a multielectronic system, now taking into account the Dirac-Breit Hamiltonian. Moreover, it is also capable of incorporating electron correlation and many QED effects not yet considered in the relativistic equation, such as the Lamb-shift, vacuum polarization, and the electron's self energy. A brief description of these contributions can be found in appendix C

## 2.4 State of the Art

### 2.4.1 Copper's characteristic X-rays

Copper is a dominant element in today's society. While most of its uses are day-to-day related, it also has a high prevalence in many physical areas, namely, copper's  $K_\alpha$  transitions [11]. While these radiative transitions have been measured countless times, with very well recorded energy values [12–16]. It is also common knowledge Copper's  $K_\alpha$  lines do not have a symmetric distribution, since both  $K_{\alpha_1}$  and  $K_{\alpha_2}$  line display a negative (left-tailed) skewness. Due to this fact, most of the fitting models used in order to analyze both Copper's  $K_{\alpha_1}$  and  $K_{\alpha_2}$  transitions involve the usage of a Lorentzian doublet, for each line [17, 18]. While many associate these asymmetries due to satellite states formed by shake processes [11, 19], with some theoretical studies having been performed [20], some authors note it could be due to X-ray resonant Raman scattering [19]. This effect occurs when a sample is exposed to energies under to near the ionization threshold, when a bound electron is excited to an upper state [21]. While there are some studies exploring this topic, most are focused on the cascade of low energy transitions that follow the

Ler bem o manual e explicar bem o método e todas as coisas diferentes que se podem fazer

This really needs a re-do



post-scattering excitation [22]. Nonetheless, a previous experimental study has been able to show that for Copper exposed to synchrotron radiation tuned to energies near K-shell's ionization energy, some  $K_{\alpha 1}$  transitions demonstrated to be narrower than expected [23].

### 2.4.2 MCDFGME capabilities

It has been noted multiple times that *MCDFGME* code excels in atomic structure calculations of super-heavy elements and highly-charged ions, where relativistic and QED effects are in prevalence [24–26]. However, it has also been proven to be an excellent tool for the calculation of less ionized and lighter atomic systems [27].

In addition, the *MCDFGME* code is able to calculate radiative and auger transition rates for the calculated configurations, which can be used in the simulation of theoretical spectra, due to being able to compute the transition's intensity and natural width. Since it is able to perform calculations, even for exotic atoms, it can be used to further understand many QED phenomena, further exploring the limits of our theory, and it's comparison to experimental data [28].

It should also be of note that there are many other code alternatives. While *MCD-FGME*, which is a close-source project, provides a very high precision in the performed calculations at a high computational cost, *FAC*, is an open source code which requires less computational time for the calculations, however, it lacks *MCDFGME*'s precision, since it only is able to consider all the spin-orb couplings, but does not mix the possible configurations originating them. It can, however, calculate other collisional processes, such as electronic impact excitation cross-sections [29]. *Grasp2k* [30], and *AUTOSTRUCTURE* [31] are other codes with some of the same capabilities.

## 2.5 Methodology

Take this out

As mentioned before, in this thesis, atomic relaxation transitions where the ejected inner shell electron, responsible for the generated hole, was not sent to the continuum, but excited to an upper state.

In order to use *MCDFGME*, since it is written in FORTRAN, there will be a multitude of input/output files, with the extension *.f0X*. The most important file extensions are *.f05*, the input file, where the calculation settings are indicated; *.f06*, where the energy results will be displayed, among others, such as some matrix element overlaps, both important to evaluate the convergence. Other file extensions, such as *.f09*, also store the final electrons' wavefunctions and can be quite useful for debugging processes, and for calculating other atomic parameters, such as shake probabilities, which will not be of relevance as of this thesis.

An example of a normal atomic structure calculation for an excited Copper atom (which has a ground state configuration of  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^1$ ) is as follows:

One starts by choosing the excitation upper orbital, and by writing in a text file all the possible 1 hole configurations from where the electron may have originated from. An example can be found in Annex I. This represents all the possible excited states where the electron is in the chosen orbital.

The same process is now performed for a 2 hole configuration, where an additional ionization occurred. This represents the possible states after the occurrence of the Auger process, which is in competition with the radiative decay.

Now, a *bash* script, *runMCDFEC.sh* is to be executed. Now, it will be possible to choose the atomic number of the element in study, the isotope mass, and, if one wishes so, to perform the calculations for an exotic atom, where, instead of electrons, other particles such as muons, negative hadrons and antiparticles can orbit the nucleus. The script will calculate all the possible spin-orbit couplings and eigenvalues, and then finally apply the *MCDFGME* method.

Depending on the excitation orbital, there will be thousands to tens of thousands of possibilities for the final configuration, with this number increasing with the quantum number of angular momentum,  $l$ , of the orbital.

However, not all the calculations performed will automatically converge. Each time, a few hundreds of computed states will fail convergence. In a file, generated for the compilation of the calculation results, these states will be missing the total system's energy, or will display a significant energy difference for the results obtained by using two different methods, summing and integrating.

The configurations that were not able to converge will now have to be converged by hand. In order to do this, their *.f05* will have to be edited, where different methods will be chosen for the calculation for each specific orbital. As there are multiple possible methods and combinations, this step can take up to a few weeks to complete.

After eliminating all the convergence problems, the radiative transition rates will be calculated, by operating the transition operator on all the possible 1 hole - 1 hole configuration pair combinations. The same will follow for all the pairs of 1 hole - 2 holes configurations, in order to compute the auger transition rates. Finally, the process is repeated for the 2 holes - 2 holes configurations, for the radiative transitions of the auger-generated satellite states.

Lastly, all the calculated transitions' widths will be summed in order to calculate each subshell's fluorescence yield (calculated by the sum of all radiative transition widths divided by the sum of all the radiative and non-radiative transition widths).

# Atomic Structure Calculations

When studying a system composed of multiple charged bodies, one must consider all the existing interactions. Whilst there are known analytical solutions for the 2-bodies Hydrogenoid systems, with the presence of more non spatially-bound particles, these become impossible to find for Schrödinger's equation, even using Born-Oppenheimer's approximation, where the nucleus is considered static, at a fixed position, making the electrons the only non spatially-bound particles in the system. That way, there was a need for the development of numerical solutions to solve this problem.

## 3.1 The non-relativistic Hamiltonian

The first approach used in order to solve the many-bodies problem used a non-relativistic consideration. This way, the Hamiltonian consisted on the sum of the system's non-relativistic momentum-related energies and the Coulomb interactions between bodies, while still considering the Born-Oppenheimer approximation.

Essentially, and in atomic units:

$$\underbrace{\sum_i^N \left( \overbrace{\frac{1}{2} \nabla_i^2}^{E_1} - \overbrace{\frac{Z}{r_i}}^{E_2} \right)}_{\text{Individual Hamiltonian}} + \underbrace{\sum_{i < j}^j \overbrace{\frac{1}{r_{ij}}}^{E_3}}_{\text{Pair repulsion}}$$

$$E_1 \rightarrow \text{Momentum} \quad E_2 \rightarrow e^- \text{ nuc. Coulomb attraction} \quad E_3 \rightarrow e^- e^- \text{ Coulomb repulsion} \quad (3.1)$$

### 3.1.1 The Hartree-Fock Method

*Note: This section used the works in [2–5] as reference.*

Hartree developed an iterative method, further enhanced by Fock and Slater, based on the field self-consistency method.

In this method, while studying a multi-electronic system, such as an atom, each electron's wavefunction is composed as a product of a spacial part,  $\psi$ , and one indicating the electron's spin,  $\chi$ , as to be able to account for relativistic effects.

$$u = \psi\chi \quad (3.2)$$

The wavefunction capable of describing the whole system,  $\Psi$ , should be somewhat of a product of all the wavefunctions describing each individual electron. However, one must not forget the need for this wavefunction to respect the antisymmetry principle, due to the electron's fermionic nature. In order to respect this,  $\Psi$  is to be composed of a Slater determinant:

$$\Psi = \frac{1}{\sqrt{N!}} \begin{vmatrix} u_1(x_1) & u_2(x_1) & \cdots & u_N(x_1) \\ u_1(x_2) & u_2(x_2) & \cdots & u_N(x_2) \\ \vdots & \vdots & \ddots & \vdots \\ u_1(x_N) & u_2(x_N) & \cdots & u_N(x_N) \end{vmatrix} \quad (3.3)$$

It is of high importance that the wavefunctions must form an orthonormal basis. These are to be initialized as trial wavefunctions.

The main goal of this method is to follow the variational principle and to minimize a functional, with the purpose of minimizing the system's energy. This optimal, yet unknown energy,  $E_0$  (calculated by Operating the Hamiltonian on the optimal wavefunctions), will try to be reached by varying the trial functions that provide a non-minimized solution, with  $\langle \Psi | H | \Psi \rangle \geq E_0$ .

The computational method consists on starting with the previously mentioned trial wavefunctions, using them to calculate Hartree-Fock's potential through the HF equations.

In a very simplified way, the self-consistent Hartree-Fock computational method can be represented by the block diagram in Figure 3.1.

## 3.2 The Dirac Equation

*Note: This section used the works in [6–9] as reference.*

It is no secret that the Schrödinger equation has some very considerable limitations. The fact that it does not account for the existence of the electron's spin and the lack of consideration of relativistic effects are some of the most impactful setbacks.

Many scientists, such as Klein, Gordon and later Fock, had already conceived a relativistic correction to Schrödinger's equation, where the free-particle energy makes use of the relativistic momentum-energy relation (3.4).

$$E = \sqrt{c^2 p^2 + m^2 c^4} \quad (3.4)$$

Which can be derived from the [Lorentz invariant Minkowski norm](#) of the momentum four-vector (3.5).




Figure 3.1: HF method's block diagram.

$$p^\mu p_\mu = m^2 c^2 \Leftrightarrow \frac{E^2}{c^2} - \mathbf{p}^2 = m^2 c^2 \Leftrightarrow \frac{E^2}{c^2} = \mathbf{p}^2 + m^2 c^2 \quad (3.5)$$

Now, inputting this new energy operator into Schrödinger's equation, yields the Klein-Gordon equation (3.6), allowing for Schrödinger's equation now to be Lorentz-invariant.

$$-\hbar^2 \frac{\partial^2}{\partial t^2} \psi = \left( -c^2 \hbar^2 \nabla^2 + m^2 c^4 \right) \psi \quad (3.6)$$

This new approach was, however, still faulty, due to only describing spin 0 particles (e.g., some mesons), and making use of a second order derivative in the time-like component.

That way, a new equation was developed by Paul Dirac, in 1928 [10], one taking into account not the classical 3 dimensional space components, but the relativistic four components.

Dirac took a spin at rewriting the energy-momentum relation, ending up with an equivalent equation (3.7), involving  $4 \times 4$  matrices, due to the 4 relativistic dimensions at play, and incorporating spins into the equation by making use of the now famous Pauli matrices (3.9).

$$E = c\boldsymbol{\alpha} \cdot \mathbf{p} + \beta mc^2, \quad \boldsymbol{\alpha} = (\alpha_1, \alpha_2, \alpha_3) \quad (3.7)$$

$$\alpha_i = \begin{pmatrix} 0 & \sigma_i \\ \sigma_i & 0 \end{pmatrix} \quad \beta = \begin{pmatrix} I_2 & 0 \\ 0 & -I_2 \end{pmatrix} \quad I_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (3.8)$$

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (3.9)$$

In order to fully comprehend this shift of notation, one should equate the square of the two equations, (3.4) and (3.7), and confirm if logic still stands.

$$c^2 \mathbf{p}^2 + m^2 c^4 = c^2 \boldsymbol{\alpha}^2 \mathbf{p}^2 + 2mc^3 \boldsymbol{\alpha} \cdot \mathbf{p} \beta + \beta^2 m^2 c^4 \quad (3.10)$$

In order for this equation to make sense, the following conditions must be true (which in fact, they are):

$$\begin{cases} c^2 \mathbf{p}^2 = c^2 \boldsymbol{\alpha}^2 \mathbf{p}^2 & \Leftrightarrow \boldsymbol{\alpha}^2 = 1 \\ 0 = 2mc^3 \boldsymbol{\alpha} \beta & \Leftrightarrow \boldsymbol{\alpha} \beta = 0 \\ m^2 c^4 = \beta^2 m^2 c^4 & \Leftrightarrow \beta^2 = 1 \end{cases} \quad (3.11)$$

With all the previous considerations taken into account, one can now construct Dirac's free-particle equation(3.12):

$$i\hbar \frac{\partial}{\partial t} \boldsymbol{\psi} = (c\boldsymbol{\alpha} \cdot \mathbf{p} + \beta mc^2) \boldsymbol{\psi} = \begin{pmatrix} mc^2 I_2 & -i\hbar c \boldsymbol{\sigma} \cdot \nabla \\ -i\hbar c \boldsymbol{\sigma} \cdot \nabla & -mc^2 I_2 \end{pmatrix} \cdot \begin{pmatrix} \psi_1 \\ \vdots \\ \psi_4 \end{pmatrix} \quad (3.12)$$

This equation, however, as mentioned above, can only describe a single particle present in a field-free region. In order to account for the existence of a field, such as the electromagnetic field, derived from the four-potential  $A^\mu$ , composed by the electric scalar potential field,  $A^0 = \phi$ , and the vector potential,  $(A^1, A^2, A^3) = \mathbf{A}$ , the following change to the momentum four-vector must be made:

$$p^\mu \rightarrow p^\mu - eA^\mu, \quad A^\mu = (\phi, \mathbf{A}) \quad (3.13)$$

The Hamiltonian can now be rewritten to account for the presence of the electromagnetic field (3.14). This way it is possible to include, for example, the electron-nucleus Coulomb attraction.

$$H_D = -e\phi + \beta mc^2 + \boldsymbol{\alpha}(c\mathbf{p} + e\mathbf{A}) \quad (3.14)$$

For a central potential, as is the one generated by the nuclear charge (assuming Born-Oppenheimer's approximation), the 3 space-like components from the four-potential

are null, and the time-like component,  $\phi = \frac{Ze}{r}$ . The Hamiltonian gains now a more recognizable form:

$$H_D = -\frac{e^2 Z}{r} + \beta mc^2 + \boldsymbol{\alpha} \cdot \mathbf{p}c \quad (3.15)$$

Something very interesting about Dirac's equation is the fact that it does not yield a single solution, but in fact, two: the large component (positive energy values), for particles, and the small component (negative energy values), for antiparticles.

### 3.3 The Dirac-Breit Equation

Once again, when considering a system composed of many bodies, one must consider all the present interactions, namely, the electron-electron repulsion in an atom. Breit, in 1929, had created a relativistic approach to treat the electron-electron interactions, consisting on a set of equations building upon the classical non-relativistic Hamiltonian from equation (3.1), which can be consulted in Appendix A. Breit's equations are able to account for angular momenta couplings and estimate level energy splittings, the change of a particle's apparent mass as a function of velocity, and even include the interaction of an applied external magnetic field [9].

It is quite obvious Breit's equations introduce a great complexity in the search of the new Hamiltonian's eigenfunctions. However, when trying to include an approximation of Breit's considerations into Dirac's equation, one must add the following operator to the one present in equation (3.15):

$$H_B = \sum_{i>j} \frac{e^2}{r_{ij}} - e^2 \left( \frac{\boldsymbol{\alpha}_i \boldsymbol{\alpha}_j}{r_{ij}} + \frac{(\boldsymbol{\alpha}_i \nabla_i)(\boldsymbol{\alpha}_j \nabla_j) r_{ij}}{2} \right) \quad (3.16)$$

This set of terms will account for the fact that Coulomb interactions, mediated by the electric field, and therefore, **virtual photons**, cannot occur at instantaneous velocities, but at the speed of light.

### 3.4 The *MCDF* Method

As previously mentioned in section 3.1.1, there is a need for a numerical method in order to compute and find the eigenfunctions for a many-body Hamiltonian. While the Hartree-Fock method was able to reasonably solve the non-relativistic problem, now, while considering the Dirac-Breit Hamiltonian from equations (3.15) and (3.16), there is a need for a new method.

Hence, the state of the art *MCDFGME* arises. This self-consistent iterative method, based on the same method present in section 3.1.1, is able to solve and find eigenfunctions for a multielectronic system, now taking into account the Dirac-Breit Hamiltonian. Moreover, it is also capable of incorporating electron correlation and many QED effects not

yet considered in the relativistic equation, such as the Lamb-shift, vacuum polarization, and the electron's self energy. A brief description of these contributions can be found in [appendix C](#)



---

## Bibliography

- [1] M. F. Vitha, R. Klockenkämper, and A. V. Bohlen. *Chemical Analysis: A Series of Monographs on Analytical Chemistry and Its Applications Total-Reflection X-ray Fluorescence Analysis and Related Methods*. 2nd Edition. Vol. 181. 2015, pp. 20–21 (cit. on pp. 4, 18, 39).
- [2] *Hartree Fock method: A simple explanation*. URL: <https://insilicosci.com/hartree-fock-method-a-simple-explanation/> (cit. on pp. 6, 20, 27).
- [3] J. P. Santos. *FÍSICA ATÓMICA Apontamentos para a UC FA 2020/21* (cit. on pp. 6, 20, 27).
- [4] R. Gabriel and T. Rosa. *The Hartree-Fock Method* (cit. on pp. 6, 20, 27).
- [5] S. M. Blinder. *Introduction to the hartree-fock method*. 2018-01. DOI: [10.1016/B978-0-12-813651-5.00001-2](https://doi.org/10.1016/B978-0-12-813651-5.00001-2) (cit. on pp. 6, 20, 27).
- [6] B. Thaller. *The Dirac Equation*. Springer Berlin Heidelberg, 1992. DOI: [10.1007/978-3-662-02753-0](https://doi.org/10.1007/978-3-662-02753-0) (cit. on pp. 7, 21, 28).
- [7] H. F. Beyer and V. P. Shevelko. *Introduction to the physics of highly charged ions*. IOP Pub, 2016, pp. 1–361. ISBN: 9781420034097. DOI: [10.1016/S0168-9002\(03\)00733-2](https://doi.org/10.1016/S0168-9002(03)00733-2) (cit. on pp. 7, 21, 28).
- [8] J. J. Sakurai and J. Napolitano. *Modern Quantum Mechanics*. Cambridge University Press, 2020-09. DOI: [10.1017/9781108587280](https://doi.org/10.1017/9781108587280) (cit. on pp. 7, 21, 28).
- [9] H. A. Bethe and E. E. Salpeter. *Quantum Mechanics of One- and Two-Electron Atoms*. Springer US, 1977. DOI: [10.1007/978-1-4613-4104-8](https://doi.org/10.1007/978-1-4613-4104-8) (cit. on pp. 7, 10, 21, 23, 28, 31).
- [10] P. A. M. Dirac. “The quantum theory of the electron”. In: *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character* 117 (778 1928-02), pp. 610–624. ISSN: 0950-1207. DOI: [10.1098/rspa.1928.0023](https://doi.org/10.1098/rspa.1928.0023) (cit. on pp. 8, 22, 29).

- [11] T. V. Nguyen et al. "Theory of copper  $K_{\alpha}$  and  $K_{\beta}$  diagram lines, satellite spectra, and ab initio determination of single and double shake probabilities". In: *Physics Letters, Section A: General, Atomic and Solid State Physics* 426 (2022-02), p. 127900. ISSN: 03759601. DOI: [10.1016/j.physleta.2021.127900](https://doi.org/10.1016/j.physleta.2021.127900). URL: <https://linkinghub.elsevier.com/retrieve/pii/S0375960121007659> (cit. on pp. 11, 24).
- [12] H. A. Melia et al. "The characteristic radiation of copper  $K_{\beta}$  including radiative Auger processes". In: *Journal of Physics B: Atomic, Molecular and Optical Physics* 53 (19 2020-10), p. 195002. ISSN: 13616455. DOI: [10.1088/1361-6455/aba3a6](https://doi.org/10.1088/1361-6455/aba3a6). URL: <https://iopscience.iop.org/article/10.1088/1361-6455/aba3a6> (cit. on pp. 11, 24).
- [13] H. A. Melia et al. "The characteristic radiation of copper  $K_{\alpha}$  1,2,3,4". In: *Acta Crystallographica Section A: Foundations and Advances* 75 (3 2019-05), pp. 527–540. ISSN: 20532733. DOI: [10.1107/S205327331900130X](https://doi.org/10.1107/S205327331900130X). URL: <http://www.ncbi.nlm.nih.gov/pubmed/31041908> (cit. on pp. 11, 24).
- [14] H. Sorum. "The  $K_{\alpha}$ 1,2 X-ray spectra of the 3d transition metals Cr, Fe, Co, Ni and Cu". In: *Journal of Physics F: Metal Physics* 17 (2 1987-02), pp. 417–425. ISSN: 03054608. DOI: [10.1088/0305-4608/17/2/011](https://doi.org/10.1088/0305-4608/17/2/011) (cit. on pp. 11, 24).
- [15] J. Bremer, T. Johnsen, and H. Sorum. "The Cu  $K_{\alpha}$ 1,2 spectrum as measured with a curved-crystal spectrometer". In: *X-Ray Spectrometry* 11 (3 1982), pp. 149–152. ISSN: 10974539. DOI: [10.1002/xrs.1300110312](https://doi.org/10.1002/xrs.1300110312) (cit. on pp. 11, 24).
- [16] M. Deutsch et al. "K and K x-ray emission spectra of copper". In: *Physical Review A* 51 (1 1995-01), pp. 283–296. ISSN: 10502947. DOI: [10.1103/PhysRevA.51.283](https://doi.org/10.1103/PhysRevA.51.283) (cit. on pp. 11, 24).
- [17] Y. Ito et al. " $K_{\alpha}$ 1,2 x-ray linewidths, asymmetry indices, and [KM] shake probabilities in elements Ca to Ge and comparison with theory for Ca, Ti, and Ge". In: *Physical Review A* 94 (4 2016-10), p. 042506. ISSN: 24699934. DOI: [10.1103/PhysRevA.94.042506](https://doi.org/10.1103/PhysRevA.94.042506) (cit. on pp. 11, 24).
- [18] H. Berger. "Study of the  $K_{\alpha}$  emission spectrum of copper". In: *X-Ray Spectrometry* 15 (4 1986-10), pp. 241–243. ISSN: 0049-8246. DOI: [10.1002/xrs.1300150405](https://doi.org/10.1002/xrs.1300150405). URL: <https://onlinelibrary.wiley.com/doi/10.1002/xrs.1300150405> (cit. on pp. 11, 24).
- [19] S. Galambosi et al. "Near-threshold multielectronic effects in the Cu  $K_{\alpha}$ 1,2 x-ray spectrum". In: *Physical Review A - Atomic, Molecular, and Optical Physics* 67 (2 2003-02), p. 5. ISSN: 10941622. DOI: [10.1103/PhysRevA.67.022510](https://doi.org/10.1103/PhysRevA.67.022510). URL: <https://link.aps.org/doi/10.1103/PhysRevA.67.022510> (cit. on pp. 11, 24).

- [20] C. T. Chantler, A. C. Hayward, and I. P. Grant. “Theoretical Determination of Characteristic X-Ray Lines and the Copper  $K_{\alpha}$  Spectrum”. In: *Physical Review Letters* 103 (12 2009-09), p. 123002. ISSN: 00319007. DOI: [10.1103/PhysRevLett.103.123002](https://doi.org/10.1103/PhysRevLett.103.123002). URL: <https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.103.123002> (cit. on pp. 11, 24).
- [21] F. Gel'mukhanov and H. Ågren. *Resonant X-ray Raman scattering*. Vol. 312. Elsevier, 1999, pp. 87–330. DOI: [10.1016/S0370-1573\(99\)00003-4](https://doi.org/10.1016/S0370-1573(99)00003-4) (cit. on pp. 11, 24).
- [22] P. Carra, M. Fabrizio, and B. T. Thole. “High resolution x-ray resonant Raman scattering”. In: *Physical Review Letters* 74 (18 1995-05), pp. 3700–3703. ISSN: 00319007. DOI: [10.1103/PhysRevLett.74.3700](https://doi.org/10.1103/PhysRevLett.74.3700) (cit. on pp. 11, 25).
- [23] P. Eisenberger, P. M. Platzman, and H. Winick. “X-ray resonant Raman scattering: Observation of characteristic radiation narrower than the lifetime width”. In: *Physical Review Letters* 36 (11 1976-03), pp. 623–626. ISSN: 00319007. DOI: [10.1103/PhysRevLett.36.623](https://doi.org/10.1103/PhysRevLett.36.623) (cit. on pp. 11, 25).
- [24] P. Indelicato, J. Bieroń, and P. Jönsson. “Are MCDF calculations 101% correct in the super-heavy elements range?” In: *Theoretical Chemistry Accounts* 129 (3-5 2011-06), pp. 495–505. ISSN: 1432881X. DOI: [10.1007/s00214-010-0887-3](https://doi.org/10.1007/s00214-010-0887-3) (cit. on pp. 11, 25).
- [25] P. Indelicato, O. Gorveix, and J. P. Desclaux. “Multiconfigurational Dirac-Fock studies of two-electron ions. II. Radiative corrections and comparison with experiment”. In: *Journal of Physics B: Atomic and Molecular Physics* 20 (4 1987-02), p. 651. ISSN: 0022-3700. DOI: [10.1088/0022-3700/20/4/007](https://doi.org/10.1088/0022-3700/20/4/007) (cit. on pp. 11, 25).
- [26] O. Gorveix, P. Indelicato, and J. P. Desclaux. “Multiconfiguration Dirac-Fock studies of two-electron ions. I. Electron-electron interaction”. In: *Journal of Physics B: Atomic and Molecular Physics* 20 (4 1987-02), pp. 639–649. ISSN: 00223700. DOI: [10.1088/0022-3700/20/4/006](https://doi.org/10.1088/0022-3700/20/4/006) (cit. on pp. 11, 25).
- [27] M. Guerra et al. “Fundamental Parameters Related to Selenium  $K_{\alpha}$  and  $K_{\beta}$  Emission X-ray Spectra”. In: *Atoms* 9 (1 2021-01), p. 8. ISSN: 2218-2004. DOI: [10.3390/atoms9010008](https://doi.org/10.3390/atoms9010008). URL: <https://www.mdpi.com/2218-2004/9/1/8> (cit. on pp. 11, 25).
- [28] N. Paul et al. “Testing Quantum Electrodynamics with Exotic Atoms”. In: *Physical Review Letters* 126 (17 2021-04), p. 173001. ISSN: 10797114. DOI: [10.1103/PhysRevLett.126.173001](https://doi.org/10.1103/PhysRevLett.126.173001) (cit. on pp. 11, 25).
- [29] M. F. Gu. *The flexible atomic code*. 2008-05. DOI: [10.1139/P07-197](https://doi.org/10.1139/P07-197) (cit. on pp. 11, 25).
- [30] P. Jönsson et al. “New version: Grasp2K relativistic atomic structure package”. In: *Computer Physics Communications* 184 (9 2013-09), pp. 2197–2203. ISSN: 00104655. DOI: [10.1016/j.cpc.2013.02.016](https://doi.org/10.1016/j.cpc.2013.02.016) (cit. on pp. 11, 25).
- [31] *ASCL.net - AUTOSTRUCTURE: General program for calculation of atomic and ionic properties*. URL: <https://ascl.net/1612.014> (cit. on pp. 11, 25).

## BIBLIOGRAPHY

---

- [32] T. D. Thomas. “Transition from Adiabatic to Sudden Excitation of Core Electrons”. In: *Phys. Rev. Lett.* 52 (6 1984), pp. 417–420. DOI: [10.1103/PhysRevLett.52.417](https://doi.org/10.1103/PhysRevLett.52.417). URL: <https://link.aps.org/doi/10.1103/PhysRevLett.52.417> (cit. on p. 17).
- [33] W. E. Lamb and R. C. Retherford. “Fine structure of the hydrogen atom by a microwave method”. In: *Physical Review* 72 (3 1947-08), pp. 241–243. ISSN: 0031899X. DOI: [10.1103/PhysRev.72.241](https://doi.org/10.1103/PhysRev.72.241) (cit. on p. 40).

## The Breit Hamiltonian Operators

Note: This operators are valid for the electrons in an atom.

The free particle energy:

$$H_0 = \sum_i^N \frac{p_i^2}{2m_e} \quad (\text{A.1})$$

The electron-nucleus Coulomb attraction:

$$H_1 = \sum_i^N -\frac{e^2 Z}{r_i} \quad (\text{A.2})$$

The electron-electron Coulomb repulsion:

$$H_2 = \sum_{i < j} \frac{e^2}{r_{ij}} \quad (\text{A.3})$$

Incorporates the relativistic apparent mass - velocity dependance:

$$H_3 = -\frac{1}{8m_e^3 c^2} \sum_i^N p_i^4 \quad (\text{A.4})$$

Electric field retardation and magnetic dipole interaction:

$$H_4 = -\frac{e^2}{2m_e^2 c^2} \sum_{i < j} \left[ \frac{\mathbf{p}_i \cdot \mathbf{p}_j}{r_{ij}} + \frac{(\mathbf{r}_{ij} \cdot \mathbf{p}_{ij})(\mathbf{r}_{ij} \cdot \mathbf{p}_j)}{r_{ij}^3} \right] \quad (\text{A.5})$$

Darwin's term, accounts for the electron's quantum fluctuation motion:

$$H_5 = \frac{\pi e \hbar^2}{2m_e^2 c^2} \sum_{i < j} \frac{Z}{2} [\delta(\mathbf{r}_i) + \delta(\mathbf{r}_j)] + \delta(\mathbf{r}_{ij}) \quad (\text{A.6})$$

And the last two operators, for the consideration of spin orbit interactions:

$$H_6 = \frac{e^2 \hbar^2}{2m_e^2 c^2} \sum_{i < j} \left( Z \frac{\mathbf{r}_i \times \mathbf{p}_i}{r_i^3} - \frac{\mathbf{r}_{ij} \times \mathbf{p}_i}{r_{ij}^3} + 2 \frac{\mathbf{r}_{ij} \times \mathbf{p}_j}{r_{ij}^3} \right) \mathbf{s}_i + \left( Z \frac{\mathbf{r}_j \times \mathbf{p}_j}{r_j^3} - \frac{\mathbf{r}_{ji} \times \mathbf{p}_j}{r_{ij}^3} + 2 \frac{\mathbf{r}_{ji} \times \mathbf{p}_i}{r_{ij}^3} \right) \mathbf{s}_j \quad (\text{A.7})$$

$$H_7 = \frac{e^2 \hbar^2}{m_e^2 c^2} \sum_{i < j} \left( -\frac{8\pi}{3} \mathbf{s}_i \cdot \mathbf{s}_j \delta(\mathbf{r}_{ij}) + \frac{1}{r_{ij}^3} \left[ \mathbf{s}_i \cdot \mathbf{s}_j - \frac{3(\mathbf{s}_i \cdot \mathbf{r}_{ij})(\mathbf{s}_j \cdot \mathbf{r}_{ij})}{r_{ij}^2} \right] \right) \quad (\text{A.8})$$

## Transition Diagram




Figure B.1: Transition notation scheme. Adapted from [1]

## QED considerations

It should be apparent by now that studying Atomic systems call not only for relativistic effects and corrections, but also for QED ones.

One of the most famous cases where QED came to light was the discovery of the Lamb Shift [33], when it was discovered Hydrogen's  $2s_{1/2}$  and  $2p_{1/2}$  levels were in fact, not degenerate (did not have the same energy), contrary to what was expected from solving Dirac's equation. This difference in energy came to be known as the Lamb Shift, only explained by QED effects.

### C.1 Self-Energy

The self-energy represents a particle's emission and reabsorption of virtual photon, present in the particle's own generated field. This interaction has the most impact in the Lamb Shift effect and when performing energy corrections. One of its [Feynman Diagrams](#) can be seen in Figure C.1a.

### C.2 Vacuum Polarization

As previously stated, electromagnetic fields, such as the Coulomb field generated by the nucleus, are mediated by virtual photons. These photons can lead to the creation of electron-positron pairs which create screening effects. Pair annihilation will follow, leading to the production of another virtual photon (Figure C.1b).



Figure C.1: QED Feynman Diagrams



## Input File .f05 Example

```

1  program_year=2019 program_version=1
2  * 1 Z=29 (1s)1 (2s)2 (2p)6 (3s)2 (3p)6 (3d)10 (4s)2 2J=1 neig=[2S1/2]#1
   LS1
3      scfmdf max :
4      mod_lightspeed=n
5      nz=29
6      mdf opt_ener=todo modfilename_ener=n modfilename_wf=n do_scf=y
7      Breit=full mag_scf=y ret_scf=y
8      qedstpg_n4=n
9      vacpol_scf=y
10     energy
11     # use_mcdfener=y
12     # opt_relax=y
13     ret_Lorentz=y
14     opt_qedel=y :
15     # ":" above is for ilams taken to be the default value
16     mod_mesh=n
17     # hx=0.012 r(1)=0.001 amesh=0.01 :
18     exotic=n
19     use_nms=y
20     mod_nuc=n
21     project=n
22     # cgt_order_vint=y order=6
23     cgt_order_vint=n
24     def_config=given
25     nbel=29 jjt=1 :
26     c 1 (1s)1 (2s)2 (2p)6 (3s)2 (3p)6 (3d)10 (4s)2 :
27     end
28     # initial state parameters
29     neigv=1 icmul=0 iprfr=0
30     norbsc=00 ndep=0 nlec=0 nec=1 :
31     nstep=0
32     lregul=n modtest=n
33     modsolv_orb=n
34     mod_odlm=n

```

## ANNEX I. INPUT FILE .F05 EXAMPLE

---

```
35      # data for uwfrdf
36      :
37      **
```



