

Lattice gauge ensembles and data management

**Yasumichi Aoki,^{a,#} Ed Bennett,^{b,1,*} Rayan Bignell,^{c,2,*} Kadir Utku Can,^{d,3,*}
Takumi Doi,^{e,4,*} Steven Gottlieb,^{f,5,*} Rajan Gupta,^{g,6,*} Georg von Hippel,^{h,7,*}
Issaku Kanamori,^{a,8,*} Andrey Kotov,^{i,9,*} Giannis Koutsou,^{j,10,#,*}
Robert Mawhinney,^{k,11,*} Agostino Patella,^{l,12,*} Giovanni Pederiva,^{i,13,*} Christian
Schmidt,^{m,14,*} Takeshi Yamazaki,^{n,15,*} and Yi-Bo Yang^{o,16,*}**

^aRIKEN Center for Computational Science (R-CCS), Kobe, 650-0047, Japan

^bSwansea Academy of Advanced Computing, Swansea University, Swansea, United Kingdom

^cSchool of Mathematics, Trinity College, Dublin, Ireland

^dCSSM, Department of Physics, The University of Adelaide, Australia

^eInterdisciplinary Theoretical and Mathematical Sciences Program (iTHEMS), RIKEN, Japan

^fDepartment of Physics, Indiana University, IN, USA

^gTheoretical Division, Los Alamos National Laboratory, NM, USA

^hPRISMA+ Cluster of Excellence and Institut für Kernphysik

ⁱJülich Supercomputing Center, Forschungszentrum Jülich, Germany

^jComputation-based Science and Technology Research Center, The Cyprus Institute, Cyprus

^kPhysics Department, Columbia University, USA

^lHumboldt Universität zu Berlin, Institut für Physik & IRIS, Germany

^mFakultät für Physik, Universität Bielefeld, Germany

ⁿInstitute of Pure and Applied Sciences, University of Tsukuba, Japan

^oCAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China

We summarize the status of lattice QCD ensemble generation efforts and their data management characteristics. Namely, this proceeding summarizes contributions to a dedicated parallel session during the 41st International Symposium on Lattice Field Theory (Lattice 2024), during which representatives of 16 lattice QCD collaborations provided details on their simulation program, with focus on plans for publication, data management, and storage requirements. The parallel session was organized by the International Lattice Data Grid (ILDG), following an open call to the lattice QCD for participation in the session.

The 41st International Symposium on Lattice Field Theory (LATTICE2024)

28 July - 3 August 2024

Liverpool, UK

¹ For the TELOS collaboration ² For the FASTSUM collaboration ³ For the CSSM/QCDSF/UKQCD collaboration

⁴ For the HAL QCD collaboration ⁵ For the MILC collaboration ⁶ For the Jlab/W&M/LANL/MIT/Marseille effort

⁷ For the CLS ⁸ For the JLQCD collaboration ⁹ For the TWEXT collaboration ¹⁰ For the ETM collaboration

(ETMC) ¹¹ For the RBC-UKQCD collaboration ¹² For the RC* collaboration ¹³ For the OPEN LAT initiative

¹⁴ For the HotQCD collaboration ¹⁵ For the PACS collaboration ¹⁶ For the CLQCD collaboration # Conveners

* Speaker

Contents

1	Introduction	2
2	Contributions	3
2.1	CLQCD	4
2.2	Jlab/W&M/LANL/MIT/Marseille	4
2.3	HotQCD	5
2.4	FASTSUM	5
2.5	TELOS	6
2.6	HAL QCD	6
2.7	TWEXT	7
2.8	CSSM/QCDSF/UKQCD	7
2.9	RBC-UKQCD	8
2.10	OPEN LAT	8
2.11	RC*	9
2.12	ETMC	9
2.13	JLQCD	10
2.14	MILC	10
2.15	CLS	11
2.16	PACS	11
3	Summary	12

1. Introduction

The simulation of Quantum Chromodynamics (QCD) via its Euclidean-time, discrete formulation on a lattice, has been one of the most compute-intensive applications in scientific computing, consuming substantial fractions of computer time at leadership HPC facilities internationally. In particular, the generation of ensembles of gauge configurations, for multiple values of the QCD parameters such as the QCD coupling, the quark masses, and the extent of the finite volume, requires multi-year simulation campaigns, coordinated by multi-member research collaborations. It is thus common that collaborations store and reuse the same gauge ensembles for multiple observables of interest, and in many cases also share the ensembles with researchers external to the collaboration that generated them.

The purpose of this proceeding is to summarize the available gauge ensembles generated by various lattice QCD collaborations internationally, with a focus on the data management practices each collaboration employs. It follows a parallel session at the 41st International Symposium on Lattice Field Theory (Lattice 2024), during which 16 collaborations provided status reports of

their simulation efforts, responding to an open call for participation addressed to the lattice QCD community prior to the conference. The first such session was during Lattice 2022 and a report of the contributions presented during that session can be found in Ref. [1].

These sessions are organized by the International Lattice Data Grid (ILDG) with the intention of obtaining, gathering, and summarizing the evolving needs of the lattice community in terms of data storage and management. The ILDG was setup in the early 2000s [2–5] by the lattice community, which realized early on the value in standardizing data management practices across the field. ILDG is organized as a federation of autonomous *regional grids*, within a single Virtual Organization [6]. It standardizes interfaces for the services, which are to be operated by each regional grid, such as storage and a searchable metadata catalog, so that the regional services are interoperable. Within ILDG, working groups specify community-wide agreed metadata schemas (QCDml) [7] to concisely mark-up the gauge configurations and develop relevant middleware tools for facilitating the use of ILDG services. The middleware and metadata specifications developed by ILDG adhere to most of the FAIR (Findable, Accessible, Interoperable, Reusable) principles [8]. A summary of recent developments in ILDG, referred to as ILDG 2.0, was presented during the same session and can be found in a separate proceeding [9].

In the remainder of this proceeding, we present the status of ensemble generation of each of the 16 collaborations that contributed to the parallel session. We restrict ourselves to simulations of QCD, and at present these are carried out using $N_f=2+1$, $N_f=2+1+1$, and $N_f=1+1+1+1$ sea quark flavors with various fermion discretizations. The contributors were asked to specify whether their data are public or if they plan on making them public, their interest in using ILDG services and tools for that purpose, as well as some overall information regarding storage requirements. This information is collected in a table and summary section that follows the individual contributions.

2. Contributions

The contributions from each collaboration follow, in the order presented during the parallel session. The original presentations can be found on the conference website [10].

2.1 CLQCD

The CLQCD collaboration focuses on the first principles QCD study of the spectrum of exotic hadrons, parton structure of the nucleon, and other traditional hadron, N-point correlation functions related to high accuracy tests of the standard model, and also QCD in extreme conditions. For this purpose, CLQCD generated a set of configurations of $N_f=2+1$ fermions using the tadpole improved clover fermion action with stout smearing and the tadpole improved Symanzik gauge action, at 5 values of the lattice spacing $\in [0.05, 0.11]$ fm, several pion masses $\in [120, 350]$ MeV, several volumes with $m_\pi L \in [2.6, 5.8]$ fm, and several temperatures $\in (0, 464]$ MeV. In addition, there are several anisotropic ensembles using clover fermions with $\xi = 5$ at different lattice spacing, pion mass, volume and flavors which concentrate on the high precision studies related to glueball properties. For the gauge generation the Chroma package [11] with QUDA [12–14] is used at present, and we plan to switch to PyQUDA [15] with QUDA in the near future. Currently the CLQCD collaboration has ~ 20 ensembles (one ensemble corresponds to one point in the space lattice spacing-spatial volume-pion mass-temperature), which occupy ~ 100 TB of disk space. Configurations are stored in the SCIDAC format. Possible collaborations on the analysis of the correlation functions are welcome and CLQCD is preparing the hardware and software for the data sharing service.

2.2 Jlab/W&M/LANL/MIT/Marseille

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

2.3 HotQCD

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

2.4 FASTSUM

The FASTSUM collaboration use $N_f=2+1$ flavor anisotropic gauge ensembles in the fixed-scale approach to study the behavior of QCD as a function of temperature in hadronic and plasma phases. Specifically we have considered the behavior of hadronic states including light, strange, charm and bottom quarks, the electrical conductivity of QCD matter, the interquark potential and properties of the chiral transition. FASTSUM gauge fields utilise an $O(a^2)$ improved Symanzik gauge action and an $O(a)$ improved spatially stout-smearred Wilson-Clover action following the parameter tuning and zero-temperature ensembles of the Hadron Spectrum collaboration [16, 17]. “Generation 2” ensembles were generated using the Chroma [11] software suite while the newer “Generation 2L” used a modification [18] of the openQCD [19, 20] package which introduces stout-link smearing and anisotropic actions. Generation 2 and 2L have an anisotropy $\xi = a_s/a_\tau \sim 3.5$ with $a_s \sim 0.12$ fm, $N_s = 24$ or 32 and a wide range of N_τ corresponding to $T \in [44, 760]$ MeV. Full details of these ensembles may be found in Refs. [21, 22]. We are in the process of production for “Generation 3” - a parameter set similar to “Generation 2” but with twice the anisotropy $\xi \sim 7$ - using openQCD-FASTSUM [18]. We maintain a centralised metadata repository detailing (among other information) who was responsible for each run, on which machine that run was produced and where copies may be found. The gauge fields are redundantly stored on two (well-separated) storage servers managed by Swansea University in the openQCD format. The “Generation 2” ensembles are publically available [23] while other ensembles will be available after an embargo period. We anticipate making ensembles available through the next incarnation of the ILDG with supplementary information also available on Zenodo [23, 24].

2.5 TELOS

The TELOS collaboration performs **T**heoretical **E**xplorations on the **L**attice with **O**rthogonal and **S**ymplectic groups. Problems of interest focus on physics beyond the Standard Model, in particular composite Higgs models. Our work to date has made use of the Wilson gauge action and Wilson fermion action. Our ensembles include studies of the $\text{Sp}(4)$ theory with two fundamental fermion flavours ($N_f=2$) [25] (five values of $\beta \in [6.9, 7.5]$, $V \leq 48 \times 42^3$, $m_{\text{PS}}/m_V \gtrsim 0.407(16)$), the $\text{Sp}(4)$ theory with three antisymmetric fermion flavours ($N_{\text{as}}=3$) [26], (six values of $\beta \in [6.6, 6.9]$, $V \leq 54 \times 36^3$, $m_{\text{PS}}/m_V \gtrsim 0.7954(44)$); and the $\text{Sp}(4)$ theory with $N_f=2$ and $N_{\text{as}}=3$ [27], (three values of $\beta \in [6.45, 6.5]$, $V \leq 56 \times 36^3$, $m_{\text{PS}}/m_V \gtrsim 0.8768(30)$, $m_{\text{PS}}/m_V \gtrsim 0.9022(27)$). In the latter two cases, the topological charge becomes slow running at small m_{as} , and at larger β . We do not retain our pure gauge ensembles, used for studies of the large- N limit of $\text{Sp}(2N)$ [28–30], since the costs of storage and data transfer are higher than those of regenerating the ensembles.

Additionally, we present ensembles generated by a subset of the collaboration, with applications to conformal and near-conformal dynamics, and to potential Walking Technicolor theories, again using the Wilson gauge and Wilson fermion actions [31]. Specifically, these are $\text{SU}(2)$ with one adjoint flavour ($N_{\text{adj}}=1$) (seven values of $\beta \in [2.05, 2.4]$, $V \leq 96 \times 48^3$, $m_{2_s^+} \gtrsim 0.28$), and $N_{\text{adj}}=2$ ($V \leq 128 \times 64^3$, $m_{2_s^+} \gtrsim 0.47$). In the former case, the majority of ensembles show ergodic topology, with $\beta = 2.4$ being marginal; in the latter, at large volumes we see significant topological freezing.

Ensembles are generated using HiRep [32, 33] and Grid [34, 35]. The above ensembles will be made available as soon as the infrastructure is in place to do so.

We are in the process of generating ensembles for $\text{Sp}(4)$ $N_f=2$ and for $\text{SU}(2)$ $N_f=1, 2$ with Möbius domain wall fermions, and for $\text{SU}(2)$ $N_f=1, 2$ with Wilson fermions and additional Pauli–Villars fields, which we aim to make available concurrently with the corresponding papers.

2.6 HAL QCD

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17

18
19

2.7 TWEXT

1 The TWEXT (Twisted Wilson @ Extreme conditions) collaboration studies the properties of
2 QCD at high temperature using Wilson Twisted Mass fermions. Problems under investigation in-
3 clude chiral properties of QCD, in particular the behavior of QCD around the chiral phase transition
4 and its scaling window [36], topological properties of QCD and QCD axion [37], hadron masses,
5 symmetries of QCD and others. For this purpose, TWEXT generated a set of configurations for
6 $N_f=2+1+1$ fermions at the physical pion mass and also uses older configurations with heavier pion
7 mass. Configurations with the physical pion mass have three lattice spacings $a \in (0.057, 0.080)$ fm
8 and cover a wide range of temperatures from ~ 120 MeV to ~ 900 MeV. It allows the TWEXT
9 collaboration to perform the continuum extrapolation for quantities of interest in this temperature
10 range. For the generation the tmLQCD software package [38–40] is used and the parameters of
11 the ensembles were taken from the zero temperature simulations of the ETM collaboration [41].
12 Currently, the TWEXT collaboration has 80 ensembles (one ensemble corresponds to one point in
13 the space temperature-pion mass-lattice spacing), which occupy ~ 80 TB of disk space. Configu-
14 rations are stored in the ILDG format. Possible collaborations are welcome and TWEXT plans to
15 make configurations public/use ILDG in the future, after performing the ongoing analysis.

2.8 CSSM/QCDSF/UKQCD

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

2.9 RBC-UKQCD

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

2.10 OPEN LAT

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

2.11 RC*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

2.12 ETMC

1 The ETM collaboration focuses on hadron spectroscopy, hadron structure, and flavor physics
2 at zero temperature. Ensembles employ the twisted mass formulation, realizing $O(a)$ -improvement
3 by tuning to maximal twist, and include a clover term to further reduce the size of lattice artifacts.
4 The Iwasaki gauge action is used. The main simulation effort is for the generation of ensembles
5 with degenerate up- and down-, strange- and charm-quarks ($N_f=2+1+1$) with lattice spacing ranging
6 between 0.049 and 0.091 fm. $M_\pi \cdot L$ varies from 2.5 up to ~ 5.5 . At the time of writing, 24 ensembles
7 are available or in the process of being generated, with 8 of these at approximately physical values of
8 the quark masses. For a recent listing of the ensembles, see [42]. Simulations are performed using
9 the Hybrid Monte Carlo (HMC) algorithm implemented in the tmLQCD software package [38–40].
10 See Ref. [41] for details on the simulation program, including the parameter tuning. The DD-
11 α AMG [43, 44] multigrid iterative solver is employed for the most poorly conditioned monomials
12 in the light sector while mixed-precision CG is used elsewhere. Multi-shift CG is used together
13 with shift-by-shift refinement using DD- α AMG [45] for a number of small shifts for the heavy
14 sector. tmLQCD has interfaces to QPhiX [46] and QUDA [12, 13]. tmLQCD automatically writes
15 gauge configurations in the ILDG format, with meta-data including creation date, target simulation
16 parameters, and the plaquette. ETMC policy is to make ensembles publicly available after a grace
17 period. Older $N_f=2$ and $N_f=2+1+1$ ensembles [47–49] have made use of ILDG storage elements.
18 The current ensembles are available upon request and the collaboration intends to use ILDG in the
19 near future. For these ensembles, we expect storage requirements to reach 3 PB.

2.13 JLQCD

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

2.14 MILC

The MILC Collaboration has been creating ensembles of gauge configurations for decades. Initial calculations used two dynamical flavors with naive staggered quarks. Second generation efforts used dynamical up, down, and strange quarks with the asqtad action, culminating with a review in Ref. [50]. Our third generation calculations use the highly improved staggered quark (HISQ) action [51]. They contain dynamical up, down, strange, and charm quarks. For most of the ensembles, up and down are degenerate. The charm quark is set near its physical value. For several ensembles, the light quarks are near their average physical value. There are also ensembles with $m_l = 0.1$, or $0.2m_s$, where m_l (m_s) is the light (strange) quark mass. We have generated a number of ensembles with m_s less than its physical value to explore low energy constants and the chiral Lagrangian. Lattice spacings are in the range of $[0.15, 0.03]$ fm. In a few cases, multiple volumes are available. Details about the generation of configurations can be found in Refs. [52–54].

The sharing policy for the MILC ensembles is available on GitHub [55]. On that web page can be found links to 1) the sharing policy, 2) a Google Sheet detailing freely available ensembles, and 3) a document summarizing which papers to cite for the use of each ensemble. Anyone wishing to use an ensemble that is not listed in the Google Sheet, but has been used in a publication or noted in a talk, is welcome to contact a member of the Fermilab Lattice or MILC Collaborations to inquire as to whether the ensemble can be made available for a specific project. Many configurations are available on USQCD resources, making access relatively easy for USQCD members. The ILDG is not operational in the US, but should it become so, we would make an effort to use it. We have assisted transfer of configurations to other researchers both within and outside the US.

2.15 CLS

The CLS (Coordinated Lattice Simulations) effort uses the openQCD code [20] to generate ensembles [56, 57] with $N_f=2+1$ non-perturbatively $O(a)$ -improved Wilson quarks and tree-level improved Symanzik glue, mostly with open boundary conditions in time to avoid topological freezing [19, 58], but also with (anti-)periodic boundary conditions in time (on some ensembles at $a \gtrsim 0.06$ fm), at six fine lattice spacings $a \in [0.039, 0.1]$ fm and quark masses from the symmetric to the physical point on three chiral trajectories ($\text{Tr}[M] = \text{const.}$, $m_s \approx \text{const.}$, $m_s = m_l$) in large volumes satisfying $M_\pi L \geq 4$ throughout, with statistics typically $\gtrsim 2,000$ MDU.

Due to the algorithm used, two reweighting factors are needed to use CLS ensembles: one to correct for the twisted-mass stabilization of the light quarks, and one to correct for the rational approximation to $\sqrt{D^\dagger D}$ for the strange quark. In the latter case, $\det D < 0$ can occur [59], so that one also needs to correct for the wrong sign of the reweighting factor; fortunately, the fraction of configurations with a negative reweighting factor is very small (or zero) for most ensembles.

At the time of Lattice 2024, there were 149,766 configurations (1.384 PB) stored on tape in the openQCD (non-ILDG) data format. Metadata regarding data provenance, simulation setup, HMC stability, and data integrity are collected automatically via automated scripts, while reweighting factors and determinant minus signs are measured separately. A first batch of ensembles has been successfully uploaded to ILDG; several more ensembles will follow soon, and the remainder will follow after an embargo period. The XML metadata are generated automatically by extraction from the existing database; the (signed) reweighting factors are included in the Config XML.

2.16 PACS

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

3. Summary

Table 1: Public: (2 = currently public, 1 = after an embargo period, 0 = no); ILDG: (N = no interest, I = interest, P = planned, U = already using); #ens: Number of ensembles; #cfg: Total number of configurations; storage: Total storage needed in TBytes.

Collaboration	Public	ILDG	#ens	#cfg	Storage (TB)
CLQCD	1	I	20	10,000	100
Jlab/W&M/LANL/MIT/Marseille					
HotQCD					
FASTSUM	2,1	I	37	37,000	65
TELOS	1	P	250	800,000	120
HAL QCD					
TWEXT	1	P	80	70,000	80
CSSM/QCDSF/UKQCD					
RBC-UKQCD					
OPEN LAT					
RC*					
ETMC	1	P,U	24	12,000	3,000
JLQCD					
MILC	1	I	50	70,000	600
CLS	1	P,U	25	150,000	1,400
PACS					

Acknowledgments

G. K. acknowledges support from EXCELLENCE/0421/0195, co-financed by the European Regional Development Fund and the Republic of Cyprus through the Research and Innovation Foundation and the AQTIVATE a Marie Skłodowska-Curie Doctoral Network GA No. 101072344. R. B. acknowledges support from a Science Foundation Ireland Frontiers for the Future Project award with grant number SFI-21/FFP-P/10186. The work of E. B. is part-funded by the UKRI Science and Technology Facilities Council (STFC) Research Software Engineer Fellowship EP/V052489/1, the UKRI Engineering and Physical Science Research Council (EPSRC) ExCALIBUR programme ExaTEPP project EP/X017168/1, and the STFC Consolidated Grant ST/T000813/1. Y.-B. Y. is supported in part by NSFC grants No. 12293060, and 12435002.

The participating collaborations acknowledge the following HPC systems, for the generation of the gauge ensembles reported here,

LUMI-C and LUMI-G at CSC, Finland,

Irène Joliot-Curie at Très Grand Centre de Calcul (TGCC) in Bruyères-le-Châtel, France,

JUGENE, JUWELS and JUWELS-Booster at Jülich Supercomputing Centre (JSC),

HAWK at Höchstleistungsrechenzentrum Stuttgart (HLRS), Germany,

SuperMUC and SuperMUC-NG at Leibniz Rechenzentrum (LRZ) in Garching, Germany,

Kay and Stokes at the Irish Centre for High-End Computing (ICHEC) in Galway, Ireland,
Leonardo, Marconi 100 and Marconi A2 at CINECA in Bologna, Italy,
Tesseract at the DiRAC Extreme Scaling Service at the University of Edinburgh, United Kingdom,
DiRAC Data Intensive 2.5 & 3 at the University of Leicester, United Kingdom,
Sunbird of Supercomputing Wales, supported by the European Regional Development Fund via the Welsh Government,
the HPC Cluster of ITP-CAS, the Southern Nuclear Science Computing Center (SNSC),
Siyuan-1 cluster, supported by the Center for High Performance Computing at Shanghai Jiao Tong University,
Frontera at TACC, TX, US,
and the DiRAC Extreme Scaling and Data Intensive services, which are part of the UKRI Digital Research Infrastructure.

References

- [1] G. Bali et al., *Lattice gauge ensembles and data management*, *PoS LATTICE2022* (2022) 203 [2212.10138].
- [2] UKQCD collaboration, C. T. H. Davies, A. C. Irving, R. D. Kenway and C. M. Maynard, *International lattice data grid*, *Nucl. Phys. B Proc. Suppl.* **119** (2003) 225 [hep-lat/0209121].
- [3] T. Yoshie, *Making use of the International Lattice Data Grid*, *PoS LATTICE2008* (2008) 019 [0812.0849].
- [4] C. M. Maynard, *International Lattice Data Grid: Turn On, Plug In, and Download*, *PoS LAT2009* (2009) 020 [1001.5207].
- [5] M. G. Beckett, B. Joo, C. M. Maynard, D. Pleiter, O. Tatebe et al., *Building the International Lattice Data Grid*, *Comput. Phys. Commun.* **182** (2011) 1208 [0910.1692].
- [6] I. L. D. Grid, “Organization of ildg activities.” <https://hpc.desy.de/ildg/organization/>. Accessed 2024-08-06.
- [7] ILDG METADATA WORKING GROUP collaboration, P. Coddington, B. Joo, C. M. Maynard, D. Pleiter and T. Yoshie, *Marking up lattice QCD configurations and ensembles*, *PoS LATTICE2007* (2007) 048 [0710.0230].
- [8] M. D. Wilkinson, M. Dumontier, I. J. Aalbersberg, G. Appleton, M. Axton et al., *The fair guiding principles for scientific data management and stewardship*, *Scientific Data* **3** (2016) 160018.
- [9] H. Matsufuru, H. Simma and C. Urbach, *ILDG 2.0*, *PoS LATTICE2024* (2024) .

- [10] “Lattice data session.” <https://conference.ippp.dur.ac.uk/event/1265/sessions/1744/#20240802>. Accessed 2024-08-02.
- [11] SciDAC, LHPC, UKQCD collaboration, R. G. Edwards and B. Joó, *The Chroma software system for lattice QCD*, *Nucl. Phys. Proc. Suppl.* **140** (2005) 832 [[hep-lat/0409003](#)].
- [12] QUDA collaboration, M. A. Clark, R. Babich, K. Barros, R. C. Brower and C. Rebbi, *Solving Lattice QCD systems of equations using mixed precision solvers on GPUs*, *Comput. Phys. Commun.* **181** (2010) 1517 [[0911.3191](#)].
- [13] QUDA collaboration, R. Babich, M. A. Clark, B. Joo, G. Shi, R. C. Brower et al., *Scaling lattice QCD beyond 100 GPUs*, in *International Conference for High Performance Computing, Networking, Storage and Analysis*, 9, 2011, [1109.2935](#), [DOI](#).
- [14] QUDA collaboration, M. A. Clark, B. Joó, A. Strelchenko, M. Cheng, A. Gambhir et al., *Accelerating lattice QCD multigrid on GPUs using fine-grained parallelization*, in *International Conference for High Performance Computing, Networking, Storage and Analysis*, 12, 2016, [1612.07873](#), [DOI](#).
- [15] X. Jiang, C. Shi, Y. Chen, M. Gong and Y.-B. Yang, *Use QUDA for lattice QCD calculation with Python*, [2411.08461](#).
- [16] R. G. Edwards, B. Joo and H.-W. Lin, *Tuning for Three-flavors of Anisotropic Clover Fermions with Stout-link Smearing*, *Phys. Rev. D* **78** (2008) 054501 [[0803.3960](#)].
- [17] HADRON SPECTRUM collaboration, H.-W. Lin et al., *First results from 2+1 dynamical quark flavors on an anisotropic lattice: Light-hadron spectroscopy and setting the strange-quark mass*, *Phys. Rev. D* **79** (2009) 034502 [[0810.3588](#)].
- [18] J. R. Glesaaen and B. Jäger, “openqcd-fastsum.” <https://gitlab.com/fastsum>, Apr., 2018. 10.5281/zenodo.2216355.
- [19] M. Lüscher and S. Schaefer, *Lattice QCD with open boundary conditions and twisted-mass reweighting*, *Comput. Phys. Commun.* **184** (2013) 519 [[1206.2809](#)].
- [20] “openQCD: Simulation programs for lattice QCD.” <https://luscher.web.cern.ch/luscher/openQCD/>.
- [21] G. Aarts, C. Allton, A. Amato, P. Giudice, S. Hands et al., *Electrical conductivity and charge diffusion in thermal QCD from the lattice*, *JHEP* **02** (2015) 186 [[1412.6411](#)].
- [22] G. Aarts et al., *Properties of the QCD thermal transition with $N_f=2+1$ flavors of Wilson quark*, *Phys. Rev. D* **105** (2022) 034504 [[2007.04188](#)].
- [23] G. Aarts, C. Allton, A. Amato, R. Bignell, T. J. Burns et al., *FASTSUM Generation 2 Anisotropic Thermal Lattice QCD Gauge Ensembles*, July, 2024. 10.5281/zenodo.8403827.

- [24] European Organization For Nuclear Research and OpenAIRE, *Zenodo*, 2013. 10.25495/7GXX-RD71.
- [25] E. Bennett, D. K. Hong, J.-W. Lee, C. J. D. Lin, B. Lucini et al., *Sp(4) gauge theories on the lattice: $N_f = 2$ dynamical fundamental fermions*, *JHEP* **12** (2019) 053 [[1909.12662](#)].
- [26] H. Hsiao, E. Bennett, D. K. Hong, J.-W. Lee, C. J. D. Lin et al., *Spectroscopy of Sp(4) lattice gauge theory with $n_f = 3$ antisymmetric fermions*, *PoS LATTICE2022* (2023) 211 [[2210.08154](#)].
- [27] E. Bennett, D. K. Hong, H. Hsiao, J.-W. Lee, C. J. D. Lin et al., *Lattice studies of the Sp(4) gauge theory with two fundamental and three antisymmetric Dirac fermions*, *Phys. Rev. D* **106** (2022) 014501 [[2202.05516](#)].
- [28] E. Bennett, J. Holligan, D. K. Hong, J.-W. Lee, C. J. D. Lin et al., *Spectrum of mesons in quenched Sp(2N) gauge theories*, *Phys. Rev. D* **109** (2024) 094517 [[2312.08465](#)].
- [29] E. Bennett, D. K. Hong, J.-W. Lee, C. J. D. Lin, B. Lucini et al., *Color dependence of the topological susceptibility in Yang-Mills theories*, *Phys. Lett. B* **835** (2022) 137504 [[2205.09254](#)].
- [30] E. Bennett, J. Holligan, D. K. Hong, J.-W. Lee, C. J. D. Lin et al., *Glueballs and strings in Sp(2N) Yang-Mills theories*, *Phys. Rev. D* **103** (2021) 054509 [[2010.15781](#)].
- [31] A. Athenodorou, E. Bennett, G. Bergner, P. Butti, J. Lenz et al., *SU(2) gauge theory with one and two adjoint fermions towards the continuum limit*, [2408.00171](#).
- [32] E. Bennett, D. K. Hong, J.-W. Lee, C.-J. D. Lin, B. Lucini et al., *Sp(4) gauge theories on the lattice: quenched fundamental and antisymmetric fermions*, *Phys. Rev. D* **101** (2020) 074516 [[1912.06505](#)].
- [33] L. Del Debbio, A. Patella and C. Pica, *Higher representations on the lattice: Numerical simulations. SU(2) with adjoint fermions*, *Phys. Rev. D* **81** (2010) 094503 [[0805.2058](#)].
- [34] E. Bennett et al., *Symplectic lattice gauge theories in the grid framework: Approaching the conformal window*, *Phys. Rev. D* **108** (2023) 094508 [[2306.11649](#)].
- [35] A. Yamaguchi, P. Boyle, G. Cossu, G. Filaci, C. Lehner et al., *Grid: OneCode and FourAPIs*, *PoS LATTICE2021* (2022) 035 [[2203.06777](#)].
- [36] A. Y. Kotov, M. P. Lombardo and A. Trunin, *QCD transition at the physical point, and its scaling window from twisted mass Wilson fermions*, *Phys. Lett. B* **823** (2021) 136749 [[2105.09842](#)].
- [37] A. Y. Kotov, A. Trunin and M. P. Lombardo, *QCD topology and axion's properties from Wilson twisted mass lattice simulations*, *PoS LATTICE2021* (2022) 032 [[2111.15421](#)].
- [38] K. Jansen and C. Urbach, *tmLQCD: A Program suite to simulate Wilson Twisted mass Lattice QCD*, *Comput. Phys. Commun.* **180** (2009) 2717 [[0905.3331](#)].

- [39] A. Deuzeman, K. Jansen, B. Kostrzewa and C. Urbach, *Experiences with OpenMP in tmLQCD*, *PoS LATTICE2013* (2014) 416 [1311.4521].
- [40] A. Abdel-Rehim, F. Burger, A. Deuzeman, K. Jansen, B. Kostrzewa et al., *Recent developments in the tmLQCD software suite*, *PoS LATTICE2013* (2014) 414 [1311.5495].
- [41] C. Alexandrou et al., *Simulating twisted mass fermions at physical light, strange and charm quark masses*, *Phys. Rev. D* **98** (2018) 054518 [1807.00495].
- [42] B. Kostrzewa, “Status of the etmc ensemble generation effort.”
<https://conference.ippp.dur.ac.uk/event/1265/contributions/7655/>.
- [43] A. Frommer, K. Kahl, S. Krieg, B. Leder and M. Rottmann, *Adaptive Aggregation-Based Domain Decomposition Multigrid for the Lattice Wilson–Dirac Operator*, *SIAM J. Sci. Comput.* **36** (2014) A1581 [1303.1377].
- [44] C. Alexandrou, S. Bacchio, J. Finkenrath, A. Frommer, K. Kahl et al., *Adaptive Aggregation-based Domain Decomposition Multigrid for Twisted Mass Fermions*, *Phys. Rev. D* **94** (2016) 114509 [1610.02370].
- [45] C. Alexandrou, S. Bacchio and J. Finkenrath, *Multigrid approach in shifted linear systems for the non-degenerated twisted mass operator*, *Comput. Phys. Commun.* **236** (2019) 51 [1805.09584].
- [46] B. Joó, D. D. Kalamkar, K. Vaidyanathan, M. Smelyanskiy, K. Pamnany et al., *Lattice QCD on Intel® Xeon Phi Coprocessors*, *Lect. Notes Comput. Sci.* **7905** (2013) 40.
- [47] R. Baron et al., *Light hadrons from lattice QCD with light (u,d), strange and charm dynamical quarks*, *JHEP* **06** (2010) 111 [1004.5284].
- [48] EUROPEAN TWISTED MASS collaboration, R. Baron et al., *Computing K and D meson masses with $N_f = 2+1+1$ twisted mass lattice QCD*, *Comput. Phys. Commun.* **182** (2011) 299 [1005.2042].
- [49] ETM collaboration, R. Baron et al., *Light Meson Physics from Maximally Twisted Mass Lattice QCD*, *JHEP* **08** (2010) 097 [0911.5061].
- [50] MILC collaboration, A. Bazavov et al., *Nonperturbative QCD Simulations with 2+1 Flavors of Improved Staggered Quarks*, *Rev. Mod. Phys.* **82** (2010) 1349.
- [51] HPQCD collaboration, E. Follana, Q. Mason, C. Davies, K. Hornbostel, G. P. Lepage et al., *Highly improved staggered quarks on the lattice, with applications to charm physics*, *Phys. Rev. D* **75** (2007) 054502 [hep-lat/0610092].
- [52] MILC collaboration, A. Bazavov et al., *Scaling studies of QCD with the dynamical HISQ action*, *Phys. Rev. D* **82** (2010) 074501 [1004.0342].
- [53] MILC collaboration, A. Bazavov et al., *Lattice QCD Ensembles with Four Flavors of Highly Improved Staggered Quarks*, *Phys. Rev. D* **87** (2013) 054505.

- [54] FERMILAB LATTICE, MILC collaboration, A. Bazavov et al., *B- and D-meson leptonic decay constants from four-flavor lattice QCD*, *Phys. Rev. D* **98** (2018) 074512 [1712.09262].
- [55] “Milc and fermilab lattice collaboration policy for sharing lattice files.”
<https://github.com/milc-qcd/sharing/wiki/LatticeSharing>.
- [56] M. Bruno et al., *Simulation of QCD with $N_f = 2 + 1$ flavors of non-perturbatively improved Wilson fermions*, *JHEP* **02** (2015) 043 [1411.3982].
- [57] D. Mohler, S. Schaefer and J. Simeth, *CLS 2+1 flavor simulations at physical light- and strange-quark masses*, *EPJ Web Conf.* **175** (2018) 02010 [1712.04884].
- [58] ALPHA collaboration, S. Schaefer, R. Sommer and F. Virotta, *Critical slowing down and error analysis in lattice QCD simulations*, *Nucl. Phys. B* **845** (2011) 93 [1009.5228].
- [59] D. Mohler and S. Schaefer, *Remarks on strange-quark simulations with Wilson fermions*, *Phys. Rev. D* **102** (2020) 074506 [2003.13359].