



# Acute effects of neonicotinoid insecticides on *Mytilus galloprovincialis*: A case study with the active compound thiacloprid and the commercial formulation calypso 480 SC

Alzbeta Stara <sup>a,b</sup>, Maria Pagano <sup>b</sup>, Gioele Capillo <sup>c</sup>, Jacopo Fabrello <sup>d</sup>, Marie Sandova <sup>a</sup>, Marco Albano <sup>b</sup>, Eliska Zuskova <sup>a</sup>, Josef Velisek <sup>a</sup>, Valerio Matozzo <sup>d</sup>, Caterina Faggio <sup>b,\*</sup>

<sup>a</sup> University of South Bohemia in Ceske Budejovice, Faculty of Fisheries and Protection of Waters, South Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, Research Institute of Fish Culture and Hydrobiology, Zatisti 728/I, 389 25, Vodnany, Czech Republic

<sup>b</sup> University of Messina, Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, Viale Ferdinando Stagno D'Alcontres 31, 98166, Messina, Italy

<sup>c</sup> University of Messina, Department of Veterinary Sciences, Polo Universitario Dell'Annunziata, 98168, Messina, Italy

<sup>d</sup> University of Padova, Department of Biology, Via Ugo Bassi 58/B, 35131, Padova, Italy

## ARTICLE INFO

### Keywords:

Pesticide  
Mussels  
Acute toxicity  
Biochemical parameters

## ABSTRACT

Pesticides can enter aquatic environments potentially affecting non-target organisms. Unfortunately, the effects of such substances are still poorly understood. This study investigated the effects of the active neonicotinoid substance thiacloprid (TH) and the commercial product Calypso 480 SC (CA) (active compound 40.4% TH) on *Mytilus galloprovincialis* after short-term exposure to sublethal concentrations. Mussels were tested for seven days to 0, 1, 5 and 10 mg L<sup>-1</sup> TH and 0, 10, 50 and 100 mg L<sup>-1</sup> CA. For this purpose, several parameters, such as cell viability of haemocytes and digestive cells, biochemical haemolymph features, superoxide dismutase (SOD) and catalase (CAT) enzymatic activity of gills and digestive gland, as well as histology of such tissues were analysed. The sublethal concentrations of both substances lead to abatement or completely stopping the byssal fibres creation. Biochemical analysis of haemolymph showed significant changes ( $P < 0.01$ ) in electrolytes ions (Cl<sup>-</sup>, K<sup>+</sup>, Na<sup>+</sup>, Ca<sup>2+</sup>, S-phosphor), lactate dehydrogenase (LDH) enzyme activity and glucose concentration following exposure to both substances. The TH-exposed mussels showed significant imbalance ( $P < 0.05$ ) in CAT activity in digestive gland and gills. CA caused significant decrease ( $P < 0.05$ ) in SOD activity in gills and in CAT activity in both tissues. Results of histological analyses showed severe damage in both digestive gland and gills in a time- and concentration-dependent manner. This study provides useful information about the acute toxicity of a neonicotinoid compound and a commercial insecticide on mussels. Nevertheless, considering that neonicotinoids are still widely used and that mussels are very important species for marine environment and human consumption, further researches are needed to better comprehend the potential risk posed by such compounds to aquatic non-target species.

## 1. Introduction

Wide range of chemicals pollutants from agriculture, industry, households etc. Get into the surface waters through various routes posing a potential risk for aquatic species (Morrissey et al., 2015; Inyinbor et al., 2018). In our study, we focused on neonicotinoid substances that are widely used insecticides in crop production and veterinary medicine since their introduction in the 1990s (Valavanidis, 2018; Craddock et al., 2019). Neonicotinoids popularity is mainly due to their

relatively low risk for vertebrates and high-target specificity to invertebrates, especially insects (Fiorenza et al., 2020). Their toxicity is primarily due to considerable structural differences between vertebrate and insect nicotinic acetylcholine receptors (nAChR). The neonicotinoids have a selective effect on the invertebrates nAChR, at first stimulating postsynaptic receptors, growing Na<sup>+</sup> ingress and K<sup>+</sup> egress and subsequently paralyzing nerve conduction, leading to rapid death (Page, 2008; Crossthwaite et al., 2017). Despite their advantages, starting from 2013 the European Commission blocked the usage of 3

\* Corresponding author.

E-mail address: [cfaggio@unime.it](mailto:cfaggio@unime.it) (C. Faggio).



neonicotinoids (imidacloprid, clothianidin, thiamethoxam) in flowering crops that appeal to honeybees and other pollinators. On April 27, 2018 the EU expanded a controversial ban on these three active substances of insecticides on all crops grown outdoors because of their unfavourable effects on pollinators (Valavanidis, 2018). Thiacloprid ([3-(6-Chloro-3-pyridinylmethyl)-2-thiazolidinylidene]cyanamide) is closely related to imidacloprid both in structure and mode of action, and it is still one of the most commonly used neonicotinoid active substance nowadays (Kammoun et al., 2019). Thiacloprid exhibits high water solubility (184 mg L<sup>-1</sup>) (Thompson et al., 2020) and it is the active substance of several insecticide products, such as Biscaya, Calypso, Proteus etc., to control a variety of chewing and sucking insects (EPA, 2003). Neonicotinoids are relatively stable in the environment. Thiacloprid shows medium persistence in the anaerobic soil and aquatic environment, with a degradation half-life of 3.4–1000 days according to environment conditions (Thompson et al., 2020). The highest measured concentration of thiacloprid in the water was 4.50 µg L<sup>-1</sup> in the Elbe river basin (Supf et al., 2006). However, for other neonicotinoids, higher concentrations up to 320 µg L<sup>-1</sup> for imidacloprid (Van Dijk et al., 2013) and 225 µg L<sup>-1</sup> for acetamiprid (Anderson et al., 2015) have been reported. Due to their properties, high concentrations of neonicotinoids can be expected in rainfall owing to runoff from agriculture treated fields (Velisek and Stara, 2018).

Studies on toxicity of thiacloprid are mainly focused on insects, especially pollinators (Iwasa et al., 2004; Morrissey et al., 2015; Brandt et al., 2016; Raby et al., 2018). However, new studies about the harmful effects of thiacloprid on macroinvertebrates and vertebrates increased in the last few years (Sekeroglu et al., 2014; Hendawi et al., 2016; Salvaggio et al., 2018; Velisek and Stara, 2018; Stara et al., 2019). European Commission adopted a draft rule that would add thiacloprid to the list of neonicotinoids to be avoided in agriculture, mainly for its risk as endocrine disruptors in mammals. However, the approval for thiacloprid is expired on April 30, 2020 (EFSA, 2019).

*Mytilus galloprovincialis* is a very important shellfish species cultured in marine temperate intertidal zones and brackish waters of Europe and is a major food resource for humans (Beyer et al., 2017; FAO, 2019; Figueras et al., 2019; Freitas et al., 2019). Given that mussels continually filter a large volume of seawater, they may be the most seriously affected organisms by various pollutants, such as pesticides (Ayad et al., 2011; Karagiannis et al., 2011; Chmist et al., 2019; Peric and Buric, 2019; Stara et al., 2020), pharmaceuticals (Falfushynska et al., 2019; Freitas et al., 2020a, 2020b, 2020c; Sehonova et al., 2018), nanoparticles (Huang et al., 2018), and UV filters (Vidal-Linan et al., 2018). They are befitting and widely used for monitoring the biological effects of environmental stressors by means of molecular and cellular biomarkers (Bocchetti and Regoli, 2006; Beyer et al., 2017; Burgos-Aceves and Faggio, 2017; Nardi et al., 2017; Figueras et al., 2019; Rahman et al., 2019). In this study, we used mussel specimens from a small brackish meromictic lagoon “Faro Lake”, locally called *Pantano Piccolo*, located on the Sicilian coast of Italy (Bottari et al., 2005; Capillo et al., 2018a, 2018b; Fazio et al., 2015).

Given the widespread use and the evidence of the toxic effects of thiacloprid on non-target organisms and little information about its effects to non-target macro-invertebrates, we have focused our study on this issue. The main objective of this study was to assess the acute effects of sub-lethal doses of the neonicotinoid active substance thiacloprid and the insecticide Calypso 480 SC on *M. galloprovincialis*. We hypothesised that these compounds can have a negative effect on i) cell viability of both haemolymph and digestive gland cells, ii) biochemical parameters of haemolymph, iii) activity of antioxidant biomarkers and iv) histopathology of the gills and digestive glands of mussels. Results of the present study are expected to provide new insights on the effects of mostly used neonicotinoids, focusing on the different sensitivity of the tissue analysed and on the main cellular pathways responsible of neonicotinoids detoxification and cellular homeostasis. Overall, the results obtained may allow a better understanding of *M. galloprovincialis* responsiveness towards neonicotinoids.

## 2. Materials and methods

### 2.1. Test organisms and chemicals

*Mytilus galloprovincialis* was supplied by a mollusc breed cooperative farm Company FARAU SRL (Frutti di Mare) Messina, Italy (Spinelli et al., 2018) from Faro Lake. Faro Lake is the deepest meromictic coastal basin in Italy reaching ~28 m depth in the central part, while it does depth from 0.50 m to 5 m around the coastal areas. Mean physical and chemical parameters of seawater are: temperature 20.3 ± 5.8 °C, pH 8.5 ± 0.3 and salinity 33.0 ± 3.5 during year (Capillo et al., 2018a c). Due to its properties, Faro Lake is a very productive environment, especially for aquaculture productions of mussels, clams or oysters, and concurrently it is part of the Naturel Oriented Reserve of Capo Peloro (Manganaro et al., 2011; Capillo et al., 2018a c). Mussels were transported to the laboratory, and acclimated for 5 days in large tanks contained continuously aerated natural brackish water with a 12 L:12D light cycle and temperature of 19.45 ± 0.98 °C.

Technical-grade thiacloprid (TH; purity 99.9%, Sigma Aldrich, Czech Republic) and pesticide product Calypso 480 SC (CA; contains active substance 40.4% thiacloprid; Bayern Crop Science Corporation) were dissolved in distilled water to obtain stock solutions.

### 2.2. Acute test conditions

Mussels, 6.85 ± 0.57 cm shell length, were randomly selected and placed into sixteen tanks each containing 20 L continuously aerated brackish water. Sixteen mussels per duplicated tank ( $n = 32$ ) were exposed for seven days to concentrations of thiacloprid: 1 mg L<sup>-1</sup> (TH1), 5 mg L<sup>-1</sup> (TH2) and 10 mg L<sup>-1</sup> (TH3); and Calypso 480 SC: 10 mg L<sup>-1</sup> (CA1), 50 mg L<sup>-1</sup> (CA2) and 100 mg L<sup>-1</sup> (CA3). Test concentrations were selected with respect to the active substance thiacloprid content of the Calypso 480 SC and based on data from the literature (Dondero et al., 2010; Stara et al., 2020). The authors describe the effect these chemicals at these concentrations on *M. galloprovincialis* but in while monitoring another kind of indicators. Control groups without tested substances (CTH, CCA) were also prepared. Natural seawater and exposure concentrations have been restored every 2 days to provide natural brackish food and preclude starvation during the exposure. Behavioural changes and mortality of the mussels, as well as water quality (temperature 18.17 ± 0.80 °C, pH 7.65 ± 0.13, salinity 34.31 ± 0.53, 1100 m Osm kg<sup>-1</sup>), were measured daily using a multiparametric probe as reported by Sanfilippo et al. (2016). Water samples were taken during the trial period (at time 0, after 24 and 48 h) for determination of actual concentrations of TH. The TH concentrations were analysed by liquid chromatography mass spectrometry (LC-MS) (Ying and Kookana, 2004) and the mean concentration of TH in the tanks was always within ±5% of the nominal concentration. Haemolymph and tissue (digestive gland, gills) samples were collected after 3 and 7 days of exposures of mussels.

### 2.3. Haemolymph collection

Pools of haemolymph (each pool consists of 3 animals) from each experimental condition were prepared. The pool was needed to obtain enough volumes for analysis. The haemolymph was collected from the anterior adductor muscle with a 23-gauge needle of a 1 ml plastic syringe. Once collected, it was placed in tubes and immediately centrifuged at 1000 rpm for 10 min (Pagano et al., 2017).

### 2.4. Isolation of digestive cells

The digestive glands of four mussels from each group were isolated to obtain digestive cells according to Pagano et al. (2016).

## 2.5. Cell viability assays

To evaluate cell damage, the cytotoxicity test was performed on both haemocytes and digestive gland cells. Cell viability was assessed by two assays: Trypan Blue (TB) exclusion test and lysosomal membrane stability by neutral red (NR), as described in [Stara et al. \(2020\)](#).

## 2.6. Haemolymph parameters

Eight pooled haemolymph samples (each pool containing three animals) for each group were centrifuged at 1000 rpm for 10 min at 4 °C and supernatant samples were stored at -80 °C. Haemolymph electrolytes composition (chloride, potassium, sodium, calcium, phosphorus, magnesium), lactate dehydrogenase (LDH) activity and glucose (GLU) levels were measured by using a multiparametric analyser (KONELAB 60 THERMO, Milano, Italy).

## 2.7. Determination of enzyme activity

Digestive gland and gill tissues of six mussels from each group were excised and frozen on -80 °C until analyses for determination of enzymatic activities (SOD and CAT). The tissues were individually homogenized on ice in 1 ml of 0.1 mM Tris-HCl buffer (pH 7, 0.15 M KCl, 0.5 M Sucrose, 1 mM EDTA, 1 mM Dithiothreitol and 40 µg L<sup>-1</sup> Aprotinin) per 1 g tissue for 1 min with an Ultra-Turrax homogeniser (model T8 basic, IKA). Homogenates were then centrifuged at 12,000 rpm for 30 min at 4 °C and supernatants were collected and used for enzymatic analyses. The SOD (EC 1.15.1.1) activity was measured using the xanthine oxidase/cytochrome c method according by [Crapo et al. \(1978\)](#). The CAT (EC 1.11.1.6) activity was made by to the method described in [Aebi \(1984\)](#). Both enzyme activities were expressed as U per mg protein. Concentration of protein in supernatant were measured according to [Bradford \(1976\)](#) using bovine serum albumin as the standard.

## 2.8. Histology

Histopathological conditions of gills and digestive glands were assessed taking fractions of tissues from each treatment (see previous sections for detailed information). Two specimens from each exposure conditions were collected. After sampling, immersion of collected samples were performed in Immunofix (paraformaldehyde 4% in phosphate saline buffer, Bio-Optica, Milan, Italy) for 8 h at room temperature. After embedded in paraffin, tissue were sectioned by a microtome (Leica, RM2235). Five µm sections were stained using Hematoxylin and Eosin stain for a qualitative histopathological examination using a light microscope (Leitz Diaplan, Germany). Detailed procedures are reported in [Pagano et al. \(2016\)](#), [Lauriano et al. \(2019\)](#) and [Zaccone et al. \(2015\)](#).

### 2.8.1. Histopathological condition indices

Semi-quantitative weighted indices approach described by [Bernet et al. \(1999\)](#) for fish, with some modifications proposed by [Costa et al. \(2013\)](#) have been applied for the evaluation of each individual histopathological indices ( $I_h$ ). Each alteration was assigned a weight (biological significance) with value ranging between 1 and 3 (maximum severity), and a score (degree of dissemination) with values between 0 (in the case that alteration is not present) and 6 (in the case alteration is diffuse). The formula for the assessment of histopathological condition indices is:

$$I_h = \frac{\sum_1^j w_j a_{jh}}{\sum_1^j M_j}$$

where  $I_h$  is the histopathological condition index for the individual  $h$ ;  $w_j$  the weight of the  $j$ th histopathological alteration;  $a_{jh}$  the score attributed to the  $h$ th individual for the  $j$ th alteration and  $M_j$  is the maximum

attributable value for the  $j$ th alteration, i.e., weight × maximum score. The equation's denominator normalizes  $I_h$  to a value between 0 and 1, this is very useful for relating different situations (ie. different organs). The indices have been generally evaluated for each organ (gills and digestive gland) and subsequently related to "reaction patterns": morphological epithelial modifications (gills), tubule and intertubular tissue alterations for digestive gland. The condition weights used have been based on observations collected in this experiment and partially on literature about both invertebrate [Costa et al. \(2013\)](#) and vertebrate histopathology. In accordance, weight assigned are shown in [Table S1](#).

## 2.9. Statistical analysis

In cell viability assays, Student's t-test was used to evaluate results obtained. The value of  $P < 0.05$  was accepted as significant. The software package Prism 5.00 (Graphpad Software Ltd., La Jolla, CA 92037, USA) was used for statistical analyses. Biomarker data of haemolymph and enzymatic activities were analysed using the STATISTICA 13.4 software package (TIBCO). Significant differences between control and treatment groups for each tissue and time were analysed using the non-parametric Mann-Whitney  $U$  test. The data of  $I_h$  were analysed by one-way ANOVA followed by a Bonferroni post-hoc test for multiple comparisons. A  $P$ -value of less than 0.05 was considered significant.

## 3. Results

### 3.1. Conditions of the test mussels during experiments

Behaviour of animals was normal in both controls and the lowest tested concentrations of TH (1 mg L<sup>-1</sup>) and CA (10 mg L<sup>-1</sup>) during the experiment. Mussels produced byssal fibres that allowed a secure attachment to the bottom of tanks or with each other mussel. Concentrations of 5 and 10 mg L<sup>-1</sup> of TH and 50 and 100 mg L<sup>-1</sup> of CA caused a reduction in byssus fibre production in mussels, inducing detachment of mussels from the bottom and no attach to each other already after 3 days. Only one mussel died at 100 mg L<sup>-1</sup> CA at the end of the experiment.

### 3.2. Cell viability assays

Both cell types (haemocytes and digestive gland cells) showed stability and normal parameters. No significant loss in haemocyte viability was observed at all the concentrations tested of TH and CA ([Table 1](#)). In the same way, no damage to the digestive cells was recorded ([Table 1](#)).

### 3.3. Haemolymph evaluation

Mussels exposed to TH showed significant differences ( $P < 0.01$ ) in haemolymph composition during the exposure period ([Table 2](#)). Concentration of TH1 (1 mg L<sup>-1</sup>) caused a significant increase ( $P < 0.01$ ) in levels of Mg<sup>2+</sup> after 3 days of exposure and glucose during the experiment, compared with control (CTH). The mussels exposed to TH2 (5 mg L<sup>-1</sup>) demonstrated a significant decrease ( $P < 0.01$ ) of haemolymph electrolytes (Cl<sup>-</sup>, K<sup>+</sup>, Ca<sup>2+</sup>) after 3 days exposure, S-phosphor after 7 days exposure, and a significant increase in levels of Mg<sup>2+</sup> and glucose after 3 days exposure, compared with control (CTH). Concentration of TH3 (10 mg L<sup>-1</sup>) caused a significant decrease ( $P < 0.01$ ) of haemolymph electrolytes Cl<sup>-</sup>, Na<sup>+</sup>, Ca<sup>2+</sup>) after 3 days exposure and S-phosphor after 7 days of exposure, together with a significant increased LDH activity after 3 day and glucose levels during the experiment compared with controls (CTH).

Results of haemolymph parameters of mussels exposed to CA are listed in [Table 2](#). Concentrations of CA (10, 50 and 100 mg L<sup>-1</sup>) led to significant decrease ( $P < 0.01$ ) of S-phosphor level compared to their control (CCA) during the experiment. The mussels exposed to CA1 (10 mg L<sup>-1</sup>) showed a significant decrease ( $P < 0.01$ ) of levels of electrolytes

**Table 1**

Percentage of viable digestive gland cells and haemocytes in *Mytilus galloprovincialis* exposed to thiacloprod (TH) and Calypso 480 SC (CA).

| Digestive gland     |             | Treatment and dose          |                             |                             |                              |                             |                              |                              |                               |
|---------------------|-------------|-----------------------------|-----------------------------|-----------------------------|------------------------------|-----------------------------|------------------------------|------------------------------|-------------------------------|
| Cell viability test | Time (days) | CTH (0 mg L <sup>-1</sup> ) | TH1 (1 mg L <sup>-1</sup> ) | TH2 (5 mg L <sup>-1</sup> ) | TH3 (10 mg L <sup>-1</sup> ) | CCA (0 mg L <sup>-1</sup> ) | CA1 (10 mg L <sup>-1</sup> ) | CA2 (50 mg L <sup>-1</sup> ) | CA3 (100 mg L <sup>-1</sup> ) |
| TB (%)              | 3           | 100.00 ± 0.00               | 99.43 ± 0.33                | 97.74 ± 0.08                | 97.86 ± 0.55                 | 100.00 ± 0.00               | 79.63 ± 0.00                 | 97.64 ± 0.00                 | 100.00 ± 0.00                 |
|                     | 7           | 100.00 ± 0.00               | 99.55 ± 0.45                | 94.34 ± 0.23                | 93.17 ± 0.73                 | 100.00 ± 0.00               | 100.00 ± 0.00                | 96.72 ± 0.00                 | 96.99 ± 0.00                  |
| NR (%)              | 3           | 100.00 ± 0.00               | 99.64 ± 0.08                | 97.84 ± 0.27                | 95.80 ± 0.80                 | 100.00 ± 0.00               | 76.99 ± 0.00                 | 96.30 ± 0.14                 | 100.00 ± 0.00                 |
|                     | 7           | 99.39 ± 0.35                | 99.02 ± 0.56                | 81.50 ± 0.75                | 91.55 ± 0.89                 | 100.00 ± 0.00               | 100.00 ± 0.00                | 98.94 ± 0.60                 | 100.00 ± 0.00                 |
| Haemocytes          |             | Treatment and dose          |                             |                             |                              |                             |                              |                              |                               |
| Cell viability test | Time (days) | CTH (0 mg L <sup>-1</sup> ) | TH1 (1 mg L <sup>-1</sup> ) | TH2 (5 mg L <sup>-1</sup> ) | TH3 (10 mg L <sup>-1</sup> ) | CCA (0 mg L <sup>-1</sup> ) | CA1 (10 mg L <sup>-1</sup> ) | CA2 (50 mg L <sup>-1</sup> ) | CA3 (100 mg L <sup>-1</sup> ) |
| TB (%)              | 3           | 98.69 ± 0.86                | 98.79 ± 0.34                | 92.13 ± 1.40                | 98.74 ± 0.48                 | 100.00 ± 0.00               | 92.43 ± 0.46                 | 97.84 ± 0.79                 | 96.90 ± 0.94                  |
|                     | 7           | 100.00 ± 0.00               | 97.85 ± 0.35                | 72.12 ± 0.21                | 80.21 ± 0.71                 | 96.26 ± 0.53                | 82.77 ± 0.63                 | 94.96 ± 0.76                 | 96.41 ± 1.24                  |
| NR (%)              | 3           | 98.45 ± 0.61                | 98.86 ± 0.74                | 98.71 ± 0.29                | 97.41 ± 1.23                 | 100.00 ± 0.00               | 76.19 ± 0.74                 | 100.00 ± 0.00                | 99.60 ± 0.40                  |
|                     | 7           | 99.70 ± 0.20                | 98.00 ± 0.26                | 93.42 ± 1.18                | 89.62 ± 0.75                 | 99.77 ± 0.15                | 100.00 ± 0.00                | 99.36 ± 0.42                 | 100.00 ± 0.00                 |

The values are mean ± SE (standard error); n = 6. Trypan blue (TB), neutral red (NR).

**Table 2**

Haemolymphatic parameters of *Mytilus galloprovincialis* exposed to thiacloprod (TH) and Calypso 480 SC (CA).

| Thiacloprod (dose)            | Time (days) | Cl <sup>-</sup> (mmol L <sup>-1</sup> ) | K <sup>+</sup> (mmol L <sup>-1</sup> ) | Na <sup>+</sup> (mmol L <sup>-1</sup> ) | Ca <sup>2+</sup> (mg dl <sup>-1</sup> ) | S-phosphor (mg dl <sup>-1</sup> ) | Mg <sup>2+</sup> (mg dl <sup>-1</sup> ) | LDH (U L <sup>-1</sup> )  | Glucose (mg dl <sup>-1</sup> ) |
|-------------------------------|-------------|---|--|---|---|-----------------------------------|---|---------------------------|--------------------------------|
| CTH (0 mg L <sup>-1</sup> )   | 3           | 473.18 ± 2.04 <sup>a</sup>              | 12.78 ± 0.15 <sup>a</sup>              | 536.50 ± 0.43 <sup>a</sup>              | 56.83 ± 0.15 <sup>a</sup>               | 0.90 ± 0.06 <sup>a</sup>          | 23.90 ± 0.07 <sup>a</sup>               | 4.50 ± 0.22 <sup>a</sup>  | 1.37 ± 0.17 <sup>a</sup>       |
|                               | 7           | 472.03 ± 2.17 <sup>a</sup>              | 11.83 ± 0.17 <sup>a</sup>              | 545.17 ± 1.94 <sup>a</sup>              | 53.55 ± 0.11 <sup>a</sup>               | 0.75 ± 0.02 <sup>ab*</sup>        | 24.85 ± 0.06 <sup>a</sup>               | 1.50 ± 0.22 <sup>a*</sup> | 2.60 ± 0.08 <sup>a*</sup>      |
| TH1 (1 mg L <sup>-1</sup> )   | 3           | 465.35 ± 1.05 <sup>a</sup>              | 12.30 ± 0.04 <sup>a</sup>              | 541.33 ± 0.67 <sup>a</sup>              | 56.33 ± 0.25 <sup>a</sup>               | 0.85 ± 0.02 <sup>a</sup>          | 25.15 ± 0.04 <sup>b</sup>               | 4.50 ± 0.22 <sup>a</sup>  | 3.07 ± 0.12 <sup>b</sup>       |
|                               | 7           | 463.03 ± 0.32 <sup>a</sup>              | 11.87 ± 0.16 <sup>a</sup>              | 543.50 ± 4.24 <sup>a</sup>              | 53.48 ± 0.23 <sup>a</sup>               | 0.87 ± 0.02 <sup>a</sup>          | 25.05 ± 0.04 <sup>a</sup>               | 1.17 ± 0.17 <sup>a*</sup> | 3.88 ± 0.47 <sup>b</sup>       |
| TH2 (5 mg L <sup>-1</sup> )   | 3           | 423.10 ± 7.48 <sup>b</sup>              | 10.77 ± 0.33 <sup>b*</sup>             | 553.50 ± 2.00 <sup>a</sup>              | 51.52 ± 1.43 <sup>*b</sup>              | 0.85 ± 0.02 <sup>a</sup>          | 24.85 ± 0.12 <sup>b</sup>               | 4.50 ± 0.22 <sup>a</sup>  | 2.15 ± 0.26 <sup>ab</sup>      |
|                               | 7           | 458.45 ± 0.52 <sup>a</sup>              | 12.25 ± 0.06 <sup>a</sup>              | 536.33 ± 1.91 <sup>a</sup>              | 52.73 ± 0.15 <sup>a</sup>               | 0.63 ± 0.06 <sup>b*</sup>         | 24.43 ± 0.18 <sup>a</sup>               | 1.17 ± 0.17 <sup>a*</sup> | 2.03 ± 0.04 <sup>a</sup>       |
| TH3 (10 mg L <sup>-1</sup> )  | 3           | 449.62 ± 10.20 <sup>b</sup>             | 12.03 ± 0.38 <sup>a</sup>              | 512.17 ± 7.56 <sup>b*</sup>             | 52.92 ± 1.66 <sup>b</sup>               | 0.87 ± 0.02 <sup>a</sup>          | 24.33 ± 0.08 <sup>a</sup>               | 7.33 ± 0.67 <sup>b</sup>  | 2.77 ± 0.14 <sup>b</sup>       |
|                               | 7           | 459.05 ± 0.28 <sup>a</sup>              | 12.65 ± 0.04 <sup>a</sup>              | 544.50 ± 2.79 <sup>a</sup>              | 54.92 ± 0.09 <sup>a</sup>               | 0.68 ± 0.05 <sup>b*</sup>         | 25.35 ± 0.13 <sup>a</sup>               | 1.50 ± 0.22 <sup>a*</sup> | 3.30 ± 0.46 <sup>b</sup>       |
| Calypso 480 SC (dose)         | Time (days) | Cl <sup>-</sup> (mmol L <sup>-1</sup> ) | K <sup>+</sup> (mmol L <sup>-1</sup> ) | Na <sup>+</sup> (mmol L <sup>-1</sup> ) | Ca <sup>2+</sup> (mg dl <sup>-1</sup> ) | S-phosphor (mg dl <sup>-1</sup> ) | Mg <sup>2+</sup> (mg dl <sup>-1</sup> ) | LDH (U L <sup>-1</sup> )  | Glucose (mg dl <sup>-1</sup> ) |
| CCA (0 mg L <sup>-1</sup> )   | 3           | 494.30 ± 0.81 <sup>a</sup>              | 12.57 ± 0.30 <sup>a</sup>              | 498.17 ± 8.03 <sup>a</sup>              | 53.25 ± 0.53 <sup>a</sup>               | 2.22 ± 0.09 <sup>a</sup>          | 24.40 ± 0.11 <sup>a</sup>               | 2.00 ± 0.37 <sup>a</sup>  | 2.93 ± 0.28 <sup>ab</sup>      |
|                               | 7           | 454.62 ± 0.78 <sup>a*</sup>             | 14.52 ± 0.06 <sup>a</sup>              | 560.00 ± 1.06 <sup>a*</sup>             | 56.75 ± 0.56 <sup>a</sup>               | 2.45 ± 0.08 <sup>a</sup>          | 25.65 ± 0.06 <sup>a</sup>               | 2.83 ± 0.54 <sup>a</sup>  | 4.42 ± 0.26 <sup>a</sup>       |
| CA1 (10 mg L <sup>-1</sup> )  | 3           | 453.35 ± 0.46 <sup>c</sup>              | 10.97 ± 0.40 <sup>b</sup>              | 450.17 ± 15.39 <sup>b</sup>             | 50.97 ± 0.70 <sup>b</sup>               | 1.20 ± 0.04 <sup>b</sup>          | 24.65 ± 0.10 <sup>a</sup>               | 4.50 ± 0.22 <sup>b</sup>  | 1.77 ± 0.32 <sup>a</sup>       |
|                               | 7           | 452.87 ± 0.27 <sup>a</sup>              | 13.07 ± 0.08 <sup>b*</sup>             | 541.33 ± 2.51 <sup>a*</sup>             | 62.85 ± 0.29 <sup>*c</sup>              | 1.25 ± 0.06 <sup>b</sup>          | 25.80 ± 0.40 <sup>a</sup>               | 3.50 ± 0.56 <sup>a</sup>  | 2.00 ± 0.18 <sup>b</sup>       |
| CA2 (50 mg L <sup>-1</sup> )  | 3           | 487.08 ± 0.57 <sup>b</sup>              | 10.23 ± 0.30 <sup>b</sup>              | 492.83 ± 4.73 <sup>a</sup>              | 56.73 ± 0.17 <sup>a</sup>               | 1.40 ± 0.15 <sup>b</sup>          | 25.15 ± 0.10 <sup>a</sup>               | 3.00 ± 0.37 <sup>a</sup>  | 2.33 ± 0.18 <sup>ab</sup>      |
|                               | 7           | 445.52 ± 0.95 <sup>b*</sup>             | 12.60 ± 0.06 <sup>b*</sup>             | 537.33 ± 0.92 <sup>a</sup>              | 54.95 ± 0.16 <sup>a</sup>               | 1.32 ± 0.06 <sup>b</sup>          | 25.70 ± 0.17 <sup>a</sup>               | 1.33 ± 0.21 <sup>b*</sup> | 1.37 ± 0.16 <sup>b</sup>       |
| CA3 (100 mg L <sup>-1</sup> ) | 3           | 498.12 ± 3.51 <sup>a</sup>              | 13.02 ± 0.29 <sup>a</sup>              | 516.67 ± 4.26 <sup>a</sup>              | 55.12 ± 0.59 <sup>a</sup>               | 1.52 ± 0.20 <sup>b</sup>          | 25.30 ± 0.06 <sup>a</sup>               | 5.50 ± 0.22 <sup>b</sup>  | 3.78 ± 0.52 <sup>b</sup>       |
|                               | 7           | 458.47 ± 1.58 <sup>a*</sup>             | 14.05 ± 0.08 <sup>a</sup>              | 549.83 ± 6.93 <sup>a</sup>              | 59.47 ± 0.19 <sup>b*</sup>              | 1.35 ± 0.06 <sup>b</sup>          | 25.08 ± 0.06 <sup>a</sup>               | 1.50 ± 0.34 <sup>b*</sup> | 3.37 ± 0.68 <sup>a</sup>       |

The values are mean ± SE (standard error); n = 6. Different superscripts indicate significant differences ( $P < 0.01$ ) among groups at the same sample time. \*Denotes significant differences among group values over time ( $P < 0.01$ ).

ions Cl<sup>-</sup>, Na<sup>+</sup>, Ca<sup>2+</sup> after 3 days, K<sup>+</sup> and glucose after 7 days and significant increase ion Ca<sup>2+</sup> the seventh day of exposure compared with control (CCA). The concentration of 50 mg L<sup>-1</sup> (CA2) caused a significant decrease ( $P < 0.01$ ) in LDH activity after 7 days of exposure and electrolytes ions (Cl<sup>-</sup>, K<sup>+</sup>) and glucose during the experiment compared to control (CCA). Mussels kept at CA3 (100 mg L<sup>-1</sup>) demonstrated significantly increased ( $P < 0.01$ ) glucose levels, LDH activity after 3 days and Ca<sup>2+</sup> after 7 days of exposure and significant decrease in LDH activity at the end of experiment, compared with control (CCA). Values of Mg<sup>2+</sup> in CA-exposed mussels did not differ significantly between the groups during the exposure.

### 3.4. Enzyme activity

In the present study, TH was shown to affect moderately CAT activity, whereas no statistically significant ( $P < 0.05$ ) effects on SOD activity were found (Fig. 1). CAT activity was significantly increased ( $P < 0.05$ ) in gills after 3 days of exposure to 10 mg L<sup>-1</sup> (TH3) and decreased in digestive gland after 7 days at 5 mg L<sup>-1</sup> (TH2), compared to controls. Calypso caused statistically significant changes ( $P < 0.05$ ) in CAT and SOD activities in mussels (Fig. 1). In digestive gland, CAT activity was significantly reduced ( $P < 0.05$ ) after 3 days at the highest concentration tested (CA3, 100 mg L<sup>-1</sup>). In gills, enzyme activity was


significantly decreased ( $P < 0.05$ ) in mussels exposed for 7 days to CA (10, 50 and 100 mg L<sup>-1</sup>), compared to control mussels. SOD activity was significantly reduced ( $P < 0.05$ ) in gills from mussels exposed for 3 days to three CA concentrations tested (10, 50 and 100 mg L<sup>-1</sup>), compared to related controls (Fig. 1).

### 3.5. Histology

Results of histological observations (Figs. 2–5) showed a considerable exposure effect to toxicants in a dose and time-dependent way for both tissues, with respect to the controls. Results of histopathological condition index (I<sub>h</sub>) are reported in Table S2 and plotted in Figs. 2–5. Statistical analyses showed significant differences among the three treated groups for both compounds; while, no statistically remarkable differences were obtained comparing the I<sub>h</sub> of different tissues (digestive tubule modifications vs intertubular tissue alterations; gills vs digestive gland).

#### 3.5.1. Digestive gland

For both toxicants, the most represented digestive gland tissue abnormalities comprised digestive tubule changes like degeneration of epithelial layer, altered anatomy and atrophy tendency and intertubular tissue modifications with massive haemocytes infiltration, presence of




**Fig. 1.** Enzymatic activity of catalase (CAT) and superoxide dismutase (SOD) in digestive gland and gill in *Mytilus galloprovincialis* exposed to thiacloprid (TH) and Calypso 480 SC (CA). The values are mean  $\pm$  SE (standard error); n = 6; Mann-Whitney U Test. \*Denotes significant differences ( $P < 0.05$ ) among groups at the same sample time.

brown cells and lipofuscin aggregates. Tubule regression, fibrosis, focal points of necrosis and nodular inflammation were the most common alterations recorded (Figs. 2–4).

### 3.5.2. Gills

Gill tissues indicate severe alterations following both CA and TH

exposures. The main histological alterations were haemocytes infiltration, epithelial detachment, vacuolation and lipofuscin deposits (Figs. 3–5). The higher values of  $I_h$  were achieved for CA3 and TH3 groups after 7 days of exposure, as also resulted for digestive gland  $I_h$  evaluation.



**Fig. 2.** *Mytilus galloprovincialis* digestive gland (DG) sections from Calypso 480 SC exposure: representative figures. a-b-c-d are respectively control, 10, 50, 100 mg L<sup>-1</sup>, 3 days exposure; e-f-g-h are respectively control, 10, 50, 100 mg L<sup>-1</sup>, 7 days exposure. Scale bars: 20 µm. Digestive tubule alteration is indicated by green circles, blue arrows indicate brown cells, orange arrows indicate granulocytoma, black arrows indicate haemocytes infiltrations, black filled circles indicate hyperplasia, asterisks indicate hypertrophy, arrowheads indicate lipofuscin aggregates, black square indicates fibrosis, nodular inflammation is widely diffused in d. “D” indicates duct, “dt” indicates digestive tubule. Insets in Figs b and d showed some additional details of haemocytes infiltrations from a different part of the related sample.  $I_h$  represents the graphical elaborations of histopathological condition index of DG from *Mytilus galloprovincialis* exposed to different concentration of Calypso 480 SC (CA): 10 mg L<sup>-1</sup> (CA1), 50 mg L<sup>-1</sup> (CA2) and 100 mg L<sup>-1</sup> (CA3). The values are mean ± SD, <sup>a</sup>P < 0.0332, <sup>b</sup>P < 0.0021, <sup>c</sup>P < 0.0002, <sup>d</sup>P < 0.0001 comparing the exposed samples (One-way ANOVA test/Bonferroni's multiple comparisons test). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

#### 4. Discussion


##### 4.1. Conditions of mussels during experiments

With increasing of exposure time, the effects of both chemicals, at concentrations of TH 5 and 10 mg L<sup>-1</sup> and CA 50 and 100 mg L<sup>-1</sup>, became more evident on physiological state of mussels. *Mytilus* mussels are sessile organisms that attach themselves to the various surfaces by the producing collagenous byssus. The number or strength of the fibres may vary depending on environment conditions (Roberts, 1975; Bell and Gosline, 1996). Mussels must be in good condition and have a large amount of energy which it is necessary to open the shells and to produce byssus. Consequently, all tissues are directly exposed to the environment and this can lead to contact with xenobiotics or molecules responsible for oxidative stress (Bell and Gosline, 1996; Burnett and Sara, 2019). Mussels protect themselves by closing the shells and reducing metabolism, helping them to cope with unfavourable conditions (Bell and Gosline, 1996; Ayad et al., 2011; Savorelli et al., 2017; Chmist et al., 2019) and simultaneously they reduce or completely stop fibres production (Bell and Gosline, 1996; Ayad et al., 2011). The apparent change fibres production under pesticide exposure has been

reported in previous studies on mussels by Ayad et al. (2011), Karagiannis et al. (2011) and Stara et al. (2020). Shells opening and associated fibres formation are first good biomarkers for monitoring weak health status of mussels (Burnett and Sara, 2019; Chmist et al., 2019; Stara et al., 2020).

##### 4.2. Cell viability assays

Viability tests are useful tools for a preliminary evaluation of the possible damage produced by pollutants to organisms. The integrity of the cell membrane assessed by Trypan Blue showed that immunocytes (haemocytes) and the cells liable for the metabolism of the whole organism (digestive gland cells) still manage to perform their tasks. This result is confirmed by the Neutral Red test. Indeed, cells were able to absorb the dye through pinocytosis and therefore reach the lysosomes in order to evaluate their stability (Aguirre-Martínez et al., 2013; Aliko et al., 2015). Lysosomes are important targets because they are the second line of defense against oxidative stress (Burgos-Aceves et al., 2017; Domouhtsidou et al., 2004). At the cellular level, both pollutants at all the concentrations tested did not affect cell viability. Generally, after exposure to xenobiotics, both haemocytes and cells of the digestive



**Fig. 3.** *Mytilus galloprovincialis* gills sections from Calypso 480 SC exposure: representative figures. a-b-c-d are respectively control, 10, 50, 100 mg L<sup>-1</sup>, 3 days exposure; e-f-g-h are respectively control, 10, 50, 100 mg L<sup>-1</sup>, 7 days exposure. Scale bars: 20  $\mu$ m. Green circles indicate vacuolation, blue arrows indicate epithelial alterations, black arrows indicate haemocytes infiltrations, and arrowheads indicate lipofuscin aggregates.  $I_h$  represents the graphical elaborations of histopathological condition index of gills from *Mytilus galloprovincialis* exposed to different concentration of Calypso 480 SC (CA): 10 mg L<sup>-1</sup> (CA1), 50 mg L<sup>-1</sup> (CA2) and 100 mg L<sup>-1</sup> (CA3). The values are mean  $\pm$  SD, <sup>a</sup>P < 0.0332, <sup>b</sup>P < 0.0021, <sup>c</sup>P < 0.0002, <sup>d</sup>P < 0.0001 comparing the exposed samples (One-way ANOVA test/Bonferroni's multiple comparisons test). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)


gland begin to be damaged and the viability percentage decreases (Faggio et al., 2016; Pagano et al., 2017; Stara et al., 2020; Torre et al., 2013). In addition, cell viability for both cell types decreased in chronic exposure to lower CA concentrations (Stara et al., 2020).

Mussels exhibit different behaviors than other aquatic organisms exposed to the same pollutants (Stara et al., 2019; Velisek and Stara, 2018; Beketov et al., 2008). This is typically evident after exposure to high levels of heavy metals (Kraak et al., 1997), mussels closing their shells and reducing the byssus production (Roberts, 1975). This therefore prevents the entry of water and both pollutants did not reach haemolymph or the digestive gland.

#### 4.3. Haemolymph biochemical composition

Nutrients, respiratory gases, metabolic wastes, enzymes, and toxicants are transported by haemolymph throughout the series of tissues body of bivalves and its biochemical investigations provide information relevant to health assessment (Gustafson et al., 2005; Matozzo et al., 2016), similarly to determination of serum biochemical parameters of fish (Qadir et al., 2014; Capillo et al., 2018a a). The biochemical composition of haemolymph varied identifiably between control and mussels exposed to TH (5 and 10 mg L<sup>-1</sup>) and CA (10, 50 and 100 mg L<sup>-1</sup>) during acute exposure. These toxicants caused most changes in

haemolymph electrolytes (Cl<sup>-</sup>, K<sup>+</sup>, Na<sup>+</sup>, Ca<sup>2+</sup>, S-phosphor), glucose concentration and LDH activity. Molluscs, as osmo-comform organisms, resist changes in the environment variation of adaptation processes, such as regulation of haemolymph ions concentration (Capillo et al., 2018a a). Glucose is the main source of energy and increased concentration is indicative of stress in aquatic organisms exposed to environment contaminants. A very marked reduction of glucose indicates acute body failure due to sudden glycogen depletion (Kulkarni and Barad, 2015). Similarly, changes in activity of the LDH is considered as a good diagnostic biomarkers of cell membrane damage caused by pollutants (Kolarova and Velisek, 2012; Ateyya et al., 2015; Faggio et al., 2016). Given the importance of haemolymph evaluation, there were many studies dealing with the evaluation of bivalve haemolymph (Lopes-Lima et al., 2012; Fritts et al., 2015; Faggio et al., 2016; Capillo et al., 2018a a), little is focused on exposure to pesticides (Ayad et al., 2011), insecticides, such as neonicotinoids (Stara et al., 2020). Stara et al. (2020) observed negative effects of CA (7.77 and 77.70 mg L<sup>-1</sup>) on composition of haemolymph *M. galloprovincialis* during 20 days exposure. Simultaneously, changes in these parameters were persistent even after 10 days of depuration in CA-free water. Although neonicotinoids are considered non-toxic to fish, similarly to the changes caused by CA in haemolymph of mussels, negative effects on serum biochemical parameters of fish have been also reported (Qadir et al., 2014; Vieira et al., 2018).




**Fig. 4.** *Mytilus galloprovincialis* digestive gland sections from thiacloprid exposure: representative figures. a-b-c-d are respectively control, 1, 5, 10 mg L<sup>-1</sup>, 3 days exposure; e-f-g-h are respectively control, 1, 5, 10 mg L<sup>-1</sup>, 7 days exposure. Scale bars: controls 25 µm, b-c-d-f-g-h 20 µm. Digestive tubule alteration is indicated by green circles, blue arrows indicate brown cells, black arrows indicate haemocytes infiltrations, orange arrows indicate granulocytoma, black filled circles indicate hyperplasia, asterisks indicate hypertrophy, arrowheads indicate lipofuscin aggregates, black square indicates fibrosis. “D” indicates duct, “dt” indicates digestive tubule.  $I_h$  represents the graphical elaborations of histopathological condition index of DG from *Mytilus galloprovincialis* exposed to different concentration of thiacloprid (TH): 1 mg L<sup>-1</sup> (TH1), 5 mg L<sup>-1</sup> (TH2) and 10 mg L<sup>-1</sup> (TH3). The values are mean  $\pm$  SD, <sup>a</sup>P < 0.0332, <sup>b</sup>P < 0.0021, <sup>c</sup>P < 0.0002, <sup>d</sup>P < 0.0001 comparing the exposed samples (One-way ANOVA test/Bonferroni's multiple comparisons test). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

#### 4.4. Determination of enzyme activities

Assessment of enzyme activities has been considered as useful biomarkers of exposure of aquatic species to pollutants (Fiorino et al., 2018; Matozzo et al., 2018; Phalova et al., 2018; et al., 2019; Vieira et al., 2018). Antioxidant enzymes play an effective antioxidant role by scavenging preferentially reactive oxygen species (ROS) and maintain a steady state in the cells (Faria et al., 2009; Vidal-Linan et al., 2016). The CAT and SOD enzymes are the first line of defense against ROS, protecting organisms from oxidative damage (Burgos-Aceves et al., 2018; Chi et al., 2019). They are found virtually in all aerobic organisms (Ewere et al., 2019). SOD catalyses the conversion of O<sub>2</sub><sup>-</sup> to O<sub>2</sub> and H<sub>2</sub>O<sub>2</sub>, whereas CAT catalyses the conversion of H<sub>2</sub>O<sub>2</sub> to H<sub>2</sub>O and O<sub>2</sub> (Somani et al., 1997). Other biotransformation enzymes are glutathione-S-transferase (GST), glutathione peroxidase (GPx), glutathione reductase (GR), and nonenzymatic biomarkers as reduced glutathion (GSH), vitamins etc. They coordinate important cellular defence mechanisms against oxidative stress which can be impacted by xenobiotic chemicals (Banni et al., 2015; Vidal-Linan et al., 2016; Vieira et al., 2018; Ewere et al., 2019; Uluturhan et al., 2019). Gill is a highly potent metabolic active tissue and is the first barrier which faced to xenobiotics in mussels (Vidal-Linan and Bellas, 2013) and they are often

used to assess the responses of mussels to environmental stressors (Banni et al., 2015). The digestive gland of molluscs is an equally important detoxifying organ aimed at metabolic regulation, providing protection through the processes of detoxification and xenobiotic elimination (Marigomez et al., 2002; Ewere et al., 2019; Faggio et al., 2018). In the present study, TH caused an increase in CAT activity in gill at TH3 (10 mg L<sup>-1</sup>) after 3 days of exposure and a decrease in digestive gland at TH2 (5 mg L<sup>-1</sup>) after 7 days. Mussels exposed to CA showed a reduction of CAT activity in digestive gland at CA3 (100 mg L<sup>-1</sup>) after 3 days and in gills at all the experimental concentrations tested (10, 50 and 100 mg L<sup>-1</sup>) after 7 days. Similarly, all concentrations of CA decreased SOD activity in gills after 3 days. Likewise, recent studies have shown that pesticides inhibit enzymatic activity in mussels. Matozzo et al. (2018) demonstrated that glyphosate (100 and 1000 µg L<sup>-1</sup>) is able to affect SOD activity in digestive gland of mussel after 7 days exposure. Similarly, Stara et al. (2020) recorded a decreased SOD activity in digestive gland of mussels exposed to CA at 7.77 mg L<sup>-1</sup> after 20 days and the activity was then balanced after 10 days in CA-free water. Ewere et al. (2019) described the negative impacts of imidacloprid and Spectrum 200SC on enzymatic activities in *Saccostrea glomerata* after two weeks of exposure at environmentally relevant concentrations (0.01 and 0.05 mg L<sup>-1</sup>). In this study, digestive gland CAT and GST activity, as well as gill



**Fig. 5.** *Mytilus galloprovincialis* gills sections from thiacloprid exposure: representative figures. a-b-c-d are respectively control, 1, 5, 10 mg L<sup>-1</sup>, 3 days exposure; e-f-g-h are respectively control, 1, 5, 10 mg L<sup>-1</sup>, 7 days exposure. Scale bars: 20 µm. Green circles indicate vacuolation, blue arrows indicate epithelial alterations, black arrows indicate haemocytes infiltrations, and arrowheads indicate lipofuscin aggregates.  $I_h$  represents the graphical elaborations of histopathological condition index of gills from *Mytilus galloprovincialis* exposed to different concentration of thiacloprid (TH): 1 mg L<sup>-1</sup> (TH1), 5 mg L<sup>-1</sup> (TH2) and 10 mg L<sup>-1</sup> (TH3). The values are mean  $\pm$  SD, <sup>a</sup>P < 0.0332, <sup>b</sup>P < 0.0021, <sup>c</sup>P < 0.0002, <sup>d</sup>P < 0.0001 comparing the exposed samples (One-way ANOVA test/Bonferroni's multiple comparisons test). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

GST activity were markedly affected by the exposure. In addition, neonicotinoids can induce antioxidant imbalance and provoke oxidative stress in fish and crayfish. Velisek and Stara (2018) described effect of TH at concentration of 4.50 µg L<sup>-1</sup> on antioxidant activity of SOD and GR during embryo-larval development of *Cyprinus carpio*. Vieira et al. (2018) observed differences in antioxidant biomarkers SOD, CAT, GSH, GPx and GST in tissues of *Prochilodus lineatus* exposed for 5 days to imidaclorpid at concentrations of 1.25–1250 µg L<sup>-1</sup>. These alterations in the antioxidants can lead to high levels of lipid peroxidation (LPO) and protein carbonyl content production, which indicates overproduction of ROS and a state of oxidative stress (Stara et al., 2019).

#### 4.5. Histology

Digestive gland is the principal site for xenobiotics detoxification in bivalve (Minier et al., 2000; Gomes et al., 2012) and it is also involved in bioaccumulation of xenobiotics and in challenging environmental and/or induced stressors (Costa et al., 2013). Gills tissue is the first contact surfaces between the organisms and the aquatic environment and are the organs responsible for respiratory function and nutrient assimilation, as for the uptake of contaminants (Azevedo et al., 2015). For these reasons, these tissues have been selected for determining of histological alteration using histopathological condition index as

suggested by Costa et al. (2013). Higher values of  $I_h$  were obtained for both tissues, as expected, in the higher concentration experimental condition at 7 days of exposure. The  $I_h$  took in account several histological abnormalities and alteration as reported in Table S1. Alterations in digestive tubule and intertubular tissues were recorded with no important differences between them.

Among the various alterations, melanosis widely occurred. Melanosis is a reported alteration in mussels as in other organisms. This condition reflects an inflammatory process related to the exposure to toxicants. Melanosis derives, in fact, from a series of events known as “pro-phenoloxidase (PO) activating systems”. The PO activating system is responsible of recognition, cytotoxicity and encapsulation of toxicants (De Vico and Carella, 2012; Luna-Acosta et al., 2010; Pagano et al., 2016). The same histological alteration was observed in *M. galloprovincialis* in a recent study that evaluated the effect of chronic exposure to CA at sub-lethal concentrations (Stara et al., 2020). The other histological abnormalities detected, confirmed a severe inflammatory response of these organisms to the pollutant. For example, it is well known that brown cells, highly present in digestive gland of treated organisms, are involved in recognition, accumulation and detoxification of pollutants (De Vico and Carella, 2012; Usheva and Frolova, 2006). Moreover, nodular inflammation has been recognized in many digestive glands of exposed mussels. Nodular inflammations occur when

haemocytes phagocytose little and abundant particles, forming different sized aggregates in haemolymph and interstices (Galloway and Depledge, 2001). Despite normally nodular formations are a consequence of microbial invasions (Rahmet-Alla and Rowley, 1989), which have shown that the aggregative properties of haemocytes are stimulated by the exposure to xenobiotics and other contaminants (Auffret and Oubella, 1997; Carella et al., 2015).

The pathological inflammatory condition has also been observed in gill tissues, with abnormalities comprehending abundant deposits of melanin, intense vacuolation, haemocytes infiltrations and presence of brown cells. Focal and/or diffuse haemocytes infiltrations in tissues, without the tendency to form aggregates, characterize the infiltrative inflammations (Kim et al., 2006; Kim and Powell, 2004, 2007). Infiltrative haemocytes phagocytose pathogens and foreign bodies in order to activate a reaction to the toxicant, activating the Multixenobiotic defence mechanism (MXDM) (Pain and Parant, 2003). The MXDM can act as protection for cells from negative actions of toxic compounds by reducing access and in promoting efflux of toxicants (Pagano et al., 2016). Haemocytes infiltrations are widely reported in gill tissues of molluscs exposed to environmental pollutants.

## 5. Conclusions

This study investigated for the first time the acute effects of the active neonicotinoid substance thiacloprid and the commercial insecticide Calypso 480 SC on *M. galloprovincialis*. Our results revealed a series of biological alterations following exposure to the compounds. Thiacloprid and also Calypso mainly affected the production of byssus fibres, biomarkers of biochemical profile of haemolymph (glucose, LDH, S-phosphor,  $\text{Ca}^{2+}$ ,  $\text{Cl}^-$ ,  $\text{K}^+$ ,  $\text{Na}^+$ ), activity of antioxidant biomarkers (SOD, CAT) and caused histopathological alterations in the digestive glands and gills. The results also show that thiacloprid in low concentrations has a stronger effect on mussels than an insecticide which containing another concomitant substances. The study of biological responses of exposed organisms is essential for understanding of the mechanisms of toxicity. To better comprehend the mechanisms of action of emerging contaminants in aquatic organisms, it is desirable to find particularly sensitive early warning signals to predict the risks posed by such substances in environment.

## Credit authors statement

Alzbeta Stara Data curation, Writing - review & editing, Maria Pagano Writing - original draft, Investigation, Gioele Capillo Writing - original draft, Jacopo Fabrello Investigation, Marco Albano Writing - original draft, Marie Sandova Data curation, Eliska Zuskova Methodology, Josef Velisek Data curation, Resources, Valerio Matozzo Visualization, Data curation, Supervision, Caterina Faggio Conceptualization, Data curation, Supervision, Writing - review & editing

## Declaration of competing interest

The authors declare that they have no conflict of interest.

## Acknowledgments

The study was financially supported by the Ministry of Education, Youth and Sports of the Czech Republic project CENAKVA (No. LM2018099) and by project Development of University of South Bohemia: International Mobility MSCA IF (No. CZ.02.2.69/0.0/0.0/17\_050/0008486). A special acknowledgement is due to the 'Farau' mussels farm for the practical support and for facilities and moreover, authors thank Dr. Fabiano Capparucci for technical support.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ecoenv.2020.110980>.

## References

- Aebi, H., 1984. Catalase in vitro. Methods Enzymol. 105, 121–126.
- Aguirre-Martínez, G.V., Buratti, S., Fabbri, E., DelValls, A.T., Martín-Díaz, M.L., 2013. Using lysosomal membrane stability of haemocytes in *Ruditapes philippinarum* as a biomarker of cellular stress to assess contamination by caffeine, ibuprofen, carbamazepine and novobiocin. J. Environ. Sci. 25 (7), 1408–1418.
- Aliko, V., Hajdaraj, G., Caci, A., Faggio, C., 2015. Copper induced lysosomal membrane destabilisation in haemolymph cells of mediterranean green crab (*Carcinus aestuarii*, nardo, 1847) from the narta lagoon (Albania). Braz. Arch. Biol. Technol. 58 (5), 750–756.
- Anderson, J.C., Dubetz, C., Palace, V.P., 2015. Neonicotinoids in the Canadian aquatic environment: a literature review on current use products with a focus on fate, exposure, and biological effects. Sci. Total Environ. 505, 409–422.
- Ateyya, H., Yosef, H., Nader, M.A., 2015. Ameliorative effect of trimetazidine on cisplatin-induced hepatotoxicity in rats. Can. J. Physiol. Pharmacol. 94, 225–230.
- Auffret, M., Oubella, R., 1997. Hemocyte aggregation in the oyster *Crassostrea gigas*: in vitro measurement and experimental modulation by xenobiotics. Comp. Biochem. Physiol. Physiol. 118, 705–712.
- Ayad, M.A., Fdil, M.A., Mouabid, A., 2011. Effects of cypermethrin (pyrethroid insecticide) on the valve activity behavior, byssal thread formation, and survival in air of the marine mussel *Mytilus galloprovincialis*. Arch. Environ. Contam. Toxicol. 60, 462–470.
- Azevedo, C.C., Guzmán-Guillén, R., Martins, J.C., Osório, H., Vasconcelos, V., da Fonseca, R.R., Campos, A., 2015. Proteomic profiling of gill GSTs in *Mytilus galloprovincialis* from the North of Portugal and Galicia evidences variations at protein isoform level with a possible relation with water quality. Mar. Environ. Res. 110, 152–161.
- Banni, M., Sforzini, S., Franzellitti, S., Oliveri, C., Viarengo, A., Fabbri, E., 2015. Molecular and cellular effects induced in *Mytilus galloprovincialis* treated with oxytetracycline at different temperatures. PLoS One 10, e0128468.
- Beketov, M.A., Schäfer, R.B., Marwitz, A., Paschke, A., Liess, M., 2008. Long-term stream invertebrate community alterations induced by the insecticide thiacloprid: effect concentrations and recovery dynamics. Sci. Total Environ. 405, 96–108.
- Bell, E.C., Gosline, J.M., 1996. Mechanical design of mussel byssus: material yield enhances attachment strength. J. Exp. Biol. 199, 1005–1017.
- Bernet, D., Schmidt, H., Meier, W., Burkhardt-Holm, P., Wahli, T., 1999. Histopathology in fish: proposal for a protocol to assess aquatic pollution. J. Fish. Dis. 22, 25–34.
- Beyer, J., Green, N.W., Brooks, S., Allan, I.J., Riis, A., Gomes, T., Brate, I.L.N., Schoeney, M., 2017. Blue mussels (*Mytilus edulis* spp.) as sentinel organisms in coastal pollution monitoring: a review. Mar. Environ. Res. 130, 338–365.
- Bocchetti, R., Regoli, F., 2006. Seasonal variability of oxidative biomarkers, lysosomal parameters, metallothioneins and peroxisomal enzymes in the Mediterranean mussel *Mytilus galloprovincialis* from Adriatic Sea. Chemosphere 65, 913–921.
- Bottari, A., Bottari, C., Carveni, P., Giacobbe, S., Spano, N., 2005. Genesis and geomorphologic and ecological evolution of the Ganzirri salt marsh (Messina, Italy). Quat. Int. 140–141, 150–158.
- Bradford, M.M., 1976. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of the protein-dye binding. Anal. Biochem. 72, 248–254.
- Brandt, A., Gorenflo, A., Siegle, R., Meixner, M., Büchler, R., 2016. The neonicotinoids thiacloprid, imidacloprid, and clothianidin affect the immunocompetence of honey bees (*Apis mellifera* L.). J. Insect Physiol. 86, 40–47.
- Burgos-Aceves, M.A., Cohen, A., Smith, Y., Faggio, C., 2017. A potential microRNA regulation of immune-related genes in invertebrate haemocytes. Sci. Total Environ. 621, 302–307.
- Burgos-Aceves, M.A., Faggio, C., 2017. An approach to the study of the immunity functions of bivalve haemocytes: physiology and molecular aspects. Fish Shellfish Immunol. 67, 513–517.
- Burgos-Aceves, M.A., Cohen, A., Smith, Y., Faggio, C., 2018. MicroRNAs and their role on fish oxidative stress during xenobiotic environmental exposures. Ecotoxicol. Environ. Saf. 148, 995–1000.
- Burnett, N.P., Sara, G., 2019. Functional responses of intertidal bivalves to repeated sub-lethal, physical disturbances. Mar. Environ. Res. 147, 32–36.
- Capillo, G., Silvestro, S., Sanfilippo, M., Fiorino, E., Giangrossi, G., Ferrantelli, V., Vazzana, I., Faggio, C., 2018a. Assessment of electrolytes and metals profile of the Faro Lake (Capo Peloro lagoon, Sicily, Italy) and its impact on *Mytilus galloprovincialis*. Chem. Biodivers. 15, e1800044.
- Capillo, G., Savoca, S., Costa, R., Sanfilippo, M., Rizzo, C., Giudice, A., Lo, Albergamo, A., Rando, R., Bartolomeo, G., Spanò, N., Faggio, C., 2018b. New insights into the culture method and antibacterial potential of *Gracilaria gracilis*. Mar. Drugs 16, 1–21.
- Capillo, G., Panarello, P., Savoca, S., Sanfilippo, M., Albano, M., Li Volsi, R., Consolo, G., Spanò, N., 2018c. Intertidal ponds of Messina's beachrock faunal assemblage, evaluation of ecosystem dynamics and communities' interactions. Atti Accad. Pelorit. Pericol. Cl. Sci. Fis. Mat. Nat. 96, 16.
- Carella, F., Feist, S.W., Bignell, J.P., De Vico, G., 2015. Comparative pathology in bivalves: aetiological agents and disease processes. J. Invertebr. Pathol. 131, 107–120.

- Chi, C., Yun, S., Giri, S.S., Kim, H.J., Kim, S.W., Kang, J.W., Park, S.C., 2019. Effect of the agicide Thiazolidinedione 49 on immune responses of bay scallop *Argopecten irradians*. *Molecules* 24, 3579.
- Chmiot, J., Szoszkiewicz, K., Drozdynski, D., 2019. Behavioural responses of *Unio tumidus* freshwater mussels to pesticide contamination. *Arch. Environ. Contam. Toxicol.* 77, 432–442.
- Costa, P.M., Carreira, S., Costa, M.H., Caeiro, S., 2013. Development of histopathological indices in a commercial marine bivalve (*Ruditapes decussatus*) to determine environmental quality. *Aquat. Toxicol.* 126, 442–454.
- Craddock, H.A., Huang, D., Turner, P.C., Quiros-Alcalá, L., Payne-Sturges, D.C., 2019. Trends in neonicotinoid pesticide residues in food and water in the United States, 1999–2015. *Environ. Health* 18, 7.
- Crapo, J.D., McCord, J.M., Fridovich, I., 1978. Preparation and assay of superoxide dismutases. *Methods Enzymol.* 53, 382–393.
- Crossthwaite, A.J., Bigot, A., Camblin, P., Goodchild, J., Lind, R.J., Slater, R., Maienfisch, P., 2017. The invertebrate pharmacology of insecticides acting at nicotinic acetylcholine receptor. *J. Pestic. Sci.* 42, 67–83.
- De Vico, G., Carella, F., 2012. Morphological features of the inflammatory response in molluscs. *Res. Vet. Sci.* 93, 1109–1115.
- Domouhtsidou, G.P., Dailianis, S., Kaloyianni, M., Dimitriadi, V.K., 2004. Lysosomal membrane stability and metallothionein content in *Mytilus galloprovincialis* (L.), as biomarkers: combination with trace metal concentrations. *Mar. Pollut. Bull.* 48, 572–586.
- Dondero, F., Negri, A., Boatti, L., Marsano, F., Mignone, F., Viarengo, A., 2010. Transcriptomic and proteomic effects of a neonicotinoid insecticide mixture in the marine mussel (*Mytilus galloprovincialis*, Lam.). *Sci. Total Environ.* 408, 3775–3786.
- EFSA (European Food Safety Authority), 2019. Peer review of the pesticide risk assessment of the active substance thiacloprid. *EFSA Journal* 17, 5595.
- EPA (Environmental Protection Agency), 2003. Name of Chemical: Thiacloprid Reason for Issuance: Conditional Registration Date Issued: September 26, 2003. United States. Office of Prevention and Toxic Substances (7501C).
- Ewere, E., Reichelt-Brushett, A., Benkendorff, K., 2019. Imidacloprid and formulated product impacts the fatty acids and enzymatic activities in tissues of Sydney rock oysters, *Saccostrea glomerata*. *Mar. Environ. Res.* 151, 104765.
- Faggio, C., Pagano, M., Alampi, R., Vazzana, I., Felice, M.R., 2016. Cytotoxicity, haemolytic parameters, and oxidative stress following exposure to sublethal concentrations of quaternium-15 in *Mytilus galloprovincialis*. *Aquat. Toxicol.* 180, 258–265.
- Faggio, C., Tsarpali, V., Dailianis, S., 2018. Mussel digestive gland as a model tissue for assessing xenobiotics: an overview. *Sci. Total Environ.* 636, 220–229.
- Falfushynska, H., Sokolov, E.P., Haider, F., Oppermann, C., Kragl, U., Ruth, W., Stock, M., Glufke, S., Winkel, E.J., Sokolova, I.M., 2019. Effects of a common pharmaceutical, atorvastatin, on energy metabolism and detoxification mechanisms of a marine bivalve *Mytilus edulis*. *Aquat. Toxicol.* 208, 47–61.
- FAO (Food and Agriculture Organization of the United Nations), 2019. National Aquaculture Sector Overview. Italy (visited online 15.11.2019). [http://www.fao.org/fishery/countrysector/naso\\_italy/en](http://www.fao.org/fishery/countrysector/naso_italy/en).
- Faria, M., Carrasco, L., diez, s., Riva, M.C., Bayona, J.M., Barata, C., 2009. Multibiomarker responses in the freshwater mussel *Dreissena polymorpha* exposed to polychlorobiphenyls and metals. *Comp. Biochem. Physiol.* C 149, 281–288.
- Fazio, F., Saoca, C., Casella, S., Fortino, G., Piccione, G., 2015. Relationship between blood parameters and biometric indices of *Sparus aurata* and *Dicentrarchus labrax* cultured in onshore tanks. *Mar. Freshw. Behav. Physiol.* 48, 289–296.
- Figueras, A., Moreira, R., Sendra, M., Novoa, B., 2019. Genomics and immunity of the Mediterranean mussel *Mytilus galloprovincialis* in a changing environment. *Fish Shellfish Immunol.* 90, 440–445.
- Fiorenza, R., Di Mauro, A., Cantarella, M., Iaria, C., Scalisi, E.M., Bruno, M.V., Gulino, A., Spitaleri, L., Nicotra, G., Dattilo, S., Carroccio, S.C., Privitera, V., Impellizzeri, G., 2020. Preferential removal of pesticides from water by molecular imprinting on TiO<sub>2</sub> photocatalysts. *Chem. Eng. J.* 379, 122309.
- Fiorino, E., Sehonova, P., Phhalova, L., Blahova, J., Svobodova, Z., Faggio, C., 2018. Effect of glyphosate on early life stages: comparison between *Cyprinus carpio* and *Danio rerio*. *Environ. Sci. Pollut. Res.* 25, 8542–8854.
- Freitas, R., Silvestro, S., Coppola, F., Meucci, V., Battaglia, F., Intorre, L., Soares, A.M.V. M., Pretti, C., Faggio, C., 2019. Biochemical and physiological responses induced in *Mytilus galloprovincialis* after a chronic exposure to salicylic acid. *Aquat. Toxicol.* (in press).
- Freitas, R., Silvestro, S., Coppola, F., Meucci, V., Battaglia, F., Intorre, L., Soares, A.M.V. M., Pretti, C., Faggio, C., 2020a. Combined effects of salinity changes and salicylic acid exposure in *Mytilus galloprovincialis*. *Sci. Total Environ.* (in press).
- Freitas, R., Silvestro, S., Coppola, F., Costa, S., Meucci, V., Battaglia, F., Intorre, L., Soares, A.M.V.M., Pretti, C., Faggio, C., 2020b. Toxic impacts induced by Sodium lauryl sulfate in *Mytilus galloprovincialis*. *Comp. Biochem. Physiol. Part A: Molecular Integrated Physiol.* (in press).
- Freitas, R., Silvestro, S., Pagano, M., Coppola, F., Meucci, V., Battaglia, F., Intorre, L., Soares, A.M.V.M., Pretti, C., Faggio, C., 2020c. Impacts of salicylic acid in *Mytilus galloprovincialis* exposed to warming conditions. *Environ. Toxicol. Pharmacol.* (in press).
- Fritts, A.K., Peterson, J.T., Hazelton, P.D., Bringolf, R.B., 2015. Evaluation of methods for assessing physiological biomarkers of stress in freshwater mussels. *Can. J. Fish. Aquat. Sci.* 72, 1450–1459.
- Galloway, T.S., Depledge, M.H., 2001. Immunotoxicity in invertebrates: measurement and ecotoxicological relevance. *Ecotoxicology* 10, 5–23.
- Gomes, T., Pereira, C.G., Cardoso, C., Pinheiro, J.P., Cancio, I., Bebianno, M.J., 2012. Accumulation and toxicity of copper oxide nanoparticles in the digestive gland of *Mytilus galloprovincialis*. *Aquat. Toxicol.* 118–119, 72–79.
- Gustafson, L.L., Stoskopf, M.K., Bogan, A.E., Showers, W., Kwak, T.J., Hanlon, S., Levine, J.F., 2005. Evaluation of a nonlethal technique for hemolymph collection in *Elliptio complanata*, a freshwater bivalve (Mollusca: unionidae). *Dis. Aquat. Org.* 65, 159–165.
- Hendawi, M.Y., Alam, R.T.M., Abdellatif, S.A., 2016. Ameliorative effect of flaxseed oil against thiacloprid-induced toxicity in rats: hematological, biochemical, and histopathological study. *Environ. Sci. Pollut. Res.* 23, 11855–11863.
- Huang, X., Liu, Y., Liu, Z., Zhao, Z., Dupont, S., Wu, F., Huang, W., Chen, J., Hu, M., Lu, W., Wang, Y., 2018. Impact of zinc oxide nanoparticles and ocean acidification on antioxidant responses of *Mytilus coruscus*. *Chemosphere* 196, 182–195.
- Inyinbor, A.A., Adebesin, B.O., Oluwori, A.P., Adelani-Akande, T.A., Dada, A.O., Oreofe, T.A., 2018. Water pollution: effect, prevention, and climatic impact, pp. 33–53 (Chapter 3), in book: *Water Challenges of an Urbanizing World*.
- Iwasa, T., Motoyama, N., Ambrose, J.T., Roe, R.M., 2004. Mechanism for the differential toxicity of neonicotinoid insecticides in the honey bee, *Apis mellifera*. *Crop Protect.* 23, 371–378.
- Kammoun, I., Sellem, I., Saad, H.B., Boudawara, T., Nasri, M., Gharsallah, N., Mallouli, L., Amara, I.B., 2019. Potential benefits of polysaccharides derived from marine alga *Ulva lactuca* against hepatotoxicity and nephrotoxicity induced by thiacloprid, an insecticide pollutant. *Environ. Toxicol.* 34, 1165–1176.
- Karagiannis, D., Vatsos, I.N., Angelidis, P., 2011. Effects of atrazine on the viability and the formation of byssus of the mussel *Mytilus galloprovincialis*. *Aquacult. Int.* 19, 103–110.
- Kim, M.K., Park, K.I., Choi, K.S., Alvarez, R.A., Cummings, R.D., Cho, M., 2006. Lectin from the manila clam *Ruditapes philippinarum* is induced upon infection with the protozoan parasite *Perkinsus olseni*. *J. Biol. Chem.* 281, 26854–26864.
- Kim, Y., Powell, E., 2004. Surfclam histopathology survey along the Delmarva mortality line. *J. Shellfish Res.* 23, 429–441.
- Kim, Y., Powell, E.N., 2007. Distribution of parasites and pathologies in sentinel bivalves: NOAA status and Trends "Mussel Watch" Program. *J. Shellfish Res.* 26, 1115–1151.
- Kolarova, J., Velisek, J., 2012. Determination and evaluation of the biochemical profile of blood fish. *FROV JU Vodnany, Methods No.* 135, 54 (In Czech).
- Kulkarni, R., Barad, V., 2015. Haematological and blood biochemical changes in the fresh water fish, *Notoperus notoperus* (Pallas) exposed to acidic medium. *Int. Lett. Nat. Sci.* 45, 27–33.
- Lauriano, E.R., Pergolizzi, S., Aragona, M., Montalbano, G., Guerrera, M.C., Crupi, R., Faggio, C., Capillo, G., 2019. Intestinal immunity of dogfish *Scyliorhinus canicula* spiral valve: a histochemical, immunohistochemical and confocal study. *Fish Shellfish Immunol.* 87, 490–498.
- Lopes-Lima, M., Freitas, S., Pereira, L., Gouveia, E., Hinzmann, M., Checa, A., Machado, j., 2012. Ionic regulation and shell mineralization in the bivalve *Anodonta cygnea* (swan mussel) following heavy-metal exposure. *Can. J. Zool.* 90, 267–283.
- Luna-Acosta, A., Rosenfeld, E., Amari, M., Fruittier-Arnaudin, I., Bustamante, P., Thomas-Guyon, H., 2010. First evidence of laccase activity in the Pacific oyster *Crassostrea gigas*. *Fish Shellfish Immunol.* 28, 719–726.
- Manganaro, A., Pulicanò, G., Sanfilippo, M., 2011. Temporal evolution of the area of Capo Peloro (Sicily, Italy) from pristine site into urbanized area. *Transitional Waters Bull* 5, 23–31.
- Marigomez, I., Soto, M., Cajaraville, M.P., Angulo, E., Giamberini, L., 2002. Cellular and subcellular distribution of metals in molluscs. *Microsc. Res. Tech.* 56, 358–392.
- Matozzo, V., Pagano, M., Spinelli, A., Caicci, F., Faggio, C., 2016. *Pinnus nobilis*: a big bivalve with big haemocytes? *Fish Shellfish Immunol.* 55, 529–534.
- Matozzo, V., Fabrello, J., Masiero, L., Ferraccioli, F., Finos, L., Pastore, P., Di Gangi, I.M., Bogialli, S., 2018. Ecotoxicological risk assessment for the herbicide glyphosate to non-target aquatic species: a case study with the mussel *Mytilus galloprovincialis*. *Environ. Pollut.* 233, 623–632.
- Minier, C., Borghi, V., Moore, M.N., Porte, C., 2000. Seasonal variation of MXR and stress proteins in the common mussel, *Mytilus galloprovincialis*. *Aquat. Toxicol.* 50, 167–176.
- Morrissey, C.A., Mineau, P., Devries, J.H., Sanchez-Bayo, F., Liess, M., Cavallaro, M.C., Liber, K., 2015. Neonicotinoid contamination of global surfacewaters and associated risk to aquatic invertebrates: a review. *Environ. Int.* 74, 291–303.
- Nardi, A., Mincarelli, L.F., Benedetti, M., Fattorini, D., d'Errico, G., Regoli, F., 2017. Indirect effects of climate changes on cadmium bioavailability and biological effects in the Mediterranean mussel *Mytilus galloprovincialis*. *Chemosphere* 169, 493–502.
- Pagano, M., Capillo, G., Sanfilippo, M., Palato, S., Trischitta, F., Manganaro, A., Faggio, C., 2016. Evaluation of functionality and biological responses of *Mytilus galloprovincialis* after exposure to Quaternium-15 (Methenamine 3-Chlorallylochloride). *Molecules* 21, 144.
- Pagano, M., Porcino, C., Briglia, M., Fiorino, E., Vazzana, M., Silvestro, S., Faggio, C., 2017. The influence of exposure of cadmium chloride and zinc chloride on haemolymph and digestive gland cells from *Mytilus galloprovincialis*. *Int. J. Environ. Res.* 11, 207–216.
- Page, S.W., 2008. Chapter 10 - antiparasitic drugs. *Small Animal Clinical Pharmacology*, second ed. Elsevier Ltd, pp. 198–260.
- Pain, S., Parant, M., 2003. Multixenobiotic defense mechanism (MDMX) in bivalves. *C. R. Biol.* 326, 659–672.
- Peric, L., Buric, P., 2019. The effect of copper and chlorpyrifos co-exposure on biomarkers in the marine mussel *Mytilus galloprovincialis*. *Chemosphere* 225, 126–134.
- Phhalova, L., Blahova, J., Divisova, L., Enevova, V., Casuscelli Di Tocco, F., Faggio, C., Tichy, F., Vecerek, V., Svobodova, Z., 2018. The effects of subchronic exposure to NeemAzal T/S on zebrafish (*Danio rerio*). *Chem. Ecol.* 34 (3), 199–210.
- Qadir, S., Latif, A., Ali, M., Iqbal, F., 2014. Effects of imidacloprid on the hematological and serum biochemical profile of *Labeo rohita*. *Pakistan J. Zool.* 46, 1085–1090.

- Raby, M., Zhao, X., Hao, C., Poirier, D.G., Sibley, P.K., 2018. Chronic toxicity of 6 neonicotinoid insecticides to *Chironomus dilutus* and *Neocloeon triangulifer*. Environ. Toxicol. Chem. 37, 2727–2739.
- Rahman, M.A., Henderson, S., Miller-Ezzy, P., Li, X.X., Qin, J.G., 2019. Immune response to temperature stress in three bivalve species: pacific oyster *Crassostrea gigas*, Mediterranean mussel *Mytilus galloprovincialis* and mud cockle *Katelysia rhytiphora*. Fish Shellfish Immunol. 86, 868–874.
- Rahmet-Alla, M., Rowley, A.F., 1989. Studies on the cellular defense reactions of the Madeira cockroach, *Leucophaea maderae*: nodule formation in response to injected bacteria. J. Invertebr. Pathol. 54, 200–207.
- Roberts, D., 1975. The effect of pesticides on byssus formation in the common mussel, *Mytilus edulis*. Environ. Pollut. 8, 241–254.
- Salvaggio, A., Antoci, F., Messina, A., Ferrante, M., Copat, C., Ruberto, C., Scalisi, E.M., Pecoraro, R., Brundo, M.V., 2018. Teratogenic effects of the neonicotinoid thiacloprid on chick embryos (*Gallus gallus domesticus*). Food Chem. Toxicol. 118, 812–820.
- Sanfilippo, M., Capillo, G., Spanò, N., Manganaro, A., 2016. Evaluation of water variables in no-take zone of ustica marine protected area (Southern Tyrrhenian Sea). Braz. Arch. Biol. Technol. 59, e16160330.
- Savorelli, F., Manfra, L., Croppo, M., Tornambe, A., Palazzi, D., Canepa, S., Trentini, P.L., Cicero, A.M., Faggio, C., 2017. Fitness evaluation of *Ruditapes philippinarum* exposed to nickel. Biol. Trace Elem. Res. 177, 384–393.
- Sehonova, P., Svobodova, Z., Dolezelova, P., Vosmerova, P., Faggio, C., 2018. Effects of waterborne antidepressants on non-target animals living in the aquatic environment: a review. Sci. Total Environ. 631–632, 789–794.
- Sekeroglu, V., Sekeroglu, Z.A., Demirhan, E.S., 2014. Effects of commercial formulations of deltamethrin and/or thiacloprid on thyroid hormone levels in rat serum. Toxicol. Ind. Health 30, 40–46.
- Somani, S.M., Husain, K., Schlorff, E.C., 1997. Response of antioxidant system to physical and chemical stress. In: Baskin, S.I., Salem, H. (Eds.), Oxidants, Antioxidants, and Free Radicals. Taylor Francis, Washington, DC, pp. 125–141.
- Spinelli, A., Capillo, G., Faggio, C., Viatale, D., Spanò, N., 2018. Returning of *Hippocampus hippocampus* (Linnaeus, 1758) (syngnathidae) in the Faro Lake – oriented natural Reserve of Capo Peloro. Italy. Nat. Prod. Res. 1–4.
- Stara, A., Bellinavia, R., Velisek, J., Strouhova, A., Kouba, A., Faggio, C., 2019. Acute exposure of common yabby (*Cherax destructor*) to the neonicotinoid pesticide. Sci. Total Environ. 665, 718–723.
- Stara, A., Pagano, M., Capillo, G., Fabrello, J., Sandova, M., Vazzana, I., Zuskova, E., Velisek, J., Matozzo, V., Faggio, C., 2020. Assessing the effects of neonicotinoid insecticide on the bivalve mollusc *Mytilus galloprovincialis*. Sci. Total Environ. 700, 134914.
- Süß, A., Bischoff, G., Mueller, A.C.W., Buhr, L., 2006. Chemisch-biologisches monitoring zu pflanzenschutzmittelbelastungen und lebensgemeinschaften in graben des alten landes. Nachrichtenbl Deut Pflanzenschutzd 58, 28–42.
- Thompson, D., Lehmler, H.J., Kolpin, D.W., Hladik, M.L., Vargo, J.D., Schilling, K.E., LeFevre, G.H., Peebles, T.L., Poch, M.C., LaDuka, L.E., Cwiertny, D.M., Field, R.W., 2020. A critical review on the potential impacts of neonicotinoid insecticide use: current knowledge of environmental fate, toxicity, and implications for human health. Environ. Sci.: Processes Impacts. <https://doi.org/10.1039/C9EM00586B>.
- Torre, A., Trischitta, F., Faggio, C., 2013. Effect of CdCl<sub>2</sub> on regulatory volume decrease (RVD) in *Mytilus galloprovincialis* digestive cells. Toxicol. Vitro 27 (4), 1260–1266.
- Uluturhan, E., Darilmaz, E., Kontas, A., Bilgin, M., Alyuruk, H., Altay, O., Sevgi, S., 2019. Seasonal variations of multi-biomarker responses to metals and pesticides pollution in *M. galloprovincialis* and *T. decussatus* from Homa Lagoon, Eastern Aegean Sea. Mar. Pollut. Bull. 141, 176–186.
- Usheva, L.N., Frolova, L.T., 2006. Morphofunctional changes of the digestive gland in the bivalve mollusk *Crenomytilus grayanus* (Dunker, 1853) in normal conditions and after parasitic invasion by trematodes. Russ. J. Mar. Biol. 32, 96.
- Valavanidis, A., 2018. Neonicotinoid insecticides. Banned by the European Union in 2018 after scientific studies concluded that harm honey bees. Scientific Reviews 37. [www.chem-tox-ecotox.org/ScientificReviews](http://www.chem-tox-ecotox.org/ScientificReviews).
- Van Dijk, T.C., Van Staalduin, M.A., Van der Sluijs, J.P., 2013. Macro-invertebrate decline in surface water polluted with imidacloprid. PloS One 8, e62374.
- Velisek, J., Stara, A., 2018. Effect of thiacloprid on early life stages of common carp (*Cyprinus carpio*). Chemosphere 194, 481–487.
- Vidal-Linan, L., Bellas, J., 2013. Practical procedures for selected biomarkers in mussels, *Mytilus galloprovincialis* – implications for marine pollution monitoring. Sci. Total Environ. 461–462, 56–64.
- Vidal-Linan, L., Bellas, J., Soriano, J.A., Concha-Grana, E., Munategui, S., Beiras, R., 2016. Bioaccumulation of PCB-153 and effects on molecular biomarkers acetylcholinesterase, glutathione-S-transferase and glutathione peroxidase in *Mytilus galloprovincialis* mussels. Environ. Pollut. 214, 885–891.
- Vidal-Linan, L., Villaverde-de-Saa, E., Rodil, R., Quintana, J.B., Beiras, R., 2018. Bioaccumulation of UV filters in *Mytilus galloprovincialis* mussel. Chemosphere 190, 267–271.
- Vieira, C.E.D., Perez, M.R., Acayaba, R.D.Á., Raimundo, C.C.M., dos Reis Martinez, C.B., 2018. DNA damage and oxidative stress induced by imidacloprid exposure in different tissues of the Neotropical fish *Prochilodus lineatus*. Chemosphere 195, 125–134.
- Ying, G.G., Kookana, R.S., 2004. Simultaneous determination of imidacloprid, thiacloprid, and thiamethoxam in soil and water by high-performance liquid chromatography with diode-array detection. J. Environ. Sci. Health B 39, 737–746.
- Zaccone, G., Fudge, D.S., Winegard, T.M., Capillo, G., Kuciel, M., Funakoshi, K., Lauriano, E.R., 2015. Confocal imaging and phylogenetic considerations of the subcutaneous neurons in the Atlantic hagfish *Myxine glutinosa*. Acta Zool. (Stockh) 96, 209–217.