



# Neurological alterations induced by formulated imidacloprid toxicity in Japanese quails

Sayed M. Rawi<sup>1,2,3</sup> · Ayed S. Al-Logmani<sup>2</sup> · Reham Z. Hamza<sup>4,5</sup> 

Received: 15 May 2018 / Accepted: 20 December 2018 / Published online: 3 January 2019  
© Springer Science+Business Media, LLC, part of Springer Nature 2019

## Abstract

Agrochemical risk assessment that takes into account only pesticide active ingredients without the spray adjuvants will miss important toxicity outcomes detrimental to non-target species including birds. In the present study toxicity of imidacloprid (IMI) pesticide was evaluated individually and in a mixture with polyethylene glycol (PEG-600) as adjuvant against Japanese quails. Oral intubation was used to obtain concentration-mortality data. Oral intubation was used to obtain concentration-mortality data. Treatments of quails for 24 h with different doses leading to the calculation of  $LC_{50}$  values. PEG enhances the pesticide efficacy and the  $LD_{50}$  value of IMI was  $17.02 \text{ mg/Kg}^1$ , and in combination with PEG it was  $15.98 \text{ mg/kg}^{-1}$ . In the second phase of the study, the effects of a single acute dose of IMI ( $1/4 LD_{50}$ ) individually or in a mixture with PEG has a potent effect on the activity of plasma AChE and brain monoamines transmitters. However, the addition of PEG-adjuvant to the selected insecticide has shown more toxic potential, more highly significant decreases in AChE activity and different changes in cortical monoamines concentration. In the present study the maximum significant inhibition of AChE activity, was recorded post 72 h exposure to IMI individually and 96 h in a mixture with PEG and exhibited  $-37.56\%$  and  $-32.65\%$  decreases, respectively. Moreover, the oral intubation of IMI individually or in a mixture with PEG caused a significant elevation in the quail cortical NE and 5-HT. The result also showed while the mixture of IMI + PEG induced the more potent effect in DA alterations, IMI individually was more effective in 5-HT changes. Our findings also indicated that PEG exposure induced remarkable changes in the studied monoamines level and the values were significant throughout the tested periods in DA. Moreover, the studied dose level was vigorously affected quail brain cerebral cortex histological structure. When administered individually or in a mixture with PEG, IMI disclosed neural congestion, neuronal degeneration, pyknosis and perivascular cuffing with glial cells.

**Keywords** PEG-600 · Imidacloprid · Histopathology · Neuroalterations · Neurotransmitters Japanese quails

## Abbreviations

5-HT Serotonin  
AChE Acetylcholinesterase

DA Dopamine  
IMI Imidacloprid  
 $LD_{50}$  Dose that induced 50% mortality  
NE Norepinephrine  
PEG-600 Polyethylene glycol-600

✉ Reham Z. Hamza  
dr\_reham\_z@yahoo.com; Reham\_z@Zu.edu.eg

<sup>1</sup> Biology Department, Faculty of Sciences, King Abdulaziz University, Jeddah, Kingdom of Saudi Arabia

<sup>2</sup> Biology Department, Faculty of Sciences and Arts, Jeddah University, Khulais, Kingdom of Saudi Arabia

<sup>3</sup> Zoology Department, Faculty of Science, Cairo University, Cairo, Egypt

<sup>4</sup> Biology Department, Faculty of Science, Taif University, Taif 888, Kingdom of Saudi Arabia

<sup>5</sup> Zoology Department, Faculty of Science, Zagazig University, Zagazig 44519, Egypt

## Introduction

Over the last 2 decades, a new class of insecticides, the neonicotinoids, has been the most remarkable and popular insecticides (Casida and Durkin 2013; Simon-Delso et al. 2015). When used for plant protection, about 5% of the active ingredient of neonicotinoid is possessed by crop plants (Wood and Goulson 2017) and is most instead disperses into the wider environment (Goulson 2013) affecting population of

multiple vertebrate and invertebrate species (Prosser and Hart 2005; Gibbons et al. 2015) directly and indirectly. There is the clear potential for ingestion of neonicotinoids by granivorous animals, specifically birds and mammals. There is an alarming trend between declines of local bird populations and neonicotinoids in the environment. Tennekes and Sánchez-Bayo (2011) investigated that neonicotinoids were acting indirectly on bird populations, by diminishing the plenty of their insect prey. Köhler and Triebkorn (2013) and Millot et al. (2017) have suggested that neonicotinoids may have a more important impact on bird populations, even used in small concentrations. In the present study, we focus on IMI, which has become one of the most widespread neonicotinoids insecticides and exert sub-lethal effects, ranging from neurotoxic, genotoxic (Khalil et al. 2017) and cytotoxic effects (Wankhede et al. 2017) to different types of birds.

After a widespread use of pesticides, including neonicotinoids several purpose pests have developed resistance (Jeschke et al. 2011; Szendrei et al. 2012; Alyokhin et al. 2007). These include sucking pests such as planthoppers, aphids, lepidopteran species and coleopteran (Bass et al. 2015). As reported by many types of researches, the mechanisms responsible for neonicotinoids resistance include behavioural and/or physiological changes, bio-degradation, sequestration and modified penetration (Puinean et al. 2010; Bass et al. 2014). Moreover, in IMI toxicity, studies of Clements et al. (2016) and Zhang et al. (2017) have suggested that cytochrome p450 is highly evolved in their detoxification. Therefore, in order to design more sustainable pest control strategies great interest has turned to pesticide formulations. Formulations are assuming improve for pesticide storage, handling, safety, application, effectiveness against pest resistance. Till now there is a gap in risk assessments for adjuvants in pesticide formulations and most studies on agrochemical risk assessment take into consideration pesticide active ingredients only without the spray adjuvants. As reported recently, exposure to environmental levels of some of these adjuvant can affect non-target organs, including birds (Edgington et al. 2004; Stills 2005). Their effects include negative impacts and provoke severe biochemical (Wieder and Davis 1983), clinical (Bendele et al. 1999), histopathological (Ishida et al. 2006) and behavior (Rodrigue et al. 2011) toxic effects. Here, we tested the acute toxicity of IMI pesticide, and its formulation with PEG-600 on the neuronal histological structure and monoamines level on the brain of quails (*Coturnix coturnix*).

## Materials and methods

### Experimental birds

Healthy adult Japanese quails (*Coturnix coturnix*) (weighing about 100–120 g) were obtained from Central Animal Facility

Cairo University. All animals were allowed to acclimatize to the experimental conditions for a period of 5 days. All the quails were housed in polyacrylic cages, not more than 3 animals per cage, and maintained under standard laboratory conditions (12 h of light and dark cycle, room temperature  $22 \pm 3$  °C). Animals were given standard diet, and tap water was provided ad libitum. The institutional animal ethics in the animal care unit in the Faculty of Science, Zoology Department, Cairo University (IACUC), (CU/I/F/60/15) committee approved the experimental protocol.

### Chemicals

#### Imidacloprid

Imidacloprid (IMI),  $C_9H_{10}ClN_5O_2$  (CAS Number 138261–41–3 5,  $\geq 98\%$  purity, molecular weight of 255.66), was donated by Novartis Egypt. Stock solutions with corn oil were prepared in accordance with the International Union of Pure and Applied Chemistry.

#### Polyethylene glycol 600 dilaurate (PEG-600)

PEG-600 was endowed by the Egyptian Company of Starch, Yeast, and Detergents in emulsified form. It is a viscous transparent liquid, of range average Molecular Weight 570–630 and diluted with corn oil.

### Pesticide formulation

Before deciding the tested dose, the insecticide formulation was prepared by mixing stock solutions of 1 g IMI and 1 ml PEG in 10 ml corn oil as described (Rawi et al. 2013). After dilutions, preparation for pesticide, PEG, and pesticide-PEG was takes place.

### Toxicity symptoms and the median lethal dose

An approximate  $LD_{50}$  is determined by a so-called “up and down” or the “staircase method” using two animals and increasing the doses of IMI and PEG, either individually or in a mixture (ratio 1:1). Six doses were given orally to 6 groups of quails (5 quails in each group) for the determination of  $LD_{50}$  starting from 0% mortality to 100% mortality (Randhawa 2009). The animals were observed for 2 h and then at 4th, 6th, and 24th h for any toxic signs and symptoms. After 24th h, the numbers of decreased quails in each group were counted and the percentage of mortality was calculated using the graphical method of Litchfield and Wilcoxon (1949).

## Dose preparation and experimental design

After LD<sub>50</sub> determinations birds were randomly divided into four groups. Group 1 served as control (Corn oil 1 ml/100 g body weight); group 2 was treated with PEG (10 ml /kg body weight); group 3 was treated with IMI (4.5 mg/kg body weight diluted in corn oil) and group 4 was treated with IMI (3.995 mg/kg) and PEG (10 ml /kg body weight) diluted with corn oil. All groups were treated once a time with the selected dose using tracheal intubation. The volume of the dose depends on the size of the animals and should not exceed 1 ml/100 g b.w. (US EPA 2012; Millot et al. 2017). At precise intervals of 1, 6, 24, 72, 96 h and, 1, 2 and 3 weeks of the oral administration, 6 animals of each group were overnight fasted, then they were sacrificed and blood samples were collected, centrifuged at 3000 rpm for 30 min and the obtained sera were used for AChE determination. At 24 and 96 h and 3 weeks, in addition to the collected blood samples, cortical brain tissue was rapidly removed and quickly divided into two parts, one was immersed in neutral buffered formalin (10%) and used for histopathological studies and the other was quickly homogenized in 0.1 M phosphate buffer (pH 7.8), using a glass-Teflon, centrifuged at 10,000 g for 60 min at 4 °C and used for the determination of monoamines.

## Aetylcholinesterase (AChE) and monoamines analysis

Serum AChE activity was measured by the method of Ellman et al. (1961) as described by Gorun et al. (1978). Norepinephrine (NE), dopamine (DA) and serotonin (5-HT) contents as monoamines were measured by the method of Pagel et al. (2000) using HPLC (Perkin-Elmer).

## Histopathology

At the end of the tested period, the cortical brain was dissected and histological examinations were done by the method of Humason (1972).

## Statistical analysis

All values were expressed as mean  $\pm$  SE. Statistical analysis was done by using SPSS 14 program. The statistical significance of differences between the two means was assessed by one way ANOVA. *P* values <0.05 were considered to be significant.

## Results

### Toxicity and mortality testing

The percentage of animals that died at each dose was transformed to probit using Finney's method (1952) and


the percentage of dead animals for 0 and 100% were corrected before the determination of probits. After that, the calculated LD<sub>50</sub> of IMI with corn oil when given orally was 17.02, mg/kg when IMI administered individually and 15.98 mg/kg when it was administered in a mixture with PEG. (Fig. 1).

### Toxicity signs recorded during the experiment

Throughout the experimental periods of toxicity testing level and the neurological effects of the acute dose, the animals exhibited general weaknesses, salivation, sometimes excitation and a variable sequence of motor symptoms involving pawing and tremor. However, the addition of PEG-adjuvant to the selected insecticide has shown more toxic potential than do the insecticide alone.


### Serum AChE

In the present study significant inhibition ( $p \leq 0.05$ ) of AChE activity, was recorded at the early exposure periods and the potency is varied in the following increasing order PEG+IMI > IMI > PEG. In the animal groups treated with PEG or IMI individually, the more highly significant decreases in the enzyme activity were recorded post 72 h exposure and exhibit, -26% and -49.38 decreases, respectively. Post one and 3 weeks of exposure, the enzyme activity showed a tendency to regain normal control level and the recorded values, however, being insignificantly affected (Fig. 2).



**Fig. 1** Lethal dose (mean  $\pm$  SD) of IMI individually and with a combination of polyethylene glycol given orally to coturnix quail

**Fig. 2** Effect of IMI exposure on AChE levels in the plasma of coturnix quail. Values represent the mean  $\pm$  SD of 6 independent samples; error bars indicate standard deviation. Statistical significance ( $*P < 0.05$ ) was analyzed using a factorial ANOVA. PEG compared to control and IMI and IMI + PEG compared to PEG



## Cortical brain monoamines


### Norepinephrine

After treatment with PEG, and as compared with the control group there were no significant changes in the quail cortical epinephrine when compared with the control group ( $p > 0.05$ ). Comparatively, the oral intubation of IMI individually or in a mixture with PEG at the tested dose levels caused a significant elevation in the quail cortical NE level. The highest increased effect ( $p < 0.01$ ) however, was attained after 24 h exposure to the tested mixture and attained 60.76% as compared to PEG-treated group (Fig. 3).

### Dopamine

The oral feeding of PEG and/or IMI to quails caused distress fluctuations in the level of DA cortical brain and the values detected being significantly affected throughout all the tested periods. Post 96 h and 3 weeks of exposure, the recorded values being significantly decreased and attained the minimum level post-exposure to PEG followed > PEG+IMI > and IMI in the increasing order (Fig. 4).

**Fig. 3** Effect of PEG and IMI exposure individually or in a mixture on cortical brain NE level of coturnix quail. Values represent the mean  $\pm$  SD of 6 independent samples; error bars indicate standard deviation. Statistical significance ( $*P < 0.05$ ) was analyzed using a factorial ANOVA. PEG compared to control and IMI and IMI + PEG compared to PEG



### Serotonin


The results given in Fig. 5 showed significant elevations in the level of quail cortical brain 5-HT after oral intubation with IMI individually or in a mixture with PEG. Comparatively, throughout all the tested periods, the IMI-treated group exhibited more increased level and attained the maximum value post 24 h and the detected value being increased more than twice the level recorded in the PEG-treated group.

## Histopathological observations

The histopathology of the cortical cerebrum of control and treated birds are given in Figs. (6 and 7). The examined sections of control brain tissue changes showed normal blood vessels, cytoplasm and nucleus in neurons (6a). In all treatment groups (PEG, IMI, and IMI + PEG) and as compared to the control group, the examined brain sections showed cerebral neuronal degeneration, pyknosis and neurophagia of some neurons due to IMI (Fig. 7a, b). In addition, relative gliosis, eosinophilic degenerated of neurons, central chromatocytolysis, demyelination of nerve fibers, and focal



**Fig. 4** Effect of PEG and IMI exposure individually or in a mixture on cortical brain DA level of cotumix quail. Values represent the mean  $\pm$  SD of 6 independent samples; error bars indicate standard deviation. Statistical significance (\*  $P < 0.05$ ) was analyzed using a factorial ANOVA. PEG compared to control and IMI and IMI + PEG compared to PEG




minute hemorrhage was also seen by IMI in a mixture with PEG (Fig. 7c, d).


## Discussion

To date, the toxicity that may induce by PEG individually and in a mixture with IMI doesn't studied in birds. In the present work, behavioral, biochemical, and histological changes have been recorded in Japanese quail administered with IMI and PEG individually or in a mixture. Histopathological findings of congestion in the blood vessels, necrosis, neurodegeneration, and hemorrhage were recognized in the cortical brain sections of animals treated with IMI individually or in a mixture with PEG. Further, cerebrum neuronal degeneration, pyknosis and neurophagia were observed in PEG-treated birds. Several investigators have shown that IMI (de Oliveira et al. 2010; Tufi et al. 2015; Rodriguez et al. 2016; Zhu et al. 2017) and PEG (Regulska et al. 2010; Pomerny et al. 2013) exerted adverse effects on the CNS cells and may contribute in pathogenesis of neurodegenerative disorders. As reported recently by Wankhede et al. (2017), congestion, degeneration of neurons, nuclear migration, vacuolation, oedema, haemorrhages, and gliosis were evident in the brain after treatments with IMI. In addition, results recorded by Kawthar

et al. (2012) showed that PEG exhibited DNA damage, vacuolar degeneration, inflammatory cellular infiltration, necrosis and pyknosis in liver and kidney cells of mice. Moreover, it is well-established that the brain histopathological injury leads to diverse behavioral consequence (Grewal et al. 2015). In the present investigations, IMI individually or in a mixture with PEG produced toxicity involves neuronal hyper-excitability, and disorders of cognition a varying degree of mild to moderate toxic symptoms and behavioral changes and the formulated form of the tested pesticide is characterized by more recognized excitability than do the insecticide alone. Indeed, Avery et al. (1994) observed ataxia in red-winged blackbirds (*Agelaius phoeniceus*) and brown-headed cowbirds (*Molothrus ater*) that fed on IMI-treated seeds. In addition, Cox (2001) reported in-coordination and inability to fly in house sparrow exposed to IMI and the toxicity often involves neuronal hyper-excitability, and disorders of cognition. Also, as reported by Webster et al. (2007) PEG can be considered as a toxic substance and the clinical signs associated with acute exposure include jumping, tremors, and convulsions in rats. Behavioral changes produced by acute IMI have been related to neuronal cell degeneration (Lang et al. 2014; Cho et al. 2017), alterations in AChE activity (Mach et al. 2004; Bonansea et al. 2016; Rodrigues et al. 2017) and regional changes in brain monoamines levels (Koprowska et al.

**Fig. 5** Effect of PEG and IMI exposure individually or in a mixture on cortical brain 5-HT level of cotumix quail. Values represent the mean  $\pm$  SD of 6 independent samples; error bars indicate standard deviation. Statistical significance (\*  $P < 0.05$ ) was analyzed using a factorial ANOVA. PEG compared to control and IMI and IMI + PEG compared to PEG






**Fig. 6** Histopathological changes of IMI toxicity (96 h). **a** Normal histology of the cerebral cortex of control group. H&E. 50  $\mu$ m, showing pyramidal neuron (arrows), blood vessel (solid arrow), axon (head arrow), and glial cell (lines). **b–c** Neuronal cells of PEG treated group showing degenerated neurons (arrows) and pyknosis of some neurons and neurophagia **c** (head arrows)

2004) and the more toxicity profile of the formulated form may be due to their synergistic toxicity on the neuronal cells. As indicated by several investigators both PEG (Herold et al. 1982) and IMI (Qadir et al. 2014) can account for the hypercalcemia, which can cause severe neurological symptoms. In addition, our data demonstrated that the tested mixture has, the more the inhibition effect of AChE activity in most of the tested period. As reported by Moser et al. (2006) the acute neurobehavioral effects of AChE-inhibiting pesticides are primarily due to overstimulation of the cholinergic system as well as different modes of action and different target sites. However, mortality data from this study indicated that IMI alone and with PEG generated different mortality, while the  $LD_{50}$  of the insecticide alone was  $17.02 \text{ mg/Kg}^{-1}$  it was  $15.98 \text{ mg/kg}^{-1}$  in the tested binary mixture.

We further examined the impact of IMI and PEG on the cortical brain monoamines level. The current study showed that exposure to PEG and IMI individually or in a mixture induced a significant decrease in DA and significantly elevated EP and 5-HT levels. There are no reports we came across in



**Fig. 7** Photomicrograph of quail cortical brain at the end of 96 h of exposure. **a–b** Cortical sections of IMI treated group showing congestion of cerebellar blood vessel (**c**), and neuronal degeneration and pyknosis of neurons (**B**-arrows). **c–d** IMI plus PEG-treated quail showing perivascular cuffing with glia cells (**C**-arrows) and necrosis of Purkinje cells (**D**-n)

Japanese quail about monoamines level affected by the tested compounds. Rodríguez et al. (2016) have reported analogous changes of brain neurotransmitters after pyrethroid cyfluthrin insecticides exposure in male rats. Moreover, Martínez et al. (2018) noted that glyphosate herbicide in a dose-dependent, altered central nervous system (CNS) monoaminergic neurotransmitters in a brain regional manner. Also, the results obtained by Tufi et al. (2015) and Martínez et al. (2018) proved the assumption that the pesticides exposure changes in dopamine (DA), noradrenaline (NE), and serotonin (5-HT). However, a number of evidence suggests that PEG included in unleaded gasoline vapours impaired the levels of monoamine neurotransmitters and other biochemical parameters in different brain areas and modulated several behavioural aspects in rats (Burbacher 1993; Kinawy 2009). In addition, as reported recently by Khalil et al. (2017) at specific dose levels IMI administration led to an alteration in the levels of cortisone and catecholamines and induced behavioral deficits, particularly in the animals exposed to the dose of  $1.0 \text{ mg/kg}$ . Our results about different exhibited changes for monoamines might be due to counter mechanism against different toxicity sign occurred due to exposure to IMI and/or PEG. It is commonly accepted that oxidative stress is responsible for the



brain damage and the excessive release of brain monoamines. Several investigators (Ahmad et al. 2010; Ge et al. 2015; Wang et al. 2016; Özdemir et al. 2018) have reported that IMI exposure induced oxidative damage which induces an increase in lipid peroxidation and protein oxidation. Consequently, the brain, with its rich lipid content, high energy demand, and weak antioxidant capacity become an easy target of excessive oxidative insult. The present data suggest that IMI individually or in a mixture with PEG induces neuroinflammation associated with monoamines imbalances and toxicity symptoms and PEG can increase the ability of IMI to affect significant toxicologic end points, including developmental neurotoxicity, and disruption of monoamines level.

**In conclusion**, there are clinical trials that have demonstrated the tolerability and the safety of PEG-600 in humans and signified the lack of central effects, but the recent study demonstrates an overview of modulation effect PEG-600 as an adjuvant on pesticides neurotoxicological effects on birds. The proposed mechanisms are potentiation at most of the studied parameters that could be assigned to neurotransmitters or their metabolism influence.

**Acknowledgments** This research was funded by King Abdulaziz University (Scientific Research (DSR) Deanship), Grant No. 857-200-D1435. Entitled: Role of Adjuvants on the physiological effects on organophosphate and pyrethroid insecticides.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## References

- Ahmad A, Rasheed N, Banu N, Palit G (2010) Alterations in monoamine levels and oxidative systems in frontal cortex, striatum, and hippocampus of the rat brain during chronic unpredictable stress. *Stress* 13(4):355–364
- Alyokhin A, Dively G, Patterson M, Castaldo C, Rogers D, Mahoney M, Wollam J (2007) Resistance and cross-resistance to imidacloprid and thiamethoxamin the Colorado potato beetle. *Pest Manag Sci* 63:32–41
- Avery ML, Decker DG, Fischer DL (1994) Cage and flight pen evaluation of avian repellency and hazard associated with imidacloprid-treated rice seed. *Crop Prot* 13:535–540
- Bass C, Puinean AM, Zimmer CT, Denholm I, Field LM, Stephen P (2014) The evolution of insecticide resistance in the peach potato aphid, *Myzus persicae*. *Insect Biochem Mol Biol* 51:41–51
- Bass C, Denholm I, Martin S, Williamson MS, Nauen R (2015) The global status of insect resistance to neonicotinoid insecticides. *Pestic Biochem Physiol* 121:78–87
- Bendele AM, McComb J, Gould T, Frazier J, Chlipala E, Seely J, Kieft G, Edwards CK (1999) Effects of PEGylated soluble tumor necrosis factor receptor type I (PEG sTNF-RI) alone and in combination with methotrexate in adjuvant arthritic rats. *Clin Exp Rheumatol* 17:553–560
- Bonansea RI, Wunderlin DA, Amé MV (2016) Behavioral swimming effects and acetylcholinesterase activity changes in *Jenynsia multidentata* exposed to chlorpyrifos and cypermethrin individually and in mixtures. *Ecotoxicol Environ Saf* 129:311–319
- Burbacher TM (1993) Neurotoxic effects of gasoline and gasoline constituents. *Environ Health Perspect* 101(6):133–141
- Casida JE, Durkin KA (2013) Neuroactive insecticides: targets, selectivity, resistance, and secondary effects. *Annu Rev Entomol* 58:99–117
- Cho Y-E, Lee M-H, Song B-J (2017) Neuronal cell death and degeneration through increased Nitroxidative stress and tau phosphorylation in HIV-1 transgenic rats. *PLoS One* 12(1):e0169945
- Clements J, Schoville S, Peterson N, Lan Q, Russell L (2016) Groves characterizing molecular mechanisms of imidacloprid resistance in select populations of *Leptinotarsa decemlineata* in the central sands region of Wisconsin. *PLoS One* 11(1):e0147844
- Cox C (2001) Insecticide factsheet: imidacloprid. *J Pestic Reform* 21:15–11
- de Oliveira IM, Nunes BV, Barbosa DR, Pallares AM, Faro LR (2010) Effects of the neonicotinoids thiametoxam and clothianidin on in vivo dopamine release in rat striatum. *Toxicol Lett* 192:294–297
- Edgington AN, Sheridan PM, Stephenson GR, Thompson DG, Boermans HJ (2004) Comparative effects of pH and vision herbicide on two life stages of four anuran amphibian species. *Environ Toxicol Chem* 23:815–822
- Ellman GL, Courtney KD, Andres V Jr, Featherstone RM (1961) A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem Pharmacol* 7:88–95
- Finney DJ (1952) Probit analysis, 2nd edn. Cambridge University Press, Cambridge
- Ge W, Yan S, Wang J, Zhu L, Chen A, Wang J (2015) Oxidative stress and DNA damage induced by imidacloprid in zebrafish (*Danio rerio*). *J Agric Food Chem* 63(6):1856–1862
- Gibbons D, Morrissey C, Mineau P (2015) A review of the direct and indirect effects of neonicotinoids and fipronil on vertebrate wildlife. *Environ Sci Pollut Res Int* 22:103–118
- Gorun V, Proinov I, Balaban V, Balaban G, Barzu O (1978) Modified Ellman procedure for assay of cholinesterase in crude enzymatic preparations. *Anal Biochem* 86:324–326
- Goulson D (2013) An overview of the environmental risks posed by neonicotinoid insecticides. *J Appl Ecol* 50:977–987
- Grewal KK, Sandhu GS, Kaur R, Brar RS, Sandhu HS (2015) Toxic impacts of cypermethrin on behavior and histology of certain tissues of albino rats. *Environ Sci Pollut Res Int* 22:103–118
- Herold DA, Rodeheaver GT, Bellamy WT, Edlich RF (1982) Toxicity of topical polyethylene glycol. *Toxicol Appl Pharmacol* 65(2):329–335
- Humason GL (1972) Animal tissue techniques. Ferdman, W. Hand Co, San Francisco
- Ishida T, Li W, Liu Z, Kiwada H (2006) Stimulatory effect of polyethylene glycol (PEG) on gene expression in mouse liver following hydrodynamics-based transfection. *Genet Med J* 8:324–334
- Jeschke P, Nauen R, Schindler M, Elbert A (2011) Overview of the status and global strategy for neonicotinoids. *J Agric Food Chem* 59:2897–2908
- Kawthar AED, Aida IE, Abd-Elmoneim OM, Sharaf HA (2012) Assessment of genotoxicity and histopathological changes induced by polyethylene glycol (PEG-6000) in male mice. *J Cytol Histol* 3(5):1000153
- Khalil SR, Awad A, Mohammed HH, Nassan MA (2017) Imidacloprid insecticide exposure induces stress and disrupts glucose homeostasis in male rats. *Environ Toxicol Pharmacol* 55:165–174
- Kinawy A (2009) Impact of gasoline inhalation on some neurobehavioural characteristics of male rats. *BMC Physiol* 9:21
- Köhler HR, Triebskorn R (2013) Wildlife ecotoxicology of pesticides: can we track effects to the population level and beyond? *Science* 341:759–765
- Koprowska M, Krotevicz M, Romaniuk A, Strzelczuk M (2004) Age-related changes in fear behavior and regional brain monoamines distribution in rats. *Acta Neurobiol Exp* 64:131–142

- Lang AE, Riherd Methner DN, Ferreira A (2014) Neuronal degeneration, synaptic defects, and behavioral abnormalities in tau<sub>45-230</sub> transgenic mice. *Neuroscience* 275:322–339
- Litchfield JT, Wilcoxon FA (1949) Simplified method of evaluating dose effect experiments. *J Pharmacol. Exp. Ther* 96:99–113
- Mach M, Grubbs RD, Price WA, Paton SJ, Lucot JB (2004) Behavioral changes after acetylcholinesterase inhibition with physostigmine in mice. *Pharmacol Biochem Behav* 79(3):533–540
- Martínez MA, Ares I, Rodríguez JL, Martínez M, Martínez-Larrañaga MR, Anadón A (2018) Neurotransmitter changes in rat brain regions following glyphosate exposure. *Environ Res* 161:212–219
- Millot F, Decors A, Mastain O, Quintaine T, Berny P, Vey D, Lasseur R, Bro E (2017) Field evidence of bird poisonings by imidacloprid-treated seeds: A review of incidents reported by the French SAGIR network from 1995 to 2014. *Environ Sci Pollut Res Int* 24(6):5469–5485
- Moser VC, Hunter DL, Marshall RS, Mcdaniel K, Phillips P, Padilla SJ (2006) Correlations of pesticide-induced cholinesterase inhibition and motor activity changes in adult rats. *Behav Toxicol Soc* 15:1006
- Özdemir S, Altun S, Arslan H (2018) Imidacloprid exposure cause the histopathological changes, activation of TNF- $\alpha$ , iNOS, 8-OHdG biomarkers, and alteration of caspase 3, iNOS, CYP1A, MT1 gene expression levels in common carp (*Cyprinus carpio* L.). *Toxicol Rep* 5:125–133
- Pagel P, Blome J, Wolf HU (2000) High-performance liquid chromatographic separation and measurement of various biogenic compounds possibly involved in the pathomechanism of Parkinson's disease. *J Chromatogr B Biomed Sci Appl* 746:297–304
- Pomierny B, Starek A, Krzyżanowska W, Starek-Swiechowicz B, Smaga I, Pomierny-Chamioło L, Regulska M, Budziszewska B (2013) Potential neurotoxic effect of ethylene glycol ethers mixtures. *Pharmacol Rep* 65(5):1415–1421
- Prosser P, Hart ADM (2005) Assessing potential exposure of birds to pesticide-treated seeds. *Ecotoxicol* 14:679–691
- Puinean AM, Foster SP, Oliphant L, Denholm I, Field LM, Millar NS, Williamson MS, Bass C (2010) Amplification of a cytochrome P450 gene is associated with resistance to neonicotinoid insecticides in the aphid *Myzus persicae*. *PLoS Genet* 6(6):e1000999
- Qadir S, Latif A, Ali M, Iqbal F (2014) Effects of Imidacloprid on the hematological and serum biochemical profile of Labeo rohita. *Pakistan J Zool* 46(4):1085–1090
- Randhawa MA (2009) Calculation of LD50 values from the method of miller and Tainter, 1944. *J Ayub Med Coll Abbottabad* 21:184–185
- Rawi SM, Al-Hazmi MA, Waggas AM (2013) The effects of polyethylene glycol (PEG-600) on the acute neurotoxicity of different types of pesticides. *Int J Manag Sci Bus* 2(7):62–72
- Regulska M, Pomierny B, Basta-Kaim A, Starek A, Filip M, Lason W, Budziszewska B (2010) Effects of ethylene glycol ethers on cell viability in the human neuroblastoma SH-SY5Y cell line. *Pharmacol Rep* 62:1243–1249
- Rodrigue JR, Balistreri W, Haber B, Jonas MM, Mohan P, Molleston JP, Murray KF, Narkewicz MR, Rosenthal P, Smith LJ, Lobritto SJ, Schwarz KB, Robuck PR, Barton B, González-Peralta RP (2011) Peginterferon with or without ribavirin has minimal effect on quality of life, behavioral/emotional, and cognitive outcomes in children. *Hepatology* 53:1468–1475
- Rodrigues KJ, Santana MB, Do Nascimento JL, Picanco-Diniz DL, Maues LA et al (2017) Behavioral and biochemical effects of neonicotinoid thiamethoxam on the cholinergic system in rats. *Ecotoxicol Environ Saf* 73:101–107
- Rodriguez JL, Ares I, Martinez M, Martinez-Larranaga MR, Anadon A, Martinez MA (2016) Effect of exposure to pyrethroid Cyfluthrin on serotonin and dopamine levels in brain region of male rats. *Environ Res* 146:388–394
- Simon-Delso N, Amaral-Rogers V, Belzunces LP, Bonmatin JM, Chagnon M, Downs C, Furlan L, Gibbons DW, Giorio C, Girolami V, Goulson D, Kreutzweiser DP, Krupke CH, Liess M, Long E, McField M, Mineau P, Mitchell EAD, Morrissey CA, Noome DA, Pisa L, Settele J, Stark JD, Tapparo A, Van Dyck H, Van Praagh J, Van der Sluijs JP, Whitehorn PR, Wiemers M (2015) Systemic insecticides (neonicotinoids and fipronil): trends, uses, mode of action and metabolites. *Environ Sci Pollut Res Int* 22:5–34
- Stills JHF (2005) Adjuvants and antibody production: dispelling the myths associated with Freund's complete and other adjuvants. *ILAR J* 46:280–293
- Szendrei Z, Grafius E, Byrne A, Ziegler A (2012) Resistance to neonicotinoid insecticides in field populations of the Colorado potato beetle (*Coleoptera: Chrysomelidae*). *Pest Manag Sci* 68:941–946
- Tennekes HA, Sánchez-Bayo F (2011) Time-dependent toxicity of neonicotinoids and other toxicants: implications for a new approach to risk assessment. *J Environ Anal Toxicol* S4:001
- Tufi S, Stel JM, de Boer J, Lamoree MH, Leonards PE (2015) Metabolomics to explore Imidacloprid-induced toxicity in the central nervous system of the freshwater snail *Lymnaea stagnalis*. *Environ Sci Technol* 49(24):14529–14536
- US EPA (2012) United States environmental protection agency ecological risk assessment
- Wang J, Wang J, Wang G, Zhu L, Wang J (2016) DNA damage and oxidative stress induced by imidacloprid exposure in the earthworm *Eisenia fetida*. *Chemosphere* 144:510–517
- Wankhede V, Hedau M, Ingole RS, Hajare SW, Wade WR (2017) Histopathological alterations induced by subacute imidacloprid toxicity in Japanese quails and its amelioration by *Butea monosperma*. *J Pharmacogn Phytochem* 6(3):252–257
- Webster R, Didier E, Harris P, Siegel N, Stadler J, Tilbury L, Smith D (2007) PEGylated proteins: Evaluation of their safety in the absence of definitive metabolism studies. *Drug Metab Dispos* 35(1):9–16
- Wieder KJ, Davis FF (1983) Enzyme therapy: II. Effect of covalent attachment of polyethylene glycol on biochemical parameters and immunological determinants of beta-glucosidase and alpha-galactosidase. *Appl Biochem Biotechnol* 5:337–347
- Wood TJ, Goulson D (2017) The environmental risks of neonicotinoid pesticides: a review of the evidence post 2013. *Environ Sci Pollut Res Int* 24(21):17285–17325
- Zhang Y, Han Y, Liu B, Yang Q, Guo H, Liu Z, Wang L, Fan J (2017) Resistance monitoring and cross-resistance role of CYP6CW1 between buprofezin and pymetrozine in field populations of *Laodelphax striatellus* (Fallén). *Sci Rep* 7:14639
- Zhu YC, Yao J, Adamczyk J, Luttrell R (2017) Feeding toxicity and impact of imidacloprid formulation and mixtures with six representative pesticides at residue concentrations on honey bee physiology (*Apis mellifera*). *PLoS One* 12(6):e0178421