

CONCLUSION ON PESTICIDE PEER REVIEW

Conclusion on the peer review of the pesticide risk assessment of confirmatory data submitted for the active substance pyriproxyfen¹

European Food Safety Authority²

European Food Safety Authority (EFSA), Parma, Italy

ABSTRACT

The conclusions of the European Food Safety Authority (EFSA) following the peer review of the initial risk assessment carried out by the competent authority of the rapporteur Member State the Netherlands for the pesticide active substance pyriproxyfen are reported. The context of the peer review was that requested by the European Commission following the submission and evaluation of confirmatory ecotoxicology data. The conclusions were reached on the basis of the evaluation of the representative uses of pyriproxyfen as an insecticide on protected tomato and aubergine and field grown cotton. The reliable endpoints concluded as being appropriate for use in regulatory risk assessment, derived from the available studies and literature in the dossier peer reviewed, are presented. A concern is identified.

© European Food Safety Authority, 2014

KEY WORDS

pyriproxyfen, peer review, risk assessment, pesticide, insecticide

¹ On request from the European Commission, Question No EFSA-Q-2014-00302, approved on 13 August 2014.

² Correspondence: pesticides.peerreview@efsa.europa.eu

Suggested citation: EFSA (European Food Safety Authority), 2014. Conclusion on the peer review of the pesticide risk assessment of confirmatory data submitted for the active substance pyriproxyfen. EFSA Journal 2014;12(8):3813, 19 pp. doi:10.2903/j.efsa.2014.3813

Available online: www.efsa.europa.eu/efsajournal

SUMMARY

Pyriproxyfen was included in Annex I to Directive 91/414/EEC on 1 July 2008 by Commission Directive 2008/69/EC, amended by Commission Directive 2010/39/EU and has been deemed to be approved under Regulation (EC) No 1107/2009, in accordance with Commission Implementing Regulation (EU) No 540/2011, as amended by Commission Implementing Regulation (EU) No 541/2011. It was a specific provision of the approval that the applicant was required to submit to the European Commission further studies on aquatic insects and pollinators by 30 June 2012.

In accordance with the specific provision, the applicant, Sumitomo Chemical Agro Europe S.A., submitted an updated dossier in July 2012, to address the risk to aquatic insects from Pyriproxyfen and from the metabolite DPH-PYR and to address the risk to pollinators, which was evaluated by the designated RMS, the Netherlands in the form of an Addendum to the Draft Assessment Report. In compliance with Guidance Document SANCO 5634/2009 rev.4.5, the RMS distributed the Addendum to Member States, the applicant and the EFSA for comments on 23 September 2013. The RMS collated all comments in the format of a Reporting Table, which was submitted to the European Commission in January 2014

Following consideration of the comments received, the European Commission requested the EFSA to organise a peer review of the RMS's evaluation of the confirmatory data submitted in relation to aquatic insects and pollinators and to deliver its conclusions on the weight of evidence approach to address the risk to aquatic insects from Pyriproxyfen and the validity of the field study on honey bees according to SANCO/10329/2002 rev. 2.

Based on the submitted confirmatory data, a low risk to aquatic organisms including insects was indicated for both pyriproxifen and the pertinent metabolite DPH-PYR for all the representative uses. The risk to pollinators from the insect growth regulator mode of action needs to be addressed further for greenhouse uses on tomato and aubergine in Southern Europe, as the submitted field study was not considered appropriate for the risk assessment for bees.

TABLE OF CONTENTS

Abstract	1
Summary	2
Table of contents	3
Background	4
The active substance and the formulated product	6
Conclusions of the evaluation	6
Concerns.....	8
1. Issues that could not be finalised	8
2. Critical areas of concern	8
3. Overview of the concerns identified for each representative use considered	8
References	9
Appendices	10
Abbreviations	16

BACKGROUND

Pyriproxyfen was included in Annex I to Directive 91/414/EEC on 1 July 2008 by Commission Directive 2008/69/EC³, amended by Commission Directive 2010/39/EU⁴ and has been deemed to be approved under Regulation (EC) No 1107/2009⁵, in accordance with Commission Implementing Regulation (EU) No 540/2011⁶, as amended by Commission Implementing Regulation (EU) No 541/2011⁷. EFSA previously finalised a Conclusion on this active substance on 21 July 2009 in the EFSA Scientific Report (2009) 336 (EFSA, 2009).

It was a specific provision of the approval that the applicant was required to submit to the European Commission further studies on aquatic insects and pollinators by 30 June 2012.

In accordance with the specific provision, the applicant, Sumitomo submitted an updated dossier in July 2012 which was evaluated by the designated rapporteur Member State (RMS), the Netherlands in the form of an Addendum to the Draft Assessment Report (Netherlands, 2013). In compliance with Guidance Document SANCO 5634/2009 rev.4.5 (European Commission, 2011), the RMS distributed the Addendum to Member States, the applicant and the EFSA for comments on 23 September 2013. The RMS collated all comments in the format of a Reporting Table, which was submitted to the European Commission in January 2014.

Following consideration of the comments received, the European Commission requested the EFSA to organise a peer review of the RMS's evaluation of the confirmatory data submitted in relation to aquatic insects and pollinators and to deliver its conclusions on the weight of evidence approach to address the risk to aquatic insects from pyriproxyfen and the validity of the field study on honey bees (Bakker F., 2011; available in: Netherlands, 2013) according to SANCO/10329/2002 rev. 2 (European Commission, 2002).

The Addendum and the Reporting Table were discussed at the Pesticides Peer Review Meeting 115 on ecotoxicology in May 2014. Details of the issues discussed, together with the outcome of these discussions were recorded in the meeting report.

A final consultation on the conclusions arising from the peer review took place with Member States via a written procedure in July 2014.

The conclusions laid down in this report were reached on the basis of the peer review of the RMS's evaluation of the confirmatory data submitted in relation to aquatic insects and pollinators. A key supporting document to this conclusion is the Peer Review Report, which is a compilation of the documentation developed to evaluate and address all issues raised in the peer review, from the compilation of comments in the Reporting Table to the conclusion. The Peer Review Report (EFSA,

³ Commission Directive 2008/69/EC/EC of 1 July 2008, amending Council Directive 91/414/EEC to include clofentezine, dicamba, difenoconazole, diflubenzuron, imazaquin, lenacil, oxadiazon, picloram and pyriproxyfen as active substances. OJ L 172, 2.7.2008, p. 9-14.

⁴ Commission Directive 2010/39/EC/EC of 22 June 2010, amending Council Directive 91/414/EEC as regards specific provisions relating to the active substances clofentezine, diflubenzuron, lenacil, oxadiazon, picloram and pyriproxyfen as active substances. OJ L 156, 23.6.2010, p. 7-11.

⁵ Regulation (EC) No 1107/2009 of the European Parliament and of the Council of 21 October 2009 concerning the placing of plant protection products on the market and repealing Council Directives 79/117/EEC and 91/414/EEC. OJ L 309, 24.11.2009, p. 1-50.

⁶ Commission Implementing Regulation (EU) No 540/2011 of 25 May 2011 implementing Regulation (EC) No 1107/2009 of the European Parliament and of the Council as regards the list of approved active substances. OJ L 153, 11.6.2011, p.1-186.

⁷ Commission Implementing Regulation (EU) No 541/2011 of 1 June 2011 amending Implementing Regulation (EU) No 540/2011 implementing Regulation (EC) No 1107/2009 of the European Parliament and of the Council as regards the list of approved active substances. OJ L 153, 11.6.2011, p.187-188.

2014) comprises the following documents, in which all views expressed during the course of the peer review, including minority views, can be found:

- the Reporting Table,
- the report of the scientific consultation with Member State experts
- the comments received on the draft EFSA conclusion.

Given the importance of the Addendum to the DAR including its final addendum (compiled version of July 2014 containing all individually submitted addenda (Netherlands, 2014)) and the Peer Review Report, these documents are considered respectively as background documents A and B to this conclusion.

It is recommended that this conclusion report and its background documents would not be accepted to support any registration outside the EU for which the applicant has not demonstrated to have regulatory access to the information on which this conclusion report is based.

THE ACTIVE SUBSTANCE AND THE FORMULATED PRODUCT

Pyriproxyfen is the ISO common name for 4-phenoxyphenyl (RS)-2-(2-pyridyloxy)propyl ether (IUPAC). Pyriproxyfen, belongs to the class of juvenile hormone mimics, other examples of this class are fenoxycarb and methoprene. The mode of action is suppression of embryogenesis, and inhibition of metamorphosis and reproduction. The representative formulated product for the evaluation was 'Pyriproxyfen 10 EC', an emulsifiable concentrate (EC).

The evaluated representative uses are as an insecticide on protected tomato and aubergine and field grown cotton. Full details of the GAP can be found in the list of end points in Appendix A.

CONCLUSIONS OF THE EVALUATION

The applicant has submitted to the Commission by the deadline of 30 June 2012 studies on aquatic insects and pollinators.

The assessment of the information was presented form of an Addendum to the Draft Assessment Report (Netherlands, 2013).

The addendum for the ecotoxicological chapter included updated risk assessment for aquatic organisms and bees using the pertinent European Commission (2002a and 2002b) which were in place at the time of the approval of pyriproxyfen.

In the previous peer review of pyriproxyfen it was concluded that the risk to aquatic insects both from pyriproxyfen and the metabolite DPH-PYR for all the representative uses could not be considered addressed by the microcosm study (Wijngaarden, 2004) included in the DAR (Netherlands, 2005) as insects were not covered. In the EFSA conclusion of pyriproxyfen issued in 2009 (EFSA, 2009) it was also concluded that the assessment factor to be applied to the microcosm endpoint should be further considered and agreed when more data on insects would become available.

To address the request for confirmatory data a chronic study with aquatic insects (Ephemeroptera) together with other four mesocosm studies from the open literature were submitted. Moreover, acute toxicity data on fish, invertebrates and algae with the metabolite DPH-PYR were also submitted.

All the available mesocosm studies including the microcosm presented in the DAR were discussed at the Pesticides Peer Review Experts' meeting 115 (May 2014). In the indoor microcosm, which is included in the DAR, the biological effects of a single application of pyriproxyfen on a plankton-dominated community consisting of algae, cladocerans, copepods, ostracods, and rotifers, were investigated. The experts noted that all the additional data showed a comparable sensitivity of crustaceans and insects to pyriproxyfen. However, it was also highlighted that all the publicly available studies were from outside Europe (3 from USA and 1 from Australia). Therefore, the endpoints from these studies were not considered suitable for risk assessment due to the uncertainties in the results extrapolation (e.g. differences in the abiotic conditions). The available additional studies, however, can be used as supportive information.

Overall, it was agreed that the endpoint to be selected for the risk assessment should be the NOEC_{population} of 1.2 µg/L instead of the NOEAEC of 5 µg/L, as previously, from the microcosm study originally included in the DAR. An assessment factor of 2 should be applied to this endpoint.

Based on the additional data and on the agreed endpoint, a low risk with FOCUS Step 3 PEC_{sw} was concluded for aquatic organisms including insects for the representative use on cotton. The risk to aquatic insects from the metabolite DPH-PYR can be also considered low, as it is likely that the

metabolite was present in the microcosm study, and therefore is considered to be covered. This is also in line with the previous conclusion on pyriproxyfen (EFSA, 2009).

For the use on tomato and aubergine (e.g. eggplant) in glasshouse a high risk to aquatic invertebrates was identified in the EFSA conclusion of 2009. For these representative uses 2 applications are proposed. However, assuming a single emission of 0.1% of the annual total application, a low risk was also concluded for aquatic organisms for the representative uses on tomato and aubergine (e.g. eggplant) using the agreed NOEC_{population} of 1.2 µg/L from the microcosm study.

In the previous conclusion on pyriproxyfen it was concluded that further studies were needed to address the risk to bee brood for the representative uses on tomato and eggplant in glasshouse in Southern Europe (1-2 applications 112.5 g a.s./ha).

A new field study at the relevant application rate for the use in glasshouse was submitted (Bakker, F., 2011). The study was conducted in *Phacelia* fields in the South-West of France in June. The study was discussed at the Pesticides Peer Review Experts' meeting 115 (May 2014). The experts agreed that the study cannot be considered appropriate for the risk assessment for bees, as it showed some limitations, i.e. lacking of a separate analysis of the results for colonies having received one or two treatments, lower average brood index for cohort 1 compared to the control but no differences for cohort 2 and 3 and deviations from the EPPO 170 (4) Bulletin 40 (2010) in the study design. Therefore, the risk to bees for the uses in tomato and aubergine (e.g. eggplant) in glasshouse in Southern Europe cannot be considered addressed. Risk mitigation measures such as covering or removing bumble bee colonies should be considered for the representative uses in glasshouse where these pollinators are used. EFSA noted that protection measures for the wild pollinators visiting the glasshouses should also be considered (e.g. keeping the glasshouses closed).

Concerns

1. Issues that could not be finalised

An issue is listed as an issue that could not be finalised where there is not enough information available to perform an assessment, even at the lowest tier level, for the representative uses in line with the Uniform Principles in accordance with Article 29(6) of Regulation (EC) No 1107/2009 and as set out in Commission Regulation (EU) No 546/2011⁸, and where the issue is of such importance that it could, when finalised, become a concern (which would also be listed as a critical area of concern if it is of relevance to all representative uses).

1. The risk to pollinators from the insect growth regulator mode of action needs to be addressed further for greenhouse uses on tomato and aubergine in Southern Europe, as the submitted field study was not considered appropriate for the risk assessment for bees.

2. Critical areas of concern

An issue is listed as a critical area of concern where there is enough information available to perform an assessment for the representative uses in line with the Uniform Principles in accordance with Article 29(6) of Regulation (EC) No 1107/2009 and as set out in Commission Regulation (EU) No 546/2011, and where this assessment does not permit to conclude that for at least one of the representative uses it may be expected that a plant protection product containing the active substance will not have any harmful effect on human or animal health or on groundwater or any unacceptable influence on the environment.

An issue is also listed as a critical area of concern where the assessment at a higher tier level could not be finalised due to a lack of information, and where the assessment performed at the lower tier level does not permit to conclude that for at least one of the representative uses it may be expected that a plant protection product containing the active substance will not have any harmful effect on human or animal health or on groundwater or any unacceptable influence on the environment.

3. Overview of the concerns identified for each representative use considered

Representative use		Cotton	Tomato, Aubergine North EU	Tomato, Aubergine South EU
	Assessment not finalised			
Risk to wild non target terrestrial organisms other than vertebrates	Risk identified			
	Assessment not finalised			X ¹
Risk to aquatic organisms	Risk identified			
	Assessment not finalised			
Comments/Remarks				

The superscript numbers in this table relate to the numbered points indicated in Section 1.1.

⁸ Commission Regulation (EU) No 546/2011 of 10 June 2011 implementing Regulation (EC) No 1107/2009 of the European Parliament and of the Council as regards uniform principles for evaluation and authorisation of plant protection products. OJ L 155, 11.6.2011, p. 127-175.

REFERENCES

- EFSA (European Food Safety Authority), 2009. Conclusion on pesticide peer review regarding the risk assessment of the active substance pyriproxyfen. EFSA Scientific Report (2009) 336 1-99 doi:10.2903/j.efsa.2009.336r. Available online: www.efsa.europa.eu
- EFSA (European Food Safety Authority), 2014. Peer Review Report to the conclusion regarding the peer review of the pesticide risk assessment of confirmatory data submitted for the active substance Pyriproxyfen. Available online: www.efsa.europa.eu
- European Commission, 2002a. Guidance Document on Terrestrial Ecotoxicology Under Council Directive 91/414/EEC. SANCO/10329/2002 rev.2 final, 17 October 2002
- European Commission, 2002b. Guidance Document on Aquatic Ecotoxicology Under Council Directive 91/414/EEC, November 2005. Available online: www.efsa.europa.eu
- Netherlands, 2005. Draft Assessment Report (DAR) on the active substance pyriproxyfen prepared by the rapporteur Member State the Netherlands in the framework of Directive 91/414/EEC, November 2005. Available online: www.efsa.europa.eu
- Netherlands, 2013. Addendum to the Draft Assessment Report, confirmatory data, September 2013. Available online: www.efsa.europa.eu
- Netherlands, 2014. Final Addendum to the Addendum to the Draft Assessment Report on pyriproxyfen., compiled by EFSA, July 2014. Available online: www.efsa.europa.eu

APPENDICES

APPENDIX A – LIST OF END POINTS FOR THE ACTIVE SUBSTANCE AND THE REPRESENTATIVE FORMULATION

Toxicity data for aquatic species (most sensitive species of each group) (Annex IIA, point 8.2, Annex IIIA, point 10.2)

Group	Test substance	Time-scale (Test type ^(D))	Endpoint	Toxicity (µg a.s./l)
Laboratory tests				
Fish				
<i>Lepomis macrochirus</i>	pyriproxyfen	96 h (f-t)	Mortality, LC ₅₀	>270 ^(A)
<i>Oncorhynchus mykiss</i>	pyriproxyfen	95 d (f-t)	NOEC	4.3 ^(A)
<i>Oncorhynchus mykiss</i>	S-71639 10EC	96 h (s)	Mortality, LC ₅₀	220 ^(A) (2100 µg form./L)
<i>Oncorhynchus mykiss</i>	4'-OH-Pyr	96 h (f-t)	Mortality, LC ₅₀	270 ^(A)
<i>Oncorhynchus mykiss</i>	PYPAC	96 h (s)	Mortality, LC ₅₀	>93000 ^(A)
<i>Oncorhynchus mykiss</i>	DPH-PYR	96 h (s)	Mortality, LC ₅₀	51000 ^(A)
Aquatic invertebrates				
<i>Daphnia magna</i>	pyriproxyfen	48 h (f-t)	Immobility, EC ₅₀	400 ^(A)
<i>Daphnia magna</i>	pyriproxyfen	21 d (f-t)	Reproduction, NOEC	0.015 ^(A)
<i>Mysidopsis bahia</i>	pyriproxyfen	28 d (f-t)	Reproduction, NOEC	0.81 ^(A)
<i>Cloeon dipterum</i>	pyriproxyfen	30 d (ss)	Nymph emergence, survival, development NOEC	≥ 0.202 ^(A)
<i>Daphnia magna</i>	S-71639 10EC	48 h (s)	Immobility, EC ₅₀	190 ^(A) (1800 µg form./L)
<i>Daphnia magna</i>	4'-OH-Pyr	48 h (f-t)	Immobility, EC ₅₀	1800 ^(A)
<i>Daphnia magna</i>	PYPAC	48 h (s)	Immobility, EC ₅₀	>95000 ^(A)
<i>Daphnia magna</i>	DPH-PYR	48 h (s)	Immobility, EC ₅₀	>9800 ^(A)
Sediment dwelling organisms				
<i>Chironomus riparius</i>	pyriproxyfen	28 d (s, spiked water)	Emergence, NOEC	10 ^(B)
Algae				
<i>Selenastrum capricornutum</i>	pyriproxyfen	72 h (s)	Biomass growth rate EbC ₅₀ ErC ₅₀	94 ^(B) 150 ^(B)
<i>Selenastrum capricornutum</i>	S-71639 10EC	72 h (s)	Biomass growth rate EbC ₅₀ ErC ₅₀	74 ^(B) 710 µg form./L) 110 ^(B) (1100 µg form./L))
<i>Pseudokirchneriella subcapitata</i>	4'-OH-Pyr	72 h (s)	Biomass growth rate EbC ₅₀ ErC ₅₀	>2500 ^(C) >2500 ^(C)
<i>Pseudokirchneriella subcapitata</i>	PYPAC	72 h (s)	Biomass growth rate EbC ₅₀ ErC ₅₀	26000 ^(A) 30000 ^(A)
<i>Pseudokirchneriella</i>	DPH-PYR	72 h (s)	Biomass EbC ₅₀	>9500 ^(A)

<i>subcapitata</i>			growth rate ErC ₅₀	>9500 ^(A)
Higher plant				
<i>Lemna gibba</i>	pyriproxyfen	14 d (s-s)	Fronds, EC ₅₀	>180 ^(A)
Microcosm or mesocosm tests				
Pyriproxyfen 10EC: NOEC _{population} 1.2 µg a.s./L ^(B)				

(A) Based on mean measured concentrations.

(B) Based on nominal concentrations (analytically confirmed for initial concentrations).

(C) Based on measured initial concentrations.

(D) f-t = flow through, s = static, s-s = semi=static

Toxicity/exposure ratios for the most sensitive aquatic organism per group for pyriproxyfen (Annex IIIA, point 10.2)

1st Tier

Tomato & Aubergine (T&A), 2*0.1125 kg as/ha: distance 1 m; 0.1% drift for greenhouse application
Cotton, 1*0.075 kg as/ha: distance 1 m; FOCUS Step 1 (2.77% drift and 10% run-off/drainage)

Crop	Organism	Test substance	Toxicity endpoint (µg a.s./L)	Time-scale	PEC _i (µg a.s./L)	TER	Annex VI Trigger
T&A	<i>O. mykiss</i>	product	220	96 h	0.0382	5765	100
Cotton	<i>O. mykiss</i>	product	220	96 h	1.5449	142	100
T&A	<i>O. mykiss</i>	a.s.	4.3	95 d	0.0382	113	10
Cotton	<i>O. mykiss</i>	a.s.	4.3	95 d	1.5449	2.8	10
T&A	<i>Daphnia</i>	product	190	48 h	0.0382	4979	100
Cotton	<i>Daphnia</i>	product	190	48 h	1.5449	123	100
T&A	<i>Daphnia</i>	a.s.	0.015	21 d	0.0382	0.39 ^(A)	10
Cotton	<i>Daphnia</i>	a.s.	0.015	21 d	1.5449	0.01	10
T&A	<i>C. riparius</i>	a.s.	10	28 d	0.0382	262	10
Cotton	<i>C. riparius</i>	a.s.	10	28 d	1.5449	6.5	10
T&A	<i>S. capricornutum</i>	product	74	72 h	0.0382	1939	10
Cotton	<i>S. capricornutum</i>	product	74	72 h	1.5449	48	10
T&A	<i>Lemna</i>	a.s.	>180	14 d	0.0382	>4717	10
Cotton	<i>Lemna</i>	a.s.	>180	14 d	1.5449	>117	10

(A) Assuming a single emission of 0.1% of the total annual application and using the NOEC_{population} of 1.2 µg/L derived from the microcosm a low risk to aquatic invertebrates can be concluded for the representative uses on tomato and aubergine.

2nd Tier

Cotton: 1*0.075 kg as/ha: distance 1 m; FOCUS Step 2 (2.77% drift and 3% run-off/drainage)

Crop	Organism	Test substance	Toxicity endpoint (µg a.s./L)	Time-scale	PEC _i (µg a.s./L)	TER	Annex VI Trigger
Cotton	<i>O. mykiss</i>	a.s.	4.3	95 d	0.6898	6.2	10
Cotton	<i>Daphnia</i>	a.s.	0.015	21 d	0.6898	0.02	1
Cotton	<i>C. riparius</i>	a.s.	10	28 d	0.6898	14	10

3rd Tier

Cotton: 1*0.075 kg as/ha: distance 1.3 m; FOCUS Step 3 (2.77% drift and substance-dependent drainage)

Crop	Organism	Test substance	Toxicity endpoint (µg a.s./L)	Time-scale	PEC _i (µg a.s./L)	TER	Annex VI Trigger
Cotton	<i>O. mykiss</i>	a.s.	4.3	95 d	0.381	11	10
Cotton	<i>Daphnia</i>	a.s.	0.015	21 d	0.381	0.04	10

4th Tier

Cotton: 1*0.075 kg as/ha: distance 1.3 m; FOCUS Step 3 (2.77% drift and substance-dependent drainage)

Crop	Organism	Test substance	Toxicity endpoint (µg a.s./L)	Time-scale	PEC _i (µg a.s./L)	TER	Annex VI Trigger
Cotton	<i>Daphnia</i>	a.s.	1.2*	56 d	0.381	3.15	2

* The endpoint is derived from a microcosm only covering zooplankton, not aquatic insects. However, it is shown with the new chronic toxicity study with *Cloeon dipterum* and with higher tier tests from public literature that that aquatic insects show a similar level of susceptibility to pyriproxyfen when compared with cladocerans. In the field studies many species are taken into account and crustaceans belong to the most sensitive organisms. Therefore, the NOEC_{population} of 1.2 µg/L derived from the microcosm is considered adequate for the higher tier risk assessment of all aquatic invertebrates. This was agreed by the experts at Pesticide Peer Review Meeting 115; they also agreed with a safety factor of 2 on this endpoint.

Effects on honeybees (Annex IIA, point 8.3.1, Annex IIIA, point 10.4)

Test substance	Acute oral toxicity (LD ₅₀ µg/bee)	Acute contact toxicity (LD ₅₀ µg/bee)
a.s. pyriproxyfen	N/A	N/A
Preparation Pyriproxyfen 10EC	LD ₅₀ 74 µg a.s./bee	LD ₅₀ >100 µg a.s./bee
Field or semi-field tests		
In a field study in Germany (dose rate 1* 75 g a.s./ha), Pyriproxyfen 10% EC did not affect mortality of adults or juvenile stages, overall colony performance or survival and development of eggs and larvae through to adult emergence.		

Hazard quotients for honey bees (Annex IIIA, point 10.4)

Application rate (g as/ha)	Crop	Route	Hazard quotient	Annex VI Trigger
Laboratory tests				
112.5	Tomato and Aubergine (e.g. eggplant)	Oral Contact	1.5 <1.1	50 50
75	Cotton	Oral Contact	1.0 <0.75	50 50

Summary of representative uses evaluated (pyriproxyfen)*


Crop and/or situation (a)	Member State or Country	Product name	F G or I (b)	Pests or Group of pests controlled (c)	Formulation		Application				Application rate per treatment			PHI (days) (l)	Remarks: (m)
					Type (d-f)	Conc. of as (i)	method kind (f-h)	growth stage & season (j)	number min max (k)	interval between applications (min)	kg as/hL min max	water L/ha min max	kg as/ha min max		
Tomato (greenhouse)	South Europe	Pyriproxyfen 10 EC	G	Greenhouse and cotton whitefly	EC	100 g/L	Foliar spray (High Volume Spraying)	BBCH 89	1-2	10 days	0.005-0.0075	1000-1500	0.05-0.1125	3	First application: as soon as adults are observed
Tomato (greenhouse)	North Europe	Pyriproxyfen 10 EC	G	Greenhouse and cotton whitefly	EC	100g/L	Foliar spray (High Volume Spraying)	BBCH 89	1-2	10 days	0.002-0.003	800-1200	0.02-0.03	3	
Aubergine (e.g. eggplant) (greenhouse)	South Europe	Pyriproxyfen 10 EC	G	Greenhouse and cotton whitefly	EC	100g/L	Foliar spray (High Volume Spraying)	BBCH 89	1-2	10 days	0.005-0.0075	1000-1500	0.05-0.1125	3	
Aubergine (e.g. eggplant) (greenhouse)	North Europe	Pyriproxyfen 10 EC	G	Greenhouse and cotton whitefly	EC	100g/L	Foliar spray (High Volume Spraying)	BBCH 89	1-2	10 days	0.002-0.003	800-1200	0.02-0.03	3	
Cotton	South Europe	Pyriproxyfen 10 EC	F	Cotton whitefly	EC	100g/L	Foliar spray (High Volume Spraying)	BBCH 78-79	1	n.a.	0.009-0.015	500-800	0.075	n.a.	

Remarks: * Uses for which risk assessment could not been concluded due to lack of essential data are marked grey

(h) Kind, e.g. overall, broadcast, aerial spraying, row, individual plant, between the plants - type of equipment used must be indicated

- | | | | |
|-----|---|-----|---|
| (a) | For crops, the EU and Codex classifications (both) should be used; where relevant, the use situation should be described (e.g. fumigation of a structure) | (i) | g/kg or g/l |
| (b) | Outdoor or field use (F), glasshouse application (G) or indoor application (I) | (j) | Growth stage at last treatment (BBCH Monograph, Growth Stages of Plants, 1997, Blackwell, ISBN 3-8263-3152-4), including where relevant, information on season at time of application |
| (c) | e.g. biting and sucking insects, soil born insects, foliar fungi, weeds | (k) | The minimum and maximum number of application possible under practical conditions of use must be provided |
| (d) | e.g. wettable powder (WP), emulsifiable concentrate (EC), granule (GR) | (l) | PHI - minimum pre-harvest interval |
| (e) | GCPF Codes - GIFAP Technical Monograph No 2, 1989 | (m) | Remarks may include: Extent of use/economic importance/restrictions |
| (f) | Method, e.g. high volume spraying, low volume spraying, spreading, dusting, drench | | |
| (g) | All abbreviations used must be explained | | |

APPENDIX B – USED COMPOUND CODE(S)

Code/Trivial name*	Chemical name/SMILES notation**	Structural formula**
DPH-PYR	4-hydroxyphenyl (RS)-2-(2-pyridyloxy)propyl ether	

* The metabolite name in bold is the name used in the conclusion.

ABBREVIATIONS

1/n	slope of Freundlich isotherm
λ	wavelength
ε	decadic molar extinction coefficient
°C	degree Celsius (centigrade)
μg	microgram
μm	micrometer (micron)
a.s.	active substance
AChE	acetylcholinesterase
ADE	actual dermal exposure
ADI	acceptable daily intake
AF	assessment factor
AOEL	acceptable operator exposure level
AP	alkaline phosphatase
AR	applied radioactivity
ARfD	acute reference dose
AST	aspartate aminotransferase (SGOT)
AV	avoidance factor
BCF	bioconcentration factor
BUN	blood urea nitrogen
bw	body weight
CAS	Chemical Abstracts Service
CFU	colony forming units
ChE	cholinesterase
CI	confidence interval
CIPAC	Collaborative International Pesticides Analytical Council Limited
CL	confidence limits
cm	centimetre
d	day
DAA	days after application
DAR	draft assessment report
DAT	days after treatment
DDD	daily dietary dose
DM	dry matter
DT ₅₀	period required for 50 percent disappearance (define method of estimation)
DT ₉₀	period required for 90 percent disappearance (define method of estimation)
dw	dry weight
EbC ₅₀	effective concentration (biomass)
EC ₅₀	effective concentration
ECHA	European Chemicals Agency
EEC	European Economic Community
EINECS	European Inventory of Existing Commercial Chemical Substances
ELINCS	European List of New Chemical Substances
EMDI	estimated maximum daily intake
ER ₅₀	emergence rate/effective rate, median
ErC ₅₀	effective concentration (growth rate)
EU	European Union
EUROPOEM	European Predictive Operator Exposure Model
f(twa)	time weighted average factor
FAO	Food and Agriculture Organization of the United Nations
fd	feed
FIR	Food intake rate
FOB	functional observation battery
FOCUS	Forum for the Co-ordination of Pesticide Fate Models and their Use

g	gram
GAP	good agricultural practice
GC	gas chromatography
GCPF	Global Crop Protection Federation (formerly known as GIFAP)
GGT	gamma glutamyl transferase
GM	geometric mean
GS	growth stage
GSH	glutathion
h	hour(s)
ha	hectare
Hb	haemoglobin
Hct	haematocrit
hL	hectolitre
HPLC	high pressure liquid chromatography or high performance liquid chromatography
HPLC-MS	high pressure liquid chromatography – mass spectrometry
HQ	hazard quotient
IEDI	international estimated daily intake
IENTI	international estimated short-term intake
ISO	International Organization for Standardization
IUPAC	International Union of Pure and Applied Chemistry
JMPR	Joint Meeting on the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Expert Group on Pesticide Residues (Joint Meeting on Pesticide Residues)
K _{doc}	organic carbon linear adsorption coefficient
kg	kilogram
K _{Foc}	Freundlich organic carbon adsorption coefficient
L	litre
LC	liquid chromatography
LC ₅₀	lethal concentration, median
LC-MS	liquid chromatography-mass spectrometry
LC-MS-MS	liquid chromatography with tandem mass spectrometry
LD ₅₀	lethal dose, median; dosis letalis media
LDH	lactate dehydrogenase
LOAEL	lowest observable adverse effect level
LOD	limit of detection
LOQ	limit of quantification (determination)
m	metre
M/L	mixing and loading
MAF	multiple application factor
MATC	maximum allowable toxicant concentration
MCH	mean corpuscular haemoglobin
MCHC	mean corpuscular haemoglobin concentration
MCV	mean corpuscular volume
mg	milligram
mL	millilitre
mm	millimetre (also used for mean measured concentrations)
mN	milli-newton
MRL	maximum residue limit or level
MS	mass spectrometry
MSDS	material safety data sheet
MTD	maximum tolerated dose
MWHC	maximum water holding capacity
NESTI	national estimated short-term intake
ng	nanogram

NOAEC	no observed adverse effect concentration
NOAEL	no observed adverse effect level
NOEAEC	no observed ecological adverse effect concentration
NOEC	no observed effect concentration
NOEL	no observed effect level
OD	oil dispersion
OECD	Organisation for Economic Co-operation and Development
OM	organic matter content
Pa	pascal
PD	proportion of different food types
PEC	predicted environmental concentration
PEC _{air}	predicted environmental concentration in air
PEC _{gw}	predicted environmental concentration in ground water
PEC _{sed}	predicted environmental concentration in sediment
PEC _{soil}	predicted environmental concentration in soil
PEC _{sw}	predicted environmental concentration in surface water
pH	pH-value
PHED	pesticide handler's exposure data
PHI	pre-harvest interval
PIE	potential inhalation exposure
pK _a	negative logarithm (to the base 10) of the dissociation constant
P _{ow}	partition coefficient between <i>n</i> -octanol and water
PPE	personal protective equipment
ppm	parts per million (10 ⁻⁶)
PT	proportion of diet obtained in the treated area
PTT	partial thromboplastin time
QSAR	quantitative structure-activity relationship
r ²	coefficient of determination
REACH	Registration, Evaluation, Authorisation of Chemicals Regulation
RPE	respiratory protective equipment
RUD	residue per unit dose
SC	suspension concentrate
SD	standard deviation
SFO	single first-order
SMILES	simplified molecular-input line-entry system
SSD	species sensitivity distribution
STMR	supervised trials median residue
t _{1/2}	half-life (define method of estimation)
TER	toxicity exposure ratio
TER _A	toxicity exposure ratio for acute exposure
TER _{LT}	toxicity exposure ratio following chronic exposure
TER _{ST}	toxicity exposure ratio following repeated exposure
TK	technical concentrate
TLV	threshold limit value
TMDI	theoretical maximum daily intake
TRR	total radioactive residue
TSH	thyroid stimulating hormone (thyrotropin)
TWA	time weighted average
UDS	unscheduled DNA synthesis
UV	ultraviolet
W/S	water/sediment
w/v	weight per volume
w/w	weight per weight
WBC	white blood cell
WG	water dispersible granule

WHO
wk
yr

World Health Organization
week
year