

# A Brief Introduction to Synthetic Differential Geometry

Gaspar Pérez Scornik

Supervised by Xavier Gràcia Sabaté

Universitat Politècnica de Catalunya  
Faculty of Mathematics and Statistics

Bachelor's Thesis, 2017



## 1 Introductory theory

- Single Variable Calculus
- A Primer on Toposes

## 2 Microlinear Objects

- The Tangent Bundle
- Vector Fields and Flows

# The Derivative

- “The rate of change, for very small changes”

$$f(x + h) = f(x) + f'(x)h + \mathcal{O}(h^2)$$

# Recreating the Derivative Axiomatically

- We begin with an object  $R$ . Naturally, this is the “synthetic version” of  $\mathbb{R}$ .

# Recreating the Derivative Axiomatically

- We begin with an object  $R$ . Naturally, this is the “synthetic version” of  $\mathbb{R}$ .

## Axiom

*$R$  is an algebra over  $\mathbb{Q}$ .*

# Recreating the Derivative Axiomatically

## The Infinitesimals

- Let  $D$  be the set

$$D = \{d \in R \mid d^2 = 0\}.$$

# Recreating the Derivative Axiomatically

## The Infinitesimals

- Let  $D$  be the set

$$D = \{d \in R \mid d^2 = 0\}.$$

### KL Axiom

*For all functions*

$$f : D \rightarrow R,$$

*there exist unique  $a, b \in R$  such that*

$$f(d) = a + bd$$

*for all  $d \in D$ .*

- *The coefficient  $a$  is easily seen to be  $f(0)$ .*

# Recreating the Derivative Axiomatically

- The derivative can now be defined.
- Let  $f : R \rightarrow R$  be a function. Fix  $x$ , define  $g : D \rightarrow R$  by

$$g(d) = f(x + d).$$



# Recreating the Derivative Axiomatically

- By the KL axiom, there exist unique  $a, b$  with

$$g(d) = f(x + d) = a + bd$$

- The first coefficient is  $g(0) = f(x)$ .

# Recreating the Derivative Axiomatically

- By the KL axiom, there exist unique  $a, b$  with

$$g(d) = f(x + d) = a + bd$$

- The first coefficient is  $g(0) = f(x)$ .

## Definition

The **derivative** of  $f$ , written  $f'(x)$ , is the unique coefficient  $b$ .

# Recreating the Derivative Axiomatically

- Thus we have

$$f(x + d) = f(x) + f'(x)d \quad \forall d \in D$$

# An Example

## Example

Consider  $f(x) = x^n$ . Then

$$\begin{aligned} f(x + d) &= \binom{n}{0} x^n + \binom{n}{1} x^{n-1} d + K_2 d^2 + \cdots + K_n d^n \\ &= x^n + nx^{n-1} d \end{aligned}$$

Therefore  $f'(x) = nx^{n-1}$ .

# How is this possible?

# How is this possible?

- We have to leave the category of sets.

# How is this possible?

- We have to leave the category of sets.
- We have to weaken logic to only constructive logic (no LEM).

# How is this possible?

- We have to leave the category of sets.
- We have to weaken logic to only constructive logic (no LEM).
- Toposes are the correct setting for SDG.



# How is this possible?

- We have to leave the category of sets.
- We have to weaken logic to only constructive logic (no LEM).
- Toposes are the correct setting for SDG.
- The objects in a topos behave like sets.

- Define “2-infinitesimals”:

$$D(2) = \{(d_1, d_2) \in D \times D \mid d_1 d_2 = 0\}$$

# Microlinear Objects

- In particular, if  $M$  is microlinear then two maps

$$f, g : D \rightarrow M$$

define a unique map

$$h : D(2) \rightarrow M$$

such that

$$h(d, 0) = f(d)$$

$$h(0, d) = g(d)$$

# Microlinear Objects

- To be thought of as “manifolds”.
- We can define tangent vectors (and they will form a vector space).

# Tangent Vectors

- Classically, they are equivalence classes of paths.

# Tangent Vectors

- Classically, they are equivalence classes of paths.

## Definition

Let  $M$  be microlinear. A **tangent vector** to  $M$  at  $p$  is a map

$$t : D \rightarrow M$$

with  $t(0) = p$ .

# Tangent Vectors

- Classically, they are equivalence classes of paths.

## Definition

Let  $M$  be microlinear. A **tangent vector** to  $M$  at  $p$  is a map

$$t : D \rightarrow M$$

with  $t(0) = p$ .

- With that, the **tangent space** at  $p$ ,  $T_p M$ , is the set of all tangent vectors at  $p$ .

# Tangent Vectors

- The tangent space  $T_p M$  is an  $R$ -vector space.
  - $0$  = the constant map  $p$ .
  - $\lambda t(d) := t(\lambda d)$
- For addition we need to use microlinearity.



# Addition of Tangent Vectors

- Microlinearity guarantees that for  $t_1, t_2$  tangent vectors, there exists a unique

$$s : D(2) \rightarrow M$$

such that

$$s(d, 0) = t_1(d)$$

$$s(0, d) = t_2(d)$$

- Define  $(t_1 + t_2)(d) = s(d, d)$

# Addition of Tangent Vectors

## The Neutral Element

- Recall we designated  $0 \equiv p$  (ct.)
- Let  $t$  be a t.v. The map  $s : D(2) \rightarrow M$  must satisfy

$$s(d, 0) = t(d, 0)$$

$$s(0, d) = 0(d) = p$$

so  $s$  must be

$$s(d_1, d_2) = t(d_1),$$

$$\text{and } (t + 0)(d) = s(d, d) = t(d)$$

# The Tangent Bundle

- It is simply the collection of all tangent spaces,  $M^D$ . We write  $TM$ .
- We have a projection  $\pi : TM \rightarrow M$  given by

$$t \mapsto t(0)$$

# Vector Fields

- They are sections of  $TM$ ; maps  $X : M \rightarrow TM$  such that  $(\pi \circ X)(p) = p$ .
- The set of vector fields is denoted  $\mathfrak{X}(M)$ .

# Vector Fields

- They are sections of  $TM$ ; maps  $X : M \rightarrow TM$  such that  $(\pi \circ X)(p) = p$ .
- The set of vector fields is denoted  $\mathfrak{X}(M)$ .
  - Each  $T_p M$  is a vector space, so by fiberwise operations so is  $\mathfrak{X}(M)$ .

# Vector Fields

- They are sections of  $TM$ ; maps  $X : M \rightarrow TM$  such that  $(\pi \circ X)(p) = p$ .
- The set of vector fields is denoted  $\mathfrak{X}(M)$ .
  - Each  $T_p M$  is a vector space, so by fiberwise operations so is  $\mathfrak{X}(M)$ .
  - $\mathfrak{X}(M)$  is also an  $M^R$ -module, by

$$(fX)(p) = f(p)X(p)$$

- Exponentials  $A^X$  have the property that

$$B \rightarrow A^X$$

is equivalent to

$$X \times B \rightarrow A$$

$$\left( \tilde{f}(x, b) = f(b)(x) \right)$$

- Vector fields are maps  $X : M \rightarrow M^D$ , therefore the same as maps

$$X : D \times M \rightarrow M.$$

We write the same  $X$  because it's comfortable.



## Proposition

*Flows are additive.*

## Proof.

Let  $f, g : D(2)$  be defined by

$$\begin{aligned}f(d_1, d_2) &= X(d_1 + d_2, p) \\g(d_1, d_2) &= X(X(d_1, p), d_2)\end{aligned}$$

we have

$$\begin{aligned}f(d, 0) &= g(d, 0) \\g(0, d) &= f(0, d)\end{aligned}$$

since  $M$  is microlinear,  $f = g$ .







# Summary

- In SDG we define geometric objects directly.
- This requires working axiomatically.
- The right context for these needs is that of topos theory.
- It's possible to recover the “same” constructs from classical geometry.

# Summary

- In SDG we define geometric objects directly.
  - This requires working axiomatically.
  - The right context for these needs is that of topos theory.
  - It's possible to recover the “same” constructs from classical geometry.
- 
- Of course, we need to construct a category which satisfies the axioms.
  - These are **models** of SDG.

# For Further Reading

-  Marta C. Bunge, Felipe Gago, and Ana María San Luis.  
*Synthetic differential topology.*  
Submitted manuscript, not yet published, 2017.
-  Anders Kock.  
*Synthetic Differential Geometry.*  
Cambridge University Press, 2nd edition, 2006.
-  Anders Kock.  
*Synthetic Geometry of Manifolds.*  
Cambridge University Press, 2010.
-  René Lavendhomme.  
*Basic Concepts of Synthetic Differential Geometry.*  
Springer Science+Business Media, B.V., 1996.