Math 60

Find the gradient  $\nabla f(\mathbf{a})$  for

$$f(x, y, z) = \cos z \ln(x + y^2)$$
$$\mathbf{a} = \left(e, 0, \frac{\pi}{4}\right)$$

Page 1 of 8 May 17, 2016

Math 60

Find the matrix  $D\mathbf{f}(\mathbf{a})$  of partial derivatives, for

$$\mathbf{f}(s,t) = (s^2, st, t^2)$$
  
 $\mathbf{a} = (-1, 1)$ 

Page 2 of 8 May 17, 2016

Math 60

Find equations for the planes tangent to  $z = x^2 - 6x + y^3$  that are parallel to the plane 4x - 12y + z = 7.

Page 3 of 8 May 17, 2016

Suppose that you have the following information concerning a differentiable function f:

$$f(2,3) = 12,$$
  $f(1.98,3) = 12.1,$   $f(2,3.01) = 12.2.$ 

(a) Give an approximate equation for the plane tangent to the graph of f at (2,3,12).

(b) Use the result of part (a) to estimate f(1.98, 2.98).

Verify the product and quotient rules (Proposition 4.2) for the pair of functions given below.

$$f(x,y) = x^{2}y + y^{3}$$
$$g(x,y) = \frac{x}{y}$$

Page 5 of 8 May 17, 2016

For the function given below determine all second-order partial derivatives (including mixed partials).

$$f(x,y) = x^2 e^y + e^{2z}$$

Page 6 of 8 May 17, 2016

Let  $f(x,y) = ye^{3x}$ . Give general formulas for  $\frac{\partial^n f}{\partial x^n}$  and  $\frac{\partial^n f}{\partial y^n}$  where  $n \ge 2$ .

Page 7 of 8 May 17, 2016

The three-dimensional heat equation is the partial differential equation

$$k\left(\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} + \frac{\partial^2 T}{\partial z^2}\right) = \frac{\partial T}{\partial t},$$

where k is a positive constant. It models the temperature T(x,y,z,t) at the point (x,y,z) and time t of a body in space.

(a) We examine a simplified version of the heat equation. Consider a straight wire "coordinatized" by x. Then the temperature T(x,t) at time t and position x along the wire is modeled by the one-dimensional heat equation

$$k\frac{\partial^2 T}{\partial x^2} = \frac{\partial T}{\partial t}$$

Show that the function  $T(x,t) = e^{-kt}\cos x$  satisfies this equation. Note that if t is held constant at value  $t_0$ , then  $T(x,t_0)$  shows how the temperature varies along the wire at time  $t_0$ . Graph the curves  $z = T(x,t_0)$  for  $t_0 = \{0,1,10\}$ , and use them to understand the graph of the surface z = T(x,t) for  $t \ge 0$ . Explain what happens to the temperature of the wire after a long period of time.

Page 8 of 8 May 17, 2016