

Data driven analysis

Giuseppe Torrisi

King's College, London

Abstract

The Haemodynamic response of different brain regions during mental arithmetic tasks is investigated.

Contents

1	Introduction	1
1.1	Measurement mechanism	1
1.2	Protocol	2
1.3	Data selection	2
2	Cusum	3

1. Introduction

Brain computer interface (BCI). are devices that aim to interpret brain activity and produce digital signal that can be read by a machine. The main challenge is to online classify the brain signal in a stable and reliable way. In particular the key point for the development in BCI is to find patterns of brain activity associated to specific tasks.

In this report I analyse the hemodynamic response to mental arithmetic operation.

1.1. Measurement mechanism

Haemoglobin (Hb) is a protein found on red cells, it binds up to four O_2 molecules, making it the main Oxygen carrier in blood. The Hb shows structural changes when bound with oxygen. In this case oxy-Hb rather than deoxy-Hb is formed. Oxy-Hb and deoxy-Hb have different absorption

spectra. Since biological tissue is relatively transparent to light in the near-infrared range between 700 nm to 1000 nm [1], near-infrared spectroscopy (NIRS) is suitable to estimate Oxy-Hb and deoxy-Hb concentration [2].

The analysis of hemodynamic response can be used to measure the activity in different areas of brain [3]. Measurement at the brain tissue level [4] have suggested that in case of higher brain activity more O_2 is required, which causes the decrease of oxy-Hb. However, the blood flow rises to satisfy the higher need of oxygen. In conclusion, after the early decrease of oxy-Hb, the long term effect is the increases of haemoglobin oxygenation. The latter effect is exploited in NIRS to assess brain activity during the performance of specific tasks, *e.g.* cognitive[5], visual [6], and motor [7]. However, the temporal resolution of hemodynamic response spans over several seconds, making any analysis of NIRS challenging and problematic in the case of BCI.

1.2. Protocol

Data have been recorded from 8 University students (three males and five females, all right-handed aged 26.0 ± 2.8 years. They were asked to perform one digit subtraction from a two digit number (*e.g.* $93 - 5$). The protocol consists in performing as many mental operation in 12 second, then a 28s rest is given. During rest person is asked to relax and not to move. This protocol is repeated 6 times. 3 or 4 run of this protocol is measured per participant.

The device records 52 channel with a sampling rate of 10 Hz

1.3. Data selection

The same dataset has already been used in [8, 9] to find pattern in the brain response to mental arithmetic tasks. They identify 3 regions of interest (ROI) each containing 3 channel, called ROI-1, ROI-2, and ROI-3 (see Table 1). They show that an antagonistic activation patterns occurs, consisting in the rise of oxy-Hb in ROI-2 and ROI-3 and decrease of oxy-Hb in ROI-1. Therefore in the following only the channels corresponding to ROI-1, ROI-2, and ROI-3 are considered. The authors point out that oxy-Hb concentration provides stronger and less noisy signal compared to deoxy-Hb concentration. Therefore in this report it has been chosen to consider the oxy-Hb concentration only.

Table 1: Channels belonging to the three Region of Interest as defined in [8]. In this report we examine the following 9 channels only.

ROI-1	46	47	48
ROI-2	18	28	29
ROI-3	13	23	24

2. Cusum

Cusum

- [1] F. F. Jobsis, Noninvasive, infrared monitoring of cerebral and myocardial oxygen sufficiency and circulatory parameters, *Science* 198 (1977) 1264–1267.
- [2] A. Villringer, B. Chance, Non-invasive optical spectroscopy and imaging of human brain function, *Trends in neurosciences* 20 (1997) 435–442.
- [3] R. B. Buxton, K. Uludağ, D. J. Dubowitz, T. T. Liu, Modeling the hemodynamic response to brain activation, *Neuroimage* 23 (2004) S220–S233.
- [4] D. Malonek, A. Grinvald, Interactions between electrical activity and cortical microcirculation revealed by imaging spectroscopy: implications for functional brain mapping, *Science* 272 (1996) 551–554.
- [5] M. A. Franceschini, S. Fantini, J. H. Thompson, J. P. Culver, D. A. Boas, Hemodynamic evoked response of the sensorimotor cortex measured non-invasively with near-infrared optical imaging, *Psychophysiology* 40 (2003) 548–560.
- [6] M. Herrmann, A.-C. Ehlis, A. Wagener, C. Jacob, A. Fallgatter, Near-infrared optical topography to assess activation of the parietal cortex during a visuo-spatial task, *Neuropsychologia* 43 (2005) 1713–1720.
- [7] S. Wriessnegger, J. Kurzmann, C. Neuper, Spatio-temporal differences in brain oxygenation between movement execution and imagery: a multichannel near-infrared spectroscopy study, *International Journal of Psychophysiology* 67 (2008) 54–63.

- [8] G. Bauernfeind, R. Scherer, G. Pfurtscheller, C. Neuper, Single-trial classification of antagonistic oxyhemoglobin responses during mental arithmetic, *Medical & biological engineering & computing* 49 (2011) 979.
- [9] G. Pfurtscheller, G. Bauernfeind, S. C. Wriessnegger, C. Neuper, Focal frontal (de) oxyhemoglobin responses during simple arithmetic, *International Journal of Psychophysiology* 76 (2010) 186–192.