

My topology exercises

Evgeny Markin

2023

Contents

I	General Topology	7
1	Set Theory and Logic	8
1.1	Fundamental Concepts	8
1.1.1	8
1.1.2	8
1.1.3	9
1.1.4	9
1.1.5	9
1.1.6	10
1.1.7	10
1.1.8	10
1.1.9	10
1.1.10	10
1.2	Functions	10
1.2.1	10
2	Topological Spaces and Continous Functions	12
2.1	Topological Spaces	12
2.2	Basis for a Topology	12
2.2.1	12
2.2.2	13
2.2.3	13
2.2.4	15
2.2.5	16
2.2.6	17
2.2.7	17
2.2.8	18
2.3	The Order Topology	19
2.4	The Product Topology on $X \times Y$	19
2.5	The Subspace Topology	19

2.5.1	19
2.5.2	20
2.5.3	20
2.5.4	21
2.5.5	21
2.5.6	22
2.5.7	22
2.5.8	23
2.5.9	23
2.5.10	24
2.6 Closed Sets and Limit Points	24
2.6.1	24
2.6.2	25
2.6.3	25
2.6.4	26
2.6.5	26
2.6.6	26
2.6.7	27
2.6.8	27
2.6.9	28
2.6.10	29
2.6.11	29
2.6.12	29
2.6.13	29
2.6.14	30
2.6.15	31
2.6.16	31
2.6.17	31
2.6.18	32
2.6.19	33
2.6.20	34
2.6.21	34
2.7 Continuous Functions	35
2.7.1	35
2.7.2	36
2.7.3	36
2.7.4	37
2.7.5	37
2.7.6	38
2.7.7	38
2.7.8	39

2.7.9	40
2.7.10	41
2.7.11	42
2.7.12	42
2.7.13	43
2.8 The Product Topology	44
2.8.1	44
2.8.2	44
2.8.3	45
2.8.4	45
2.8.5	45
2.8.6	45
2.8.7	46
2.8.8	47
2.8.9	47
2.8.10	47
2.9 The Metric Topology	48
2.9.1	48
2.9.2	49
2.9.3	49
2.9.4	52
2.9.5	55
2.9.6	55
2.9.7	56
2.9.8	61
2.9.9	62
2.10 The Metric Topology (continued)	62
2.10.1	62
2.10.2	63
2.10.3	63
2.10.4	64
2.11 The Quotient Topology	64
3 Connectedness and Compactness	65
3.1 Connected Spaces	65
3.1.1	65
3.1.2	65
3.1.3	65
3.1.4	66
3.1.5	66
3.1.6	66

3.1.7	67
3.1.8	67
3.1.9	67
3.1.10	68
3.2	Connected Subspaces of the Real Line	69
3.2.1	69
3.2.2	69
3.2.3	70
3.2.4	71
3.2.5	72
3.2.6	72
3.2.7	73
3.2.8	74
3.2.9	75
3.2.10	75
3.2.11	76
3.3	Compact Spaces	76
3.3.1	76
3.3.2	77

Preface

Those are my solutions for the James Munkres' "Topology", 2nd edition.

Majority of the notation that is used here migrated from my course on the set theory. In my very personal opinion, notation that is used there is far superior that whatever is happening in Munkres' book. Sometimes I use some abusive notation when it is painfully clear what's going on.

If you decide to persue the study of topology yourself, then I highly recommend firstly to go through a course on axiomatic set theory and logic, because first chapter of this book is highly insufficient in this regard. My personal recommendations are the combo by Cunningham, which includes "Set theory: A first course" and "A Logical Introduction to Proof", or "A first course in Mathematical Logic and Set Theory" by Michael L. O'Leary for both subjects.

Notation

Sometimes I use specific notation, that migrated from my previous endeavours in pure maths. This notation includes:

$$V_{\epsilon}(x) = (x - \epsilon, x + \epsilon)$$

Part I

General Topology

Chapter 1

Set Theory and Logic

1.1 Fundamental Concepts

1.1.1

*Check distributive and DML laws
GOTO set theory book*

1.1.2

Determine which of the following are true.

- (a) - impl
- (b) - impl
- (c) - true
- (d) - rimpl
- (e) - \subseteq , true if $B \subseteq A$.
- (f) - \supseteq ; $A - (B - A) = A$.
- (g) - true
- (h) - \supseteq
- (i) - true
- (j) - true
- (k) - false
- (l) - true
- (m) - \subseteq
- (n) - true
- (o) - true
- (p) - true
- (q) - \supseteq

1.1.3

(a) Write a contrapositive and converse of the following statement: "If $x < 0$, then $x^2 - x > 0$ " and determine which ones are true

Contrapositive:

$$x^2 - x \leq 0 \Rightarrow x \geq 0$$

Converse

$$x^2 - x > 0 \Rightarrow x < 0$$

Contrapositive is correct, converse is incorrect ($2^2 - 2 > 0$)

(b) Do the same for the statement $x > 0 \Rightarrow x^2 - x > 0$

Contrapositive:

$$x^2 - x \leq 0 \Rightarrow x \leq 0$$

Converse

$$x^2 - x > 0 \Rightarrow x > 0$$

Contrapositive is false ($1^2 - 1 = 0$); Converse is also false ($(-2)^2 - (-2) = 6$).

1.1.4

Let A and B be the sets of real numbers. Write the negation of each of the following statements:

(a)

$$(\exists a \in A)(a^2 \notin B)$$

(b)

$$(\forall a \in A)(a^2 \notin B)$$

(c)

$$(\exists a \in A)(a^2 \in B)$$

(d)

$$(\forall a)(a \notin A \Rightarrow a^2 \notin B)$$

1.1.5

Let A be a nonempty collection of sets. Determine the truths of each of the following and their converses

(a)

$$x \in \bigcup A \Leftrightarrow (\exists B \in A)(x \in B)$$

(b)

$$x \in \bigcup A \Leftrightarrow (\forall B \in A)(x \in B)$$

(c)

$$x \in \bigcap A \Rightarrow (\exists B \in A)(x \in B)$$

(d)

$$x \in \bigcap A \Leftrightarrow (\forall B \in A)(x \in B)$$

1.1.6

Skip

1.1.7

skip

1.1.8

GOTO set theory book

1.1.9*Formulate DML for arbitrary unions and intersections*

$$A \setminus \bigcap (B) = \bigcup (A \setminus B)$$

$$A \setminus \bigcup (B) = \bigcap (A \setminus B)$$

For the proof goto set theory or real analysis book

1.1.10

(a, b, d) are true

1.2 Functions**1.2.1***Let $f : A \rightarrow B$. Let $A_0 \subseteq A$ and $B_0 \subseteq B$.**(a) Show that $A_0 \subseteq f^{-1}[f[A_0]]$ and that equality holds if f is injective.*

Suppose that $x \in A_0$. We follow that there exists $\langle x, y \rangle \in f$ for some $y \in f[A_0]$. Therefore there exists $\langle y, x \rangle \in f^{-1}$. Because $y \in f[A_0]$, we follow that $x \in f^{-1}[f[A_0]]$. Therefore $A_0 \subseteq f^{-1}[f[A_0]]$.

Suppose that f is injective. Suppose that there exists $x_0 \in f^{-1}[f[A_0]]$ such that $x_0 \notin A_0$. We follow that $\langle y, x_0 \rangle, \langle y, x \rangle \in f^{-1}$, therefore $\langle x_0, y \rangle, \langle x, y \rangle \in f$, and because $x_0 \neq x$ we follow that we've got a contradiction.

((b)

pretty similar to (a)

This chapter practicly mirrors the content of my set theory course . Gonna skip it for now, and will come back if the need arises.

Chapter 2

Topological Spaces and Continuous Functions

2.1 Topological Spaces

I want to state here that if $\mathcal{T} \subseteq \mathcal{P}(X)$ satisfies properties

$$\{X, \emptyset\} \subseteq \mathcal{T}$$

$$(\forall Y \in \mathcal{P}(\mathcal{T}))(\bigcup U \in \mathcal{T})$$

$$(\forall Y \in \mathcal{P}(\mathcal{T}))(Y \neq \emptyset \wedge |Y| <_c |\omega| \rightarrow \bigcap U \in \mathcal{T})$$

then \mathcal{T} is a topology on X .

2.2 Basis for a Topology

Let $Y \subseteq \mathcal{P}(X)$. If

$$(\forall x \in X)(\exists y \in Y)(x \in y)$$

and

$$(\forall x \in X)(\exists y_1, y_2, y_3 \in Y)(x \in y_1 \cap y_3 \rightarrow x \in y_3 \wedge y_3 \subseteq y_1 \cap y_2)$$

then Y is a basis for a topology on X .

2.2.1

Let X be a topological space; Let A be a subset of X . Suppose that for each $x \in A$ there is an open set U containing x such that $U \subseteq A$. Show that A is open in X .

Let $U : A \rightarrow \mathcal{P}(A)$ be an indexed function such that

$$x \in U(x) \wedge U(x) \subseteq A \wedge U(x) \in \mathcal{T}(X)$$

We want to show that $A = \bigcup \text{ran}(U)$. Suppose that $x \in A$. We follow that $x \in U(x)$. Thus $x \in \bigcup \text{ran}(U)$. Therefore $A \subseteq \bigcup \text{ran}(U)$.

Suppose that $z \in \bigcup \text{ran}(U)$. We follow that

$$(\exists Y \in \text{ran}(U))(z \in Y) \Rightarrow (\exists x \in A)(z \in U(x))$$

Since $(\forall x \in A)(U(x) \subseteq A)$, we follow that $z \in A$. Thus $\bigcup \text{ran}(U) = A$.

Because $(\forall x \in A)(U(x) \in \mathcal{T}(X))$, we follow that

$$\text{ran}(U) \subseteq \mathcal{T}(A)$$

, therefore by definition of topology we follow that

$$\bigcup \text{ran}(U) \in \mathcal{T}(X)$$

as desired.

2.2.2

Too tedious, skip

2.2.3

Show that the collection \mathcal{T}_c given in Example 4 of p. 12 is a topology on the set X . Is the collection

$$\mathcal{T}_\infty = \{U \in \mathcal{P}(X) : |X \setminus U| \leq_c |\omega| \vee X \setminus U = \emptyset \vee X \setminus U = X\}$$

a topology on X ?

We firstly state that

$$\mathcal{T}_c = \{U \in \mathcal{P}(X) : |X \setminus U| \leq_c |\omega| \vee X \setminus U = X\}$$

We can follow that $X \setminus X = \emptyset$, which is countable, thus $X \in \mathcal{T}_c$. $X \setminus \emptyset = X$, therefore $\emptyset \in \mathcal{T}_c$.

Suppose that $U' \subseteq \mathcal{T}_c$. If $U' = \{\emptyset\}$, then $X \setminus \bigcap U' = X$ and $X \setminus \bigcup U' = X$. Thus assume that $U' \neq \{\emptyset\}$.

We follow that

$$(\forall u \in U')(|X \setminus u| \leq_c |\omega| \vee X \setminus u = X)$$

We follow that if $\emptyset \in U'$, then $\bigcup U' = \bigcup (U' \setminus \{\emptyset\})$. Then we follow by DML that

$$X \setminus \bigcup \{U'\} = X \setminus \bigcup \{U' \setminus \{\emptyset\}\} = \bigcap_{U' \setminus \{\emptyset\}} X \setminus u$$

we know that $(\forall u \in U')(|X \setminus u| \leq_c |\omega|)$. For any $u \in U'$ we follow that

$$\bigcap_{u \in U' \setminus \{\emptyset\}} X \setminus u \subseteq X \setminus u'$$

and given that $X \setminus u'$ is countable, we follow that $\bigcap_{u \in U'} X \setminus u$ is countable as well, thus $\bigcup U' \in \mathcal{T}_c$.

Now let $U' \subseteq \mathcal{T}_c$ and $|U'| < |\omega|$ and $U' \neq \{\emptyset\}$. We follow that if $\emptyset \in U'$, then $\bigcap U' = \emptyset$, and therefore $X \setminus \bigcap U' = X$. Therefore assume that $\emptyset \notin U'$.

Then we can follow that

$$X \setminus \bigcap U' = \bigcup_{u \in U'} X \setminus u$$

Given that U' is countable and $X \setminus u$ is countable we follow that $\bigcup_{u \in U'} X \setminus u$ is countable, thus $X \setminus \bigcap U'$ is countable.

Therefore we conclude that \mathcal{T}_c is a topology on X .

Now let us consider T_∞ . We can state that $X \in T_\infty$ because $X \setminus X = \emptyset$. Because $X \setminus \emptyset = X$, we follow that $\emptyset \in T_\infty$.

Suppose that X is not infinite and $T_\infty \neq \{\emptyset, X\}$. Then there exists $u \in T_\infty$ such that $u \neq \emptyset$ and $u \neq X$. Therefore $X - u$ is nonempty finite set, therefore $u \notin T_\infty$, which is a contradiction. Therefore we conclude that if X is finite, then T_∞ is a trivial topology.

If X is infinite, then we follow that we can have an injection $f : \omega \rightarrow X$. Let O be the set of odd naturals and E be the set of evens. Then we follow that

$$|X \setminus f[O]| = |f[E]| \geq_c |\omega|$$

and

$$|X \setminus f[E]| =_c |f[O]| \geq_c |\omega|$$

which tells us that $f[O]$ and $f[E]$ are both in X . We can also follow that

$$|X \setminus f[O \cup \{0\}]| \geq |\omega|$$

thus $f[O \cup \{0\}] \in \mathcal{T}_\infty$. This gives us that

$$f[E] \cap f[O \cup \{0\}] = \{f(0)\} \in \mathcal{T}_\infty$$

but $\{f(0)\}$ is a finite nonempty set for which none of the conditions of \mathcal{T}_∞ hold. Therefore we conclude that if X is infinite, then \mathcal{T}_∞ is not a topology.

Therefore we conclude that if X is a finite set, then T_∞ is equal to a trivial topology; if X is infinite, then T_∞ is not a topology at all, since it is not closed under finite intersections.

2.2.4

(a) if $\{\mathcal{T}_\alpha\}$ is a family of topologies on X , show that $\bigcap \mathcal{T}_\alpha$ is a topology on X . Is $\bigcup \mathcal{T}_\alpha$ a topology on X ?

Since every topology on X has X and \emptyset as elements, we follow that

$$\{X, \emptyset\} \subseteq \bigcap \mathcal{T}_\alpha$$

If $Y \subseteq \bigcap \mathcal{T}_\alpha$, then we follow that

$$(\forall Z \in \{\mathcal{T}_\alpha\})(\bigcap \mathcal{T}_\alpha \subseteq Z)$$

$$(\forall Z \in \{\mathcal{T}_\alpha\})(Y \subseteq Z)$$

since every Z is a topology, we follow that

$$(\forall Z \in \{\mathcal{T}_\alpha\})(\bigcup Y \in Z)$$

$$\bigcup Y \in \bigcap \mathcal{T}_\alpha$$

If Y is finite and nonempty, we can also follow that

$$(\forall Z \in \{\mathcal{T}_\alpha\})(Y \in Z) \Rightarrow (\forall Z \in \{\mathcal{T}_\alpha\})(\bigcap Y \in Z) \Rightarrow \bigcap Y \in \bigcap \mathcal{T}_\alpha$$

thus we conclude that $\bigcap \mathcal{T}_\alpha$ is a topology.

$\bigcup \mathcal{T}_\alpha$ is not necessarily a topology. Although $\{X, \emptyset\} \in \bigcup \mathcal{T}_\alpha$, we cannot follow that the topology is closed under unions. Case in point: Let $X = \{a, b, c\}$ and

$$\mathcal{T}_1 = \{\emptyset, X, \{a\}\}, \mathcal{T}_2 = \{\emptyset, X, \{b\}\}$$

then $Y = \mathcal{T}_1 \cup \mathcal{T}_2$ does not contain $\{a, b\}$, which would be necessary for this case. Thus we conclude that in general we can't have implications for $\bigcup \mathcal{T}_\alpha$.

(b) Let $\{\mathcal{T}_\alpha\}$ be a family of topologies on X . Show that there is a unique smallest topology on X containing all the collections \mathcal{T}_α and a unique largest topology contained in all \mathcal{T}_α .

Let us take $\bigcup \{\mathcal{T}_\alpha\}$. We cannot follow that presented set is a topology on X , nor can we state that it is a basis of a topology. Former is followed from the discussion in the previous section of this exercise, and the latter cannot be followed because we don't necessarily satisfy the second point of the definition of the basis. Namely, we don't have that

$$(\forall x \in X)(\exists y_1, y_2, y_3 \in \bigcup \{\mathcal{T}_\alpha\})(x \in y_1 \cap y_3 \rightarrow x \in y_3 \wedge y_3 \subseteq y_1 \cap y_2)$$

Let Q be a set of all of the intersections of finite nonempty subsets of $\bigcup \{\mathcal{T}_\alpha\}$. We follow that $(\forall x \in \bigcup \{\mathcal{T}_\alpha\})(x = \bigcap \{x\})$, therefore $\bigcup \{\mathcal{T}_\alpha\} \subseteq Q$. Thus we follow that Q satisfies

the first requirement for the basis of X . Now let $x \in X$ be such that there exist $y_1, y_2 \in Q$ such that $x \in y_1 \cap y_2$. We follow that there exist finite subsets $Y_1, Y_2 \subseteq \bigcup \{\mathcal{T}_\alpha\}$ such that

$$y_1 = \bigcap Y_1 \wedge y_2 = \bigcap Y_2$$

therefore

$$y_1 \cap y_2 = \bigcap Y_1 \cap \bigcap Y_2$$

which is an intersection of a finite subset of $\bigcup \mathcal{T}_\alpha$. Thus we follow that there exists $y_3 \in Q$ such that $x \in y_3 \wedge y_3 \subseteq y_1 \cap y_2$. Therefore we can follow that the set Q is indeed a basis for a topology on X . Let us name the topology generated by this set as \mathcal{T}_q .

Suppose that there is a topology, which contains all of the topologies $\{\mathcal{T}_\alpha\}$. Then we follow that it contains $\bigcup \{\mathcal{T}_\alpha\}$, therefore we follow that it contains all of the unions of $\bigcup \{\mathcal{T}_\alpha\}$, and finite intersections of subsets of $\bigcup \{\mathcal{T}_\alpha\}$, and thus it contains \mathcal{T}_q . Therefore we follow that \mathcal{T}_q is the smallest topology, which contains all the topologies of $\{\mathcal{T}_\alpha\}$.

Suppose that \mathcal{T}_p is a topology, which is contained in all of the $\{\mathcal{T}_\alpha\}$. Then we follow that $\mathcal{T}_p \subseteq \bigcap \mathcal{T}_\alpha$. Because $\bigcap \mathcal{T}_\alpha$ is a topology itself, we follow that it is the largest topology, which is contained in all of the $\{\mathcal{T}_\alpha\}$.

(c) If $X = \{a, b, c\}$, let

$$\mathcal{T}_1 = \{\emptyset, X, \{a\}, \{a, b\}\}$$

$$\mathcal{T}_2 = \{\emptyset, X, \{a\}, \{b, c\}\}$$

Find the smallest topology containing \mathcal{T}_1 and \mathcal{T}_2 , and the largest topology contained in $\mathcal{T}_1, \mathcal{T}_2$.

We can follow from previous discussions that largest contained topology is

$$\{\emptyset, X, \{a\}\}$$

and the smallest containing topology is

$$\{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{b, c\}\}$$

2.2.5

Show that if A is a basis for a topology on X , then the topology generated by A equals the intersection of all topologies on X that contains A . Prove the same if A is a subbasis.

Let A be a subbasis. Let $\{\mathcal{T}_\alpha\}$ be a set of topologies, that contain A and \mathcal{T}_A is a topology generated by A . We can follow that $\mathcal{T}_A \in \{\mathcal{T}_\alpha\}$, therefore $\bigcap \{\mathcal{T}_\alpha\} \subseteq \mathcal{T}_A$. If $x \in \mathcal{T}_A$, then we follow that there exists a subset $B \subseteq A$ such that x is equal to some union of some finite intersections of B . Since $B \subseteq A$, we follow that $(\forall y \in \mathcal{T}_\alpha)(B \subseteq y)$. Therefore all of the finite intersections of B are in any topology of \mathcal{T}_α . Therefore all of the unions of those intersections are in any \mathcal{T}_α . Therefore we conclude that $(\forall y \in \mathcal{T}_\alpha)(x \in y)$.

and thus $x \in \bigcap \mathcal{T}_\alpha$. Therefore we conclude that $\mathcal{T}_A \subseteq \bigcap \mathcal{T}_\alpha$, and by double inclusion we get that $\mathcal{T}_A = \bigcap \mathcal{T}_\alpha$, as desired.

Since every basis of a topology is a subbasis by first clause of the definition, we follow that the desired result holds for bases as well.

2.2.6

Show that the topologies of R_l and R_k are not comparable.

Let $[0, 1)$ be an element of a basis of topology R_l . Then we follow that there are no elements of basis of standard topology on R that contains 0 and lies inside $[0, 1)$. We can follow this by contradiction

Suppose that $0 \in (x, y)$ and $(x, y) \subseteq [0, 1)$. Since $0 \in (x, y)$, we follow that $x < 0$. Thus we conclude that there exists $n \in \mathbb{Z}_+$ such that $1/n < |x|$. Therefore $-1/n \in (x, y)$ and $-1/n \notin [0, 1)$ which gives us that $(x, y) \not\subseteq [0, 1)$, which is a contradiction. The same logic applies to any element of basis of R_k .

Now let us look at the basis element $(-1, 1) \setminus K$ and the point 0. We can follow that $0 \in (-1, 1) \setminus K$ and suppose that there exists basis element of R_l $[a, b)$ that has point 0 and is contained within $(-1, 1) \setminus K$. Since $0 \in [a, b)$, we follow that $a \leq 0 < b$. Thus we conclude that there exists $n \in \mathbb{Z}_+$ such that $0 < 1/n < b$. Thus we conclude that $1/n \in [a, b)$ and $1/n \notin (-1, 1) \setminus K$, since $1/n \in K$ for all $n \in \mathbb{Z}_+$. Thus we conclude that R_k and R_l are not comparable, as desired.

2.2.7

Consider the following topologies on R :

$\mathcal{T}_1 = \text{the standard topology on } R$

$\mathcal{T}_2 = \text{the topology of } R_k$

$\mathcal{T}_3 = \text{the finite complement topology}$

$\mathcal{T}_4 = \text{the upper limit topology, having all sets } (a, b] \text{ as basis}$

$\mathcal{T}_5 = \text{the topology having all sets } (-\infty, a) = \{x : x < a\} \text{ as a basis}$

Determine, for each of these topologies, which of the others it contains

We can follow that \mathcal{T}_2 contains \mathcal{T}_1 , since it's finer, as proven in the chapter. The reverse is not true, as proven in the chapter.

We can follow that \mathcal{T}_3 does not contain \mathcal{T}_1 , because if it is, then we follow that $(-\infty, a] \cup [b, \infty)$ has finite number of points. The reverse is true, since we can divide each element of a finite complement into a union of open intervals. For example, if $x \in \mathcal{T}_3$ is such that $x = R \setminus \{x_1, x_2, x_3\}$ and $x_1 < x_2 < x_3$, then we can state that $x = (-\infty, x_1) \cup (x_1, x_2) \cup (x_2, x_3) \cup (x_3, \infty)$. We can follow that middle 2 intervals are in the basis of standard

topology, and two infinite intervals are unions of infinite set of intervals of basis. Thus \mathcal{T}_1 is strictly finer than \mathcal{T}_3 .

We can follow that the same logic, that worked with lower limit, works with upper limit as well. thus we conclude that \mathcal{T}_4 is strictly finer than \mathcal{T}_1 .

We can follow that for $(-\infty, a) \in \mathcal{T}_5$ we can get a sequence $(x_n) = a - n$, then get a set of intervals $\{(a, a - 1), (x_{n+1}, x_n)\}$, all of which are in the basis of standard topology, get another set $\{V_{0,1}(x_n)\}$ to patch the holes in this set, and take union of unions of both sets to get that $(-\infty, a) \in \mathcal{T}_1$.

For (a, b) - a set in the basis of standard topology we follow that every set in the basis of \mathcal{T}_5 contains $a - 1$, thus we conclude that $(a, b) \notin \mathcal{T}_5$. Thus we conclude that \mathcal{T}_1 is strictly finer than \mathcal{T}_5 .

Topology \mathcal{T}_2 is strictly finer than \mathcal{T}_1 , therefore we follow that topologies that are finer than \mathcal{T}_1 are a subset of \mathcal{T}_2 . This includes \mathcal{T}_3 and \mathcal{T}_5 . (Almost) the same reasoning that worked with R_k and R_l can be applied to show that \mathcal{T}_2 is not finer than \mathcal{T}_4 . On the other hand, suppose that $x \in X$ and $y \in \mathcal{T}_2$ is such that $x \in y$. We follow that if $y \in \mathcal{T}_1$, then there exists an element of \mathcal{T}_4 that is finer than y . Thus assume that $y \notin \mathcal{T}_1$ and therefore is in the form $y = (a, b) \setminus K$ for some $a, b \in R$. If $x \leq 0$, then we can have set $(a, x] \subseteq y$ that will satisfy. Thus assume that $x > 0$. We follow that there exists $n \in \mathbb{Z}_+$ such that $1/n < x$. By well-ordering properties of \mathbb{Z}_+ we follow that there exists lowest $n \in \mathbb{Z}_+$ such that $1/nx$. Therefore we follow that there are no elements $z \in K$ such that $1/n < z < x$. Since $x \in (a, b) \setminus K$, we follow that $x \notin K$, therefore $(\forall y \in (1/n, x))(y \in x \in (a, b) \setminus K)$. Therefore we conclude that \mathcal{T}_4 is strictly finer than \mathcal{T}_2 , which is neat.

\mathcal{T}_3 is strictly coarser than $\mathcal{T}_1, \mathcal{T}_2$. Since \mathcal{T}_4 is strictly finer than \mathcal{T}_2 , we follow that \mathcal{T}_3 is coarser than \mathcal{T}_4 . Suppose that $a < x < b$ and let $y = R \setminus \{a, b\}$ be an element of \mathcal{T}_3 . Then we follow that no element of basis of \mathcal{T}_5 has x and does not have a . If $(-\infty, a)$ is an element of \mathcal{T}_5 , then we follow that every element of topology \mathcal{T}_3 has numbers greater than a in it (since there are infinitely many of them). Thus we conclude that no element of \mathcal{T}_3 is a subset of $(-\infty, a)$. Thus we conclude that \mathcal{T}_3 and \mathcal{T}_5 are not comparable.

And after all of the discussion, we can conclude that

$$[\mathcal{T}_3 | \mathcal{T}_5] \subset \mathcal{T}_1 \subset \mathcal{T}_2 \subset \mathcal{T}_4$$

is the desired conclusion.

2.2.8

(a) Apply Lemma 13.2 to show that the countable collection

$$B = \{(a, b) : a < b \wedge a, b \in \mathbb{Q}\}$$

is a basis that generates the standard topology on R .

Denote \mathcal{T} as a standard topology on R . Let $x \in \mathcal{T}$. We follow that there exists an interval (a, b) in basis of standard topology such that $x \in (a, b)$. We can follow that there

exist $a', b' \in Q$ such that $a < a' < x < b' < b$ (otherwise we run into some problem with density of rationals in reals). Therefore we follow that $x \in (a', b')$. Lemma 13.2 tells us that the presented result implies that B is a basis for standard topology, as desired.

(b) Show that the collection

$$C = \{[a, b) : a < b \wedge a, b \in Q\}$$

is a basis that generates a topology different from the lower limit topology on R .

Proof that C is a basis is trivial. Let us look at $[\sqrt{2}, 2)$ - an element of R_l . Suppose that $c = [a, b) \in C$ is such that $\sqrt{2} \in c$. Because $\sqrt{2} \notin Q$, we follow that $a \neq \sqrt{2}$, therefore $a < \sqrt{2} < b$. Therefore we can conclude that C is not finer than R_l . Proving that C is a subset of R_l is trivial, thus we conclude that R_l is strictly finer than C , and thus C generates a topology different than R_l , as desired.

2.3 The Order Topology

2.4 The Product Topology on $X \times Y$

2.5 The Subspace Topology

2.5.1

Show that if Y is a subspace of X , and A is a subspace of Y , then the topology A inherits as a subspace of Y is the same as the topology it inherits as a subspace of X .

Suppose that Q is an open set in A with respect to topology, inherited from X . We follow that there exists an open set in X $Q_x \subseteq X$ such that $Q = Q_x \cap A$ by definition of a subspace topology. We follow that there exists open in Y set $Q_y \subseteq Y$ such that $Q_y = Q_x \cap Y$. With respect to Q_y there exists an open in A set $Q' = Q_y \cap A$. Thus

$$Q' = Q_y \cap A$$

$$Q' = Q_x \cap Y \cap A$$

Since $A \subseteq Y$, we follow that $Y \cap A = A$. Thus

$$Q' = Q_x \cap (Y \cap A)$$

$$Q' = Q_x \cap A$$

$$Q' = Q$$

Therefore we conclude that if Q is in topology of A inherited from X , then Q is also in a topology of A inherited from Y . Proof of the converse is pretty much the same proof

Here's another, more logical and rigorous proof. Denote topology of A inherited from Y by \mathcal{T}_A and topology of A inherited from X by \mathcal{T}'_A . Also denote topology of X by \mathcal{T}_X and topology of Y inherited from X by \mathcal{T}_Y . Then we can state that

$$\begin{aligned} Q \in \mathcal{T}_A &\Leftrightarrow (\exists Q_y \in \mathcal{T}_Y)(Q = Q_y \cap A) \Leftrightarrow (\exists Q_X \in \mathcal{T}_X)(Q_y = Q_x \cap Y \wedge Q = Q_y \cap A) \Leftrightarrow \\ &\Leftrightarrow (\exists Q_X \in \mathcal{T}_X)(Q = Q_x \cap Y \cap A) \Leftrightarrow (\exists Q_X \in \mathcal{T}_X)(Q = Q_x \cap (Y \cap A)) \Leftrightarrow \\ &\Leftrightarrow (\exists Q_X \in \mathcal{T}_X)(Q = Q_x \cap A) \Leftrightarrow Q \in \mathcal{T}'_A \end{aligned}$$

thus $\mathcal{T}'_A = \mathcal{T}_A$ by extensionality axiom.

2.5.2

if \mathcal{T} and \mathcal{T}' are topologies on X and \mathcal{T}' is strictly finer than \mathcal{T} , what can you say about the corresponding topologies on the subset Y of X .

Denote corresponding topologies by \mathcal{T}'_Y and \mathcal{T}_Y . There're three plausible cases:

- 1 - we can't say nothing
- 2 - $\mathcal{T}'_Y \supset \mathcal{T}_Y$
- 3 - $\mathcal{T}'_Y \supseteq \mathcal{T}_Y$

I'm betting on the second case, so let us try to prove that. In order to do that, let us firstly prove the third case, which is a "subcase" of the second.

Suppose that $Q \in \mathcal{T}_Y$. We follow that there exists $Q_X \in \mathcal{T}$ such that $Q = Q_X \cap Y$. Since $Q_X \in \mathcal{T}$, we follow by $\mathcal{T} \subset \mathcal{T}'$ that $Q_X \in \mathcal{T}'$. Thus $Q = Q_X \cap Y$ implies that $Q \in \mathcal{T}'_Y$. Therefore we follow that $\mathcal{T}'_Y \supseteq \mathcal{T}_Y$.

Although I'm betting on the second case, it seems that I'm not getting my money back. We can follow that second case is not always true, if we substitute \emptyset for Y . Then $\mathcal{T}_Y = \mathcal{T}'_Y = \emptyset$. If we look into topologies of some almost-trivial set, such as $X = \{a, b, c\}$, then I think that we can come up with a more persuasive case as well. Therefore we conclude that presented conditions imply that $\mathcal{T}_Y \subseteq \mathcal{T}'_Y$.

2.5.3

Consider the set $Y = [1, 1]$ as a subspace of R . Which of the following sets are open in Y ? Which are open in R ?

$$A = \{x : \frac{1}{2} < |x| < 1\}$$

$$B = \{x : \frac{1}{2} < |x| \leq 1\}$$

$$C = \{x : \frac{1}{2} \leq |x| < 1\}$$

$$D = \{x : \frac{1}{2} \leq |x| \leq 1\}$$

$$E = \{x : 0 < |x| < 1 \wedge 1/x \notin Z_+\}$$

We can follow that $A = (-1, -1/2) \cup (1/2, 1)$ is open in both Y and R .

$B = [-1, -1/2) \cup (1/2, 1]$ is a union of two rays in Y , therefore we follow that it is open in Y . For R we've got that there is no open interval, that contains a point 1 and does not contain anything larger than 1. Therefore we conclude that given set is not a union of open intervals, and therefore it is not open in R .

We can follow pretty easily that C and D are not open in both Y and R since there is no open interval/ray that contains $1/2$ and does not contain anything in the interval $(-1/2, 1/2)$.

We can represent E as

$$E = (-1, 0) \cup ((0, 1) \setminus K)$$

We follow that $(-1, 0)$ is an element of a basis of both Y and R . Suppose that $x \in (0, 1) \setminus K$. Then we follow that there exist lowest $n_1 \in Z_+$ such that $1/n_1 < x < 1/(n_1 + 1)$. Therefore we can conclude that if $x \in E$, then there exist a basis element Q of both Y and R such that $x \in Q \subseteq Y, R$. Therefore we follow that E is an open set in both Y and R .

2.5.4

A map $f : X \rightarrow Y$ is said to be an open map if for every open set U of X , the set $f(U)$ is open in Y . Show that $\pi_1 : X \times Y \rightarrow X$ and $\pi_2 : X \times Y \rightarrow Y$ are open maps.

Suppose that $Q \in X \times Y$ is an open set. Therefore we follow that it is a union of some element of a basis of $X \times Y$, therefore there exist a subset R of a basis of $X \times Y$ such that $Q = \bigcup R$. From a set theory course we know that

$$U[\bigcup G] = \bigcup \{R[C] : C \in G\}$$

for any relation U . Therefore we can follow that the same result holds for functions π_1, π_2 . We can follow that for any $r \in R$ we've got that both $\pi_1(r)$ and $\pi_2(r)$ are open by the definition of a basis for the product topology. Therefore we conclude that $\pi_1[Q] = \pi_1[\bigcup R] = \bigcup \{\pi_1[\bigcup r] : r \in R\}$. Therefore we conclude that $\pi_1[Q]$ is a union of open sets of X , therefore we conclude that it is in topology of X . We can follow the similar result for π_2 using similar logic.

2.5.5

Let X and X' denote a single set in the topologies \mathcal{T} and \mathcal{T}' respectively; let Y and Y' denote a single set in the topologies U and U' , respectively. Assume that these sets are nonempty.

There're a couple of ways to deconstruct the text of this exercise: we can assume that $X = X', Y = Y', X \in \mathcal{T}, X' \in \mathcal{T}', Y \in U$ and $Y' \in U'$, or we can assume that $X \in \mathcal{T}, X' \in \mathcal{T}', Y \in U$ and $Y' \in U'$ without $X = X'$ and $Y = Y'$. The latter case will obviously

present some problems in the proofs, therefore we will assume that the author intended to use the former case.

(a) Show that if $\mathcal{T}' \supseteq \mathcal{T}$ and $U' \supseteq U$, then the product topology on $X' \times Y'$ is finer than the product topology on $X \times Y$.

Let \mathcal{B} denote the basis for $\mathcal{T}_{X \times Y}$. Let $q \in X \times Y$. Because \mathcal{B} is a basis for $\mathcal{T}_{X \times Y}$ we follow that there exists $b \in \mathcal{B}$ such that $q \in b$. Since $b \in \mathcal{B}$, we follow that there exist $x \in \mathcal{T}$ and $y \in U$ such that $b = x \times y$. Since $\mathcal{T} \subseteq \mathcal{T}'$ and $U \subseteq U'$, we follow that $x \in \mathcal{T}'$ and $y \in U'$. Therefore $x \times y \in \mathcal{B}'$ where \mathcal{B}' denotes the basis for $\mathcal{T}_{X' \times Y'}$. Therefore we conclude that for every $q \in X \times Y$ and every basis element $q \in \mathcal{B}$ there exists $q' \in \mathcal{B}'$ such that $q' \subseteq q$ and $q \in q'$. Therefore we conclude that $\mathcal{T}_{X \times Y} \subseteq \mathcal{T}_{X' \times Y'}$, as desired.

(b) Does the converse of (a) hold? Justify your answer.

Let \mathcal{T} and \mathcal{T}' be defined on a set $Q = \{a, b\}$ and U and U' be defined on $W = \{c, d\}$. Let $X = X' = \{a\}$, $Y = Y' = \{c\}$, $\mathcal{T} = \{\emptyset, \{a\}, \{b\}, Q\}$, $\mathcal{T}' = \{\emptyset, \{a\}, Q\}$, and $U = U'$. Then we follow that topology defined on $X \times Y$ is finer than the topology defined on $X' \times Y'$ (and vice versa), but \mathcal{T}' is not finer than \mathcal{T} .

2.5.6

Show that the countable collection

$$\{(a, b) \times (c, d) : a < b \wedge c < d \wedge a, b, c, d \in \mathbb{Q}\}$$

is a basis for \mathbb{R}^2 .

Let us denote this set by L . Suppose that $x \in \mathbb{R}^2$. We follow that there exist $x_1, x_2, y_1, y_2 \in \mathbb{Q}$ such that $x_1 < x < x_2$ and $y_1 < y < y_2$, therefore $(\exists l \in L)(x \in l)$. Thus we follow that the first condition of a definition of a basis is satisfied. The last condition can be satisfied by through the argument about the density of rationals in reals.

We can follow that topology, that is presented by given basis is a subset of the standard topology on \mathbb{R}^2 , and we can follow though pretty much the same argument that given topology is finer than the standard topology. Therefore I'm pretty sure that we can state that given basis generates the standard topology (I'll not provide any proof of that, just stating what I think).

2.5.7

Let X be an ordered set. If Y is a proper subset of X that is convex in X , does it follow that Y is an interval or a ray in X ?

Don't think so. I think that the author tries to give us a hint to what's going to come afterwards (probably something about the completeness and whatnot).

Pretty sure, that we don't need to prove that \mathbb{Q} is a totally ordered set, so we're going to take it as a given. Let

$$M = \{x \in \mathbb{Q} : x^2 < 2 \wedge x \geq 0\}$$

(I've added the latter condition in order not to be bogged down by several cases, depending on the sign). Let $x < y \in M$. Suppose that $z \in Q$ is such that $x < z < y$. Then we follow that $z > x \geq 0$, thus $z > 0$. Since all of the numbers are positive, we're justified to square them and get that $x^2 < z^2 < y^2$. Given that $y^2 < 2$, we conclude that $z^2 < 2$ as well. Therefore we follow that $z \in M$. Thus we can follow that $z \in (x, y) \Rightarrow z \in M$. Therefore we can state that presented set is convex.

Given that M is bounded above and below, we follow that it is not a ray. Suppose that it is an interval. Then we follow that there exists $k \in Q$ such that $M = [0, k)$. Therefore we follow that k is a least lower bound of M , which is not the case, as proven in numerous real analysis books. Thus we conclude that M is not an interval.

2.5.8

If L is a straight line in the plane, describe the topology L inherits as a subspace of $R_l \times R$ and as a subspace of $R_l \times R_l$. In each case it is a familiar topology.

Let \mathcal{B} be the basis for $R_l \times R$ and \mathcal{B}' be the basis for $R_l \times R_l$. Let $q \in \mathcal{B}$ and suppose that $b = L \cap q \neq \emptyset$. From plotting elements of the basis and the line itself on the graph, we can conclude that b is some sort of an interval on the plane (either closed or open), and it might as well be a ray (once again, open or closed). In case with $R_l \times R_l$ we conclude that the topology here is once again open or closed intervals on the plane.

2.5.9

Show that the dictionary order topology on the set $R \times R$ is the same as the product topology $R_d \times R$, where R_d denotes r in the discrete topology. Compare this topology with the standard topology on R^2 .

Let $\langle x, y \rangle \in R^2$. We follow that there exists q - element of basis of $R_d \times R$ such that $\langle x, y \rangle \in q$. Because q is an element of a basis, we follow that $q = w \times r$, where $w \in \mathcal{T}_{R_d}$ and $r \in \mathcal{T}_R$. Because $r \in \mathcal{T}_R$, we follow that there exists (a, b) such that $(a, b) \subseteq r$. Therefore we follow that element $\{x\} \times (a, b)$ is an element of a basis of dictionary order such that $\{x\} \times (a, b) \subseteq q$. Therefore we follow that $R_d \times R$ is coarser than dictionary order topology.

Suppose that $\langle x, y \rangle$ is in R^2 and q is in basis of dictionary topology of R^2 such that $\langle x, y \rangle \in q$. By definition (and immediate implications of thereof) we follow that the set $q \cap \{x\} \times R$ is nonempty. Since $\langle x, y \rangle$ is in q , and q is a basis element, we follow that there exist $a, b \in R$ such that $\{x\} \times (a, b) \subseteq q$ (follows from definitions and maybe some trivial manipulations of definition of dictionary order). Since $\{x\} \times (a, b)$ is an element of a basis of $R_d \times R$, we conclude that dictionary order topology is coarser than $R_d \times R$, and thus by double inclusion we conclude that topology over $R_d \times R$ and dictionary order topologies are equal, as desired.

We can follow that topology in R_d is strictly finer than standard topology of R since $R_d = \mathcal{P}(R)$, and thus it is the largest possible topology. Strictness follows from the fact

that $\{0\} \in \mathcal{T}_{R_d}$ and $\{0\} \notin \mathcal{T}_R$.

Thus we can be pretty sure that there is no element of basis of standard topology on $R \times R$ that contains $\{0\} \times R$. Since every element of basis of $R \times R$ is also contained in $R_d \times R$, we conclude that standard topology on R^2 is coarser than $R_d \times R$.

2.5.10

Let $I = [0, 1]$. Compare the product topology on $I \times I$, the dictionary order topology on $I \times I$, and the topology $I \times I$ inherits as a subspace of $R \times R$ in the dictionary order topology.

Wanted to skip this one, since I've solved it incorrectly the first time, but instead of skipping I'll just present the half-assed proof here for completeness' sake.

Important thing to remember: sets such as $\{x\} \times [0, 0.1)$ are not elements of basis of dictionary topology on $I \times I$

Let us look at the point $\langle 0.5, 1 \rangle$ and basis element of standard topology $[0, 1] \times (0.5, 1]$. We can follow that since the elements of dictionary bases cannot just stop at the corners and must wrap around, we follow that there is no element of basis of dictionary order topology, that is contained in presented element of basis and contains the desired point.

We can follow also that $\{0.5\} \times (0, 1)$ cannot be presented in standard topology as well. Thus the first two are not comparable

Suppose that $\langle x, y \rangle \in I \times I$ and q is the basis element with respect to the dictionary order topology. We follow that we can take a "strand" from dictionary (i.e. take a set $\{x\} \times R \cap q$, where q is an element of the basis, that contains point $\langle x, y \rangle$) and get that dictionary order over $I \times I$ is coarser than the topology $I \times I$ inherits as a subspace of $R \times R$ in the dictionary order topology, since the strand is the element of the basis of the latter. We can pull the same trick that we've used in the previous paragraph to show that the inherited topology is strictly coarser than the dictionary topology. Using the "strand" method (i.e. taking basis elements in form $\{x\} \times (a, b)$ or its closed analogs) we can prove that the last topology is strictly finer than the standard topology on $I \times I$.

2.6 Closed Sets and Limit Points

2.6.1

let C be a collection of subsets of the set X . Suppose that \emptyset and X are in C , and that finite unions and arbitrary intersections of elements of C are in C . Show that the collection

$$T = \{X \setminus C : C \in C\}$$

is a topology on X .

We're gonna use the definition of topology on this one. We follow that since X and \emptyset are in C that

$$X \setminus X = \emptyset \in T$$

$$X \setminus \emptyset = X \in T$$

Assume that J is an arbitrary subset of T . We follow that for every $j \in J$ there exists a unique $k \in C$ such that $j = X \setminus k$. Thus we follow that there exists $C' \subseteq C$ such that

$$\{j : j \in J\} = \{X \setminus k : k \in C'\}$$

thus

$$\bigcup J = \bigcup \{j : j \in J\} = \bigcup \{X \setminus k : k \in C'\} = X \setminus \bigcap \{k : k \in C'\} = X \setminus \bigcap C'$$

where we've used DML to justify one of the equations. Since C' is an arbitrary subset of C we follow that $\bigcap C' \in C$. Thus we follow that $X \setminus \bigcap C' \in T$. Thus we follow that $J \subseteq T \Rightarrow \bigcup J \in T$. Therefore we've got second property of topology.

If J is a finite subset of T , then we follow that we can define C' by the same definition and that C' is finite as well. Thus

$$\bigcap J = \bigcap \{j : j \in J\} = \bigcap \{X \setminus k : k \in C'\} = X \setminus \bigcup \{k : k \in C'\} = X \setminus \bigcup C'$$

thus we follow that if J is a finite subset of T , then $\bigcap J \in T$, therefore we've got the third and final condition of topology. Thus we follow that T is a topology, as desired.

2.6.2

Show that if A is closed in Y and Y is closed in X , then A is closed in X

Since Y is closed in X we follow that $Y \subseteq X$. Assuming that the topology on Y is a subset topology, we follow that if A is a closed set in Y , then there exists $A' \subseteq X$ such that A' is closed and $A = A' \cap Y$. Since both A' and Y are closed in X we follow that $A = A' \cap Y$ is closed in X as well by definition of topology, as desired.

2.6.3

Show that if A is closed in X and B is closed in Y , then $A \times B$ is closed in $X \times Y$.

If A is closed in X , then we follow that there exists A' such that $A = X \setminus A'$, where A' is an open set in X . Same goes for $B = Y \setminus B'$. Thus we follow that $A' \times B'$ is an open set in $X \times Y$. one of the exercises in chapter 1 gives us that

$$(A \times B) = (X \setminus A') \times (Y \setminus B') = (X \times Y \setminus A' \times Y) \setminus X \times B'$$

We follow that $A' \times Y$ is an open set, then $(X \times Y \setminus A' \times Y)$ is a closed set. We follow also that $X \times B'$ is an open set, thus $X \times Y \setminus X \times B'$ is a closed set. Thus

$$(X \times Y \setminus A' \times Y) \setminus X \times B' = (X \times Y \setminus A' \times Y) \cap (X \times Y \setminus X \times B')$$

is an intersection of closed sets and therefore is closed itself. Thus we conclude that $A \times B$ is a closed set, as desired. (we can also follow the same thing by the following exercise)

2.6.4

Show that if U is open in X and A is closed in X , then $U \setminus A$ is open in X and $A \setminus U$ is closed in X .

Firstly I want to prove that if $A, B \subseteq X$, then

$$A \setminus B = A \cap (X \setminus B)$$

We follow that by

$$x \in A \setminus B \Leftrightarrow x \in A \wedge x \notin B \Leftrightarrow x \in A \wedge x \in X \wedge x \notin B \Leftrightarrow x \in A \wedge (x \in X \setminus B) \Leftrightarrow x \in A \cap (X \setminus B)$$

We can follow by the fact that $U, A \subseteq X$ that

$$U \setminus A = U \cap (X \setminus A)$$

and

$$A \setminus U = A \cap (X \setminus U)$$

In the former case we've got finite intersection of two open sets, and in the latter we've got finite intersection of two closed sets, thus proving that $U \setminus A$ is open and $A \setminus U$ is closed, as desired.

2.6.5

Let X be an ordered set in the order topology. Show that $\overline{(a, b)} \subseteq [a, b]$. Under what conditions does equality hold?

Let us firstly state that $a, b \in X$. We follow that $(a, b) \subseteq [a, b]$, thus $[a, b]$ is a closed set that contains (a, b) , therefore by definition of closure we follow that $\overline{(a, b)} \subseteq [a, b]$.

We follow that $\overline{(a, b)} = [a, b]$ if and only if a, b are limit points of (a, b) .

2.6.6

Let A, B and A_α denote subsets of a space X . Prove the following

(a) If $A \subseteq B$, then $\overline{A} \subseteq \overline{B}$

Assume that $A \subseteq B$. Let $x \in \overline{A}$. We follow that every neighborhood of x intersects A . Thus every neighborhood of x intersects B by the fact that $A \subseteq B$. Therefore $x \in \overline{B}$. Therefore $\overline{A} \subseteq \overline{B}$.

(b) $\overline{A \cup B} = \overline{A} \cup \overline{B}$

Let $x \in \overline{A \cup B}$. We follow that every neighborhood of x intersects $A \cup B$. Thus every neighborhood of x intersects A or B . Assume that $x \notin \overline{A}$ and $x \notin \overline{B}$. Then we follow that there exists a neighborhood U of x such that $U \cap A = \emptyset$. There also exists neighborhood U' of x such that $U' \cap B = \emptyset$. Thus we follow that $U \cap U'$ is a neighborhood of x such that it

does not intersect A nor B . Thus $x \notin \overline{A \cup B}$, which is a contradiction. Thus we conclude that $x \in \overline{A \cup B} \Rightarrow x \in \overline{A} \cup \overline{B}$.

If $x \in \overline{A} \cup \overline{B}$, then $x \in \overline{A}$ or $x \in \overline{B}$. Assume that the former is true. Then we follow that $x \in \overline{A}$. Thus we follow that every neighborhood of x intersects A . Since $A \subseteq A \cup B$, we follow that every neighborhood of x intersects $A \cup B$. Thus $x \in \overline{A \cup B}$, as desired.

(c) $\overline{\bigcup A_\alpha} \supseteq \bigcup \overline{A_\alpha}$.

I think that we need to assume here that $A_\alpha \subseteq \mathcal{P}(X)$ and what we actually need to prove is that

$$\overline{\bigcup A_\alpha} \supseteq \bigcup \{\overline{a} : a \in A_\alpha\}$$

if that's the case, then we follow that if $x \in \bigcup \{\overline{a} : a \in A_\alpha\}$, then there exists $x \in a \in A_\alpha$ such that $x \in \overline{a}$. This means that every neighborhood of x intersects a at some point. Since $a \subseteq \bigcup A_\alpha$, we follow that every neighborhood of x intersects $\bigcup A_\alpha$ at some point and thus $x \in \overline{\bigcup A_\alpha}$.

We can follow that if $A_\alpha = \{\{1/n\} : n \in \mathbb{Z}_+\}$ and we've got standard topology on reals, then $0 \in \overline{\bigcup A_\alpha}$, but $0 \notin \bigcup \{\overline{a} : a \in A_\alpha\}$, since there is no $a \in A_\alpha$ such that $0 \in \overline{a}$.

2.6.7

Critsize the following "proof" that $\overline{\bigcup A_\alpha} \subseteq \bigcup \overline{A_\alpha}$: if $\{A_\alpha\}$ is a collection of sets in X and if $x \in \bigcup \overline{A_\alpha}$, then every neighborhood of x intersects $\bigcup \overline{A_\alpha}$. Thus x must intersect some A_α , so that x must belong to the closure of some A_α . Therefore $x \in \bigcup \overline{A_\alpha}$.

We don't have implication "Thus x must intersect some A_α , so that x must belong to the closure of some A_α ", as it was just made up. Although every neighborhood of x indeed intersects some A_α , there's no implication that there exists A_α such that every neighborhood of x intersects A_α .

2.6.8

Let A, B and A_α denote subsets of a space X . Determine whether the following equations hold; if an equality fails, determine whether one of the inclusions holds.

(a) $\overline{A \cap B} = \overline{A} \cap \overline{B}$.

If $x \in \overline{A \cap B}$, then we follow that every neighborhood of x intersects $A \cap B$ at some point. Thus every neighborhood of x intersects A and B . Thus $x \in \overline{A}$ and $x \in \overline{B}$. Thus we've got forward inclusion.

If $x \in \overline{A} \cap \overline{B}$, then we follow that every neighborhood of x intersects A and every neighborhood of x intersects B . Thus every neighborhood of x intersects both A and B . This does not mean that every neighborhood of x intersects $A \cap B$ since points of intersection can be different. We can come up with some counterexample for this claim: for example we can set $A = \{1/2n : n \in \mathbb{Z}_+\}$ and $B = \{1/(2n+1) : n \in \mathbb{Z}_+\}$. We follow that $A \cap B = \emptyset$ and thus $\overline{A \cap B} = \emptyset$, but $\overline{A} \cap \overline{B} = \{0\}$.

Therefore we follow only the forward inclusion.

$$(b) \overline{\bigcap A_\alpha} = \bigcap \overline{A_\alpha}$$

We're once again struck with this awful notation, so let's change that

$$\overline{\bigcap A_\alpha} = \bigcap_{a \in A_\alpha} \bar{a}$$

we follow that reverse inclusion is not true, since that would imply the correctness of counterexample in previous point.

We follow the forward inclusion by pretty much the same logic as in previous one

If $x \in \overline{\bigcap A_\alpha}$, then we follow that every neighborhood of x intersects $\bigcap A_\alpha$ at some point. Thus every neighborhood of x intersects every $a \in A_\alpha$. Thus $(\forall a \in A_\alpha)(x \in \bar{a})$. Therefore we follow that $a \in \bigcap_{a \in A_\alpha} \bar{a}$. Thus we've got forward inclusion.

$$(c) \overline{A \setminus B} = \bar{A} \setminus \bar{B}$$

We've got a case against $\overline{A \setminus B} \subseteq \bar{A} \setminus \bar{B}$ by setting once again $A = \{1/2n : n \in \mathbb{Z}_+\}$, $B = \{1/(2n+1) : n \in \mathbb{Z}_+\}$. We follow that $A \setminus B = A$ since they are disjoint. Therefore we follow that $0 \in \overline{A \setminus B}$ but $0 \notin \bar{A} \setminus \bar{B}$, since $0 \in \bar{B}$.

If $x \in \bar{A} \setminus \bar{B}$, then we follow that every neighborhood of x intersects A , but x is not in \bar{B} . Assume that some neighborhood of x intersects A only at a points $A \cap B$. Then we follow that $x \in \overline{A \cap B}$ and thus $x \in \bar{A} \cap \bar{B}$. Therefore $x \in \bar{B}$, which is a contradiction. Therefore we follow that if $x \in \bar{A} \setminus \bar{B}$ then every neighborhood of x intersects $A \setminus B$. Therefore $x \in \overline{A \setminus B}$, which gives us reverse inclusion.

2.6.9

Let $A \subseteq X$ and $B \subseteq Y$. Show that in the space $X \times Y$

$$\overline{A \times B} = \bar{A} \times \bar{B}$$

Let $x \in X$ and $y \in Y$ be such that $\langle x, y \rangle \in \overline{A \times B}$. Let U be an arbitrary neighborhood of x and V be an arbitrary neighborhood for y . We follow that $U \times V$ is an element of basis of $X \times Y$ that contains $\langle x, y \rangle$, and thus it intersects $A \times B$. Therefore there exists $\langle q, w \rangle \in A \times B \cap U \times V$. Thus we follow that $\langle q, w \rangle \in (A \cap U) \times (B \cap V)$. Therefore we follow that U intersects A and V intersects B . Since U and V are arbitrary, we follow that every neighborhood of x intersects A and every neighborhood of y intersects B . Thus $x \in \bar{A}$ and $y \in \bar{B}$ and thus $\langle x, y \rangle \in \bar{A} \times \bar{B}$.

Suppose that $x \in X, y \in Y$ are such that $\langle x, y \rangle \in \bar{A} \times \bar{B}$. We follow that $x \in \bar{A}$ and $y \in \bar{B}$. Assume that $U \times V$ is a basis element of $X \times Y$ that contains $\langle x, y \rangle$. We follow that U intersects A and V intersects B . Thus there exist $u \in U \cap A$ and $v \in V \cap B$. Therefore $\langle u, v \rangle \in A \times B \cap U \times V$. Therefore $U \times V$ intersects $A \times B$. Thus we follow that every basis element that contains $\langle x, y \rangle$ intersects $A \times B$, and thus $\langle x, y \rangle \in \overline{A \times B}$.

Double inclusion produces the desired equality, as desired.

2.6.10

Show that every order topology is Hausdorff.

Let X be a toset, and let \mathcal{T} be respective order topology. I think that definition, that is presented in the book implies that if there is only one element in X , then the topology is vacuously Hausdorff (same goes for empty set). Thus assume that X contains at least two elements, and let $x_1, x_2 \in X$ be such that $x_1 \neq x_2$. Since $x_1, x_2 \in X$ and $x_1 \neq x_2$, we follow that $x_1 \prec x_2$ or $x_2 \prec x_1$. Assume the former.

Essentially we want to produce two open sets, that will prove that the space is Hausdorff, and those two sets will be just plain old intervals. But we've got two cases: there could exist x_3 such that $x_1 \prec x_3 \prec x_2$, or there might not. If such an element exists, then set $b_1 = a_2 = x_3$. If there is no such element, then set $b_1 = x_2$ and $a_2 = x_1$. Let a_1 be either the lowest element of X if such exists, or some element that is less than x_1 in case that it does not exist. Let b_2 be either the largest element of X if such exists, or some element that is larger than x_2 in case that it does not exist.

Then we can follow that $x_1 \in (a_1, b_1)$ and $x_2 \in (a_2, b_2)$ and $(a_1, b_1) \cap (a_2, b_2) = \emptyset$ by their respective definitions. Thus we follow that the space is Hausdorff, as desired.

2.6.11

Show that the product of two Hausdorff spaces is Hausdorff.

Let X be the first space and let Y be the second. Let $\langle x_1, y_1 \rangle, \langle x_2, y_2 \rangle \in X \times Y$ be distinct. We follow that $x_1 \neq x_2$ or $y_1 \neq y_2$. If we've got $A_1, A_2 \subseteq X$ and $B_1, B_2 \subseteq Y$ such that $A_1 \cap A_2 = \emptyset$ or $B_1 \cap B_2 = \emptyset$, then we follow that

$$A_1 \times B_1 \cap A_2 \times B_2 = (A_1 \cap A_2) \times (B_1 \cap B_2) = \emptyset$$

the desired result follows easily from that.

2.6.12

Show that a subspace of a Hausdorff space is Hausdorff.

Assume that X is a Hausdorff space and let $Y \subseteq X$. Let $y_1, y_2 \in Y$. We follow that there exist open sets $U, V \subseteq X$ such that $y_1 \in U, y_2 \in V, U \cap V = \emptyset$ by the fact that X is Hausdorff. We follow that $U \cap Y$ and $V \cap Y$ are open sets in Y , and since $U \cap V = \emptyset$, we follow that $(U \cap Y) \cap (V \cap Y) = \emptyset$ by commutativity and associativity of \cap . Therefore we conclude that subspace generated by Y is a Hausdorff. Since Y is arbitrary, we follow that any subspace of X is Hausdorff, as desired.

2.6.13

Show that X is Hausdorff iff diagonal $\Delta = \{x \times x : x \in X\}$ is closed in $X \times X$.

Since X is Hausdorff, we follow that for all $x \in X$ we've got that $\{\langle x, x \rangle\}$ is closed.

Assume that $\Delta \neq \overline{\Delta}$ and thus there exists $\langle y_1, y_2 \rangle \in \overline{\Delta} \setminus \Delta$. We follow that since $\langle y_1, y_2 \rangle \notin \Delta$ that $y_1 \neq y_2$. We also follow that $y_1, y_2 \in X$. Since X is Hausdorff, we follow that $y_1, y_2 \in X$ implies that there exist neighborhoods $U, V \subseteq X$ of y_1 and y_2 respectively, such that $U \cap V = \emptyset$. Thus we follow that $U \times V$ is an open set in $X \times X$. Then we follow that since $\langle y_1, y_2 \rangle \in \overline{\Delta}$ that there exists $d \in \Delta$ such that $d \in U \cap V$. Thus we follow that there exists $d_1 \in X$ such that $\langle d_1, d_1 \rangle = d$. Therefore $\langle d_1, d_1 \rangle \in U \times V$ and therefore $d_1 \in U \wedge d_1 \in V$, which implies that $d_1 \in U \cap V$, which is a contradiction, since $U \cap V = \emptyset$ by the fact that X is Hausdorff, as desired.

Therefore we conclude that $\overline{\Delta} \setminus \Delta = \emptyset$, therefore $\Delta = \overline{\Delta}$ and Δ is a closed set, thus giving us forward implication.

Now assume that Δ is a closed set and X is not Hausdorff. Since X is not Hausdorff, we follow that there exists $x_1, x_2 \in X$ such that $x_1 \neq x_2$ and for all $U \in \mathcal{T}$ we've got that $x_1 \in U \Rightarrow x_2 \in U$.

Let B be an arbitrary basis element that contains $\langle x_1, x_2 \rangle$. We follow that since it is a basis element, that $B = U \times V$, where U, V are open subsets of X . We follow that $x_1 \in U$ and $x_2 \in V$, that implies that $x_1, x_2 \in U$ and $x_1, x_2 \in V$. Thus we follow that $\langle x_1, x_2 \rangle \in B$. Therefore we follow that if B is a basis element, that contains $\langle x_1, x_2 \rangle$, then B intersects Δ by the fact that $\langle x_1, x_1 \rangle$ and $\langle x_2, x_2 \rangle$ are both in B . Thus we follow that $\langle x_1, x_2 \rangle \in \overline{\Delta}$. Since $x_1 \neq x_2$ we follow that $\langle x_1, x_2 \rangle \notin \Delta$, which implies that $\Delta \neq \overline{\Delta}$, which implies that Δ is not closed, which is a contradiction. Thus we follow that if Δ is a closed set, then X is Hausdorff, which gives us reverse implication, as desired.

2.6.14

In the finite complement topology on R , to what point or points does the sequence $x_n = 1/n$ converge?

We can follow pretty easily that finite topology on R is not Hausdorff by the fact that if U, V are nonempty open sets in finite complement topology, then $U = R \setminus S_1, V = R \setminus S_2$ for some finite subsets S_1, S_2 of R . Thus we follow that $U \cap V = R \setminus S_1 \cap R \setminus S_2 = R \setminus (S_1 \cup S_2)$, where we follow that $S_1 \cup S_2$ is finite and therefore $R \setminus (S_1 \cup S_2)$ is infinite and therefore nonempty.

Assume that U is a neighborhood of 0. We follow that $U = R \setminus S_1$ such that $0 \notin S_1$ and S_1 is finite. Since x_n is an infinite sequence, we follow that there must exist n such that $n_0 > n \Rightarrow x_n \notin S_1 \Rightarrow x_n \in U$. Thus we conclude that x_n converges to 0.

By the same logic we follow that if U is a neighborhood of any other number, then the same logic applies. Therefore we follow that x_n converges to any number in finite complement topology.

2.6.15

Show that T_1 axiom is equivalent to the condition that for each pair of points of X , each has a neighborhood not containing the other.

Let $x_1, x_2 \in X$ are such that $x_1 \neq x_2$. T_1 axiom implies that $\{x_1\}$ and $\{x_2\}$ are closed and thus $X \setminus \{x_1\}$ and $X \setminus \{x_2\}$ are open. Since $x_1 \neq x_2$, we follow that $x_2 \in X \setminus \{x_1\}$ and $x_1 \in X \setminus \{x_2\}$, thus implying that each point has a neighborhood, that does not contain the other point.

2.6.16

Consider the five topologies on R given in Exercise 7 of paragraph 13.

(a) Determine the closure of the set $K = \{1/n : n \in \mathbb{Z}_+\}$ under each of these topologies.

Under standard topology we follow that $\overline{K} = K \cup \{0\}$ from real analysis course.

Under K topology we follow that R is open by default, and thus $R \setminus K$ is a basis element, therefore it is open and thus $K = R \setminus (R \setminus K)$ is closed.

Under finite complement topology we follow that all closed sets that are not R are finite, and thus only R contains K . Therefore $\overline{K} = R$.

Since the upper limit topology is finer than the K -topology, we follow that $R \setminus K$ is an open set in upper limit topology, and thus $\overline{K} = K$.

For the topology that has sets $(-\infty, a)$ as a basis, we follow that closed sets there are either R or $[a, \infty)$ for some $a \in R$, and thus $\overline{K} = [0, \infty)$.

(b) Which of these topologies satisfy the Hausdorff axiom? The T_1 axiom?

We follow that standard topology satisfies the Hausdorff axiom, as proven in the chapter or in the real analysis course. Thus we follow that since standard topology is coarser than K -topology and upper limit topology, that both of them satisfy the axiom as well. In exercise 14 of this section we've shown that finite complement topology is definitely not Hausdorff, and for the last topology we follow that if $0 \in U$, then $-1 \in U$, therefore it is not Hausdorff as well.

In the chapter we've discussed that T_1 axiom is weaker than Hausdorff, thus we follow that standard topology, K -topology and upper limit topology all satisfy the T_1 axiom. We can follow that finite complement topology satisfies T_1 by some trivial implications and the last topology is definitely does not satisfy T_1 .

2.6.17

Consider the lower limit topology on R and the topology given by the basis C of exercise 8 of paragraph 13. Determine the closures of the intervals $A = (0, \sqrt{2})$ and $B = (\sqrt{2}, 3)$ in these two topologies.

Let's talk about lower limit topology first.

Suppose that $0 \in [a, b)$. We follow that $a \leq 0 < b$, and by density of reals we follow that there exists $d \in R$ such that $0 < d < b$ and thus $d \in [a, b)$. Thus we follow that every

basis element that contains 0 also intersects A , and thus $0 \in \bar{A}$. We follow that $[\sqrt{2}, 2)$ is a neighborhood of $\sqrt{2}$ such that it does not intersect A at any point. Thus we follow that $\sqrt{2} \notin \bar{A}$. There're plenty of methods to state that if $x < 0$ or $x > \sqrt{2}$, then it is not in closure of A , therefore we conclude that $[0, \sqrt{2})$ is clousre of A under lower limit topology.

For B we follow that pretty much the same logic holds with respect to different numbers, and thus $\bar{B} = [\sqrt{2}, 3)$.

Now let's talk about the weird topology.

We can follow that pretty much the same argument holds for $0 \in \bar{A}$. Now assume that $a < \sqrt{2} < b$. We follow that since $a \notin I$, that $a \neq \sqrt{2}$, and thus we follow that $a < \sqrt{2} < b$. Thus we follow that $\sqrt{2} \in \bar{A}$. If $\sqrt{2} < c$, then we follow that there exist rational a, b such that $\sqrt{2} < a < c < b$, and thus $c \notin \bar{A}$. If $c < 0$, then we have pretty much the same result. Therefore we follow that $\bar{A} = [0, \sqrt{2}]$.

For the case of $B = (\sqrt{2}, 3)$, we follow that $[3, 5) \cap B = \emptyset$, thus we follow that $3 \notin \bar{B}$. We can easily follow that $\sqrt{2} \in \bar{B}$, and the rest is followed by pretty much the same logic as in the previous paragraph. Therefore we conclude that $\bar{B} = [\sqrt{2}, 3)$.

2.6.18

Determine the closures of the following subsets of the ordered square:

We're talking about the lexicographical order on the set $I \times I$ for $I = [0, 1]$. We're also not gonna use the dumb notation.

$$A = \{\langle 1/n, 0 \rangle : n \in \mathbb{Z}_+\}$$

We follow that $A \subseteq \bar{A}$. Let $\langle x_1, x_2 \rangle \in I \times I$.

Assume that $x_1 \neq 0$. If $x_1 = 1/n$ for some $n \in \mathbb{Z}_+$ and $x_2 = 0$, then we follow that $\langle x_1, x_2 \rangle \in A$, thus assume that $x_2 \neq 0$. We follow that if $x_2 > 0$ and $x_2 < 1$, then there exist $a, b \in \mathbb{R}$ such that $0 < a < x_2 < b < 1$ and thus $\langle x_1, x_2 \rangle \in (\langle x_1, a \rangle, \langle x_1, b \rangle)$, where we follow that $(\langle x_1, a \rangle, \langle x_1, b \rangle)$ does not intersect A at any point.

If $x_1 = 1/n$ and $x_2 = 1$ then we follow that there exists some space between x_1 and the previous point in the sequence, therefore we can have interval $[x_1, y)$ such that $A \cap [x_1, y) = \{x_1\}$. Thus if we take open interval $(\langle x_1, 0.5 \rangle, \langle y_1, 0.5 \rangle)$, then $(\langle x_1, 0.5 \rangle, \langle y_1, 0.5 \rangle) \cap A = \emptyset$. Therefore we follow that $\{1/n\} \times I \cap A \subseteq A$. We can follow pretty much the same thing for $x_1 \neq 0$ in a more general case.

Now assume that $x_1 = 0$. For the point $\langle 0, 0 \rangle$ we've got that $(\langle 0, 0 \rangle, \langle 0, 0.5 \rangle) \cap A = \emptyset$, therefore we conclude that $\langle 0, 0 \rangle \notin \bar{A}$. Simple logic implies that for $x_2 \neq 1$ we've got pretty much the same result.

The case with $\langle 0, 1 \rangle$ is an interesting one. Suppose that we've got an element of the basis $(\langle j_1, j_2 \rangle, \langle k_1, k_2 \rangle)$ such that

$$\langle j_1, j_2 \rangle < \langle 0, 1 \rangle < \langle k_1, k_2 \rangle$$

We can follow that $j_1 = 0$ by the definition of order. We also follow that $0 < k_1$. Thus

$$\langle 0, j_2 \rangle < \langle 0, 1 \rangle < \langle k_1, k_2 \rangle$$

we can follow that for any $k_1 > 0$ there exists $n \in \mathbb{Z}_+$ such that $0 < 1/n < k_1$, and therefore

$$\langle 0, j_2 \rangle < \langle 1/n, 0 \rangle < \langle k_1, k_2 \rangle$$

Therefore we follow that if B is an element of the basis such that $\langle 0, 1 \rangle \in B$, then it intersects A , and thus $\langle 0, 1 \rangle$ is the only point outside of A that is in the closure of A . Thus we follow that

$$\overline{A} = A \cup \{\langle 0, 1 \rangle\}$$

$$B = \{1 - 1/n \times 1/2 : n \in \mathbb{Z}_+\}$$

$$\overline{B} = B \cup \{\langle 1, 1 \rangle\}$$

$$C = \{x \times 0 : 0 < x < 1\}$$

$$\overline{C} = C \cup \{\langle x, 1 \rangle : 0 \leq x \leq 1\}$$

$$D = \{x \times 1/2 : 0 < x < 1\}$$

$$\overline{D} = D \cup \{\langle x, 1 \rangle : 0 \leq x \leq 1\} \cup \{\langle x, 0 \rangle : 0 < x < 1\}$$

$$E = \{1/2 \times x : 0 < x < 1\}$$

$$\overline{E} = E \cup \{\langle 1/2, 0 \rangle, \langle 1/2, 1 \rangle\}$$

last answers are probably wrong, but I just want to move on.

2.6.19

If $A \subseteq X$, we define the boundary of A by the equation

$$BdA = \overline{A} \cap \overline{(X - A)}$$

(a) Show that $IntA$ and BdA are disjoint and $\overline{A} = IntA \cup BdA$.

Suppose that $x \in IntA$. We follow that there exists open set U such that $x \in U$ and $U \subseteq A$. We follow that U and $X \setminus A$ are disjoint. Suppose that $x \in \overline{X \setminus A}$. Then we follow that every neighborhood of x intersects $X \setminus A$. Since U is a neighborhood of x we follow that $U \cap X \setminus A \neq \emptyset$. Therefore there exists $j \in U$ such that $j \in X \setminus A$. Therefore $j \in U$ and $j \notin A$. This contradicts the fact that $U \subseteq A$. Thus we follow that if $x \in IntA$ then $x \notin \overline{X \setminus A}$. Thus we follow that $IntA \cap \overline{X \setminus A} = \emptyset$. Now we can follow that

$$BdA \cap IntA = \overline{A} \cap \overline{X \setminus A} \cap IntA = \overline{A} \cap \emptyset = \emptyset$$

We can also follow that $BdA \subseteq \overline{A}$ by definition and $IntA \subseteq \overline{A}$ by the fact that $IntA \subseteq A \subseteq \overline{A}$. Therefore we follow that $IntA \cup BdA \subseteq \overline{A}$.

Now assume that $x \in \bar{A}$ and $x \notin \text{Int}A$. Since $x \notin \text{Int}A$ we follow that there is no neighborhood U of x such that $U \subseteq A$. We follow that every neighborhood U of x has an element $y \in U$ such that $y \notin A$. And since $U \subseteq X$, we follow that $U \cap (X \setminus A) \neq \emptyset$ for every neighborhood of x . Thus we follow that $x \in \overline{(X \setminus A)}$. And since $x \in \bar{A}$, we follow that $x \in \text{Bd}A$ and thus

$$(\forall x \in \bar{A})(x \notin \text{Int}A \Rightarrow x \in \text{Bd}A)$$

Thus we follow that $\bar{A} \setminus \text{Bd}A = \text{Int}A$ and since both of $\text{Bd}A$ and $\text{Int}A$ are subsets of A we conclude that $\bar{A} = \text{Bd}A \cup \text{Int}A$, as desired.

(b) Show that $\text{Bd}A = \emptyset \Leftrightarrow A$ is both open and closed.

If $\text{Bd}A = \emptyset$ we follow that $A = \text{Int}A$ and thus it is open. We also follow that $\bar{A} = \text{Bd}A \cup \text{Int}A = \text{Int}A = A$ since $\text{Int}A \subseteq A \subseteq \bar{A}$, and thus A is closed as well, as desired.

If A is both opened and closed we follow that $\text{Int}A = A$ and $\bar{A} = A$, thus $\text{Bd}A = \emptyset$.

(c) Show that U is open $\Leftrightarrow \text{Bd}U = \bar{U} \setminus U$.

If U is open, then we follow that $X \setminus U$ is closed and thus $\overline{X \setminus U} = X \setminus U$. This implies that $\text{Bd}U = \bar{U} \cap (X \setminus U)$ and since $\bar{U} \subseteq X$ and $U \subseteq \bar{U}$ we follow by some identity with a \setminus that $\text{Bd}U = \bar{U} \setminus U$, as desired.

Suppose that $\text{Bd}U = \bar{U} \setminus U$. Since $\bar{U} = \text{Bd}U \cup \text{Int}U$ and $\text{Int}U \cap \text{Bd}U = \emptyset$, we follow that $\text{Int}U = \bar{U} \setminus \text{Bd}U$. Thus $\text{Int}U = \bar{U} \setminus (\bar{U} \setminus U)$ and since $U \subseteq \bar{U}$ we follow that $\text{Int}U = U$, thus proving that U is open, as desired.

(d) If U is open, is it true that $U = \text{Int}\bar{U}$? Justify your answer.

Suppose that $x \in U$. We follow that there exists open $V \subseteq U$ such that $x \in V \subseteq U$. Therefore $V \subseteq \bar{U}$. Thus we follow that $V \subseteq \text{Int}\bar{U}$ and thus $x \in \text{Int}\bar{U}$. Therefore we follow that $U \subseteq \text{Int}\bar{U}$.

If $x \in \text{Int}(\bar{U})$. We follow that there exists V such that $x \in V \subseteq \bar{U}$. If $x \notin U$, then we follow nothing.

Let $U = R \setminus \{0\}$ and assume stanadrd topology. We follow that $\bar{U} = R = \text{Int}\bar{U} \neq U$, which gives us a solid contradiction of the reverse inclusion.

2.6.20

Skip

2.6.21

(Kuratowski) Consider the collection of all subsets A of the topological space X . The operations of closure $A \rightarrow \bar{A}$ and complementation $A \rightarrow X \setminus A$ are functions from this collection to itself.

(a) Show that starting with a given set A , ine can form no more than 14 distinct sets by applying therse two operations successively.

We follow that if $X \neq \emptyset$, then $A \neq X \setminus A$. We also know that $A = X \setminus (X \setminus A)$

Let $A = [0, 2] \setminus \{1\}$ so that it is neither open nor closed.

$$\overline{A} = [0, 2]$$

$$X \setminus \overline{A} = (-\infty, 0) \cup (2, \infty)$$

$$\overline{X \setminus \overline{A}} = (-\infty, 0] \cup [2, \infty)$$

$$X \setminus \overline{X \setminus \overline{A}} = (0, 2)$$

$$\overline{X \setminus \overline{X \setminus \overline{A}}} = [0, 2]$$

$$A = [0, 1) \cup \{2\} \cup (3, 4]$$

$$\overline{A} = [0, 1] \cup \{2\} \cup [3, 4]$$

$$X \setminus \overline{A} = (-\infty, 0) \cup (1, 2) \cup (2, 3) \cup (4, \infty)$$

$$\overline{X \setminus \overline{A}} = (-\infty, 0] \cup [1, 3] \cup [4, \infty)$$

$$X \setminus \overline{X \setminus \overline{A}} = (0, 1) \cup (3, 4)$$

$$\overline{X \setminus \overline{X \setminus \overline{A}}} = [0, 1] \cup [3, 4]$$

$$X \setminus \overline{X \setminus \overline{X \setminus \overline{A}}} = (-\infty, 0) \cup (1, 3) \cup (4, \infty)$$

$$\overline{X \setminus \overline{X \setminus \overline{X \setminus \overline{A}}}} = (-\infty, 0] \cup [1, 3] \cup [4, \infty)$$

SKIP for now, probably will come back later

2.7 Continuous Functions

2.7.1

Prove that for functions $f : R \rightarrow R$, the $\epsilon - \delta$ definition of continuity implies the open set definition.

Let f be continuous with respect to the $\epsilon - \delta$ definition of continuity. Let (a, b) be an element of basis of R . If $(a, b) \cap \text{ran}(f) = \emptyset$, then we follow that $f^{-1}[(a, b)] = \emptyset$, which is an open set.

Suppose that $(a, b) \cap \text{ran}(f) \neq \emptyset$. Let $y \in (a, b) \cap \text{ran}(f)$. Since $y \in (a, b)$, we follow that there exists ϵ such that $V_\epsilon(y) \subseteq (a, b)$.

Let $x \in f^{-1}[\{y\}]$. For that $V_\epsilon(y)$ there exists $\delta > 0 \in R$ with corresponding $V_\delta(x)$ such that $z \in V_\delta(x) \Rightarrow f(z) \in V_\epsilon(y)$. By AC (not sure that we actually need AC at this point, but we're doing topology, so why not) we follow that for each y we can pick exclusive δ such that everything holds. Define $K : (a, b) \rightarrow \mathcal{P}(R)$ by $K(y) = \bigcup_{x \in f^{-1}[\{y\}]} V_\delta(x)$ in case

that $y \in \text{ran}(f)$ and empty set otherwise. Since $V_\delta(x)$ are all open intervals and empty sets are also open, we follow that for all $y \in (a, b)$ $K(y)$ is an open set. Moreover, we follow that $\bigcup \text{ran}(K)$ is an open set as well. By definition of K and $\epsilon - \delta$ continuity of f we follow that $x \in \bigcup \text{ran}(K) \Rightarrow f(x) \in (a, b)$

Now let $x \in R$ be such that $x \in f^{-1}[(a, b)]$. We follow that $x \in \bigcup \text{ran}(K)$ by definition. Therefore we conclude that $f^{-1}[(a, b)] = \bigcup \text{ran}(K)$. As proven earlier, $\bigcup \text{ran}(K)$ is an open set, and therefore we conclude that if f is $\epsilon - \delta$ -continuous, then for arbitrary interval (a, b) we've got that $f^{-1}[(a, b)]$ is open, and thus f is continuous according to our definition. Taking into account stuff that we were given in the chapter, we follow that f is $\epsilon - \delta$ -continuous if and only if f is continuous, as desired.

2.7.2

Suppose that $f : X \rightarrow Y$ is continuous. If x is a limit point of the subset A of X , is it necessary true that $f(x)$ is a limit point of $f(A)$?

Short answer: no.

Let $U \subseteq Y$ be a neighborhood of $f(x)$. We follow that $f^{-1}[U]$ is open and $x \in f^{-1}[U]$. Thus we follow that there exists a point $u \in f^{-1}[U]$ such that $u \neq x$ and $u \in A$. Thus $f(u) \in f[A]$. This means that we can't follow crap on the account that f might not be injective.

Let $f : R \rightarrow R$ be defined by $f(x) = 5$ and let us assume standard topology. For $A = (0, 1)$ we follow that $x = 0$ is a limit point of A , but $\text{ran}(f) = \{5\}$ and it doesn't have no limit points. Thus we've got a contradiction of our conjecture.

2.7.3

Let X and X' denote a single set in the two topologies \mathcal{T} and \mathcal{T}' respectively. Let $X' \rightarrow X$ be the identity function.

Just to be clear: we've assumed that we've got two topological spaces $\langle X, \mathcal{T} \rangle$ and $\langle X', \mathcal{T}' \rangle$ such that $X = X'$, but \mathcal{T} might be different to \mathcal{T}' .

(a) *Show that i is continuous $\Leftrightarrow \mathcal{T} \subseteq \mathcal{T}'$*

Suppose that i is continuous. Let $U \in \mathcal{T}$. Thus U is an open set in X . We follow by continuity of i that $i^{-1}[U] = U$ is an open set in \mathcal{T}' . Thus we follow that $U \in \mathcal{T}'$. Thus we follow that $\mathcal{T} \subseteq \mathcal{T}'$, as desired.

Same logic in reverse gets us the reverse implication.

(b) *Show that i is a homeomorphism $\Leftrightarrow \mathcal{T}' = \mathcal{T}$.*

If i is a homeomorphism, then we follow that $i^{-1} : X \rightarrow X'$ is continuous. Previous point implies that $\mathcal{T}' = \mathcal{T}$.

If $\mathcal{T}' = \mathcal{T}$, then previous point implies that both i and i^{-1} are continuous and thus i is an homeomorphism.

2.7.4

Given $x_0 \in X$ and $y_0 \in Y$, show that the maps $f : X \rightarrow X \times Y$ and $g : Y \rightarrow X \times Y$ defined by

$$f(x) = x \times y_0$$

$$g(y) = x_0 \times y$$

are imbeddings

We follow that $\text{ran}(f) = X \times \{y_0\}$ and $\text{ran}(g) = \{x_0\} \times Y$. Let $U_x \subseteq X$ and $U_y \subseteq Y$ be open sets. We follow that

$$f[U_x] = U_x \times \{y_0\} = U_x \times Y \cap X \times \{y_0\} = U_x \times Y \cap \text{ran}(f) =$$

$$g[U_y] = \{x_0\} \times U_y = X \times U_y \cap \{x_0\} \times Y = U_x \times Y \cap \text{ran}(g)$$

since $X \times U_y$ and $U_x \times Y$ are open sets in $X \times Y$ by the fact that U_x and U_y are open, we follow that $f[U_x]$ and $g[U_y]$ are open as well.

Let $W \subseteq X \times Y \cap \text{ran}(f)$ be an element of basis. We follow that $W = U \times V \cap \text{ran}(f)$ for some U, V - open sets in X and Y respectively. Since $\text{ran}(f) = X \times \{y_0\}$, we follow that $W = U \times \{y_0\}$. We follow that $W = f[U]$. Since U is an open set and taking into account previous paragraph, we conclude that U is open if and only if $f[U]$ is open and thus f is an embedding. Similar argument for reverse implication also holds for g , thus we follow that g is an imbedding as well, as desired.

2.7.5

Show that the subspace (a, b) of R is homeomorphic with $(0, 1)$ and the subspace $[a, b]$ of R is homeomorphic with $[0, 1]$.

In order to be clear I'll state here that we assume standard topology, and also assume that $a, b \in R$ are constants such that $a < b$.

Let $f_p : R \rightarrow R$ be defined by

$$f_p(x) = \frac{x - a}{b - a}$$

By the fact that this function is a constant shift of linear and nonconstant function, we can follow that this function is bijective (we can also follow that through standard methods in order to be more precise, but I'll skip that proof because it's pretty obvious). We can also follow that this function is continuous by the fact that identity function is continuous and by algebraic properties of continuity, that were proven in the course of real analysis (previous exercises and text in the chapter implies that $\epsilon - \delta$ -continuity is equivalent to topological continuity with respect to standard topology, so that we're clear with using that stuff here). Since f_p is bijective we follow that it's got an inverse, and basic algebra implies that

$$f_p^{-1}(x) = (b - a)x + a$$

which is continuous as well by algebraic properties of continuity.

We can follow that $f_p[(a, b)] = (0, 1)$ and $f_p[[a, b]] = [0, 1]$ by some basic algebra. Thus we follow that we can define bijections $f : (a, b) \rightarrow (0, 1)$ and $g : [a, b] \rightarrow [0, 1]$ by restriction of domain, i.e.

$$f = f_p|_{(a,b)}$$

$$g = f_p|_{[a,b]}$$

previous discussions imply that both f, g are continuous and bijective. We also follow that

$$f^{-1} = f_p^{-1}|_{(0,1)}$$

$$g^{-1} = f_p^{-1}|_{[0,1]}$$

thus implying that inverses of f and g are also continuous. Thus we follow the desired result.

2.7.6

Find a function $f : R \rightarrow R$ that is continuous at precisely one point.

We've handled that in real analysis course, but I'll state this function here anyways.

We can modify Dirichlet's function to get

$$f(x) = \begin{cases} x \notin Q \Rightarrow x \\ x \in Q \Rightarrow 0 \end{cases}$$

proof of the fact that this function is continuous exclusively at zero was handled using $\epsilon - \delta$ definition in real analysis course.

2.7.7

(a) Suppose that $f : R \rightarrow R$ is "continuous from the right", that is

$$\lim_{x \rightarrow a^+} f(x) = f(a)$$

for each $a \in R$. Show that f is continuous when considered as a function from $R_l \rightarrow R$

Let (a, b) be a basis element of R . Assume that $f^{-1}[(a, b)]$ is nonempty and let $y \in f^{-1}[(a, b)]$. Assume that $x \in (a, b)$ is such that $f(x) = y$. Let ϵ be such that $V_\epsilon(y) \subseteq (a, b)$. We follow that there exists $\delta > 0$ such that $j \in (x, x + \delta)$ implies that $f(j) \in V_\epsilon(y)$. We follow that $j \in [x, x + \delta)$ implies that $f(j) \in V_\epsilon(y)$ since $f(j) = y \in V_\epsilon(y)$. Thus we follow that for every $y \in (a, b)$ there exists interval $[x, x + \delta) \subseteq f^{-1}[(a, b)]$. Now we can define a function $K : (a, b) \rightarrow \mathcal{P}(R)$ such that

$$K(f(x)) = [x, x + \delta) \subseteq f^{-1}[(a, b)]$$

We follow that every value of K is a basis element of R_l and thus union of its range is the open set in topology R_l . From the construction of K we follow that $\text{ran}(K) = f^{-1}[(a, b)]$ and thus we conclude that $f^{-1}[(a, b)]$ is the open set for all intervals (a, b) . Thus we follow that f is continuous, as desired.

(b) Can you conjecture what functions $f : R \rightarrow R$ are continuous when considered as maps from R to R_l ? As maps from $R_l \rightarrow R_l$? We shall return to this question in chapter 3.

We're required only to conjecture, so there're no wrong answers here.

Since R_l is finer than R , we follow that any function, that is continuous in $R \rightarrow R$ is continuous in $R_l \rightarrow R_l$. Same goes for $R \rightarrow R_l$.

For $R_l \rightarrow R_l$ i think that we're gonna have something that has only jump discontinuities. For $R \rightarrow R_l$ I don't think that we're going to have more continuous functions, since the continuity is mainly governed by the domain set.

I wonder if the uniform continuity is just a continuity with respect to some crazy topology, or maybe it is some sort of a new form of continuity. Maybe it has something to do with metrics, or maybe even measures.

2.7.8

Let Y be an ordered set in the order topology. Let $f, g : X \rightarrow Y$ be continuous.

(a) Show that the set $\{x : f(x) \leq g(x)\}$ is closed in X .

Let $z \in X$ be such that $f(z) > g(z)$. Since Y is ordered, we follow that it is Hausdorff, and thus there exist disjoint neighborhoods V_p of $f(z)$ and U_p of $g(z)$. By the fact that open intervals constitute a basis for an ordered topology, we follow that those respective neighborhoods have intervals $V \subseteq V_p$ and $U \subseteq U_p$ such that $f(z) \in V$ and $g(z) \in U$. Since the original neighborhoods are disjoint, we follow that V and U are also disjoint, and thus we follow that

$$(\forall v \in V)(\forall u \in U)(v > u)$$

(in order to follow it faithfully we need to delve deeper into the boundaries and whatnot, but the point is pretty obvious and trivial). Since f and g are continuous, we follow that $f^{-1}[V]$ and $g^{-1}[U]$ are open. Thus $f^{-1}[V] \cap g^{-1}[U]$ is open. We also have that $z \in f^{-1}[V] \cap g^{-1}[U]$. Let $j \in f^{-1}[V] \cap g^{-1}[U]$. We follow that $f(j) \in V$ and $g(j) \in U$ and thus we follow that $f(j) > g(j)$. Therefore we follow that $f^{-1}[V] \cap g^{-1}[U] \subseteq \{x \in X : f(x) > g(x)\}$. Therefore we conclude that for all $j \in X$ if $f(j) > g(j)$ then there is a neighborhood $f^{-1}[V] \cap g^{-1}[U] \subseteq \{x \in X : f(x) > g(x)\}$ such that $j \in f^{-1}[V] \cap g^{-1}[U]$. Thus we follow that we can once again create a function $K : X \rightarrow f^{-1}[V] \cap g^{-1}[U]$ by setting

$$K(j) = f^{-1}[V] \cap g^{-1}[U]$$

if $j \in f^{-1}[V] \cap g^{-1}[U]$ or

$$K(j) = \emptyset$$

otherwise. Since $j \in \{x \in X : f(x) > g(x)\} \Rightarrow j \in K(j)$ we follow that $\{x \in X : f(x) > g(x)\} \subseteq \bigcup \text{ran}(K)$ and we follow that $\bigcup \text{ran}(K) \subseteq \{x \in X : f(x) > g(x)\}$ by construction of the function. Therefore we conclude that $\bigcup \text{ran}(K) = \{x \in X : f(x) > g(x)\}$, and since all values of J are open sets we conclude that $\{x \in X : f(x) > g(x)\}$ is an open set.

We can follow that

$$\{x : f(x) \leq g(x)\} = X \setminus \{x \in X : f(x) > g(x)\}$$

and since $\{x \in X : f(x) > g(x)\}$ is an open set we conclude that $\{x : f(x) \leq g(x)\}$ is closed, as desired.

(b) Let $h : X \rightarrow Y$ be the function

$$h(x) = \min f(x), g(x)$$

Show that h is continuous.

We know from a previous point that

$$A = \{x \in X : f(x) \geq g(x)\}$$

is closed. Also by this point we follow that

$$B = \{x \in X : f(x) \leq g(x)\}$$

is closed. We can follow that functions $g_p : A \rightarrow Y$, $f_p : B \rightarrow Y$ are continuous with respect to the subset topology (pretty trivial point). Thus we follow that by the pasting lemma that the function $h : X \rightarrow Y$, which is produced as in the text of lemma is continuous. And by construction of f_p and g_p we follow that

$$h(x) = \min f(x), g(x)$$

thus h is continuous, as desired.

2.7.9

Let $\{A_\alpha\}$ be a collection of subsets of X ; let $X = \bigcup_\alpha A_\alpha$. Let $f : X \rightarrow Y$; Suppose that $f|_{A_\alpha}$ is continuous for each α .

(a) Show that if the collection $\{A_\alpha\}$ is finite and each set A_α is closed, then f is continuous.

If $A, B \in \{A_\alpha\}$, then we follow by pasting lemma that $f|_{A \cup B} : A \cup B \rightarrow Y$ is continuous. Thus we follow by induction that since $\{A_\alpha\}$ is finite that $f|_{\bigcup \{A_\alpha\}} = f$ is continuous, as desired.

(b) Find an example where the collection $\{A_\alpha\}$ is countable and each A_α is closed, but f is not continuous.

Let $f : [0, 1] \rightarrow \mathbb{R}$ and let $A_0 = \{0\}$ and $A_n = [1/n + 1, 1/n]$. Define

$$f(x) = \begin{cases} 1 & \text{if } x = 0 \\ 0 & \text{otherwise} \end{cases}$$

(c) An indexed family of sets $\{A_\alpha\}$ is said to be locally finite if each point x of X has a neighborhood that intersects A_α for only finitely many values of α . Show that if the family $\{A_\alpha\}$ is locally finite and each A_α is closed, then f is continuous.

Let $x \in X$. We follow that there exists $U \subseteq X$ - neighborhood of x such that U intersects finitely many A_α 's. Let C be a union of A_α 's that intersect U . Let $u \in U$. We follow that since $\bigcup A_\alpha = X$ that u is in some A_α , and thus u is the point of intersection between U and C . Thus we follow that $u \in C$, and therefore $U \subseteq C$.

We follow that $f|_C$ is continuous by the first point. Thus $f|_U : U \rightarrow \mathbb{R}$ is also continuous. Let $K = f[U]$ with subspace topology. We follow that $g : U \rightarrow K$ is continuous.

Let V be a neighborhood of $f(x)$. We follow that $V \cap K$ is open in the subspace topology with respect to K . We also have that $V \cap K \subseteq K$. Thus we follow that $g^{-1}[V \cap K]$ is open by continuity of g . We also follow that $x \in g^{-1}[V \cap K]$. Since U is open in X , we follow that $g^{-1}[V \cap K]$ is open in X as well. Thus we conclude that for all $x \in X$ and each neighborhood V of $f(x)$ there exists neighborhood $g^{-1}[V \cap K]$ of x such that $f[g^{-1}[V \cap K]] \subseteq V$. Therefore f is continuous, as desired.

2.7.10

Let $f : A \rightarrow B$ and $g : C \rightarrow D$ be continuous functions. Let us define a map $f \times g : A \times C \rightarrow B \times D$ by

$$(f \times g)(a \times c) = f(a) \times g(c)$$

Show that $f \times g$ is continuous.

Let $f_1 : A \times C \rightarrow B$ be defined by

$$f_1(\langle a, c \rangle) = f(a)$$

let $f_2 : A \times C \rightarrow D$ be defined by

$$f_2(\langle a, c \rangle) = g(c)$$

Let U be an open set in B . Then we follow that

$$f_1^{-1}[U] = f^{-1}[U] \times C$$

Since f is continuous, we follow that $f^{-1}[U]$ is open. Thus $f^{-1}[U] \times C$ is also open. Thus we follow that for all open $U \subseteq B$ we've got that $f_1^{-1}[U]$ is also open. Thus f_1 is continuous. Same logic can be followed to show that f_2 is also continuous.

Thus we conclude that we can define $h : A \times C \rightarrow B \times D$ by

$$h(\langle a, c \rangle) = \langle f_1(a), f_2(c) \rangle$$

by one of the theorems we follow that h is continuous. We can follow that $h = f \times g$, and thus we conclude that $f \times g$ is continuous, as desired.

2.7.11

Let $F : X \times Y \rightarrow Z$. We say that F is continuous in each variable separately if for each y_0 in Y , the map $h : X \rightarrow Z$ defined by $h(x) = F(x \times y_0)$ is continuous, and for each $x_0 \in X$, the map $k : Y \rightarrow Z$ defined by $k(y) = F(x_0 \times y)$ is continuous. Show that if F is continuous, then F is continuous in each variable separately.

Let $j : X \rightarrow X \times Y$ be defined by

$$j(x) = \langle x, y_0 \rangle$$

Let $U \times V$ be a basis element of $X \times Y$. We follow that if $y_0 \notin V$, then $j^{-1}[U \times V] = \emptyset$, and if $y_0 \in V$, then $j^{-1}[U \times V] = U$. Thus we follow that j is a continuous function.

We can follow that $h = F \circ j$, and thus it is a composition of continuous functions, therefore making it itself a continuous function. Similar logic can show that k is a continuous function as well.

2.7.12

Let $F : R \times R \rightarrow R$ be defined by the equation

$$F(x \times y) = \begin{cases} x \neq 0 \wedge y \neq 0 \Rightarrow \frac{xy}{x^2+y^2} \\ 0 \text{ otherwise} \end{cases}$$

Show that F is continuous in each variable separately

Let $y_0 = 0$. Then we follow that if $x \neq 0$, then

$$F(\langle x, y_0 \rangle) = \frac{0}{x^2 + 0} = 0$$

and if $x = 0$, then $F(\langle x, y_0 \rangle) = 0$. Thus we follow that if $y_0 = 0$, then F is a constant function. Same goes for $x_0 = 0$.

If $y_0 \neq 0$, then we follow that

$$F(\langle x, y_0 \rangle) = \frac{xy_0}{x^2 + y^2}$$

which is continuous by algebraic properties of continuity. Same thing holds for $x_0 \neq 0$. Thus we follow that F is continuous in each variable separately

(b) Compute the function $g : \mathbb{R} \rightarrow \mathbb{R}$ defined by $g(x) = F(x \times x)$.

We follow that if $x \neq 0$, then

$$g(x) = F(x, x) = \frac{x^2}{2x^2} = 1/2$$

and

$$g(0) = 0$$

thus we follow that

$$g(x) = \begin{cases} x = 0 \Rightarrow 0 \\ 1/2 \text{ otherwise} \end{cases}$$

therefore it is obviously not continuous.

(c) Show that F is not continuous

Let $c = \langle 0, 0 \rangle$. We follow that $F(c) = 0$. We follow that $(-0.1, 0.1)$ is a neighborhood of 0. Let C be a neighborhood of c . We follow that there is a basis element $(a_1, b_1) \times (a_2, b_2) \subseteq C$ such that that basis element is a neighborhood of c . Since $c \in (a_1, b_1) \times (a_2, b_2)$, we follow that $a_1, a_2 < 0 < b_1, b_2$. We can follow that there is an element $d \in (a_1, b_1) \times (a_2, b_2)$ such that $d = \langle x_1, x_1 \rangle$ for some $x_1 \neq 0$ (albeit tedious, this little fact can be followed from the properties of intervals). Thus we follow that $f(d) = 1/2$, and thus $f(d) \notin (-0.1, 0.1)$, which contradicts one of the equivalent statements of continuity.

2.7.13

Let $A \subseteq X$; let $f : A \rightarrow Y$ be continuous; let Y be Hausdorff. Show that if f may be extended to a continuous function $f : \bar{A} \rightarrow Y$, then g is uniquely determined by f .

Let $f : A \rightarrow Y$ be defined as in the exercise. Suppose that there exist two functions $g_1, g_2 : \bar{A} \rightarrow Y$ such that both of them are continuous and $f \subseteq g_1 \wedge f \subseteq g_2$.

If $g_1 \neq g_2$, then we follow that there exists $x \in \bar{A}$ such that $g_1(x) \neq g_2(x)$. Since $f \subseteq g_1, g_2$, we follow that $x \notin A$. We follow that $g_1(x), g_2(x) \in Y$, and because Y is Hausdorff, we follow that there exist two disjoint neighborhoods $U, V \subseteq Y$ of $g_1(x)$ and $g_2(x)$ respectively. Since both g_1, g_2 are continuous, we follow that $g_1^{-1}[U]$ and $g_2^{-1}[V]$ are neighborhoods of x . We also follow that since U and V are disjoint, we've got that

$$g_1^{-1}[U] \cap g_1^{-1}[V] = \emptyset$$

$$g_2^{-1}[U] \cap g_2^{-1}[V] = \emptyset$$

but $x \in g_1^{-1}[U] \cap g_2^{-1}[V]$ and thus $g_1^{-1}[U] \cap g_2^{-1}[V]$ is nonempty. By the fact that $g_1^{-1}[U]$ and $g_2^{-1}[V]$ are open, we follow that $g_1^{-1}[U] \cap g_2^{-1}[V]$ is a finite intersection of open sets and thus it is open as well. Thus we follow that $g_1^{-1}[U] \cap g_2^{-1}[V]$ is a neighborhood of x . Thus we follow that it intersects A at some point $a \in A$. Since $x \notin A$, we follow that

$a \neq x$. Therefore we conclude that there's a point $a \in A$ such that $a \in g_1^{-1}[U] \cap g_2^{-1}[V]$. Since $a \in A$, we follow that $g_1(a) = g_2(a) = f(a)$. Thus we can conclude that

$$a \in g_1^{-1}[U] \Rightarrow g_1(a) \in U$$

$$a \in g_2^{-1}[V] \Rightarrow g_2(a) \in V$$

and since $g_1(a) = f(a) = g_2(a)$, we follow that $f(a) \in U \cap V$, which contradicts our assumption that $U \cap V = \emptyset$.

Therefore we conclude that if Y is Hausdorff, $f : A \rightarrow Y$ and there exist continuous functions $g_1, g_2 : \bar{A} \rightarrow Y$ such that $f \subseteq g_1, g_2$, then $g_1 = g_2$, which is not the exact wording of the conclusion of the exercise, but its somewhat more exact equivalence.

2.8 The Product Topology

2.8.1

Prove Theorem 19.2

Let X_α be an original indexed set of topological spaces and B_α be an arbitrary element of the basis. We follow that since each B_α is open that $\prod B_\alpha$ is an element of a basis of the box topology and therefore it's open.

Let x be an arbitrary point in the original space with corresponding neighborhood U . We follow that there is an element of original basis $\prod U_\alpha$ that is a subset of U and contains x . We follow by properties of bases that for each U_α there is an element of basis B_α such that $x_\alpha \in B_\alpha \subseteq U_\alpha$. Therefore we can conclude that there is a set $\prod B_\alpha$ such that $x \in \prod B_\alpha \subseteq \prod U_\alpha$. Therefore we conclude by Lemma 13.2 that collection of products of elements of bases indeed constitutes a basis for a box topology, as desired. Pretty much the same logic with minor tweaks will imply similar result for product topologies.

2.8.2

Prove Theorem 19.3

Firstly, we state that

$$\prod U_\alpha \cap \prod A_\alpha = \prod (U_\alpha \cap A_\alpha)$$

the proof for this statement is somewhat redundant, but I like FOL, so I'll state it anyways:

$$\begin{aligned} x \in \prod U_\alpha \cap \prod A_\alpha &\Leftrightarrow x \in \prod U_\alpha \wedge x \in \prod A_\alpha \Leftrightarrow (\forall \alpha \in J)(x_\alpha \in U_\alpha) \wedge (\forall \alpha \in J)(x_\alpha \in A_\alpha) \Leftrightarrow \\ &\Leftrightarrow (\forall \alpha \in J)(x_\alpha \in U_\alpha \wedge x_\alpha \in A_\alpha) \Leftrightarrow (\forall \alpha \in J)(x_\alpha \in U_\alpha \cap A_\alpha) \Leftrightarrow x \in \prod (U_\alpha \cap A_\alpha) \end{aligned}$$

Lemma 16.1 with our definition and logic in Lemma 16.3 now implies the desired result.

2.8.3

Prove Theorem 19.4

Statement in the previous exercise and logic from exercise 17.11 (2.6.11) imply the desired result.

2.8.4

Show that $(X_1 \times \dots \times X_{n-1}) \times X_n$ is homeomorphic with $X_1 \times \dots \times X_n$.

In set theory course we've defined cartesian product for finite amount of elements recursively, so that exercise boils down to basically nothing. I think that here we need to show that with our new definition of cartesian product everything works.

Let $f : (X_1 \times \dots \times X_{n-1}) \times X_n \rightarrow X_1 \times \dots \times X_n$ be defined by $f(\langle u, v \rangle) = \langle u_1, u_2, \dots, u_{n-1}, v \rangle$. By common sense we follow that f is bijective. Let V be an open set in $X_1 \times \dots \times X_n$. Let B be an element of a basis of $X_1 \times \dots \times X_n$. We follow that $B = \prod U_\alpha$ for $\alpha \in \{1, \dots, n\}$ and some open sets U_α . We follow that

$$f^{-1}[B] = \left(\prod_{\alpha \in \{1, \dots, n-1\}} U_\alpha \right) \times U_n$$

and thus it is open. Thus f is continuous. We also follow that

$$f \left[\left(\prod_{\alpha \in \{1, \dots, n-1\}} U_\alpha \right) \times U_n \right] = \prod U_\alpha$$

thus we follow that f^{-1} is also continuous, thus f is homeomorphism and both spaces are homeomorphic.

2.8.5

One of the implications stated in Theorem 19.6 holds for the box topology. Which one?

Example 2 gives a contradiction for reverse implication, therefore the forward implication must be true.

2.8.6

Let x_1, x_2, \dots be a sequence of points of the product space $\prod X_\alpha$. Show that this sequence converges to the point x if and only if the sequence $\pi_\alpha(x_1), \pi_\alpha(x_2), \dots$ converges to $\pi_\alpha(x)$ for each α . Is this fact true for if one uses the box topology instead of the product topology?

Suppose that the sequence

$$x_1, x_2, \dots$$

converges to a point x . Then we follow that for each neighborhood U of x there is a $n \in \omega$ such that $m \geq n \Rightarrow x_m \in U$.

Let V be a neighborhood of $\pi_\alpha(x)$. Just to be clear here, we're gonna state here that $\pi_\theta : \prod X_\alpha \rightarrow X_\theta$, and thus $\pi_\theta(x) \in X_\theta$. It can be followed easily (if it wasn't done already) that π_θ is a continuous function and thus $\pi_\theta^{-1}(V)$ is an open set in $\prod X_\alpha$. Thus we follow that there is $n \in \omega$ such that $m \geq n \Rightarrow x_m \in \pi_\theta^{-1}(V)$. Since $x_m \in \pi_\theta^{-1}(V)$, we follow that $\pi_\theta(x_m) \in V$. Since everything here was chosen arbitrarily, we follow that for each neighborhood V of $\pi_\theta(x)$ there is $n \in \omega$ such that $m \geq n \Rightarrow x_m \in V$.

Suppose that $\pi_\alpha(x_n)$ converges for every α . Let U be a neighborhood of x . Since U is a neighborhood in box topology, we follow that it is an arbitrary union of finite intersection of finite elements in form $\pi_\theta^{-1}[U_\theta]$. Thus we follow that there exists a finite set C , whose elements are in form $\pi_\theta^{-1}[U_\theta]$, and $x \in \bigcap C \subseteq U$ (trivial proof). Thus we follow that there exists a finite set $D \subseteq \omega$ such that

$$n \in D \Rightarrow (\exists \mu)(\pi_\mu[U_\mu] \in C)$$

Since D is finite, we follow that there is a maximal element of it, and thus there is $n \in \omega$ such that

$$m \in \omega \Rightarrow x_m \in \bigcap C \subseteq U$$

thus we can conclude that our original sequence converges, which proves reverse direction.

We can follow that box topology will work just fine in forward direction, but we can make a whacky open set (like one in the example 2) in $\prod X_n$ so that reverse direction does not work with infinite "dimentions".

2.8.7

Let R^ω be the subset of R^ω consisting of all sequences that are "eventually zero", that is, all sequences (x_1, x_2, \dots) such that $x_i \neq 0$ for only finitely many values of i . What is the closure of R^ω in R^ω in the box and product topologies? Justify your answer.

Let $x \in R^\omega$ and let U be an arbitrary neighborhood of x with respect to product topology. Since U is an open set, we follow that there's a finite set C that consists of elements in form $\pi_\alpha^{-1}[U_\alpha]$ and such that $x \in \bigcap C \subseteq U$. Since C is finite, we follow that we can make a function $f : \omega \rightarrow R$ such that

$$\begin{cases} \pi_\alpha^{-1}[U_\alpha] \in C \Rightarrow f(\alpha) \in \pi_\alpha^{-1}[U_\alpha] \\ 0 \text{ otherwise} \end{cases}$$

we then follow that $f \in R^\omega$, and thus we conclude that every neighborhood of x intersects R^ω at some point. Thus we conclude that with respect to product topology, $\overline{R^\omega} = R^\omega$.

Now assume box topology and let $x \in R^\omega \setminus R^\omega$. We follow that there is an infinite set $D \subseteq \omega$ such that $d \in D \Rightarrow x_d \neq 0$. Assuming that topology of R is standart, we follow that for every $d \in D$ there is an open set (a, b) such that $0 \notin (a, b)$ and $x \in (a, b)$. We now can

define an indexed set (which is a fancy name for a function) $A : \omega \rightarrow \mathcal{P}(R)$ by

$$\begin{cases} a \in D \Rightarrow A(a) = (a, b) \\ A(a) = R \text{ otherwise} \end{cases}$$

which won't have no elements of R^ω on the account that D is infinite and $0 \notin (a, b)$. Thus we conclude that for every $x \in R^\omega \setminus R^\infty$ there's a neighborhood of x that doesn't contain no elements of R^∞ . Thus we conclude that $\overline{R^\infty} = R^\infty$ in box topology.

2.8.8

Given sequences (a_1, a_2, \dots) and (b_1, b_2, \dots) of real numbers with $a_i > 0$ for all i define $h : R^\omega \rightarrow R^\omega$ by the equation

$$h((x_1, x_2, \dots)) = (a_1x_1 + b_1, a_2x_2 + b_2, \dots)$$

Show that if R^ω is given the product topology, h is homeomorphism of R^ω with itself. What happens if R^ω is given the box topology?

Since $a_i \neq 0$ for all i , we follow that h is a bijection (trivial proof).

Let A be a basis element of R^ω with respect to product topology. We follow that it is an indexed set A_α such that $A_\alpha = (c, d)$ or $A_\alpha = R$, and there are finitely many α 's such that $A_\alpha = (c, d)$. Let $n \in \omega$. If $n = \alpha$ for some $\alpha \in J$, then there's an interval (e, f) such that $h_\alpha^{-1}[(e, f)] = (c, d)$ (trivial proof, for clarifications use common sense). We also follow that $h_\alpha[R] = R$, and thus there is an open set C , that consists of product of those intervals and R 's such that $h^{-1}[C] = U$. Similar logic holds in reverse direction. Thus we conclude that h is homeomorphism.

Since we haven't used the fact that a basis element consists of finitely many intervals, we conclude that h is a homeomorphism with respect to a box topology as well.

2.8.9

Show that the choice axiom is equivalent to the statement that for any indexed family $\{A_\alpha\}_{\alpha \in J}$ of nonempty sets, with $J \neq \emptyset$, the cartesian product is not empty.

Forward direction is directly implied by AC (or its lemma with choice function). Reverse implication is almost rewording of AC (or its lemma with choice function). For more GOTO set theory course (2nd and 5th chapter if I remember it correctly).

2.8.10

Skip for now, but by looking at it, we can probably expect some Zorn Lemma action coming

2.9 The Metric Topology

2.9.1

(a) In R^n , define

$$d'(x, y) = \sum |x_i - y_i|$$

Show that d' is a metric that induces the usual topology of R^n . Sketch the basis element under d' when $n = 2$.

Since it's a sum of nonnegative numbers, we follow that $d'(x, y) \geq 0$. We also follow that if $x \neq y$, then there exists $i < n$ such that $x_i \neq y_i$, thus $|x_i - y_i| > 0$ and thus $d'(x, y) > 0$. We also follow that $x = y$, then $d'(x, x) = 0$. Since $|x_i - y_i| = |y_i - x_i|$, we follow that $d'(x, y) = d'(y, x)$. By triangle inequality of normal absolute value function we follow the triangle inequality for this metric.

We can follow that

$$\max\{|x_1 - y_1|, \dots, |x_n - y_n|\} \leq \sum |x_i - y_i|$$

and thus if we set $\delta = \epsilon$ then

$$B_{d'}(x, \delta) \subseteq B_d(x, \epsilon)$$

thus we follow that this topology is finer standard this topology. We also follow that

$$\frac{\sum |x_i - y_i|}{n} \leq \max\{|x_1 - y_1|, \dots, |x_n - y_n|\}$$

and thus we can set $\delta = \frac{\epsilon}{n}$ to get

$$B_d(x, \delta) \subseteq B_{d'}(x, \epsilon)$$

thus standard topology is finer than given topology. Therefore we conclude that the topologies are equal, as desired.

I think that this metric is also called a Manhattan distance. For this one an open set is a diamond in R^2 .

(b) More generally, given $p \geq 1$, define

$$d'(x, y) = \left[\sum_{i=1}^n |x_i - y_i|^p \right]^{1/p}$$

for $x, y \in R^n$. Assume that d' is a metric. Show that it induces the usual topology on R^n .

We follow that

$$\max\{|x_1 - y_1|, \dots, |x_n - y_n|\} \leq \left(\left[\sum_{i=1}^n |x_i - y_i|^p \right]^{1/p} \right)^p$$

and

$$\frac{\left(\sum_{i=1}^n |x_i - y_i|\right)^{1/p}}{n} \leq \max\{|x_1 - y_1|, \dots, |x_n - y_n|\}$$

thus we can set $\delta = \epsilon^p$ in the former case and $\delta = \frac{\epsilon^{1/p}}{n}$ in the latter case to get the desired result (or some variation of those, but the point is clear).

2.9.2

Show that $R \times R$ in the dictionary order is metrizable.

We can define a metric by

$$d(\langle x_1, x_2 \rangle, \langle y_1, y_2 \rangle) = \begin{cases} x_1 = y_1 \Rightarrow \min(|x_2 - y_2|, 1) \\ 1 \text{ otherwise} \end{cases}$$

this is a somewhat modified bounded metric, from which we follow that it is a metric. Proof that this metric indeed generates a dictionary topology is somewhat trivial.

2.9.3

Let X be a metric space with metric d .

(a) Show that $d : X \times X \rightarrow R$ is continuous.

Let us firstly assume that R has a standard topology, X has a metric topology with respect to d and $X \times X$ has a product topology. Let U be a basis element of R . We follow that $U = V_\epsilon(x)$ for some $x, \epsilon \in R$. Let $W = d^{-1}[U]$. Suppose that $\langle z, y \rangle \in W$. We follow that $d(z, y) \in V_\epsilon(x)$, and thus

$$x - \epsilon < d(z, y) < x + \epsilon$$

which a syntactic sugar for

$$x - \epsilon < d(z, y) \wedge d(z, y) < x + \epsilon$$

Let $\langle z', y' \rangle \in X \times X$ be such that

$$d(z', y') < x + \epsilon$$

Let

$$\delta = \frac{x + \epsilon - d(z', y')}{2}$$

Let $B_\delta(z')$ and $B_\delta(y')$ be basis elements. Let $k \in B_\delta(z')$ and $l \in B_\delta(y')$. Then we follow that

$$d(k, l) \leq d(k, z') + d(z', y') + d(y', l) < d(z', y') + 2\delta =$$

$$= d(z', y') + 2 \frac{x + \epsilon - d(z', y')}{2} = d(z', y') + x + \epsilon - d(z', y') = x + \epsilon$$

thus we follow that if $\langle k, l \rangle \in B_\delta(z') \times B_\delta(y')$ then

$$d(k, l) < x + \epsilon$$

Since $B_\delta(z')$ and $B_\delta(y')$ are basis elements, we follow that they are open, thus $B_\delta(z') \times B_\delta(y')$ is an open element in $X \times X$ and thus we follow that for every

$$w \in \{q \in X \times X : d(q) < x + \epsilon\}$$

there is an open subset U of $X \times X$ such that $w \in U$

$$U \subseteq \{q \in X \times X : d(q) < x + \epsilon\}$$

Thus we follow that $\{q \in X \times X : d(q) < x + \epsilon\}$ is an open set.

Let $\langle z', y' \rangle \in X \times X$ be such that

$$x - \epsilon < d(z', y')$$

from triangular inequality we follow that we can set

$$\delta = \frac{d(z', y') - (x - \epsilon)}{2}$$

define k, l and such as in previous paragraph. From triangular inequality we follow that

$$d(z', y') \leq d(k, z') + d(k, l) + d(l, y')$$

$$d(z', y') - d(k, z') - d(l, y') \leq d(k, l)$$

and since $d(k, z'), d(l, y') < \delta$, we follow that

$$d(z', y') - 2\delta \leq d(k, l)$$

thus

$$\begin{aligned} d(k, l) &\geq d(z', y') - 2\delta = d(z', y') - 2 \frac{d(z', y') - (x - \epsilon)}{2} = d(z', y') - d(z', y') + (x - \epsilon) = \\ &= x - \epsilon \end{aligned}$$

thus by the same logic as in the previous paragraph we follow that

$$\{q \in X \times X : d(q) > x - \epsilon\}$$

is open.

Now we can follow that

$$\{q \in X \times X : d(q) < x + \epsilon\} \cap \{q \in X \times X : d(q) > x - \epsilon\}$$

is a finite intersection of open sets, and therefore it is open. By doing some set algebra we get

$$\begin{aligned} & \{q \in X \times X : d(q) < x + \epsilon\} \cap \{q \in X \times X : d(q) > x - \epsilon\} = \\ &= \{q \in X \times X : d(q) > x - \epsilon \wedge d(q) < x + \epsilon\} = \\ &= \{q \in X \times X : x - \epsilon < d(q) \wedge d(q) < x + \epsilon\} = \\ &= \{q \in X \times X : x - \epsilon < d(q) < x + \epsilon\} \end{aligned}$$

is an open set, which is precisely equal to W . Therefore we conclude that d^{-1} maps open sets to open sets, and therefore d is continuous, as desired.

(b) Let X' denote a space having the same underlying set as X . Show that if $d : X' \times X' \rightarrow R$ is continuous, then the topology of X' is finer than the topology of X .

Let $x \in X$ and let U be a basis element of X such that $x \in U$. We follow that since U is a basis element, there's $k \in R$ and $y \in X$ such that $U = B_k(y)$. We now can follow that $(-1, k)$ is open in R and thus $d^{-1}[(-1, y)]$ is open in both $X \times X$ and $X' \times X'$. Now we want to follow that

$$d^{-1}[(-1, y)] \cap \{y\} \times R = \{y\} \times B_k(y)$$

Let us firstly state and prove a theorem:

Strand theorem: Let X, Y be topological spaces and let $x \in X$ and $y \in Y$. We follow that if U is an open set in $X \times Y$, then

$$U \cap X \times \{y\} = V \times \{y\}$$

and

$$U \cap \{x\} \times Y = \{x\} \times W$$

where V is an open set in X and W is an open set in Y .

Suppose that U is an open set in $X \times Y$. We follow that there is an indexed collection (i.e. a function) A_j of elements of basis of $X \times Y$ such that

$$U = \bigcup_{j \in J} A_j$$

We follow that since A_j is a basis element for all $j \in J$ we've got that

$$A_j = X_j \times Y_j$$

such that X_j and Y_j are open sets in X and Y respectively. Let $K \subseteq J$ be such that if $k \in K$ then $y \in Y_k$. Thus we follow that

$$\begin{aligned} \bigcup_{j \in J} A_j \cap X \times \{y\} &= \left(\bigcup_{k \in K} A_k \cup \bigcup_{j \in J \setminus K} A_j \right) \cap X \times \{y\} = \\ &= \left(\bigcup_{k \in K} A_k \cap X \times \{y\} \right) \cup \left(\bigcup_{j \in J \setminus K} A_j \cap X \times \{y\} \right) = \left(\bigcup_{k \in K} A_k \cap X \times \{y\} \right) \cup \emptyset = \\ &= \bigcup_{k \in K} A_k \cap X \times \{y\} \end{aligned}$$

Now we want to follow that

$$\bigcup_{k \in K} A_k \cap X \times \{y\} = \bigcup_{k \in K} X_k \times \{y\}$$

$$q \in \bigcup_{k \in K} A_k \cap X \times \{y\} \Leftrightarrow q = \langle x, y \rangle \wedge x \in \bigcup_{k \in K} X_k \wedge y \in Y \Leftrightarrow q = \langle x, y \rangle \wedge x \in \bigcup_{k \in K} X_k \Leftrightarrow q \in \bigcup_{k \in K} X_k \times \{y\}$$

since X_k is a collection of open sets in X , we follow that we can set $V = \bigcup_{k \in K} X_k$ to get the desired result. The same logic holds for the later case as well. ■

From the strand theorem we follow that $B_k(y)$ is open in X' . Thus we follow that topology on X is coarser than X' , as desired.

2.9.4

Consider the product, uniform and box topologies on R^ω .

Let U_p, U_u and U_b be basis elements of product, uniform and box topologies respectively.

(a) In which topologies are the following functions from R to R^ω continuous.

$$f(t) = (t, 2t, 3t, \dots)$$

$$g(t) = (t, t, t, \dots)$$

$$h(t) = (t, \frac{1}{2}t, \frac{1}{3}t, \dots)$$

Consider the product topology. Let U be a basis element such that $f^{-1}[U], g^{-1}[U], h^{-1}[U] \neq \emptyset$ (not necessarily at the same time, but the point is clear). We follow that

$$U = \prod_{j \in J} A_j$$

We follow that there're finitely many intervals A_j such that $A_j \neq R$ and such that A_j is an open interval in R . Let $K = \bigcap_{j \in J} A_j$. We follow by the fact that there are finitely many

j 's such that $A_j \neq R$ that $f^{-1}(U)$ is an open set, restriction on which comes from the last A_j such that $A_j \neq R$. We can follow that

$$g^{-1}(U) = K$$

and we can follow that h is restricted by the first A_j such that $A_j \neq R$. Therefore we conclude that all of the functions are continuous in product topology.

When it comes to the uniform topology, then we follow that we can create basis elements

$$U = B((1/2 - 1/3, 1/3 - 1/3, 1/4 - 1/3, \dots), 1/3)$$

such that $f^{-1}[U] = g^{-1}[U] = \{0\}$. Thus not g nor f are continuous in uniform topology. We can do a set, similar to the one in box topology set, so that h is $h^{-1}[U]$ is not open as well. Therefore we conclude that no given functions are continuous under uniform topology.

Consider the box topology. We follow that there's a basis element of box topology

$$U = (-1/2, 1/2) \times (-1/3, 1/3) \dots$$

we can follow that the only $V \subseteq R$ such that $f(t) \in U$ is $V = \{0\}$. We can follow that V is not an open set in R , and thus f and g are not continuous with respect to box topology.

We can also create an open set

$$K = \prod (1/n - 1/n^2, 1/n + 1/n^2)$$

We follow that $h(1) \in K$. If $j \neq 1$ thought, then let $k = j - 1$. We follow that if $k > 0$, then

$$h(j) = h(k + 1) = (k + 1, \frac{1}{2}k + \frac{1}{2}, \dots)$$

Since $k \in R$, we follow that there is $l \in Q$ such that $0 < l < k$. We follow that there's $o, p \in N$ such that $l = o/p$ and thus Thus we follow that

$$\frac{1}{n}l + \frac{1}{n} = \frac{o}{np} + \frac{1}{n}$$

and thus there's some $n \in N$ (namely some $n > p$) such that

$$\frac{o}{np} > \frac{1}{n^2}$$

and since $h(l) < h(k)$ we conclude that $h(k) \notin K$. Thus we follow that $h^{-1}[K] = \{1\}$, which proves that h is not continuous as well.

(b) *In which of topologies do the following sequences converge?*

Firtly, we must state that all of the sequences either converge to $(0, 0, \dots)$ or don't converge at all. For the simplicity of notation I'm gonna use 0 as a shorthand for this vector.

Consider the box topology. We follow that there's

$$U = \prod_{j \in \omega} (1, -1)$$

we follow that $0 \in U$ but every element of w_n such that $n > 1$ has the property that $0 \notin w_n$. Thus we conclude that w_n does not converge in box topology.

If we set

$$U = \prod_{j \in \omega} (1/j^2, -1)$$

then the fact that $1/j^2 < 1/j$ for all $j \in \omega$ will give us that x_n does not converge in box topology. Same U will suffice to show that y_n does not converge in box topology because for every $n \in \omega$ there's $j \in \omega$ such that $1/n > 1/j$, thus giving us that at n 'th position both y_n and x_n won't be confined inside U .

Case with z_n is different though. We follow that if U is a basis element in box topology, then there's two open intervals A_1, A_2 such that $A_1, A_2 \in \text{ran } U_j$. We follow that $A_1 \cap A_2$ is an interval as well, and thus we follow that there's $n \in \omega$ such that $1/n \in A_1 \cap A_2$. Thus we conclude that z_n will converge in box topology. Given that both uniform and product topologies are coarser than box topology, we can follow that z_n converges in those topologies as well.

Consider uniform topology. We follow that

$$0 \in B(0, 1)$$

but $j > 2 \Rightarrow w_j \notin B(0, 1)$. Thus we follow that w_n does not converge in uniform topology. We can use a method of sliding interval (just as in part (a)) to conclude that x_n does not converge in uniform topology.

We can follow that if $x \neq (k, 0, 0, \dots)$ for some $k \in R$, then $0 \notin B(x, \epsilon)$. Thus we conclude that if $0 \in B(x, \epsilon)$, then $x = (k, 0, 0, \dots)$. We can also follow pretty easily that $\epsilon > k$. Thus we can follow that y_n converges in uniform topology. Since product topology is coarser than uniform, we follow that y_n converges in product topology as well.

Consider product topology. Let U be a basis element such that $0 \in U$. We follow that

$$U = \prod_{j \in \omega} U_j$$

and by the nature of basis elements in product topologies we follow that there's $n \in \omega$ such that $U_j \neq R \Rightarrow j < n$. Thus we follow that there's $n \in \omega$ such that $w_n \in U$. Therefore we conclude that w_n converges in product topology. Slightly modified logic can tell us that x_n converges in product topology as well.

2.9.5

Let R^∞ be the subset of R^ω consisting of all sequences that are eventually zero. What is the closure of R^∞ in R^ω in the uniform topology? Justify your answer.

I want to say that it's all the sequences that converge to 0.

Suppose that $z_n \in R^\omega$ is such that it does converge to 0. Let $B(x_n, \epsilon)$ be a basis element of uniform topology such that $z_n \in B(x, \epsilon)$. We follow that there's $n \in \omega$ such that $m \geq n \Rightarrow |x_n| < \epsilon$. This gives us that there's an element y_n of R^∞ , specifically

$$\begin{cases} m < n \Rightarrow y_m = x_n \\ m \geq n \Rightarrow y_m = 0 \end{cases}$$

such that $y_n \in B(x_n, \epsilon)$. Thus we conclude that if z_n converges to 0, then $z_n \in \overline{R^\infty}$ in uniform topology.

Let $x_n \in R^\omega$ be such that it does not converge to 0. We follow that there exists $\epsilon > 0$ such that for all $n \in \omega$

$$|x_n| > \epsilon$$

moreover, we follow that there's $\epsilon \in (0, 1)$ that will do the job. Thus we conclude that there's a basis element $B(x_n, \epsilon)$ such that there's no element z of R^∞ such that $z \in B(x_n, \epsilon)$. Therefore we conclude that if x_n does not converge to 0, then $x_n \notin \overline{R^\infty}$.

Thus we can conclude that $x_n \in \overline{R^\infty}$ if and only if x_n converges to 0 (or so I think).

2.9.6

Let ρ be the uniform metric on R^ω . Given $x = (x_1, x_2, \dots) \in R^\omega$ and given $0 < \epsilon < 1$, let

$$U(x, \epsilon) = (x_1 - \epsilon, x_1 + \epsilon) \times (x_2 - \epsilon, x_2 + \epsilon) \times \dots$$

(a) Show that $U(x, \epsilon)$ is not equal to ϵ -ball $B_\rho(x, \epsilon)$.

Let

$$y = (x_1 + \frac{1}{2}\epsilon, x_2 + \frac{2}{3}\epsilon, \dots)$$

we follow that $y \in U(x, \epsilon)$. At the same time we follow that

$$\rho(x, y) = \sup\{\bar{d}(x_n, y_n) : n \in \omega\} = \sup\{\frac{n-1}{n}\epsilon : n \in \omega\} = \epsilon$$

since

$$y \in B_\rho(x, \epsilon) \Leftrightarrow \rho(x, y) < \epsilon$$

we follow that $y \notin B_\rho(x, \epsilon)$.

(b) Show that $U(x, \epsilon)$ is nor even open in the uniform topology

We can follow that there's no basis element of uniform topology such that it contains y and is a subset of $U(x, \epsilon)$.

(c) Show that

$$B_\rho(x, \epsilon) = \bigcup_{\delta < \epsilon} U(x, \delta)$$

Suppose that $y \in B_\rho(x, \epsilon)$. We follow that

$$\rho(y, x) < \epsilon$$

thus we follow that there's $\delta \in R$ such that $\rho(y, x) < \delta < \epsilon$. We follow that $y \in U(x, \delta)$ and thus $y \in \bigcup_{\delta < \epsilon} U(x, \delta)$. Therefore

$$B_\rho(x, \epsilon) \subseteq \bigcup_{\delta < \epsilon} U(x, \delta)$$

If $y \in \bigcup_{\delta < \epsilon} U(x, \delta)$, then there's $\delta < \epsilon$ such that $y \in U(x, \delta)$. We follow that $\rho(y, x) \leq \delta < \epsilon$ and thus $y \in B_\rho(x, \epsilon)$. Thus

$$B_\rho(x, \epsilon) \supseteq \bigcup_{\delta < \epsilon} U(x, \delta)$$

Double inclusion gives us the desired result.

2.9.7

Consider the map $h : R^\omega \rightarrow R^\omega$ defined in Exercise 8 of paragraph 19, give R^ω the uniform topology. Under what conditions on the numbers a_i and b_i is h continuous? a homeomorphism?

$h : R^\omega \rightarrow R^\omega$ is defined as

$$h((x_1, x_2, \dots)) = (a_1x_1 + b_1, a_2x_2 + b_2, \dots)$$

We can also use some notation from linear algebra (where we denote piecewise vector multiplication by \cdot) to get

$$h((x_1, x_2, \dots)) = ((x_1, x_2, \dots) \cdot (a_1, a_2, \dots)) + (b_1, b_2, \dots)$$

and if none of a_i 's are zeroes, then

$$\begin{aligned} h^{-1}((x_1, x_2, \dots)) &= ((x_1, x_2, \dots) - (b_1, b_2, \dots)) \cdot \left(\frac{1}{a_1}, \frac{1}{a_2}, \dots\right) = \\ &= ((x_1, x_2, \dots) \cdot \left(\frac{1}{a_1}, \frac{1}{a_2}, \dots\right) - (b_1, b_2, \dots) \cdot \left(\frac{1}{a_1}, \frac{1}{a_2}, \dots\right)) = \left(\frac{x_1}{a_1}, \frac{x_2}{a_2}, \dots\right) - \left(\frac{b_1}{a_1}, \frac{b_2}{a_2}, \dots\right) \end{aligned}$$

although the notation is somewhat abusive, we only want to show the idea; the same conclusion can be drawn with ol' reliable FOL and stuff like that.

The conjecture is that set of a_i 's need to be bounded. Let's expand our idea a bit and let a_i and b_i 's be arbitrary for now. We follow that if $B(x, \epsilon)$ is a basis element and $B(x, \epsilon) \cap h^{-1}(R^\omega) \neq \emptyset$, then there exists $y \in R^\omega$ such that

$$h(y) \in B(x, \epsilon)$$

Suppose that there's a basis element around y such that its image is contained in $B(x, \epsilon)$. We follow that there's $\gamma \in R$ such that $\gamma > 0$ and

$$h(B(y, \gamma)) \subseteq B(x, \epsilon)$$

We follow that there's $\beta \in R$ such that $0 < \beta < \gamma$ and thus we follow that

$$y + (\beta, \beta, \dots) \in B(y, \gamma)$$

as proven in previous exercise. We follow then that

$$h(y + (\beta, \beta, \dots)) \subseteq B(x, \epsilon)$$

we thus follow that for all i 's we've got that

$$(y_i + \beta)a_i + b_i \in (x_i - \gamma, x_i + \gamma)$$

thus

$$x_i - \gamma < (y_i + \beta)a_i + b_i < x_i + \gamma$$

$$x_i - \gamma - b_i < (y_i + \beta)a_i < x_i + \gamma - b_i$$

$$x_i - \gamma - b_i < y_i a_i + \beta a_i < x_i + \gamma - b_i$$

$$x_i - \gamma - b_i - y_i a_i < \beta a_i < x_i + \gamma - b_i - y_i a_i$$

$$x_i - y_i a_i - b_i - \gamma < \beta a_i < x_i - b_i - y_i a_i + \gamma$$

for all i 's. Let

$$z = x - ay - b$$

i.e.

$$(z_1, z_2, \dots) = (x_1, x_2) - (a_1, a_2, \dots) \cdot (y_1, y_2, \dots) - (b_1, b_2, \dots)$$

we follow then that thus we follow that

$$\beta a \in U(z, \gamma)$$

and by using pretty much the same logic we can follow that

$$(-\beta)a \in U(z, \gamma)$$

thus for all i 's we've got that

$$\begin{cases} z_i - \gamma < \beta a_i < z_i + \gamma \\ z_i - \gamma < -\beta a_i < z_i + \gamma \end{cases}$$

let's do some algebra on the second inequality

$$-z_i + \gamma > \beta a_i > -z_i - \gamma$$

$$-z_i - \gamma < \beta a_i < -z_i + \gamma$$

thus we get

$$\begin{cases} z_i - \gamma < \beta a_i < z_i + \gamma \\ -z_i - \gamma < \beta a_i < -z_i + \gamma \end{cases}$$

thus we follow that we can sum up two inequalities to get

$$z_i - \gamma - z_i - \gamma < 2\beta a_i < z_i + \gamma - z_i + \gamma$$

$$-2\gamma < 2\beta a_i < 2\gamma$$

$$-\gamma < \beta a_i < \gamma$$

$$-\gamma/\beta < a_i < \gamma/\beta$$

thus we follow that

$$|a_i| < \gamma/\beta$$

which means that a_i 's ought to be bounded, otherwise we don't have a basis element around y whose image is contained in the original basis element.

Now suppose that a_i 's are bounded by M . If M is zero, then we follow that a_i 's are certainly are bounded by some nonzero M , thus let us follow that in those circumstances M is a positive real number. Let $B(x, \epsilon)$ be a basis element. Now we follow that there's two cases, namely

$$B(x, \epsilon) \cap \text{ran}(h) = \emptyset$$

or

$$B(x, \epsilon) \cap \text{ran}(h) \neq \emptyset$$

if the former case is true, then $h^{-1}[B(x, \epsilon)] = \emptyset$, which is an open set and we're done. Thus assume the latter case. Let $y \in B(x, \epsilon) \cap \text{ran}(h)$. We want to show now that there's a basis element around y that is contained in $y \in B(x, \epsilon) \cap \text{ran}(h)$, which would imply that $h^{-1}[B(x, \epsilon)]$ is an open set. Previous exercise implies that

$$B_\rho(x, \epsilon) = \bigcup_{\delta < \epsilon} U(x, \delta)$$

which means that there's $\delta \in R$ such that $0 < \delta < x$ and

$$h(y) \in U(x, \delta)$$

we also follow that since $\delta, x \in R$ and $\delta \neq x$ there's $\gamma \in R$ such that

$$\delta < \gamma < x$$

and thus

$$U(x, \delta) \subseteq U(x, \gamma)$$

therefore

$$h(y) \in U(x, \gamma)$$

and

$$h(y) \in U(x, \delta)$$

Therefore

$$h(y) \in \prod [(x_i - \gamma, x_i + \gamma)]$$

$$h(y) \in \prod [(x_i - \delta, x_i + \delta)]$$

we follow that for all i 's we've got that

$$x_i - \delta < a_i y_i + b_i < x_i + \delta$$

$$-\delta < a_i y_i + b_i - x_i < \delta$$

$$|a_i y_i + b_i - x_i| < \delta$$

Let $z \in U(y, \frac{\gamma - \delta}{M})$. We follow that for all i 's we've got

$$|y_i - z_i| < \frac{\gamma - \delta}{M}$$

$$M|y_i - z_i| < \gamma - \delta$$

given that $|a_i| < M$ for all a_i 's, we follow that

$$|a_i||y_i - z_i| < \gamma - \delta$$

$$|a_i y_i - a_i z_i| < \gamma - \delta$$

$$|a_i y_i - a_i z_i| + \delta < \gamma$$

using earlier proven inequality $|a_i y_i + b_i - x_i| < \delta$ we get

$$|a_i y_i - a_i z_i| + |a_i y_i + b_i - x_i| < \gamma$$

now we need to use some absolute value properties to get

$$|a_i y_i - a_i z_i| + |-a_i y_i - b_i + x_i| < \gamma$$

let us use the magestic triangle ($|a + b| \leq |a| + |b|$) now

$$|a_i y_i - a_i z_i - a_i y_i - b_i + x_i| < \gamma$$

$$|-a_i z_i - b_i + x_i| < \gamma$$

$$|a_i z_i + b_i - x_i| < \gamma$$

$$-\gamma < a_i z_i + b_i - x_i < \gamma$$

$$x_i - \gamma < a_i z_i + b_i < x_i + \gamma$$

thus

$$h(z) \in U(x, \gamma)$$

and therefore

$$h(z) \in B(x, \epsilon)$$

therefore we conclude that if $z \in U(y, \frac{\gamma - \delta}{M})$ then $h(z) \in B(x, \epsilon)$. Given that

$$B(y, \frac{\gamma - \delta}{M}) \subseteq U(y, \frac{\gamma - \delta}{M})$$

(proof of that is pretty straightforward, so I'll omit it) we follow that

$$z \in B(y, \frac{\gamma - \delta}{M}) \Rightarrow h(z) \in B(x, \epsilon)$$

thus we conclude that around every point of $h^{-1}[B(x, \epsilon)]$ there's a basis element that is contained in $h^{-1}[B(x, \epsilon)]$. Thus we conclude that for all $B(x, \epsilon)$ we've got that $h^{-1}[B(x, \epsilon)]$ is open and thus h is continous, as desired.

Now to the question of homeomorphism. We can follow pretty easily that if some a_i 's are equal to zero, then h is not bijective, and if a_i 's are all nonzero, then the function is bijective (if the proof of that is not included in my linear algebra course, then it's pretty straightforward anyways). Thus we conclude that all a_i 's ought to be nonzero in order to function to be homeomorphism. If none of the a_i 's are zero, then we can follow that

$$h^{-1}((x_1, x_2, \dots)) = (x_1 \frac{1}{a_1}, x_2 \frac{1}{a_2}, \dots) + (-\frac{b_1}{a_1}, -\frac{b_2}{a_2}, \dots)$$

as shown in the beginning of this solution. Thus we follow that h^{-1} is continous if and only if $\frac{1}{a_i}$'s are bounded by some number. This statement is equivalent to saying that there's $\gamma \in \mathbb{R}$ such that $\gamma < |a_i|$ for all a_i 's. This also takes care of the nonzero clause.

Therefore we conclude that h is a homeomorphism if and only if there exist $M, \gamma \in \mathbb{R}$ such that

$$\gamma < |a_i| < M$$

for all a_i 's.

2.9.8

Let X be the subset of R^ω consisting of all sequences x such that $\sum x^2$ converges. Then the formula

$$d(x, y) = \left[\sum_{i=1}^{\infty} (x_i - y_i)^2 \right]^{1/2}$$

defines a metric on X . On X we have the three topologies it inherits from the box, uniform and product topologies on R^ω . We have also the topology given by the metric d , which we call the l^2 topology

(a)) Show that on X , we have the inclusions

$$\text{uniform topology} \subseteq l^2\text{-topology} \subseteq \text{box topology}$$

Suppose that $B_\rho(x, \epsilon)$ is a basis element of uniform topology such that $B_\rho(x, \epsilon) \cap X$ is not empty. We follow that $B_\rho(x, \epsilon) \cap X$ is a basis element of inherited topology on X from uniform topology. Let $y \in B_\rho(x, \epsilon) \cap X$. We follow that

$$\rho(x, y) < \epsilon$$

thus

$$\sup\{\bar{d}(x_i, y_i)\} < \epsilon$$

Now let $B_d(x, \epsilon/2)$ be a basis element in l^2 -topology. We follow that for all $z \in B_d(x, \epsilon/2)$ we've got that

$$\left[\sum_{i=1}^{\infty} (x_i - z_i)^2 \right]^{1/2} < \epsilon/2$$

$$\sum_{i=1}^{\infty} (x_i - z_i)^2 < (\epsilon/2)^2$$

we thus follow that for all x_i and z_i we've got

$$(x_i - z_i)^2 < (\epsilon/2)^2$$

thus

$$|x_i - z_i| < \epsilon/2$$

therefore $z \in U(x, \epsilon/2)$. Therefore $z \in B_\rho(x, \epsilon)$. Thus we conclude that topology inherited from uniform topology on R^ω is coarser than l^2 -topology on X , which gives us the first inclusion.

Now let $B_l(x, \epsilon)$ be a basis element in l^2 topology. We follow that we can make a basis element in box topology

$$U = \prod (x_i - \epsilon * 2^{-i}, x_i + \epsilon * 2^{-i})$$

we then follow that if $y \in U \cap X$, then

$$d(x, y) < \epsilon$$

thus we conclude that l^2 -topology is coarser than the box topology, as desired.

(b) The set R^∞ of all sequences that are eventually zero is contained in X . Show that the four topologies that R^∞ inherits as a subspace of X are all distinct.

Let

$$U = \prod (-1/n, 1/n)$$

be a basis element in the box topology. We follow that $U \cap R^\infty \neq \emptyset$, by the fact that 0 vector is in both of those elements. Thus we conclude that $U \cap R^\infty$ is a nonempty open set in inherited topology with respect to the box topology.

Let $B_d(x, \epsilon)$ be a basis element around 0 with respect to the given metric (the one, that defines l^2 topology). Since $\epsilon > 0$ we can follow that $\epsilon/2 > 0$ and thus there's $n \in \mathbb{N}$ such that $1/n < \epsilon/2$. We follow that element

$$f_x = \begin{cases} x = n \Rightarrow \epsilon/2 \\ 0 \text{ otherwise} \end{cases}$$

is an element in R^∞ , and we can also conclude that $f \in B_d(x, \epsilon)$ by the fact that

$$\sqrt{\sum (x_i - f_i)^2} = \sqrt{(\epsilon/2)^2} = \epsilon/2$$

thus $d(x, f) < \epsilon$. We can also follow that $f \notin U$, because we've specifically engineered it not to be in it. Given that ϵ is arbitrary, we can conclude that there's an x such that there's no basis neighborhood in l^2 -topology that is a subset of $U \cap R^\infty$ and thus we conclude that $U \cap R^\infty$ is not open in l^2 -topology.

We can define

$$V = \prod (0, 1/n)$$

to be an open set in l^2 -topology, which is

TODO: later

2.9.9

This one is handled in the linear algebra course.

2.10 The Metric Topology (continued)

2.10.1

Let $A \subseteq X$. If d is a metric for the topology of X , show that $d|_{A \times A}$ is a metric for the subspace topology on A .

We follow that if some metric property does not hold for a pair of points in A , then it doesn't hold in X as well, which produces contradiction, and thus $d|_A \times A$ is a metric on set A .

Let U be an open set in A . We follow that there's an open set $V \subset X$ such that $U = V \cap A$. Since V is an open set, it is a union of balls, and thus U is union of open balls in A . Thus we conclude that $d|_A \times A$ imposes the same topology, as desired.

2.10.2

Let X and Y be metric spaces with metrics d_X and d_Y respectively. Let $f : X \rightarrow Y$ have the property that for every pair of points x_1, x_2 of X

$$d_Y(f(x_1), f(x_2)) = d_X(x_1, x_2)$$

Show that f is an imbedding. It is called isometric imbedding of X in Y .

Suppose that $x_1, x_2 \in X$ and $x_1 \neq x_2$. We follow that

$$d_X(x_1, x_2) \neq 0$$

and thus

$$d_Y(f(x_1), f(x_2)) \neq 0$$

thus $f(x_1) \neq f(x_2)$. Thus we conclude that f is injective.

Let $x \in X$ and $\epsilon \in \mathbb{R}$ be arbitrary. Then we follow that if we set $\delta = \epsilon$ and get a point $x' \in X$ such that

$$d_X(x, x') < \delta$$

then

$$d_X(x, x') < \epsilon$$

and since $d_X(x, x') = d_Y(f(x), f(x'))$ we follow that

$$d_Y(f(x), f(x')) < \epsilon$$

thus we conclude that f is continuous. By the same logic we can prove that $f^{-1} : \text{ran}(f) \rightarrow X$ is continuous as well.

Thus we follow that f is an imbedding by definition, as desired.

2.10.3

Let X_n be a metric space with metric d_n for $n \in \mathbb{Z}_+$.

(a) Show that

$$\rho(x, y) = \max\{d_1(x_1, y_1), \dots, d_n(x_n, y_n)\}$$

is a metric for the product space $X_1 \times \dots \times X_n$

We follow that $\rho(x, y) \geq 0$ for all $x \in R$ by the virtue of the fact that all of the numbers under the max are non-negative. If $x = y$, then $d_j(x, y) = 0$ for all j 's, and thus $\rho(x, y) = \max\{0\} = 0$. If $x \neq y$, then there is a component j such that $d_j(x, y) \neq 0$, and thus we've got the first property of the metric.

We can follow commutativity of this thing by common sense.

Suppose that $x, y, z \in \prod X_i$. We follow that for all j 's we've got

$$d_j(x_j, z_j) \leq d_j(x_j, y_j) + d_j(y_j, z_j)$$

Then we can follow the triangle by either induction or contradiction.

Now suppose that U is a basis element of $\prod X_i$ and $x \in U$. We follow that there are open sets V_j such that $U = \prod V_j$. Given that any of the V_j 's are open, we follow that for each one of them there's a basis element B_j such that B_j is centered on x_j and $B_j \subseteq V_j$ and we follow that $\prod(B_j) \subseteq \prod V_j$. Each B_j has ϵ_j such that $B_j = B_j(x_j, \epsilon_j)$ and thus we can follow that if $\rho(x, y) < \max\{\epsilon_j\}$, then $y \in \prod(B_j)$. Thus we follow that

$$B_\rho(x, \max\{\epsilon_j\}) \subseteq U$$

, therefore product topology is a subset of topology, that is induced by this metric. We can also follow that every basis element of topology, that is induced by this metric is a basis element in product topology, and thus we conclude that given metric induces the product topology, as desired.

(b) Let $\bar{d}_i = \min\{d_i, 1\}$. Show that

$$D(x, y) = \sup \bar{d}_i(x_i, y_i) / i$$

is a metric for the product space $\prod X_i$.

This proof is practically the same as in theorem 20.5. If it ain't, then I skip it anyways

2.10.4

Show that R_l and the ordered square satisfy the first countability axiom

Let $x \in R$. We follow that $[x, x + 1/n)$ is the countable collection, that satisfies the countability axiom.

For the ordered square we've got either $(\langle x_1, x_2 - 1/n \rangle, \langle x_1, x_2 + 1/n \rangle)$ or some similar stuff for the edges.

the rest

the rest of the exercises were taken care of in the real analysis course. Maybe later I'll repeat the proofs for s 'n'g

2.11 The Quotient Topology

Chapter 3

Connectedness and Compactness

3.1 Connected Spaces

3.1.1

Let \mathcal{T} and \mathcal{T}' be two topologies on X . If $\mathcal{T} \subseteq \mathcal{T}'$, what does connectedness of X in one topology imply about connectedness in the other?

We follow that if $\mathcal{T} \subseteq \mathcal{T}'$ and U is a connected subspace of \mathcal{T}' , then there's no separations in \mathcal{T}' , and thus there're no separations in \mathcal{T} as well.

If U is open in \mathcal{T} , then and space X has some sane amount of elements (i.e. not zero or one), then we can't follow nothing from it. For example, every topology is a subset of a discrete topology on a set, and every subspace of discrete topology is disconnected.

3.1.2

Let $\{A_n\}$ be a sequence of connected subspaces of X , such that $A_n \cap A_{n+1} \neq \emptyset$ for all n . Show that $\bigcup A_n$ is connected.

We firstly assume that $|\{A_n\}| \leq \omega$ because of the word "sequence"

Assume that $\bigcup A_n$ is disconnected and has a separation U and V . We thus follow that for all $\{A_n\}$ we've got that A_n is in either U or V . Assume that A_1 is in U . Thus we follow that there's a minimal j such that $A_j \subseteq V$ (if there isn't one, then V is empty, which is a contradiction). Therefore $A_{j-1} \subseteq U$ and $A_j \subseteq V$, which means that $A_{j-1} \cap A_j = \emptyset$, which is a contradiction.

3.1.3

Let A_α be a collection of connected subspaces of X ; let A be a connected subspace of X . Show that if $A \cap A_\alpha \neq \emptyset$ for all α , then $A \cup (\bigcup A_\alpha)$ is connected.

Unlike the previous exercise, we can't reasonably assume nothing about the collection A_α . We firstly state that

$$A \cup \left(\bigcup A_\alpha \right) = \bigcup (A \cup A_\alpha)$$

which is either handled in the set theory course, or is justified by some trivial FOL.

Since $A \cap A_\alpha \neq \emptyset$ for all α , we follow that $A \cap A_\alpha$ is connected for all α , thus 23.3 implies that $\bigcup (A \cup A_\alpha)$ is connected as well.

3.1.4

Show that if X is an infinite set, it is connected in the finite complement topology

Suppose that it ain't. We follow that sets U and V form separation. Given that U and V are open, we follow that $X \setminus U = V$ is finite and thus V is not open, which is a contradiction.

3.1.5

A space is totally disconnected if its only connected subspaces are one point sets. Show that if X has the discrete topology, then X is totally disconnected. Does the converse hold?

I'm pretty sure that we don't need to prove that one point sets are connected. If U is a subspace with more than one point in it, then we follow that there's $u \in U$ and thus $U \setminus \{u\}$ and $\{u\}$ make a separation of U .

We've seen already that Q in R is totally disconnected (if there're 2 or more points in $U \subseteq Q$, then we can take $i \in (\inf(U), \sup(U)) \cap I$, where I is irrational, and then $(-\infty, i) \cap U$ and $(i, \infty) \cap U$ form the desired separation) and it just inherits the standard topology, so the converse doesn't hold.

3.1.6

Let $A \subseteq X$. Show that if C is a connected subspace of X that intersects both A and $X \setminus A$, then C intersects $Bd(A)$

We firstly state here that

$$Bd(A) = \overline{A} \setminus Int(A)$$

and

$$Bd(A) = \overline{A} \cap \overline{X \setminus A}$$

Suppose that C does not intersect $Bd(A)$. We firstly want to follow that $X \setminus \overline{A}$ and \overline{A} make a partition of set X , and since $Bd(A)$ and $Int(A)$ are partitions of \overline{A} , we follow that $X \setminus \overline{A}$, $Bd(A)$ and $Int(A)$ form a partition of X . We also note that $X \setminus \overline{A}$ and $Int(A)$ are open sets. Since C does not intersect $Bd(A)$, we follow that there're sets $X \setminus \overline{A} \cap C$ and $Int(A) \cap C$ that form a separation of C , which means that C is not connected, which is a contradiction.

3.1.7

Is the space R_l connected? Justify your answer.

We follow that

$$(-\infty, 0), [0, \infty)$$

are both open in R_l since

$$(-\infty, 0) = \bigcup_{n \in \mathbb{Z}_+} [-n, -n + 1)$$

$$[0, \infty) = \bigcup_{n \in \mathbb{Z}_+} [n - 1, n)$$

and

$$(-\infty, 0) \cup [0, \infty) = R$$

thus we follow that those two sets constitute a separation.

3.1.8

Determine whether or not R^ω is connected in the uniform topology

We can follow that R^ω can be partitioned in the same manner as the box topology. We can follow that $B(x, \epsilon) \subseteq U(x, \epsilon)$, and then proceed with the same logic as in the example 6 to show that bounded and unbounded sequences form a separation on R^ω .

3.1.9

Let A be a proper subset of X , and let B be a proper subset of Y . If X and Y are connected, show that

$$(X \times Y) \setminus (A \times B)$$

is connected.

The idea here is to plant a cross somewhere in the desired set, and then add strands to this set, until we prove that the desired set is connected.

Let $z \notin A$ and $q \notin B$ (those points exist since both A and B are proper subsets). Thus $\langle z, q \rangle \in (X \times Y) \setminus (A \times B)$. Now we can follow that set

$$\langle z, q \rangle \in \{z\} \times Y \cup X \times \{q\}$$

is a subset of

$$(X \times Y) \setminus (A \times B)$$

and the set $\{z\} \times Y \cup X \times \{q\}$ is connected. Now let us use the identity

$$(X \times Y) \setminus (A \times B) = [X \times (Y \setminus B)] \cup [(X \setminus A) \times Y]$$

we now follow that for all $j \in (Y \setminus B)$ it is true that every set $X \times \{j\}$ is connected and also that

$$X \times \{j\} \cap \{z\} \times Y \cup X \times \{q\}$$

is nonempty. Therefore we conclude that

$$[X \times (Y \setminus B)] \cup \{z\} \times Y \cup X \times \{q\}$$

is connected. By the same logic we follow that

$$[(X \setminus A) \times Y] \cup \{z\} \times Y \cup X \times \{q\}$$

is connected. Since both of those sets have a point $\langle z, q \rangle$ in common, we conclude that the desired set is connected as well, as desired.

3.1.10

Let $\{X_\alpha\}_{\alpha \in J}$ be an indexed family of connected spaces; let X be a product space

$$X = \prod_{\alpha \in J} X_\alpha$$

Let $a = (a_\alpha)$ be a fixed point of X .

Important point: in this exercise we assume the product topology on X .

(a) Given any finite subset K of J , let X_K denote the subspace of X consisting of all points $x = (x_\alpha)$ such that $x_\alpha = a_\alpha$ for $\alpha \notin K$. Show that X_K is connected.

We can prove the desired result by stating that $\prod_{j \in K} X_j$ is homeomorphic to X_K , and since the former is a cartesian product of finite amount of connected spaces we follow that both of those spaces are connected.

(b) Show that the union Y of the spaces X_K is connected

All of X_K 's have point a in common, and thus their union is connected.

(c) Show that X equals the closure of Y ; conclude that X is connected.

Suppose that $b \in X$. Let U be a basis neighborhood of b (in the product topology of course). Since U is a basis element, we follow that there's a finite set K and function $f : J \rightarrow \mathcal{P}(\bigcup X_j)$ such that

$$f(j) = \begin{cases} j \notin K \Rightarrow X_k \\ j \in K \Rightarrow V_j \end{cases}$$

and

$$U = \prod_{j \in J} f(j)$$

since K is finite, we follow that U intersects X_K for that particular K , and since Y is a union of X_K 's, we follow that U intersects Y . Since U is arbitrary, we follow that b is in a closure of Y . Therefore

$$X \subseteq \overline{Y}$$

We can also state that $\overline{Y} \subseteq X$ since we use the product topology on X , and thus we conclude that $\overline{Y} = X$ by double inclusion, as desired.

Last two exercises are left for the time, when I'm finished with quotient products.

3.2 Connected Subspaces of the Real Line

3.2.1

(a) *Show that no two spaces $(0, 1)$, $(0, 1]$, and $[0, 1]$ are homeomorphic.*

Suppose that $h : (0, 1] \rightarrow (0, 1)$ is a homeomorphism. Since both of those sets have the same cardinality, we follow that there's a bijection between the two, and therefore we won't dispute that.

Let $u = h(1)$. We follow that $u \in (0, 1)$ and thus $(0, 1) \setminus \{h(1)\}$ is a disconnected set. Thus we follow that

$$h[(0, 1)] = h[(0, 1]] \setminus h(1) = (0, 1) \setminus \{h(1)\}$$

is a disconnected set, which contradicts the fact that h is a homeomorphism.

By pretty much the same logic, but applied to two points we can show that the resulting two pairs aren't homeomorphic.

(b) *Suppose that there exists imbeddings $f : X \rightarrow Y$ and $g : Y \rightarrow X$. Show by means of an example that X and Y need not be homeomorphic.*

We've shown earlier that any two closed intervals are homeomorphic, and also that any two open intervals are homeomorphic as well. Thus we can follow that we can create imbeddings out of homeomorphisms

$$f : [0, 1] \rightarrow [1/3, 2/3]$$

and

$$g : (0, 1) \rightarrow (1/3, 2/3)$$

by expanding their respective codomains. Thus we've got the desired functions, and previous point shows that given sets aren't homeomorphic, as desired.

(c) *Show that R^n and R aren't homeomorphic if $n > 1$.*

Since R^n is Hausdorff, we follow that $\{0\}$ is open and thus punctured euclidian space $R^n \setminus \{0\}$ is connected, which means that if there's a bijection $h : R^n \rightarrow R$, then the set $h[R^n \setminus \{0\}]$ is always disconnected, which means that no bijection between those two sets can be continuous.

3.2.2

Let $f : S^1 \rightarrow R$ be a continuous map. Show there exists a point x of S^1 such that $f(x) = f(-x)$.

Firstly, we want to assume that $f(-x)$ means that we apply f to the x 's additive inverse in vector notation, because no other meaning makes sense. Since scalar multiplication of vectors is continuous, we follow that

$$f(-x) = (f \circ T)(x)$$

where $T(x) = -x$ is continuous as well. Thus we follow that we can define $g : S \rightarrow R$ by

$$g(x) = f(x) - f(-x)$$

is a composition of continuous operator $-$ and two continuous functions, and thus g itself is continuous. We also follow that

$$g(-x) = f(-x) - f(-(-x)) = f(-x) - f(x) = -g(x)$$

Now let $s \in S$. If $g(s) = 0$, then

$$f(x) - f(-x) = 0 \Leftrightarrow f(x) = f(-x)$$

and we're done. If $g(s) > 0$, then we follow that

$$g(-s) = -g(s) < 0$$

therefore IVT implies together with the connectedness of S imply that there's a point $s' \in S$ such that $g(s') = 0$. If $s < 0$, then we can set $s' = -s$ and default to the previous case. Therefore we derive the desired result.

3.2.3

Let $f : X \rightarrow X$ be continuous. Show that if $X = [0, 1]$, there is a point x such that $f(x) = x$. The point x is called a fixed point of f . What happens if X equals $[0, 1)$ or $(0, 1)$?

Set $g : X \rightarrow R$ by

$$g(x) = f(x) - x$$

since the identity function, $f(x)$ and $- : R \times R \rightarrow R$ are all continuous functions, we follow that g is a continuous function as well. We can also argue that condition that $f(x) = x$ is equivalent to condition $g(x) = 0$; thus we need to show that there's a $x \in X$ such that $g(x) = 0$.

We can follow that

$$g(0) = f(0) - 0 = f(0)$$

and thus $g(0) \in X$. If $g(0) = 0$, then we're done, thus assume that $g(0) \neq 0$. We then conclude that $g(0) \in (0, 1]$.

Let us consider also $g(1)$. We follow that

$$g(1) = f(1) - 1 \in [-1, 0]$$

We follow that if $g(x) = 0$, then we're done, and thus $g(x) \in [-1, 0)$.

Now we can combine two arguments to follow that $g(0) > 0$ and $g(1) < 0$, which means that there's $x \in [0, 1]$ such that $g(x) = 0$, as desired.

We can also follow that we can define function $h : \mathbb{R} \rightarrow \mathbb{R}$

$$h(x) = \frac{x+1}{2}$$

we follow that we can restrict domain of h to either $[0, 1)$ or $(0, 1)$ and in those cases we'll have that

$$\text{ran}(h|_{[0, 1)}) \subseteq [0, 1)$$

$$\text{ran}(h|_{(0, 1)}) \subseteq (0, 1)$$

Some trivial analysis can show us that $h(x) \neq x$ for all $[0, 1)$. Thus we conclude that if we restrict domain and codomain to presented values, then we can't say anything about the existence of the fixed point.

3.2.4

Let X be an ordered set in the order topology. Show that if X is connected, then X is a linear continuum.

Suppose that X is not a linear continuum. If the first condition does not hold, then we follow that there are $x, y \in X$ such that $x < y$ and there's no $z \in X$ such that $x < z < y$. Thus we follow that $(-\infty, y)$ is an open set, as well as (x, ∞) , and they constitute separation of X , which is a contradiction.

Suppose that X does not have a least upper boundary property. Thus we follow that there's a subset U of X such that U is bounded above and does not have a least upper bound. Define

$$V = \bigcup_{u \in U} (-\infty, u)$$

Since U is bounded, we follow that $V \neq X$. Let $i \in X \setminus V$. We follow that i is an upper bound, but because there's no least upper bound, we follow that there's $i' \in X$ such that $i' < i$ and i' is also an upper bound for V . AC implies that we've got a function on our hands, and thus we follow that

$$\bigcup_{i \in X \setminus V} (f(i), \infty)$$

is an open set, that is disjoint from V , and is equal to $X \setminus V$. Since U is nonempty, we follow that V is also nonempty, and thus V is both open and closed, which implies that X is disconnected, which is a contradiction.

Therefore we follow that if X is connected, then it satisfies both conditions of linear continuum, as desired.

3.2.5

Consider the following sets in the dictionary order. Which are linear continua?

(a) $Z_+ \times [0, 1)$

Z_+ is a woset, and thus the next exercise will imply that the desired set is linear continuum. The intuitive conclusion is that this thing is homeomorphic to R .

(b) $[0, 1) \times Z_+$

We follow that there's no elements between $\langle 0, 1 \rangle$ and $\langle 0, 2 \rangle$ on the account that Z_+ does not hold neither of the preproperties of linear continuum.

(c) $[0, 1) \times [0, 1]$

We can follow that $[0, 1) \times [0, 1] = \{i \in I^2 : i < \langle 1, 0 \rangle\}$, thus we follow that given set is convex, and thus by sub-theorem of 24.1 we conclude that it is connected. Since it is a connected set in order topology, we follow that it is a linear continuum.

(d) $[0, 1] \times [0, 1)$

We follow that the subset

$$U = \{\langle 0, 1/n \rangle : n \in Z_+\}$$

does not have a supremum, and thus we conclude that given set cannot be a linear continuum.

3.2.6

Show that if X is a well-ordered set, then $X \times [0, 1)$ in the dictionary order is a linear continuum.

Let $x, y \in X \times [0, 1)$ be such that $x < y$. We follow that

$$x = \langle a, b \rangle$$

$$y = \langle c, d \rangle$$

If $a = c$, then we follow that there's $q \in [0, 1)$ such that $b < q < d$, and thus

$$x < \langle a, q \rangle < y$$

If $a < c$, then we follow that there are two cases: $c \neq a^+$ or $c = a^+$. If former is the case, then we follow that

$$x < \langle a^+, 0 \rangle < y$$

if the latter is the case, then we follow that there's $q \in (b, 1)$ such that

$$x < \langle a, q \rangle < y$$

thus we conclude that $x < y$ implies that there's $z \in X \times [0, 1)$ such that

$$x < z < y$$

which proves the second requirement of the linear continuum.

Now let U be a bounded set of $X \times [0, 1)$. We can set

$$s = \sup\{x \in X : \exists y \in [0, 1) : \langle x, y \rangle \in U\}$$

i.e. the supremum of the first parts of elements of a given set. Now define

$$V = \{u \in U : \exists y \in [0, 1) : \langle s, y \rangle \in U\}$$

i.e. set of elements of with the supremum in the first part.

Now we can define an element j by cases on V . If $V = \emptyset$, then we can set $j = \langle s, 0 \rangle$. Construction of j implies that j is indeed an upper bound, and we follow that if $k < j$, then there's $l \in X, q \in [0, 1)$ such that $k = \langle l, q \rangle$ and thus $l < s$. Now construction of s implies that k is not an upper bound. Thus we can conclude that j is indeed a least upper bound.

Let y be equal to the supremum of the second parts of V . If $y \neq 1$, then we set $j = \langle s, y \rangle$, and if $y = 1$, then we can set $j = \{s^+, 0\}$. Some trivial logic shows that j is indeed the least upper bound for U .

Thus we conclude that $X \times [0, 1)$ is indeed a linear continuum, as desired.

3.2.7

(a) Let X be ordered sets in the order topology. Show that if $f : X \rightarrow Y$ is order preserving and surjective, then f is a homeomorphism.

We firstly note that X is a toset (implied by the fact that X is in the order topology, which was defined so far exclusively on tosets), and thus it's got a trichotomy. Thus we follow that if $x \neq y$, then we can assume that $x < y$, thus $f(x) < f(y)$, and thus f is injective. Now surjectivity of f with respect to the chosen codomain makes it a bijection.

Now let $a, b \in Y$. We follow that

$$\forall x \in X : x \in f^{-1}[(a, b)] \Leftrightarrow f^{-1}(a) < x < f^{-1}(b)$$

(all of this stuff can be derived pretty easily). Thus we follow that if U is a basis oin Y , then $f^{-1}[U]$ is open, thus making f continous. Check that f^{-1} is continous is well is the same as this one. Thus we conclude that f is a homeomorphism, as desired.

(b) Let $X = Y = \overline{R_+}$. Given a positive inveger n , show that the function $f(x) = x^n$ is order preserving and surjective. Conclude that its inverse, the n th root function is continous.

We follow that if $n = 1$, then f is an identity, thus it's order preserving and surjective. Suppose that $f(x) = x^{n-1}$ is order preserving and surjective. Let $z, y \in \overline{R_+}$ are such that $z < y$. We follow that z, y, z^j, y^j are all nonnegative for all $j \in \omega$. Thus we follow that

$$z < y \iff f(z) < f(y) \iff z^{n-1} < y^{n-1}$$

now $z < y$ and $z^{n-1} < y^{n-1}$ and nonnegativity of those things imply that

$$z * z^{n-1} < y * y^{n-1}$$

$$z^n < y^n$$

given that z and y are arbitrary, we conclude that $f(x) = x^n$ is order preserving. Thus we conclude that

$$f(x) = x^n$$

is order preserving for all $n \in \omega$ by induction.

Now by induction we can also prove that f is continuous and nonnegative. Thus we can follow that IVT implies that f is surjective. Therefore we conclude that f is order-preserving and surjective, which means that it's a homeomorphism. Thus we conclude that f^{-1} is indeed continuous.

(c) *blah blah blah*

Yes, it is, but not with respect to the subset topology, but with respect to the order topology defined on $(-\infty, 1) \cup [0, \infty)$. goto page 90 of original book for examples and explanation

3.2.8

(a) *Is a product of path-connected spaces necessarily path connected?*

It seems to be so. Let X and Y be path-connected spaces. Now let $q, w \in X \times Y$. We follow that

$$q = \langle a, b \rangle$$

$$w = \langle c, d \rangle$$

We now follow that there exist continuous functions $f : [x_1, x_2] \rightarrow X$ and $g : [y_1, y_2] \rightarrow Y$ such that

$$f(x_1) = a, f(x_2) = c, g(y_1) = b, g(y_2) = d$$

We can follow that spaces X and $X \times \{b\}$ with respect to the subset topology of a product topology in $X \times Y$ are homeomorphic. Thus we follow that we can define continuous $f' : [x_1, x_2] \rightarrow X \times Y$ by

$$f'(x) = \langle f(x), b \rangle$$

and $g' : [x_2, (y_2 - y_1) + x_2] \rightarrow X \times Y$ by

$$g'(y) = \langle c, g(y - x_2 + y_1) \rangle$$

We can follow that $f'(x_2) = \langle f(x_2), b \rangle = \langle c, b \rangle = g'(x_2)$. Thus we follow that we can concatenate those two functions to get the desired path.

By induction we can follow that finite product of path-connected spaces is path connected.

(b) If $A \subseteq X$ and A is path connected, is \overline{A} necessarily path connected?

Let S be the topologist's sine curve. If $x, y \in S$, then we follow that obviously there is a continuous function such that everything connects, but \overline{S} is not path connected, and thus we've got the contradiction.

(c) If $f : X \rightarrow Y$ is continuous and X is path connected, is $f[X]$ necessarily path connected?

Suppose that $y_1, y_2 \in f[X]$. We follow that there are $x_1, x_2 \in X$ (maybe not unique, but still) such that $f(x_1) = y_1, f(x_2) = y_2$. Since X is path connected, we follow that there's a continuous $g : [a, b] \rightarrow X$ such that $g(a) = x_1$ and $g(b) = x_2$. Since f and g are both continuous, we follow that $f \circ g$ is also continuous, and thus we've got the desired path.

(d) If $\bigcup A_\alpha$ is a collection of path-connected subspaces of X and if $\bigcap A_\alpha \neq \emptyset$, is $\bigcup A_\alpha$ necessarily path-connected?

Suppose that $x, y \in \bigcup A_\alpha$. We follow that there's $q \in \bigcap A_\alpha$. Thus we follow that there's a path between x and q and a path between q and y , which means that there's a concatenated path between x and y . (Although it feels like a somewhat liberal proof, we can follow all the rigorous stuff pretty easily)

3.2.9

Assume that R is uncountable. Show that if A is a countable subset of R^2 , then $R^2 \setminus A$ is path connected.

Let $x, y \in R^2$. We follow that there are uncountable amount of lines that pass through each one of those points. We can also follow that there's no point $a \in A$ such that there are two lines from a single point, that intersect a . Thus we follow that the set of lines from x that does not intersect A is uncountable. Same goes for y . We can follow now that there are uncountably many unparallel pairs of lines that go through the x and y , which gives us the desired path.

3.2.10

Show that if U is an open connected subspace of R^2 , then U is path connected.

Let $x_0 \in U$ and let A denote the set of points in U , from which there's a path to x_0 . We follow that A is a subset of U , and we also follow that it's nonempty, since there's always a path from x_0 to x_0 .

Let $y \in A$. Since $y \in U$ and U is open, we follow that there's a basis element B inside U such that $y \in B$. Since y is quite literally the center of the basis element B , we follow that there's a path to any given element of B from y . Thus we follow that $B \subseteq A$, which allows us to conclude that A is open.

Suppose that $l \in U$ is a limit point of A . We follow that there's a basis neighborhood around l that intersects A and lies fully within U . Thus we conclude that once again we can connect l to some point inside A , and thus we conclude that $l \in A$. Therefore A contains

all of its limit points, that are in U , and thus we conclude that A is both open and closed inside subset topology of U . Thus we conclude that $A = \emptyset$ or $A = U$, where former is impossible, as we've shown before. Thus we conclude that U is path connected, as desired.

3.2.11

If A is a connected subspace of X , does it follow that $\text{Int}(A)$ and $\text{Bd}(A)$ are connected?

We can define a set

$$\{\langle x, y \rangle \in \mathbb{R}^2 : 0 \leq y \leq x \mid 0 \geq y \geq x\}$$

(i.e. two triangular regions, that meet at the origin). Then we follow that the origin is not in the interior of A , since no basis neighborhood of A is contained within A . But A in general is connected, which is trivially proven. Thus we conclude that A being connected does not imply anything about connectedness of its interior.

We can also punch two holes through \mathbb{R}^2 to get a boundary that will consist of two points in \mathbb{R}^2 , which will obviously be disconnected. Thus we conclude that connectedness of A does not imply anything about connectedness of $\text{Int}(A)$ or $\text{Bd}(A)$.

Suppose that both $\text{Int}(A)$ and $\text{Bd}(A)$ are connected. We follow that $\text{Int}(A)$ or $\text{Bd}(A)$ can be both open and closed, which would imply that A is disconnected. It can be connected as well. So we can't conclude much from it.

The last exercise is left for later

3.3 Compact Spaces

3.3.1

(a) Let \mathcal{T} and \mathcal{T}' be two topologies on the set X ; suppose that $\mathcal{T} \subseteq \mathcal{T}'$. What does compactness of X under one of these topologies imply about the other?

We follow that if X is compact under \mathcal{T}' , and U is a collection of open sets from \mathcal{T} , that form open cover for X , we can conclude that U is an open cover in \mathcal{T}' as well, thus there's a finite subcover, and thus we follow that X is compact in \mathcal{T}' as well. If X is compact only in \mathcal{T} , then we can't conclude nothing. Example: standard and discrete topologies on \mathbb{R} .

(b) Show that if X is compact Hausdorff under both \mathcal{T} and \mathcal{T}' , then either \mathcal{T} or \mathcal{T}' are equal or they are not compatible.

We essentially want to prove that \mathcal{T} cannot be a proper subset of \mathcal{T}' and vice versa. We follow that they can be equal (duh), so assume that $\mathcal{T} \subset \mathcal{T}'$.

We know that X is a compact space under \mathcal{T}' . Let U be an open set of \mathcal{T}' such that $U \notin \mathcal{T}$. We can state now that $U \neq \emptyset$, since $U \notin \mathcal{T}$ and $\emptyset \in \mathcal{T}$ by definition.

We can follow that $X \setminus U$ is a closed set. Since $U \subseteq X$, we follow that $X \setminus U \subseteq X$. Thus $X \setminus U$ is a closed subspace of X , and since X is compact we follow that $X \setminus U$ is compact. Since $U \notin \mathcal{T}$, we follow that $X \setminus U$ is not closed in X with respect to \mathcal{T} , and thus

we follow that there's an open cover C for $X \setminus U$ in \mathcal{T} , that does not have a finite subcover for $X \setminus U$. Since $C \subseteq \mathcal{T}$, we follow that $C \subseteq \mathcal{T}'$, which means that C is an open cover for $X \setminus U$ for which there's no finite subcover, which implies that $X \setminus U$ is not compact, which is a contradiction.

3.3.2

(a) *Show that in the finite complement topology on R , every subspace is compact.*

Let $U \subseteq R$ and let \mathcal{V} be its open cover.