

# My real analysis exercises

Evgeny Markin

2022

Exercises are from UTM-040 Understanding analysis by Stephen Abbott. Edition is unknown, but the date in the preface is August 2000. I don't own any rights for the book and I am not planning to acquire them anytime soon.

Also it wouldn't hurt to mention, that some (or maybe even most) of exercises were not proof-read after they were written, and that none of them were evaluated by anyone who matters. There are a lot of mistakes out there, both in mathematical and logical parts, and in the language they were written as well. I am doing this kind of stuff for fun, so cut me some slack.

### 1.2.1

(a) *Prove that  $\sqrt{3}$  is irrational. Does a similar argument work to show  $\sqrt{6}$  is irrational?*  
Suppose that  $\sqrt{3}$  is a rational number; then it is true that

$$\exists m \in \mathbf{Z}, n \in \mathbf{N} : \frac{m}{n} = \sqrt{3}$$

where  $m$  and  $n$  are at their lowest possible terms. Then

$$\sqrt{3}n = m$$

$$3n^2 = m^2$$

Therefore we can state, that  $m \% 3 = 0$ . Therefore  $\exists k : 3k = m$ . Thus we can reformulate formula as

$$3n^2 = (3k)^2$$

$$n^2 = 3k^2$$

Therefore  $n \% 3 = 0$  as well. Therefore  $n$  and  $k$  are not in their possible terms, which contradicts our initial assumptions. Therefore we can state that  $\sqrt{3} \notin \mathbf{Q}$ .

Let's try the same argument for  $\sqrt{6}$ .

$$\exists m \in \mathbf{Z}, n \in \mathbf{N} : \frac{m}{n} = \sqrt{6}$$

$$\sqrt{6}n = m$$

$$6n^2 = m^2$$

then  $m$  has as their dividers both 2 and 3. Therefore  $m \% 2 = 0$  and  $m \% 3 = 0$ . Therefore we can proceed with the same argument as earlier

$$6n^2 = (6k)^2$$

$$n^2 = 6k^2$$

Therefore  $n$  is divided by 6, etc., etc.,  $\sqrt{6} \notin \mathbf{Q}$ .

(b) Where does the proof of Theorem 1.1.1 break down if we try to use it to prove  $\sqrt{4}$  is irrational?

Suppose that  $\sqrt{4}$  is a rational number; then it is true that

$$\exists m \in \mathbf{Z}, n \in \mathbf{N} : \frac{m}{n} = \sqrt{2}$$

where  $m$  and  $n$  are at their lowest possible terms. Then

$$\sqrt{4n} = m$$

$$4n^2 = m^2$$

$n$  can still be odd and  $m$  can still be even. In other words,  $m$  is divisible by a prime, and the number under the radical consists of two primes. Therefore if a number decomposes to two equal sets of primes, then its square root is a rational number. Otherwise it isn't.

## 1.2.2

Decide which of the following represent true statements about the nature of sets. For any that are false, provide a specific example where the statement in question does not hold.

(a) if  $A_1 \supseteq A_2 \supseteq A_3 \supseteq \dots$  are all sets containing an infinite number of elements, then the intersection  $\cap_{n=1}^{\infty} A_n$  is infinite as well. -

$\cap_{n=1}^{\infty} A_n = (0, 1/n)$  has no numbers in it.

Proof is easy -

$$\forall x \in \mathbf{R} > 0 : \exists n \in \mathbf{N} : 1/n < x$$

(b) if  $A_1 \supseteq A_2 \supseteq A_3 \supseteq \dots$  are all finite, nonempty sets of real numbers, then the intersection  $\cap_{n=1}^{\infty} A_n$  is finite and nonempty. -

True.

There is no need for the proof, but I'll supply one anyways. If all  $A_n$  are finite and nonempty, then  $\exists j \in \mathbf{N} : |A_1| = j$ . Therefore, because of the same reasons, there are only  $j - 1$  times when

$$A_k \supset A_{k+1}$$

can happen, because after  $j - 1$  times the set will be empty. Therefore, because it is finite, their intersection will have finite number of elements and will be non-empty.

(c)  $A \cap (B \cup C) = (A \cap B) \cup C$

False: let

$$x \notin A, x \notin B, x \in C$$

Then

$$x \in A \cap (B \cup C); x \notin (A \cap B) \cup C$$

$$(c) A \cap (B \cap C) = (A \cap B) \cap C$$

True. Kinda goes without a proof; if you imagine a Vien diagram, then it's obvious.

$$(c) A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

True. For the same reason as before.

I'm sure that there exist more concrete versions of those proofs, but I'm not required to provide any. My suspicion on why is it so, is because it's a little more complicated and requires more knowlege in set theory and/or logic.

### 1.2.3 (De Morgan's Laws).

Let  $A$  and  $B$  be subsets of  $\mathbf{R}$

(a) If  $x \in (A \cap B)^c$ , explain why  $x \in A^c \cup B^c$ . This shows that  $(A \cap B)^c \subseteq (A \cap B)^c$ .

If we have two sets  $A$  and  $B$ , then  $\mathbf{R}$  desintegrates into 4 different sets:  $A$ ,  $B$ ,  $A^c$ ,  $B^c$ .

Therefore there must exists sets

$$S_1 = A \cap B$$

$$S_2 = A^c \cap B$$

$$S_3 = A \cap B^c$$

$$S_4 = A^c \cap B^c$$

An element cannot be in the set and not in the set at the same time. Therefore, there does not exist an element, which is in two of  $S_n$ 's.

For any  $x \in \mathbf{R} \rightarrow x \in A$  or  $x \notin A$ . Therefore an element of  $\mathbf{R}$  needs to be in at least one of those sets. It is easily seen by

$$A \cap \mathbf{R} = A$$

$$A \cap (B \cup B^c) = A$$

$$(A \cap B) \cup (A \cap B^c) = A$$

Therefore  $\cup_{n=1}^4 S_n = \mathbf{R}$  and  $\cap_{n=1}^4 S_n = \emptyset$ .

Suppose  $x \in (A \cap B)^c$ . Then  $x \notin A \cap B$ . Therefore  $x \in S_2 \cup S_3 \cup S_4$ .

Suppose that  $x \in A^c \cup B^c$ . Then  $x \in S_2 \cup S_3 \cup S_4$ .

Therefore  $(A \cap B)^c \subseteq A^c \cup B^c$ .

(b) Prove the reverse inclusion

As seen in part (a)

$$(A \cap B)^c = S_2 \cup S_3 \cup S_4 = A^c \cup B^c$$

(c) Show  $(A \cup B)^c = A^c \cap B^c$  by demonstrating inclusion both ways.  
No need to do both ways.

$$(A \cup B)^c = S_4 = A^c \cap B^c$$

## 1.2.4

Verify the triangle inequality in the special cases where

(a)  $a$  and  $b$  have the same sign

Suppose  $a \geq 0$ ,  $b \geq 0$ . Then  $|a| = a$  and  $|b| = b$ . Therefore

$$|a + b| = a + b = |a| + |b| \leq |a| + |b|$$

Suppose  $a < 0$ ,  $b < 0$ . Then  $|a| = -a$  and  $|b| = -b$ ; also  $a + b < 0 \rightarrow |a + b| = -(a + b) = -a - b$ . Therefore

$$|a + b| = -a + (-b) = |a| + |b| \leq |a| + |b|$$

.

(b)  $a \geq 0$ ,  $b < 0$  and  $a + b \geq 0$ .

$$a + b \geq 0 \rightarrow a + b = |a + b|$$

Also,  $|a| = a$  and  $|b| = -b$ . Therefore

$$a + b \geq 0 \rightarrow a \geq -b \rightarrow a \geq |b| \rightarrow |a| \geq |b|$$

$$b < 0$$

$$b \leq 0$$

$$2b \leq 0$$

$$b + b \leq 0$$

$$b \leq (-b)$$

$$a + b \leq a + (-b)$$

$$|a + b| \leq |a| + |b|$$

## 1.2.5

Use the triangle inequality to establish the inequalities

$$(a) |a - b| \leq |a| + |b|;$$

$$|a - b| = |a + (-b)| \leq |a| + |-b| = |a| + |b|$$

$$(b) ||a| - |b|| \leq |a - b|;$$

let  $a = a + b - b$ . Then

$$|a| = |a - b + b| \leq |a - b| + |b|$$

$$|a| - |b| \leq |a - b|$$

$$|b| = |b - a + a| \leq |b - a| + |a| = |a - b| + |a|$$

$$|b| - |a| \leq |a - b|$$

$$|a| - |b| \geq -|a - b|$$

$$-|a - b| \leq |a| - |b| \leq |a - b| \rightarrow ||a| - |b|| \leq |a - b|$$

## 1.2.6

Given a function  $f$  and a subset  $A$  of its domain, let  $f(A)$  represent the range of  $f$  over the set  $A$ ; that is,  $f(A) = \{f(x) : x \in A\}$ .

(a) Let  $f(x) = x^2$ . if  $A = [0, 2]$  (the closed interval  $\{x \in \mathbf{R} : 0 \leq x \leq 2\}$ ) and  $B = [1, 4]$ , find  $f(A)$  and  $f(B)$ . Does  $f(A \cap B) = f(A) \cap f(B)$  in this case? Does  $f(A \cup B) = f(A) \cup f(B)$ ?

First things first:  $f(A) = [0, 4]$ ;  $f(B) = [1, 16]$  (without any proof because if we don't go with axiomatic stuff, then it is obvious).

$$f(A \cap B) = f([1, 2]) = [1, 4]$$

$$f(A) \cap f(B) = [1, 4]$$

Therefore in this case  $f(A) \cap f(B) = f(A \cap B)$ .

$$f(A \cup B) = f([0, 4]) = [0, 16] = f(A) \cup f(B)$$

(b) Find two sets  $A$  and  $B$  for which  $f(A \cap B) \neq f(A) \cap f(B)$ .

Let  $A = [-1, 0]$  and  $B = [0, 1]$ . Then

$$f(A \cap B) = f(\{0\}) = \{0\}$$

$$f(A) \cap f(B) = [0, 1] \cap [0, 1] = [0, 1] \neq f(A \cap B)$$

(c) Show that, for an arbitrary function  $g : \mathbf{R} \rightarrow \mathbf{R}$ , it is always true that  $g(A \cap B) \subseteq g(A) \cap g(B)$  for all sets  $A, B \subseteq \mathbf{R}$ .

$$x \in g(A \cap B) \rightarrow x \in g(A)$$

$$x \in g(A \cap B) \rightarrow x \in g(B)$$

Therefore

$$x \in g(A \cap B) \rightarrow x \in g(A) \cap g(B)$$

Thus

$$g(A \cap B) \subseteq g(A) \cap g(B)$$

(d) Form and prove a conjecture about the relationship between  $g(A \cup B)$  and  $g(A) \cup g(B)$  for an arbitrary function  $g$ .

$$x \in g(A) \rightarrow x \in g(A \cup B)$$

$$x \in g(B) \rightarrow x \in g(A \cup B)$$

Therefore

$$x \in g(A) \cup g(B) \rightarrow x \in g(A \cup B)$$

Thus

$$g(A) \cup g(B) \subseteq g(A \cup B)$$

Suppose that

$$\exists y \in \mathbf{R} : y \in g(A \cup B); y \notin g(A) \cup g(B)$$

Then  $\exists q_1 \in A \cup B : g(q_1) = y$  but

$$\forall q_2 \in A, q_3 \in B : g(q_2) \neq y; g(q_3) \neq y$$

Therefore  $q_1 \notin A$  and  $q_1 \notin B$ . Therefore  $q_1 \in A^c \cap B^c$ . Using De Morgan rule

$$q_1 \in A^c \cap B^c \rightarrow q_1 \in (A \cup B)^c$$

therefore

$$q_1 \notin g(A \cup B)$$

which is a contradiction. Therefore

$$y \in g(A \cup B) \rightarrow y \in g(A) \cup g(B)$$

Thus

$$g(A \cup B) \subseteq g(A) \cup g(B)$$

Therefore if we take into account previous conclusion

$$g(A \cup B) = g(A) \cup g(B)$$

for any  $g$ .

### 1.2.7

Given a function  $f : D \rightarrow \mathbf{R}$  and a subset  $B \subseteq \mathbf{R}$ , let  $f^{-1}(B)$  be the set of all points from the domain  $D$  that get mapped into  $B$ ; that is,  $f^{-1}(B) = \{x \in D : f(x) \in B\}$ . This is called the preimage of  $B$ .

(a) Let  $f(x) = x^2$ . If  $A$  is the closed interval  $[0, 4]$  and  $B$  is the closed interval  $[-1, 1]$ , find  $f^{-1}(A)$  and  $f^{-1}(B)$ . Does  $f^{-1}(A \cap B) = f^{-1}(A) \cap f^{-1}(B)$  in this case? Does  $f^{-1}(A \cup B) = f^{-1}(A) \cup f^{-1}(B)$ ?

$$f^{-1}(A) = [-2, 2]$$

$$f^{-1}(B) = [-1, 1]$$

$$f^{-1}(A \cap B) = f^{-1}([0, 1]) = [-1, 1] = f^{-1}(A) \cap f^{-1}(B)$$

$$f^{-1}(A \cup B) = f^{-1}([-1, 4]) = [-2, 2] = f^{-1}(A) \cup f^{-1}(B)$$

(b) The good behaviour of preimages demonstrated in (a) is completely general. Show that for an arbitrary function  $g : \mathbf{R} \rightarrow \mathbf{R}$ , it is always true that  $g^{-1}(A \cap B) = g^{-1}(A) \cap g^{-1}(B)$  and  $g^{-1}(A \cup B) = g^{-1}(A) \cup g^{-1}(B)$  for all sets  $A, B \subseteq \mathbf{R}$ .

By definition

$$x \in g^{-1}(A \cap B) \rightarrow \exists y \in A \cap B : y = g(x)$$

Therefore if we use the fact  $y \in A \cap B \rightarrow y \in A$  and  $y \in A \cap B \rightarrow y \in B$

$$x \in g^{-1}(A \cap B) \rightarrow \exists y \in A : y = g(x) \rightarrow x \in g^{-1}(A)$$

$$x \in g^{-1}(A \cap B) \rightarrow \exists y \in B : y = g(x) \rightarrow x \in g^{-1}(B)$$

therefore  $x \in g^{-1}(A \cap B)$  implies that  $x \in g^{-1}(A)$  and  $x \in g^{-1}(B)$ , or in other words

$$g^{-1}(A \cap B) \subseteq g^{-1}(A) \cap g^{-1}(B)$$

In other direction:

$$x \in g^{-1}(A) \rightarrow \exists y_1 \in A : y_1 = g(x)$$

$$x \in g^{-1}(B) \rightarrow \exists y_2 \in B : y_2 = g(x)$$

$x \in g^{-1}(A) \cap g^{-1}(B)$  implies that  $\exists y_1 \in A : g(x) = y_1$  and  $\exists y_2 \in B : y_2 = g(x)$ . Because  $g$  is a function we know, that for every  $x$  there exists only one  $y = g(x)$ . Therefore  $y_1 = y_2 = g(x)$ . Thus we can state that  $y \in A \cap B$ .

thus

$$x \in g^{-1}(A) \cap g^{-1}(B) \rightarrow \exists y \in A \cap B : y = g(x) \rightarrow x \in g^{-1}(A \cap B)$$

Therefore



$$g^{-1}(A) \cap g^{-1}(B) \subseteq g^{-1}(A \cap B)$$

If we take previous conclusion into account, then it follows that

$$g^{-1}(A) \cap g^{-1}(B) = g^{-1}(A \cap B)$$

as desired.

Now let's prove that  $g^{-1}(A \cup B) = g^{-1}(A) \cup g^{-1}(B)$ :

If  $x \in g^{-1}(A) \cup g^{-1}(B)$  then  $\exists y \in A : y = g(x)$  or  $\exists y \in B : y = g(x)$ . If we take into account that  $y \in A \rightarrow y \in A \cup B$  then we can conclude that

$$x \in g^{-1}(A) \cup g^{-1}(B) \rightarrow \exists y \in A \cup B : y = g(x) \rightarrow x \in g^{-1}(A \cup B)$$

Thus

$$g^{-1}(A) \cup g^{-1}(B) \subseteq g^{-1}(A \cup B)$$

In other direction:

$$x \in g^{-1}(A \cup B) \rightarrow \exists y \in A \cup B : y = g(x) \rightarrow y = g(x)$$

As proven before  $g(A \cup B) = g(A) \cup g(B)$  and therefore  $y \in g(A \cup B)$  implies that either  $y \in g(A)$  or  $y \in g(B)$ . Therefore

$$x \in g^{-1}(A \cup B) \rightarrow x \in g^{-1}(A) \cup g^{-1}(B)$$

$$g^{-1}(A \cup B) \subseteq g^{-1}(A) \cup g^{-1}(B)$$

And if we combine this fact with previous conclusion:

$$g^{-1}(A \cup B) = g^{-1}(A) \cup g^{-1}(B)$$

as desired.

### 1.2.8

*Form the logical negation of each claim. One way to do this is to simply add "It is not the case that ..." in front of each assertion, but for each statement, try to embed the word "not" as deeply into the resulting sentence as possible (or avoid using it altogether).*

(a) *For all real numbers satisfying  $a < b$ , there exist an  $n \in \mathbf{N}$  such that  $a + 1/n < b$ .*  
 There exist real numbers  $a < b$  such that  $a + 1/n \geq b$  for all  $n \in \mathbf{N}$ .

(b) *Between every two distinct real numbers, there is a rational number*

There exist two real numbers, such that there are only irrational numbers between them.

(c) For all natural numbers  $n \in \mathbf{N}$ ,  $\sqrt{n}$  is either a natural number or an irrational number

There exist a natural number  $n \in \mathbf{N}$ , such that  $\sqrt{n}$  is a rational number, that is not a natural number. (This one is a little bit weird if we try to negate this, but "not" is stuffed as deep as possible)

(d) Given any real number  $x \in \mathbf{R}$ , there exist  $n \in \mathbf{N}$  satisfying  $n > x$

There exist a real number  $x \in \mathbf{R}$ , such that for all  $n \in \mathbf{N}$  it is true that  $n \leq x$ .

## 1.2.9

Show that the sequence  $(x_1, x_2, x_3, \dots)$  defined in Example 1.2.7 is bounded above by 2; that is, prove that  $x_n \leq 2$  for every  $n \in \mathbf{N}$ .

The mentioned sequence is defined by

$$x_1 = 1$$

$$x_{n+1} = (1/2)x_n + 1$$

A great advice about induction states, that when you hear words "prove" and "sequence" in the same sentence, then the word "induction" should pop up in your head. So here we go

Base case:  $x_1 \leq 2$ .

Inductive proposition:  $x_n \leq 2$

Inductive step:

$$x_n \leq 2$$

$$(1/2)x_n \leq 1$$

$$(1/2)x_n + 1 \leq 2$$

$$x_{n+1} = (1/2)x_n + 1 \leq 2$$

$$x_{n+1} \leq 2$$

as desired.

## 1.2.10

Let  $y_1 = 1$ , and for each  $n \in \mathbf{N}$  define  $y_{n+1} = (3 * y_n + 4)/4$ .

(a) Use induction to prove that the sequence satisfies  $y_n < 4$  for all  $n \in \mathbf{N}$ .

Base case:  $y_1 = 1 < 4$

Inductive proposition:  $y_n < 4$

Inductive step:

$$y_n < 4$$

$$3 * y_n < 12$$

$$3 * y_n + 4 < 16$$

$$(3 * y_n + 4)/4 < 4$$

$$y_{n+1} = (3 * y_n + 4)/4 < 4$$

$$y_{n+1} < 4$$

as desired.

(b) Use another induction to show the sequence  $(y_1, y_2, y_3, \dots)$  is increasing

We need to show that  $y_{n+1} - y_n \geq 0$

As shown earlier

$$y_n < 4$$

therefore

$$y_n < 4$$

$$\frac{y_n}{4} < 1$$

$$1 > \frac{y_n}{4}$$

$$1 - \frac{y_n}{4} > 0$$

$$1 + \left(\frac{3y_n}{4} - y_n\right) > 0$$

$$\frac{3y_n}{4} + 1 - y_n > 0$$

$$(3 * y_n + 4)/4 - y_n > 0$$

$$y_{n+1} - y_n > 0$$

as desired.

### 1.2.11

If a set  $A$  contains  $n$  elements, prove that the number of different subsets of  $A$  is equal to  $2^n$ . (Keep in mind that the empty set  $\emptyset$  is considered to be a subset of every set.)

This proof is dumb, but intuitive:

Every subset is corresponding to a number in binary system: 0 for excluded, 1 for included. Therefore there exist  $2^n$  possible combinations.

For a more concrete proof let's resort to induction.

Base case(s): subsets of  $\emptyset$  are  $\emptyset$  itself ( $2^0 = 1$  in total). Subsets of set with one element are  $\emptyset$  and set itself ( $2^1 = 2$  in total).

Proposition is that set with  $n$  elements has  $2^n$  subsets.

Inductive step is that for set with  $n+1$  elements can either have or not have the  $n+1$ 'th element. Therefore there exist  $2^n + 2^n = 2 * 2^n = 2^{n+1}$  subsets, as desired.

### 1.2.12

For this exercise, assume Exercise 1.2.3 has been successfully completed (as it was)

(a) Show how induction can be used to conclude that

$$(A_1 \cup A_2 \cup \dots \cup A_n)^c = A_1^c \cap A_2^c \cap \dots \cap A_n^c$$

First of all, base case

$$(A_1 \cup A_2)^c = A_1^c \cap A_2^c$$

Proposition

$$(A_1 \cup A_2 \cup \dots \cup A_n)^c = A_1^c \cap A_2^c \cap \dots \cap A_n^c$$

Step

$$(A_1 \cup A_2 \cup \dots \cup A_n \cup A_{n+1})^c = A_1^c \cap A_2^c \cap \dots \cap A_n^c \cap A_{n+1}^c$$

Let us denote  $Q = A_1^c \cap A_2^c \cap \dots \cap A_n^c$ . Then by inductive proposition

$$A_1^c \cap A_2^c \cap \dots \cap A_n^c \cap A_{n+1}^c = Q \cap A_{n+1}^c = (Q \cup A_{n+1})^c = (A_1 \cup A_2 \cup \dots \cup A_n \cup A_{n+1})^c$$

as desired.

(b) Explain why induction cannot be used to conclude

$$(\cup_{n=1}^{\infty} A_n)^c = \cap_{n=1}^{\infty} A_n^c$$

It might be useful to consider part (a) of Exercise 1.2.2.

Induction cannot be used on this one, because for induction we need finite set of elements. This stands on the fact, that if induction works for some case, then it works for

$n - 1$ 'th element, and for  $n - 2$ 'th element and this way all the way down to the base case. As an example of why it doesn't work, we can take into account exercise 1.2.2, where all of the elements have infinite number of elements, and their intersection would have infinite amount of elements for any finite number of elements, but it is not true for the infinite amount.

(c) *Is the statement in part (b) valid? If so, write a proof that does not use induction.*

Suppose that

$$x \in (U_{n=1}^{\infty} A_n)^c$$

Then  $x \notin A_n$  for every  $n \in \mathbf{N}$ . Therefore  $x \in A_n^c$  for every  $n \in \mathbf{N}$ . Therefore  $x \in \cap_{n=1}^{\infty} A_n^c$ . Thus

$$(U_{n=1}^{\infty} A_n)^c \subseteq \cap_{n=1}^{\infty} A_n^c$$

Suppose that  $x \in \cap_{n=1}^{\infty} A_n^c$ . Then  $x \notin A_n$  for every  $n \in \mathbf{N}$ . Therefore  $x \notin (U_{n=1}^{\infty} A_n)$  for every  $n \in \mathbf{N}$ . Therefore  $x \in (U_{n=1}^{\infty} A_n)^c$  for every  $n \in \mathbf{N}$ . Therefore

$$\cap_{n=1}^{\infty} A_n^c \subseteq (U_{n=1}^{\infty} A_n)^c$$

Thus if we combine two statements

$$(U_{n=1}^{\infty} A_n)^c = \cap_{n=1}^{\infty} A_n^c$$

as desired.

### 1.3.1

Let  $\mathbf{Z}_5 = \{0, 1, 2, 3, 4, 5\}$  and define addition and multiplication modulo 5. In other words, compute the integer remainder when  $a + b$  and  $ab$  are divided by 5, and use this as the value for the sum and product, respectively.

(a) Show that, given any element  $z \neq 0$  in  $\mathbf{Z}_5$ , there exists an element  $y$  such that  $z + y = 0$ . The element  $y$  is called the additive inverse of  $z$ .

It is true, that for every of those elements we can set  $y = 5 - z$ , for which it is true that  $z + y = 5$  and  $(z + y) \% 5 = 0$  as desired.

(b) Show that, given any element  $z \neq 0$  in  $\mathbf{Z}_5$ , there exists an element  $x$  such that  $zx = 1$ . The element  $x$  is called the multiplicative inverse of  $z$ .

$$(1 * 1) \% 5 = 1$$

$$(2 * 3) \% 5 = 1$$

$$(3 * 2) \% 5 = 1$$

$$(4 * 4) \% 5 = 1$$

as desired.

(c) *The existence of additive and multiplicative inverses is part of the definition of a field. Investigate the set  $\mathbf{Z}_4 = \{0, 1, 2, 3\}$  (where addition and multiplication are defined modulo 4) for the existence of additive and multiplicative inverses. Make a conjecture about the values of  $n$  for which additive inverses exist in  $\mathbf{Z}_n$ , and then form another conjecture about the existence of multiplicative inverses.*

For  $\mathbf{Z}_4$  we define additive inverse the same way we defined it in the part (a) ( $4 - z = y$ ). For multiplicative inverse we have the way with 1, but any for 2 we don't have multiplicative inverse.

Therefore the conjecture about the additive inverse is that every  $\mathbf{Z}_n$  with addition defined as addition modulo  $n$  we have additive inverse.

Proof of it is that every element of  $\mathbf{Z}_n$  is less than  $n$ , and that because of this there exists  $s = n - x \in \mathbf{Z}_n$

For multiplicative inverse the conjecture is that it exists only when  $n$  is a prime number.

### 1.3.2

(a) *Write a formal definition in the style of Definition 1.3.2 for the infimum or greatest lower bound of a set*

A real number  $s$  is the *greatest lower bound* for a set  $A \subseteq \mathbf{R}$  if it meets the following two criteria:

(i)  $s$  is a lower bound

(ii) if  $b$  is any lower bound for  $A$ , then  $s \geq b$ .

(b) *Now, state and prove a version of Lemma 1.3.7 for greatest lower bounds.*

#### **Lemma for greatest lower bounds**

Assume  $s \in \mathbf{R}$  is a lower bound for a set  $A \subseteq \mathbf{R}$ . Then,  $s = \inf(A)$  if and only if, for every choice of  $\epsilon > 0$ , there exists an element  $a \in A$  satisfying  $s + \epsilon > a$ .

Proof:

In one direction: Suppose  $s = \inf(A)$ . Then, by definition of greatest lower bound, there does not exist a lower bound, greater than  $s$ . In other words, suppose that there exist  $\epsilon > 0$  such that there is no element  $a \in A$  such that  $a < s + \epsilon$ . Then  $s + \epsilon$  is a lower bound, which is greater than  $s$ , therefore  $s$  is not a greatest lower bound, which is a contradiction. Therefore there does not exist  $\epsilon > 0$  for which there exist no  $a \in A$  such that  $a < s + \epsilon$ . Therefore for every  $\epsilon > 0$  there exist an  $a \in A$  such that  $a < s + \epsilon$ , as desired.

In other direction: suppose that  $s$  is a lower bound and for every  $\epsilon > 0$  there exist  $a \in A$  such that  $a < s + \epsilon$ . Then any number  $s + \epsilon$  is not a lower bound. Therefore any number, which is greater than  $s$  is not a lower bound. Therefore any lower bound is less or equal to  $s$ . Therefore  $s$  is a greatest lower bound.

Maybe this proof is a little bit more complicated, than it should be, but at least every step is followed properly.

### 1.3.3

(a) Let  $A$  be bounded below, and define  $B = \{b \in \mathbf{R} : b \text{ is a lower bound for } A\}$ . Show that  $\sup(B) = \inf(A)$ .

$B$  is a set, therefore Axiom of Completeness states that there exist real number  $k = \sup(B)$ . Therefore all lower bounds are less or equal to  $k$ .

In order to prove that  $k$  is infimum, we need to show that it is a lower bound.

We do it by contradiction: suppose that  $k$  is not a lower bound. Therefore there exists  $a \in A$  such that  $k > a$ . Let  $\epsilon = k - a > 0$ . Then, because  $k = \sup(B)$  there exist  $b \in B$  such that  $k - \epsilon < b$ . Therefore  $a < b$ . Therefore there exist an element of  $A$ , that is less than lower bound of  $A$ . Therefore  $b$  is not a lower bound. Therefore we have a contradiction. Thus  $k$  is a lower bound.

Because  $k$  is a lower bound,  $k \in B$  by definition of  $B$ . Therefore it is a lower bound, that is greater or equal to any other lower bounds, because it is a supremum of  $B$ . Therefore it is an infimum of  $A$  by definition of infimum.

(b) Use (a) to explain why there is no need to assert that greatest upper bound exist as part of the Axiom of Completeness.

Proof of part (a) does not take into account the fact, that  $A$  (that is bounded below) has an infimum. We prove its existence of the infimum by the fact, that we have a set of lower bounds (which is in its turn has a supremum by Axiom of Completeness), and setting the fact, that its supremum is lower bound itself and therefore proving that it is in the set of lower bounds and therefore setting the fact, that it exists.

(c) Propose another way to use the Axiom of Completeness to prove that sets bounded below have greatest lower bounds.

The only idea, that goes into my mind is to create set  $B = \{-a : a \in A\}$ . Then it'll have a supremum, for which the inverse will be the infimum of the set. We can polish this idea with some theorems and axioms, but I'm satisfied with the current proof already, and nobody is requiring it.

### 1.3.4

Assume that  $A$  and  $B$  are nonempty, bounded above, and satisfy  $B \subseteq A$ . Show  $\sup(B) \leq \sup(A)$ .

We prove it by contradiction: let  $B \subseteq A$  and

$$\sup(B) > \sup(A)$$

Then let  $\epsilon = \sup(B) - \sup(A) > 0$ . Then by lemma we have

$$\begin{aligned} b \in B : b &> \sup(B) - \epsilon \\ b \in B : b &> \sup(B) - \sup(B) + \sup(A) \end{aligned}$$

$$b \in B : b > \sup(A)$$

Therefore  $b > \sup(A)$ , which is an upper bound for  $A$  and by extension  $\forall a \in A : b > a$ . Therefore  $b \notin A$  and  $b \in B$ . Therefore  $B \not\subseteq A$ , which is a contradiction. Therefore  $\sup(B) \leq \sup(A)$ .

### 1.3.5

Let  $A \subseteq \mathbf{R}$  be bounded above, and let  $c \in \mathbf{R}$ . Define the sets  $c + A$  and  $cA$  by  $c + A = \{c + a : a \in A\}$  and  $cA = \{ca : a \in A\}$

(a) Show that  $\sup(c + A) = c + \sup(A)$ .

We gonna prove it by contradiction

Suppose  $c + \sup(A)$  is not an upper bound of  $c + A$ . Then

$$\exists n \in c + A : n > c + \sup(A)$$

let us call such element  $l$ ; also, by the definition of  $c + A$

$$\forall n \in c + A : \exists a \in A : c + a = n$$

therefore

$$\exists a \in A : c + a = l$$

because  $l > c + \sup(A)$

$$c + a > c + \sup(A)$$

$$a > \sup(A)$$

Which is a contradiction. Therefore  $c + \sup(A)$  is an upper bound for  $c + A$ .

Suppose  $c + \sup(A) \neq \sup(c + A)$ . Then  $\sup(c + A)$  is less than  $c + \sup(A)$ . Let  $\epsilon = c + \sup(A) - \sup(c + A)$ . Then

$$\exists k \in A : k > \sup(A) - \epsilon$$

$$k > \sup(A) - c - \sup(A) + \sup(c + A)$$

$$k > -c + \sup(c + A)$$

Therefore

$$\exists h \in c + A : h = k + c$$

$$h - c = k$$

$$h - c > -c + \sup(c + A)$$

$$h > \sup(c + A)$$

therefore



$$\exists h \in c + A : h > \sup(c + A)$$

which is a contradiction. Therefore  $c + \sup(A) = \sup(c + A)$ , as desired.

(b) If  $c \geq 0$ , show that  $\sup(cA) = c * \sup(A)$

If  $c = 0$ , then it  $cA = \{0\}$ , and the case is trivial. Therefore let's discuss further case when  $c > 0$ .

Suppose  $c * \sup(A)$  is not an upper bound for  $cA$ .

Then

$$\exists q \in cA : q > c * \sup(A)$$

by the definition of  $cA$

$$\exists j \in A : q = cj$$

therefore

$$q > c * \sup(A)$$

$$cj > c * \sup(A)$$

$$j > \sup(A)$$

Which is a contradiction, because  $j \in A$ . Therefore  $c * \sup(A)$  is an upper bound for  $cA$ .

Suppose  $c * \sup(A) \neq \sup(cA)$ . Then  $\sup(cA)$  is less than  $c * \sup(A)$ .

$$c * \sup(A) > \sup(cA)$$

$$c * \sup(A) - \sup(cA) > 0$$

$$\frac{c * \sup(A) - \sup(cA)}{c} > 0$$

Let  $\epsilon = \frac{c * \sup(A) - \sup(cA)}{c}$ . Then

$$\exists k \in A : k > \sup(A) - \epsilon$$

$$k > \sup(A) - \frac{c * \sup(A) - \sup(cA)}{c}$$

$$k > \sup(A) - \sup(A) - \sup(cA)/c$$

$$k > \sup(cA)/c$$

Therefore

$$\exists h \in cA : h = ck$$

$$h/c = k$$

$$h/c > \sup(cA)/c$$

$$h > \sup(cA)$$

therefore

$$\exists h \in cA : h > \sup(cA)$$

Which is a contradiction. Therefore  $\sup(cA) = c * \sup(A)$  as desired.

(c) *Postulate a similar type of statement for  $\sup(cA)$  for the case  $c < 0$*

Proposition: suppose that  $A$  is bounded below; if  $c < 0$  then  $\sup(cA) = c * \inf(A)$

Suppose  $c * \inf(A)$  is not an upper bound for  $cA$ .

Then

$$\exists q \in cA : q > c * \inf(A)$$

by the definition of  $cA$

$$\exists j \in A : q = cj$$

therefore

$$q > c * \inf(A)$$

$$cj > c * \inf(A)$$

$$j < \inf(A)$$

Which is a contradiction, because  $j \in A$ . Therefore  $c * \sup(A)$  is an upper bound for  $cA$ .

Suppose  $c * \inf(A) \neq \sup(cA)$ . Then  $\sup(cA)$  is less than  $c * \inf(A)$ .

$$\sup(cA) < c * \inf(A)$$

$$\sup(cA) - c * \inf(A) < 0$$

$$\frac{\sup(cA) - c * \inf(A)}{c} > 0$$

Let  $\epsilon = \frac{\sup(cA) - c * \inf(A)}{c} > 0$ . Then

$$\exists k \in A : k < \inf(A) + \epsilon$$

$$k < \inf(A) + \epsilon$$

$$k < \inf(A) + \frac{\sup(cA) - c * \inf(A)}{c}$$

$$k < \inf(A) + \sup(cA)/c - \inf(A)$$

$$k < \sup(cA)/c$$

Therefore

$$\exists h \in cA : h = ck$$

$$\begin{aligned}
h/c &= k \\
h/c &= k < \sup(cA)/c \\
h/c &< \sup(cA)/c \\
h &> \sup(cA)
\end{aligned}$$

therefore

$$\exists h \in cA : h > \sup(cA)$$

Which is a contradiction. Therefore  $\sup(cA) = c * \inf(A)$  as desired.

### 1.3.6

Compute, without proofs, the suprema and infima of the following sets:

- (a)  $\{n \in \mathbf{N} : n^2 < 10\}$   
 $\sup = 3, \inf = 1.$
- (b)  $\{n/(m+n) : m, n \in \mathbf{N}\}$   
 $\sup = 1/2, \inf = 0.$
- (c)  $\{n/(2n+1) : n \in \mathbf{N}\}$   
 $\sup = 1/2, \inf = 1/3.$
- (d)  $\{n/m : m, n \in \mathbf{N} \text{ with } m+n \leq 10\}$   
 $\sup = 9, \inf = 1/9.$

### 1.3.7

Prove that if  $a$  is an upper bound for  $A$ , and if  $a$  is also an element  $A$ , then it must be that  $a = \sup(A)$

Let's prove this one by contradiction

Suppose that  $a \neq \sup(A)$ . Because  $a$  is still an upper bound,  $\sup(A) < a$ , Therefore  $a \in A$ , but  $\sup(A) < a$ , which is a contradiction. Therefore  $a = \sup(A)$ .

### 1.3.8

If  $\sup(A) < \sup(B)$ , then show that there exists an element  $b \in B$ , that is an upper bound for  $A$ .

Let  $\epsilon = \sup(B) - \sup(A)$ . Then

$$\begin{aligned}
\exists b \in B : b &> \sup(B) - \epsilon \\
b &> \sup(B) - \epsilon
\end{aligned}$$

$$b > \sup(B) - \sup(B) + \sup(A)$$

$$b > \sup(A)$$

therefore  $b$  is an upper bound for  $A$ .

### 1.3.9

Without worrying about formal proofs for the moment, decide if the following statements about suprema and infima are true or false. For any that are false, supply an example where the claim in question does not appear to hold.

(a) A finite, nonempty set always contains its supremum

True

(b) If  $a < L$  for every element  $a$  in the set  $A$ , then  $\sup(A) < L$ . False.  $\sup((0, 1)) = 1$ ;  $\forall a \in (0, 1) : a < 1$ , therefore  $\sup(A) = L$ .

(c) If  $A$  and  $B$  are sets with the property that  $a < b$  for every  $a \in A$  and  $b \in B$ , then it follows that  $\sup(A) < \inf(B)$

False.  $\sup((0, 1)) = \inf((1, 2))$

(d) If  $\sup(A) = s$  and  $\sup(B) = t$ , then  $\sup(A + B) = s + t$ . The set  $A + B$  is defined as  $A + B = \{a + b : a \in A \text{ and } b \in B\}$ .

True

(e) If  $\sup(A) \leq \sup(B)$ , then there exists an element  $b \in B$  that is an upper bound for  $A$ .

False.  $\sup([1, 2]) = \sup((1, 2))$

### 1.4.1

Without doing too much work, show how to prove Theorem 1.4.3 in the case where  $a < 0$  by converting this case into the already proven.

First of all, let's state the theorem itself.

**Theorem 1.4.3 (Density of  $\mathbb{Q}$  in  $\mathbb{R}$ ).** For every two real numbers  $a$  and  $b$  with  $a < b$ , there exists a rational number  $r$  satisfying  $a < r < b$ .

Then let's talk about the possible cases for  $a < 0$ . Then  $b > a$  by the assumptions of the theorem. Therefore  $b \geq 0$  or  $b < 0$ . If  $b > 0$  then there exist 0 between them. If  $b = 0$ , then there exist a rational number  $b = 0 < 1/n < -a$ , and by extension  $a < 1/n < b$  as desired. Therefore we will have some work to do only with case  $b < 0$ .

It is proven, that  $a_1 < r < b_1$  if  $0 \leq a_1 < b_1$ . Therefore for  $a < b < 0$ . Therefore  $-a > -b > 0$ . Therefore if we set  $a_1 = -b$  and  $b_1 = -a$  then by previously stated theorem, there exist

$$a_1 < r < b_1$$

$$-b < r < -a$$

$$b > r > a$$

$$a < r < b$$

as desired.

## 1.4.2

Recall that  $\mathbf{I}$  stands for the set of irrational numbers.

(a) Show that if  $a, b \in \mathbf{Q}$ , then  $ab$  and  $a + b$  are elements of  $\mathbf{Q}$  as well.

Because  $a, b \in \mathbf{Q} \exists m_1, m_2 \in \mathbf{Z}, \exists n_1, n_2 \in \mathbf{N}$  such that

$$a = \frac{m_1}{n_1}$$

$$b = \frac{m_2}{n_2}$$

therefore

$$a + b = \frac{m_1}{n_1} + \frac{m_2}{n_2} = \frac{m_1 n_2 + m_2 n_1}{n_1 n_2}$$

$\mathbf{Z}$  is presumed closed under addition and  $\mathbf{N}$  is presumed closed under  $\mathbf{N}$ , therefore

$$a + b = \frac{m_1 n_2 + m_2 n_1}{n_1 n_2} \in \mathbf{Q}$$

also  $\mathbf{Z}$  is closed under multiplication, and therefore

$$ab = \frac{m_1 m_2}{n_1 n_2} \in \mathbf{Q}$$

(b) Show that if  $a \in \mathbf{Q}$  and  $t \in \mathbf{I}$ , then  $a + t \in \mathbf{I}$  and  $at \in \mathbf{I}$  as long as  $a \neq 0$ .

We prove both things by contradiction.

Suppose  $a + t \in \mathbf{Q}$ . Then  $\exists b \in \mathbf{Q} : a + t = b$ . Also,  $a \in \mathbf{Q} \rightarrow -a \in \mathbf{Q}$ . Therefore

$$a + t = b$$

$$t = b - a$$

Therefore, because  $\mathbf{Q}$  is closed under addition (as we discussed previously),  $t \in \mathbf{Q}$ , which is a contradiction. Therefore  $a + t \in \mathbf{I}$ .

Suppose  $at \in \mathbf{Q}$  for  $a \neq 0$ . Then

$$\exists b \in \mathbf{Q} : at = b$$

therefore if  $a = m/n \in Q$ , then  $1/a = n/m \in Q$ . Therefore

$$t = b/a$$

Therefore  $t \in Q$ , which is a contradiction. Therefore  $at \in I$  for  $a \neq 0$ .

(c) Part (a) can be summarized by saying that  $Q$  is closed under addition and multiplication. Is  $I$  closed under addition and multiplication? Given two irrational numbers  $s$  and  $t$ , what can we say about  $s + t$  and  $st$ .

$I$  is not closed under addition, nor under multiplication. Proof is  $\sqrt{2} + 1$  and  $0 - \sqrt{2}$  are both irrational, but

$$\sqrt{2} + 1 - \sqrt{2} = 1 \in Q$$

and

$$\sqrt{2} * \sqrt{2} = 2 \in Q$$

### 1.4.3

Using Exercise 1.4.2, supply a proof for Corollary 1.4.4 by applying Theorem 1.4.3 to the real numbers  $a - \sqrt{2}$  and  $b - \sqrt{2}$ .

First, let's state Corollary 1.4.4.

Given any two real numbers  $a < b$ , there exists an irrational number  $t$  satisfying  $a < t < b$ .

We know, that between two numbers  $a_1 < b_1$  there exists a rational number  $r$ , for which it is true

$$a_1 < r < b_1$$

Let  $a_1 = a + \sqrt{2}$  and  $b_1 = b + \sqrt{2}$ . Then

$$a_1 < r < b_1$$

$$a + \sqrt{2} < r < b + \sqrt{2}$$

$$a < r - \sqrt{2} < b$$

As we know,  $r - \sqrt{2}$  is an irrational number, therefore between  $a$  and  $b$  there exists an irrational number.

### 1.4.4

Use the Archimedean Property of  $R$  to rigorously prove that  $\inf\{1/n : n \in N\} = 0$

It is true, that  $\forall n \in N : n > 0$ . Therefore

$$n > 0$$

$$1/n > 0$$

Therefore 0 is a lower bound for  $1/n$ . Also, because of Archimedean Property, if we take any  $\forall \epsilon > 0 : \exists 1/n : 1/n < \epsilon$

$$\forall \epsilon > 0 : \exists 1/n : 1/n < \epsilon$$

$$\forall \epsilon > 0 : \exists 1/n : 0 + \epsilon > 1/n$$

Therefore for every  $\epsilon$  there exists an element of a set such that  $0 + \epsilon$  is greater than this set. Therefore 0 is an infimum of this set, as desired.

### 1.4.5

*Prove that  $\cap_{n=1}^{\infty} (0, 1/n) = \emptyset$ . Notice that this demonstrates that the Nested Interval Property must be closed for the conclusion of the theorem to hold.*

First of all,  $1/n > 0$  implies, that if  $y \leq 0$  then  $y \notin (0, 1/n)$  for any  $n \in N$ .

Because of the Archimedean Property, for every  $y \in R > 0$  there exists  $n \in N$  such that  $1/n < y$ . Therefore there does not exist  $y > 0$  such that  $y \geq 1/n$  for every  $n \in N$ . Therefore for every  $y \geq 0$  there exist  $n \in N$  such that  $y \notin (0, 1/n)$ .

Therefore there are no real numbers in  $\cap_{n=1}^{\infty} (0, 1/n)$ . Therefore

$$\cap_{n=1}^{\infty} (0, 1/n) = \emptyset$$

as desired.

This conclusion proves, that if we have use an open interval for nested interval property, then we'll have a problem.

### 1.4.6

*(a) Finish the proof of Theorem 1.4.5 by showing that the assumption  $\alpha^2 > 2$  leads to a contradiction of the fact that  $\alpha = \sup(T)$ .*

First, Theorem 1.4.5 states that there exists a real number  $\alpha \in R$  satisfying  $\alpha^2 = 2$ .

Our assumption is that

$$T = \{t \in R : t^2 < 2\} \text{ and } \alpha = \sup(T)$$

Our strategy is to state that if  $a^2 > 2$ , then  $a$  is not a least upper bound.

If  $a$  is an least upper bound for  $T$ , then it is true, that

$$\forall \epsilon > 0 : \exists t \in T : t > \alpha - \epsilon$$

Therefore for every  $n \in N$

$$\exists t \in T : t > \alpha - 1/n$$

Now let us follow with the proof. For all  $n \in N$ .

$$t > \alpha - 1/n$$

$$\alpha - 1/n < t$$

$$(\alpha - 1/n)^2 < t^2$$

$$(\alpha - 1/n)^2 < t^2$$

$$\alpha^2 - \frac{2\alpha}{n} + \frac{1}{n^2} < t^2$$

$$\alpha^2 - \frac{2\alpha}{n} + \frac{1}{n^2} < t^2$$

Let's justify something now.  $1^2 = 1 < 2$ , therefore  $1 < \alpha$ . Therefore  $2\alpha - 1 > 0$ . Also, because  $\alpha^2 > 2$ ,  $\alpha^2 - 2 > 0$ . Therefore

$$\frac{\alpha^2 - 2}{2\alpha - 1} > 0 \in R$$

Now, let us pick  $n \in N$  such that

$$1/n < \frac{\alpha^2 - 2}{2\alpha - 1}$$

then

$$1/n < \frac{\alpha^2 - 2}{2\alpha - 1}$$

$$1/n^2 < \frac{\alpha^2 - 2}{2\alpha - 1}$$

$$n^2 > \frac{2\alpha - 1}{\alpha^2 - 2}$$

$$\frac{2\alpha - 1}{n^2} < \alpha^2 - 2$$

$$\frac{2\alpha}{n^2} - \frac{1}{n^2} < \alpha^2 - 2$$

$$-\alpha^2 + \frac{2\alpha}{n^2} - \frac{1}{n^2} < -2$$

$$\alpha^2 - \frac{2\alpha}{n^2} + \frac{1}{n^2} > 2$$

$$\alpha^2 - \frac{2\alpha}{n} + \frac{1}{n^2} > 2$$



but

$$\alpha^2 - \frac{2\alpha}{n} + \frac{1}{n^2} < t^2 < 2$$

Therefore we have a contradiction. Therefore  $\alpha \leq 2$ . Therefore  $\alpha = 2$ , as desired. Phew.

(b) *Modify the argument to prove the existence of  $\sqrt{B}$  for any real number  $b \geq 0$ .*

Let's discuss the case  $a^2 < b$ .

$$(\alpha + 1/n)^2 = \alpha^2 + \frac{2\alpha}{n} + \frac{1}{n^2} < \alpha^2 + \frac{2\alpha}{n} + \frac{1}{n} = \alpha + \frac{2\alpha + 1}{n}$$

Let

$$\frac{1}{n} < \frac{b - \alpha}{2\alpha + 1}$$

then

$$(\alpha + \frac{1}{n})^2 = \alpha^2 + (b - \alpha^2) = b$$

Therefore  $\alpha + 1/n \in T$ . Therefore  $a^2 \geq b$ .

Now let

$$1/n < \frac{\alpha^2 - b}{2\alpha - 1}$$

. Then by reasoning in the last part of exercise we can state, that  $\alpha^2 \leq b$ . Therefore  $\alpha^2 = b$ , as desired.

## 1.4.7

*Finish the following proof for Theorem 1.4.12.*

First of all, let us state Theorem 1.4.12

If  $A \subseteq B$  and  $B$  is countable, then  $A$  is either countable, finite or empty.

Assume  $B$  is a countable set. Thus, there exists  $f : N \rightarrow B$ , which 1-1 and onto. Let  $A \subseteq B$  be an infinite subset of  $B$ . We must show that  $A$  is countable.

Let  $n_1 = \min\{n \in N : f(n) \in A\}$ . As a start to a definition of  $g : N \rightarrow A$ , set  $g(1) = f(n_1)$ . Show how to inductively continue this process to produce a 1-1 function  $g$  from  $N$  onto  $A$ .

Proposition:  $g(n) = f(n_n)$ .

Step: let  $n_{n+1} = \min\{k > n \in N : f(k) \in A\}$ . Then  $f(n) < n_{n+1} \notin n_{n+1}$ . Therefore  $g(n+1) = f(n_{n+1})$ . Therefore

$$\forall n_1 \neq n_2, k_1, k_2 \in N \exists g(n_1) = f(k_1) \neq g(n_2) = f(k_2)$$

and

$$\forall l \in g(N) \exists k \in N : g(k) = f(n_k) = l$$

Therefore there exist a bijective function  $g : N \rightarrow A$ . Therefore  $A$  is countable, as desired.

If  $A$  is not infinite, then it's finite (duh).

Same with empty.

## 1.4.8

Use the following outline to supply for the statements in Theorem 1.4.13.

First of all, let's state Theorem 1.4.13

(i) If  $A_1, A_2, \dots, A_m$  are each countable sets, then the union  $A_1 \cup A_2 \cup \dots \cup A_m$  is countable.

(ii) If  $A_n$  is countable set for each  $n \in N$ , then  $\bigcup_{n=1}^{\infty} A_n$  is countable.

(a) First, prove statement (i) for two countable sets,  $A_1$  and  $A_2$ . Example 1.4.8 (ii) may be a useful reference. Some technicalities can be avoided by first replacing  $A_2$ , with the set  $B_2 = A_2 \setminus A_1 = \{x \in A_2 : x \notin A_1\}$ . The point of this is that the union  $A_1 \cup B_2$  is equal to  $A_1 \cup A_2$  and the sets  $A_1$  and  $B_2$  are disjoint. (What happens if  $B_2$  is finite?)

Let us first set  $B_2 = A_2 \setminus A_1$ . We will do it in order to have two useful properties:

$$a \in B_2 \rightarrow a \in A_1^c \cup A_2 \rightarrow a \notin A_1$$

$$A_1 \cup B_2 = A_1 \cup (A_1^c \cap A_2) = (A_1 \cap A_1^c) \cup (A_1 \cap A_2) = A_1 \cap B_2$$

Let's finally begin with the proof.  $B_2 \subseteq A_2$ . By using previous theorem we can state, that  $B_2$  is either countable, finite, or empty. If it is empty, then  $A_1 \cup A_2 = A_1$ , and therefore is countable.

For the finite case we'll need function, that returns the  $n$ 'th smallest element of the set. We can argue, that this function does not need the rigorous definition, but we'll give it anyways:

$$s_F(1) = \{x \in F : \forall y \in F : x \leq y\}$$

$$s_F(n) = \{x \in F \setminus \{s(1), \dots, s(n-1)\} : \forall y \in F \setminus \{s(1), \dots, s(n-1)\} : x \leq y\}$$

for the finite set  $F \subseteq R$ .

$$\forall n_1 > n_2 \in N : s_F(n_2) \notin F \setminus \{s(1), \dots, s(n_1-1)\} \rightarrow s_F(n_1) \neq s_F(n_2)$$

therefore the function is injective

We remove one element at every iteration from  $F$  every iteration, therefore  $\{s_F(1), \dots, s_F(|F|)\}$  spans the whole set  $F$ . Therefore the function is surjective. Therefore the function is bijective.

Now let us define  $q : N \rightarrow A_1 \cup B_2$ . Let  $g_1 : N \rightarrow A_1$  be a bijective function, that exists, because the set  $A_1$  is countable. Then

$$q(x) = \begin{cases} s_{B_2}(x) & \text{if } x \leq |B_2| \\ g_1(x - |B_2|) & \text{if } x > |B_2| \end{cases} \quad (1)$$

Let's analyse this function. If  $x_1 < x_2 \leq |B_2|$  or  $x_1 > x_2 > |B_2|$ , then  $q(x_1) \neq q(x_2)$  by injectivity of  $s_{B_2}$  or  $g_1(x)$  respectively. If  $x_1 \leq |B_2| < x_2$ , then  $q(x_1) \in B_2$  and  $q(x_2) \in A_1$ . Those two sets are disjoint, and therefore  $q(x_1) \neq q(x_2)$ . Therefore for all possible  $x_1, x_2 \in N \rightarrow q(x_1) \neq q(x_2)$ . Therefore the function is injective.

Let  $b \in N$  and  $N_b = \{x \in N : x > b\}$ . Also let  $N_a = \{x \in N_b : x - b\}$ . Then

$$\forall c \in N_a : \exists v \in N_b : c = v - b$$

$$\forall v \in N_b : \exists n \in N : v = n + b$$

$$\forall c \in N_a : \exists n \in N : c = n$$

therefore  $N \subseteq N_a$ .

$$\forall n \in N : \exists v \in N_b : v = n + b$$

$$\forall v \in N_b : \exists c \in N_a : c = v - b$$

$$\forall n \in N : \exists c \in N_a : c = n$$

therefore  $N_a \subseteq N$ . Thus  $N = N_a$ .

Therefore

$$\forall x > |B_2| : \exists n \in N : g_1(x - |B_2|) = g_1(n)$$

Therefore  $g_1$  spans the whole set  $A_1$ .  $s_{B_2}$  spans the whole  $B_2$  by surjectivity of  $s$ . Therefore  $q$  spans  $A_1 \cup B_2$ . Therefore it is surjective.

Therefore  $A_1 \cup B_2$  is bijective, therefore  $A_1 \cup B_2$  is countable if  $B_2$  is finite. One last thing, that I want to add before finishing this case is to acknowledge the fact, that this theorem is true not for real numbers only. This fact throws our minimal element part out of the proof. This misfortune can be avoided through the usage of lists and by converting sets into them. I did not use this fact here because of the need to axiomatically define lists, which I can, but don't want to do.

Now let us proceed with the case when  $B_2$  is countable. Let  $g_1 : N \rightarrow A_1$  be a bijective function for  $A_1$  and  $g_2 : N \rightarrow B_2$  be a bijective function for  $B_2$  (both of those exist because of the fact, that both of the sets are countable). Let  $q : N \rightarrow A_1 \cup B_2$  be defined as

$$q(x) = \begin{cases} g_1((x+1)/2) & \text{if } x \text{ is odd} \\ g_2(x/2) & \text{if } x \text{ is even} \end{cases} \quad (2)$$

Then  $\forall n_1 \neq n_2 \in N \rightarrow q(n_1) \neq q(n_2)$  because of either injectivity of both functions, or because of the fact, that both sets are disjoint. Therefore  $q$  is injective. Also

$$x \text{ is odd} : (x+1)/2 = x \text{ is even} : x/2 = N$$

Therefore  $q$  spans  $A_1 \cup B_2$ . Therefore the function is surjective.

Thus  $q$  is bijective and  $A_1 \cup B_2 = A_1 \cup B_2$  is countable if  $B_2$  is countable.

Therefore for any countable sets  $A_1$  and  $A_2$  it is true, that their union is countable as well.

*Now, explain how the more general statement in (i) follows*

We prove the first part of the theorem by induction.

Base:  $A_1 \cup A_2$  is countable.

Proposition:  $\cup_{n=1}^m A_n$  is countable for  $m \in N$ .

Step:

$$\cup_{n=1}^{m+1} A_n = \cup_{n=1}^m A_n \cup A_{m+1}$$

$\cup_{n=1}^m A_n$  is a countable set because of the proposition.  $A_{m+1}$  is countable by assumption of the theorem. Therefore  $\cup_{n=1}^{m+1} A_n$  is a union of two countable sets, and therefore is itself countable, as desired.

*(b) Explain why induction cannot be used to prove part (ii) of Theorem 1.4.13 from part (i)*

Induction cannot be used to prove part (ii) because of the fact, that induction relies of finality of the given set.

*(c) Show how arranging  $N$  into the two-dimentional array*

$$\begin{array}{ccccccc} 1 & 3 & 6 & 10 & 15 & \dots \\ & 2 & 5 & 9 & 14 & \dots \\ & & 4 & 8 & 13 & \dots \\ & & & 7 & 12 & \dots \\ & & & & 11 & \dots \\ & & & & & \dots \end{array}$$

*leads to a proof of a Theorem 1.4.13 (ii)*

This proof will not be as rigorous as the ones before that, but i'll try them anyways. Let us convert all of the given sets onto lists. Then let us construct lists

$$\begin{aligned} l_1 &= A_{1_1} \\ l_2 &= A_{2_1}, A_{1_2} \\ l_3 &= A_{3_1}, A_{2_2}, A_{3_1} \\ &\dots \\ l_n &= A_{n_1}, A_{n-1_2}, A_{n-2_3} \dots \end{aligned}$$

if  $A_{n_n}$  already has been included into the lists  $l_{1..n-1}$ , then we don't include it in the list. Then if we concatenate all of those lists together, then we will have function  $q : N \rightarrow \cup_{n=1}^{\infty} A_n$  which is defined as

$$q(x) = x\text{'th element of the final list}$$

This function will be injective, because we threw out already included elements, and will be surjective, because all of the elements of  $\cup_{n=1}^{\infty} A_n$  are eventually in the list. Thus, we have a bijective function between  $N$  and  $\cup_{n=1}^{\infty} A_n$ . Therefore  $\cup_{n=1}^{\infty} A_n$  is countable, as desired.

### 1.4.9

(a) *Given sets  $A$  and  $B$ , explain why  $A \sim B$  is equivalent to asserting  $B \sim A$ .*

Short answer: bijectivity implies inversability.

Medius answer: injectivity leads us to the fact, that for every element of codomain there exists only one element of the domain and by extension lets us define inverse function in the first place; surjectivity guarantees, that the inverse function is defined for every element of the domain.

Long answer:

$A \sim B$  implies, that there exists a bijective function  $g : A \rightarrow B$ . Bijective means that

$$\forall a_1 \neq a_2 \in A \rightarrow g(a_1) \neq g(a_2)$$

$$\forall b \in B : \exists a \in A : g(a) = b$$

This implies that for all  $b \in B$  there exist only one  $a \in A$  such that  $g(a) = b$ . This fact lets us define  $g^{-1} : B \rightarrow A$  as

$$g^{-1}(b) = a \text{ such that } g(a) = b$$

Then

$$\forall b_1 \neq b_2 \in B \rightarrow g^{-1}(b_1) \neq g^{-1}(b_2)$$

because  $g$  is a function and therefore  $g^{-1}(b_1) \neq g^{-1}(b_2) \rightarrow g(a_1) \neq g(a_2) \rightarrow a_1 \neq a_2$ . Therefore  $g^{-1}$  is injective.

$$\forall a \in A : \exists b \in B : g^{-1}(b) = a$$

and therefore

$$\forall a \in A : \exists b \in B : g^{-1}(b) = a$$

therefore function is surjective.

Therefore the function is bijective and therefore  $B \sim A$ .

Therefore  $A \sim B \iff B \sim A$ , as desired.

(b) *For three sets  $A, B$  and  $C$ , show that  $A \sim B$  and  $B \sim C$  implies  $A \sim C$ . These two properties are what is meant by saying that  $\sim$  is an equivalence relation.*

Because  $A \sim B$  and  $B \sim C$  there exist two bijective functions  $g_1 : A \rightarrow B$  and  $g_2 : B \rightarrow C$ . Therefore

$$\forall a_1 \neq a_2 \in A \rightarrow g_1(a_1) \neq g_1(a_2)$$

$$\forall b_1 \neq b_2 \in B \rightarrow g_2(b_1) \neq g_2(b_2)$$

therefore

$$\forall a_1 \neq a_2 \in A \rightarrow g_1(a_1) \neq g_1(a_2) \rightarrow g_2(g_1(a_1)) \neq g_2(g_1(a_2))$$

Also

$$\forall a \in A : \exists b \in B : g_1(a) = b$$

$$\forall b \in B : \exists c \in C : g_2(b) = c$$

and therefore

$$\forall a \in A : \exists b \in B : g_1(a) = b \rightarrow \exists c \in C : g_2(g_1(a)) \in C$$

Therefore  $g_2 \circ g_1 : A \rightarrow C$  is a bijective function. Therefore  $A \sim C$ , as desired.

### 1.4.10

*Show that the set of all finite subsets of  $N$  is a countable set. (It turns out that the set of all subsets of  $N$  is not a countable set. This is a title of Section 1.5)*

Our strategy here will be to show that for every  $n \in N$  set of sets of length  $n$  is countable, therefore their union is countable.

Maybe I should proceed with induction. Base: First of all, set  $N$  is countable. Therefore the set of sets of length 1 is countable.

Proposition: Suppose that set of sets of length  $n \in N$  is countable.

Step: Because set of sets of length  $N$  is countable, let us take a set  $S_m$ . Then, let us add a number into it, which is not already in this set. To be more precise, let us define

$$N_m = N \setminus S_m$$

$$S_{m_k} = S_m \cup \{k\text{'th number of } N_m\}$$

Then each of  $S_{m_k}$  is a union of countable sets, and therefore countable. Therefore  $\bigcup_{k=1}^{\infty} S_{m_k} = S_{m+1}$  is countable.

Therefore each set of sets of length  $n \in N$  is countable. Therefore their union is countable. Therefore set of finite subsets of  $N$  is countable, as desired.

### 1.4.11

*Consider the open interval  $(0,1)$ , and let  $S$  be the set of points in the open unit square; that is,  $S = \{(x,y) : 0 \leq x, y \leq 1\}$*

*(a) Find a 1-1 function, that maps  $(0, 1)$  into, but not necessarily onto,  $S$ . (This is easy.)*

Yeah, it is. Let  $g(x) = (x, 0.5)$ .



$$x^3 + 0 - 2 = 0$$

$$x^3 = 2$$

$$x = \sqrt[3]{2}$$

Therefore  $\sqrt[3]{2}$  is algebraic.

Let  $a_0 = 1$ ,  $a_1 = 0$ ,  $a_2 = -10$ ,  $a_3 = 0$  and  $a_4 = 1$ . Then

$$x^4 - 10x^2 + 1 = 0$$

$$x^4 - 10x^2 + 25 = 24$$

$$(x^2 - 5)^2 = 4 * 6$$

$$x^2 - 5 = 2\sqrt{6}$$

$$x^2 = 5 + 2\sqrt{6}$$

$$x^2 = 2 + 3 + 2\sqrt{2}\sqrt{3}$$

$$x = \sqrt{2} + \sqrt{3}$$

Therefore  $\sqrt{2} + \sqrt{3}$  is algebraic.

(b) Fix  $n \in N$ , and let  $A_n$  be the algebraic numbers obtained as roots of polynomials with integer coefficients that have degree  $n$ . Using the fact that every polynomial has a finite number of roots, show that  $A_n$  is countable. (For each  $m \in N$ , consider polynomials  $a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$  that satisfy  $|a_n| + |a_{n-1}| + \dots + |a_1| + |a_0| \leq m$ .)

Let us fix  $m \in M$ , then there exist a finite number of  $a_n, \dots, a_1, a_0 \in Z$  such that  $|a_n| + |a_{n-1}| + \dots + |a_1| + |a_0| \leq m$ . Therefore for each  $m \in M$  there are finitely many combinations of coefficients and thus finitely many numbers of roots. Therefore let us correspond each root, with a number  $n \in N$  in ascending order (keeping in mind, we should check, that this number is not already in the function). Then we'll get a bijective function from  $N$  to all the possible roots of polynomials with integral coefficients. Therefore  $A_n$  is countable.

(c) Now, argue that the set of all algebraic numbers is countable. What may we conclude about the set of transcendental numbers?

Because  $A_n$  is countable for each  $n \in N$  we know, that  $\cup_{n=1}^{\infty} A_n$  is countable as well. By definition, algebraic number is a root of one of such polynomials, and therefore algebraic numbers are a countable set.

The fact, that the set of algebraic numbers is countable presents us with a fact, that the set of transcendental numbers is not countable (because if it was, then the set of real numbers would be a union of two countable sets and therefore countable as well).



### 1.4.13 (Schroder-Bernstein Theorem)

Assume that there exists a 1-1 function  $f : X \rightarrow Y$  and another 1-1 function  $g : Y \rightarrow X$ . Follow the steps to show that there exists a 1-1, onto function  $h : X \rightarrow Y$  and hence  $X \sim Y$ .

(a) The range of  $f$  is defined by  $f(X) = \{y \in Y : y = f(x) \text{ for some } x \in X\}$ . Let  $y \in f(X)$ . (Because  $f$  is not necessarily onto, the range  $f(X)$  may not be all of  $Y$ .) Explain why there exists a unique  $x \in X$  such that  $f(x) = y$ . Now define  $f^{-1}(y) = x$ , and show that  $f^{-1}$  is a 1-1 function from  $f(X)$  onto  $X$ .

Because  $f$  is given to be injective, it is true that

$$\forall x_1 \neq x_2 \in X \rightarrow f(x_1) \neq f(x_2)$$

$f$  is also a function. By definition of a function we know, that for every element of domain there exists only one element of codomain. Therefore

$$\forall f(x_1) \neq f(x_2) \in f(X) \rightarrow x_1 \neq x_2$$

By plugging  $y$  into the equation we have

$$\forall y_1 \neq y_2 \in f(X) \rightarrow x_1 \neq x_2$$

Thus we can define  $f^{-1} : f(X) \rightarrow X$ :

$$f^{-1}(y) = \{x \in X : f(x) = y\}$$

This function will be injective by the fact that

$$\forall y_1 \neq y_2 \in f(X) \rightarrow x_1 \neq x_2 \rightarrow f^{-1}(x_1) \neq f^{-1}(x_2)$$

In a similar way, we can also define the 1-1 function  $g^{-1} : g(X) \rightarrow Y$

Same logic applies to  $g(x)$ . Therefore we have

$$g^{-1}(x) = \{y \in Y : g(y) = x\}$$

which by the same logic is injective.

(b) Let  $x \in X$  be arbitrary. Let the chain  $C_x$  be the set consisting of all elements of the form

$$(1) \dots, f^{-1}(g^{-1}(x)), g^{-1}(x), x, f(x), g(f(x)), f(g(f(x))), \dots$$

Explain why the number of elements to the left of  $x$  in the above chain may be zero, finite, or infinite

The number of elements on the left is 0, if there does not exist  $y \in Y$  such that  $g(y) = x$ .

The number of elements on the left is finite number, if after some element  $l$  there does not exist an element  $y \in Y$  or  $x \in X$ , such that  $g(y) = l$  or  $f(x) = l$ .

The number of elements on the left is infinite, if there exist both of those numbers for every element (for example, if both functions are onto).

(c) Show that any two chains are either identical, or completely disjoint

Suppose that  $x_1 \neq x_2$ . Then  $f(x_1) \neq f(x_2)$  by injectivity of  $f$ . Also,  $f(x) \in Y$ , and therefore  $g(f(x_1)) \neq g(f(x_2))$  and so on.

$f^{-1}$  and  $g^{-1}$  are injective as well. Therefore the same logic applies. Thus we can state that

$$\forall x_1 \neq x_2 \rightarrow C_{x_1} \cap C_{x_2} = \emptyset$$

and

$$\forall x_1 = x_2 \rightarrow C_{x_1} \cap C_{x_2} = C_{x_1} = C_{x_2}$$

(d) Note that the terms of the chain in (1) alternate between elements of  $X$  and elements of  $Y$ . Given a chain  $C_x$ , we want to focus on  $C_x \cap Y$ , which is just a part of the chain that sits in  $Y$ .

Define the set  $A$  to be the union of all chains  $C_x$  satisfying  $C_x \cap Y \subseteq f(X)$ . Let  $B$  consist of the union of the remaining chains not in  $A$ . Show that any chain contained in  $B$  must be of the form

$$y, g(y), f(g(y)), g(f(g(y))), \dots$$

where  $y$  is an element of  $Y$  that is not in  $f(X)$

Let  $A$  and  $B$  be defined as in the exercise text. Then

$$\forall C_x \in B : \exists c \in C_x : c \in Y \text{ and } c \notin f(X)$$

. Thus

$$\nexists x \in X : f(x) = c$$

at the same time

$$c \in Y \rightarrow \exists x \in X : g(y) = x$$

.

Because  $g : Y \rightarrow g(Y) \subseteq X$  and  $f : X \rightarrow f(X) \subseteq Y$  we can write the chain in the form

$$c, g(c), f(g(c)), \dots$$

Substituting  $c$  with  $y$  we get

$$y, g(y), f(g(y)), \dots$$

where  $y \notin f(X)$ , as desired.

(e) Let  $X_1 = A \cap X$ ,  $X_2 = B \cap X$ ,  $Y_1 = A \cap Y$ , and  $Y_2 = B \cap Y$ . Show that  $f$  maps  $X_1$  onto  $Y_1$  and that  $g$  maps  $Y_2$  onto  $X_2$ . Use this information to prove  $X \sim Y$ .

Let  $X_1, X_2, Y_1$  and  $Y_2$  be defined as in exercise text. Then let us discuss  $f(X_1)$

$$A = \{C_x : C_x \cap Y \subseteq f(X)\}$$

$$X_1 = \{C_x \cap X : C_x \cap Y \subseteq f(X)\}$$

$$Y_1 = \{C_x \cap Y : C_x \cap X \subseteq g(Y)\}$$

therefore

$$\forall y \in Y_1 : \exists x \in X_1 : f(x) = y$$

and

$$\forall x \in X_2 : \exists y \in Y_2 : g(y) = x$$

Or in other words,  $f : X_1 \rightarrow Y_1$  is surjective(onto).

By the same logic,  $g : Y_2 \rightarrow X_2$  is also surjective.

Let us define function  $h : X \rightarrow Y$

$$h(x) = \begin{cases} f(x) & \text{if } x \in X_1 \\ g^{-1}(x) & \text{if } x \in X_2 \end{cases} \quad (3)$$

Functions  $f$  and  $g^{-1}$  are both injective.  $X_1 \cap X_2 = \emptyset$ . Therefore  $h$  is injective as well. Also,  $X_1 \cup X_2 = X$ , and both  $f$  and  $g$  are surjective. Therefore  $h$  is surjective as well.

Therefore this function is bijective. Therefore we have a bijective function from  $X$  to  $Y$ . Therefore  $X \sim Y$ , as desired.

### 1.5.1

Show that  $(0, 1) = \{x \in R : 0 < x < 1\}$  is uncountable if and only if  $R$  is uncountable.

Let

$$f(x) = \frac{2x - 2}{x^2 - 2x}$$

Calculus shows that this function maps  $(0, 1) \rightarrow R$ . Also, it shows, that it is increasing and therefore it is bijective. Thus there exist a bijective. Thus  $(0, 1) \sim R$ . Therefore it is uncountable if and only if  $R$  is uncountable.

### 1.5.2

(a) Explain why the real number  $x = b_1b_2b_3\dots$  cannot be  $f(1)$ .

If  $f(1) = b$  then

$$b_1 = 2 \rightarrow b_1 \neq 2$$

$$b_1 \neq 2 \rightarrow b_1 = 2$$

Therefore we have a contradiction.

(b) Now, explain why  $x \neq f(2)$ , and in general why  $x \neq f(n)$  for any  $n \in N$ .

If  $f(1) = b$  then

$$b_n = 2 \rightarrow b_n \neq 2$$

$$b_n \neq 2 \rightarrow b_n = 2$$

Therefore we have a contradiction for  $n \in N$ .

(c) Point out the contradiction that arises from these observations and conclude that  $(0, 1)$  is uncountable.

If  $(0, 1)$  is countable, then  $b \in (0, 1)$ , but it cannot be correspondent to  $n \in N$  for any  $f : N \rightarrow (0, 1)$ . Therefore either  $b \notin (0, 1)$  or  $(0, 1)$  is not countable. Because  $b \in (0, 1)$  we conclude that  $(0, 1)$  is uncountable.

### 1.5.3

Supply rebuttals to the following complaints about the proof of Theorem 1.5.1

(a) Every rational number has a decimal expansion so we could apply this same argument to show that the set of rational numbers between 0 and 1 is uncountable. However, because we know that any subset of  $Q$  must be countable, the proof of Theorem 1.5.1 must be flawed

We can try to apply the same argument to  $Q$ , but now we have a problem with the fact, that every rational number in decimal expansion repeats itself after some point or another. Therefore  $b \notin Q$ , therefore we cannot conclude anything.

(b) A few numbers have two different decimal representations. Specifically, any decimal expansion that terminates can also be written with repeating 9's. For instance,  $1/2$  can be written as  $.5$  or as  $.4999\dots$ . Doesn't this cause some problems?

No, it doesn't. And the reason on why it doesn't cause any problems is because our argument stems on the fact, that given  $b \in R$  is not in the set, if the set is countable. Therefore different representations problem is irrelevant.

### 1.5.4

Let  $S$  be the set consisting of all sequences of 0's and 1's. Observe that  $S$  is not a particular sequence, but rather a large set whose elements are sequences; namely,

$$S = \{(a_1, a_2, a_3, \dots) : a_n = 0 \text{ or } 1\}$$

As an example, the sequence  $(1, 0, 1, 0, 1, 0, \dots)$  is an element of  $S$ , as the sequence  $(1, 1, 1, 1, \dots)$ .

Give a rigorous argument, that  $S$  is uncountable

I don't know, if it counts, as rigorous, but here we go.

Each  $s \in S$  corresponds to a binary representation of a number in  $(0, 1)$ . Therefore  $S \sim (0, 1) \sim \mathbb{R}$ , therefore it is uncountable.

We can work around sets and NIP to show the same thing if we want to, but I don't want to.

### 1.5.5

(a) Let  $A = \{a, b, c\}$ . List the eight elements of  $P(A)$ . (Do not forget that  $\emptyset$  is considered to be a subset of every set.)

$$\emptyset, \{a\}, \{b\}, \{b, a\}, \{c\}, \{c, a\}, \{c, b\}, \{c, b, a\}$$

(b) If  $A$  is finite with  $n$  elements, show that  $P(A)$  has  $2^n$  elements. (Constructing a particular subset of  $A$  can be interpreted as making a series of decisions about whether or not to include each element of  $A$ .)

Repeat of 1.2.11

This proof is dumb, but intuitive:

Every subset is corresponding to a number in binary number: 0 for excluded, 1 for included. Therefore there exist  $2^n$  possible combinations.

For a more concrete proof let's resort to induction.

Base case(s): subsets of  $\emptyset$  are  $\emptyset$  itself ( $2^0 = 1$  in total). Subsets of set with one element are  $\emptyset$  and set itself ( $2^1 = 2$  in total).

Proposition is that set with  $n$  elements has  $2^n$  subsets.

Inductive step is that for set with  $n+1$  elements can either have or not have the  $n+1$ 'th element. Therefore there exist  $2^n + 2^n = 2 * 2^n = 2^{n+1}$  subsets, as desired.

### 1.5.6

(a) Using the particular set  $A = \{a, b, c\}$ , exhibit two different 1-1 mappings from  $A$  to  $P(A)$

$$f_1(x) = \begin{cases} a \rightarrow \{a\} \\ b \rightarrow \{b\} \\ c \rightarrow \{c\} \end{cases} \quad (4)$$

$$f_2(x) = \begin{cases} a \rightarrow \emptyset \\ b \rightarrow \{b\} \\ c \rightarrow \{c\} \end{cases} \quad (5)$$

(b) Letting  $B = \{1, 2, 3, 4\}$ , produce an example of a 1-1 map  $g : B \rightarrow P(B)$ .

$$f_3(x) = \begin{cases} 1 \rightarrow 1 \\ 2 \rightarrow 2 \\ 3 \rightarrow 3 \\ 4 \rightarrow 4 \end{cases} \quad (6)$$

(c) Explain why, in parts (a) and (b), it is impossible to construct mappings, that are onto

$$|A| < |P(A)|$$

### 1.5.7

Return to the particular functions constructed in Exercise 1.5.6 and construct subset  $B$  that results using the preceding rule. In each case, note that  $B$  is not in the range of the function used.

For  $f_1$ :

$$B = \{\emptyset\}$$

For  $f_2$ :

$$B = \{a\}$$

For  $f_3$ :

$$B = \{\emptyset\}$$

### 1.5.8

(a) First, show that the case  $a' \in B$  leads to a contradiction.

Suppose that  $a' \in B$ . By definition of  $B$ ,  $a' \notin f(a')$ . Therefore we have a contradiction.

(b) Now, finish the argument by showing that the case  $a' \notin B$  is equally unacceptable.

Suppose that  $a' \notin B$ . Therefore, by definition of  $B$ ,  $a' \notin B \rightarrow a' \in B$ . Therefore we have a contradiction. Thus, we cannot construct a surjective map from  $A$  to  $P(A)$ . Therefore  $A \not\sim P(A)$ .

### 1.5.9

As a final exercise, answer each of the following by establishing a 1-1 correspondence with a set of known cardinality.

(a) Is the set of all function from  $\{0, 1\}$  to  $N$  countable or uncountable?

Examples of such functions

$$f_1 = \begin{cases} \{0\} \rightarrow 5 \\ \{1\} \rightarrow 123 \end{cases} \quad (7)$$

$$f_2 = \begin{cases} \{0\} \rightarrow 7 \\ \{1\} \rightarrow 7 \end{cases} \quad (8)$$

Each of those functions we can correspond to a set of  $Q^+ = \{q \in Q : q > 0\}$ , which is countable (infinite subset of a countable set  $\rightarrow$  countable) .

To clarify my result, think of  $f(0)$  as of numerator, and  $f(1)$  as denominator.

(b) *Is the set of all function from  $N$  to  $\{0, 1\}$  countable or uncountable.*

Uncountable. Each of those function we can correspond to a sequence of 0's and 1's from Exercise 1.5.4.

(c) *Given a set  $B$ , a subset  $A$  of  $P(B)$  is called an antichain, if no element of  $A$  is a subset of any other element of  $A$ . Does  $P(N)$  contain an uncountable antichain?*

Yes, it does. We can correspond a function from  $N$  to  $\{0, 1\}$  to an antichain by adding element  $2n$  to the set, if the  $n$ 'th element of a sequence is 0, and adding  $2n - 1$  to the set, if the  $n$ 'th position is 1. Therefore, for 2 different chains we will have two different sets, each of which will be different by at least two numbers, and therefore not a subset of each other.

## 2.2.1

*Verify, using the definition of convergence of a sequence, that the following sequences converge to the proposed limit.*

(a)  $\lim_{n \rightarrow \infty} \frac{1}{6n^2+1} = 0$

Let  $\epsilon$  be arbitrary. Choose  $N \in \mathbf{N} : N > \sqrt{\frac{1}{6\epsilon} - \frac{1}{6}}$ . Let  $n \in N \geq N$ . Then

$$n > \sqrt{\frac{1}{6\epsilon} - \frac{1}{6}}$$

$$n^2 > \frac{1}{6\epsilon} - \frac{1}{6}$$

$$6n^2 > 1/\epsilon - 1$$

$$6n^2 + 1 > 1/\epsilon$$

$$\frac{1}{(6n^2 + 1)} < \epsilon$$

$$\left| \frac{1}{(6n^2 + 1)} \right| < \epsilon$$

as desired.

(b)  $\lim_{n \rightarrow \infty} \frac{3n+1}{2n+5} = \frac{3}{2}$

Let  $\epsilon$  be arbitrary. Choose  $N \in \mathbf{N} : N > \frac{13}{4\epsilon} - 10/4$ . Let  $n \in N \geq N$ . Then

$$n > \frac{13}{4\epsilon} - 10/4$$

$$4n > \frac{13}{\epsilon} - 10$$

$$4n + 10 > \frac{13}{\epsilon}$$

$$\frac{13}{4n + 10} < \epsilon$$

$$|\frac{13}{4n + 10}| < \epsilon$$

$$|\frac{-13}{4n + 10}| < \epsilon$$

$$|\frac{6n + 2 - 6n - 15}{4n + 10}| < \epsilon$$

$$|\frac{2(3n + 1) - 3(2n + 5)}{2(2n + 5)}| < \epsilon$$

$$|\frac{3n + 1}{2n + 5} - \frac{3}{2}| < \epsilon$$

as desired.

$$(c) \lim_{\sqrt{n+3}} \frac{2}{\sqrt{n+3}} = 0$$

Let  $\epsilon$  be arbitrary. Choose  $N \in \mathbf{N} : N > \frac{2^2}{\epsilon} - 3$ . Let  $n \in \mathbf{N} \geq N$ . Then

$$n > \frac{2^2}{\epsilon} - 3$$

$$n + 3 > \frac{2^2}{\epsilon}$$

$$\sqrt{n + 3} > \frac{2}{\epsilon}$$

$$\frac{2}{\sqrt{n + 3}} < \epsilon$$

$$|\frac{2}{\sqrt{n + 3}}| < \epsilon$$

as desired.



## 2.2.2

What happens if we reverse the order of the quantifiers on Definition 2.2.3?

*Definition:* A sequence  $(x_n)$  verconges to  $x$  if there exists an  $\epsilon > 0$  such that for all  $N \in \mathbf{N}$  it is true that  $n \geq N$  implies  $|x_n - x| < \epsilon$ .

Give an example of a vercongent sequence. Can you give an example of a vercongent sequence, that is divergent? What exactly is being described in this strange definition?

An example of a vercongent sequence:

$$(x_n) = 5$$

An example of a vercongent sequence, that is divergent:

$$(x_n) = (-1)^n$$

Here described a bounded sequence (i.e.  $|(x_n)| < M$  for some  $M > 0 \in \mathbf{R}$ )

## 2.2.3

Describe what we would have to demonstrate in order to disprove each of the following statements.

(a) At every college in the United States, there is a student who is at least seven feet tall

There exist a college, where every student is shorter than 7 feet.

(b) For all colleges in the United States, there exists a professor who gives every student a grade of either A or B

There exist a college, where every professor gives C or less every time.

(c) There exist a college in the United States where every student is at least six feet tall

In all colleges across US there exists a student, who is shorter than 6 feet.

## 2.2.4

Argue that the sequence

$$1, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 0, 0, 0, 1, (5 \text{ zeroes}), 1, \dots$$

does not converge to zero. For what values of  $\epsilon > 0$  does there exist a response  $N$ ? For which values of  $\epsilon > 0$  is there no suitable response?

If we set  $\epsilon = 0.5$ , then from time to time elements will get out of the desired range.

For values  $\epsilon > 1$  there always exist a suitable  $N$ .

For values  $0 < \epsilon \leq 0$  there exists no response.

## 2.2.5

Let  $\lfloor x \rfloor$  be the greatest integer less than or equal to  $x$ . For example,  $\lfloor \pi \rfloor = 3$  and  $\lfloor 3 \rfloor = 3$ . Find  $\lim a_n$  and supply proofs for each conclusion if

$$(a) a_n = \lfloor 1/n \rfloor$$

$$\begin{aligned} \forall n \in N \rightarrow 0 < 1/n \leq 1 \\ n \in N \geq 2 \rightarrow 0 < 1/n < 1 \end{aligned}$$

Therefore for  $n > 2$

$$\forall \epsilon > 0 \in R : \lfloor \lfloor 1/n \rfloor \rfloor = \lfloor 1/n \rfloor = 0 < \epsilon$$

as desired.

$$(b) a_n = \lfloor (10 + n)/2n \rfloor$$

Let  $n \geq 10$ . Then

$$\begin{aligned} n &> 10 \\ 2n &> 20 \\ \frac{10}{2n} &< \frac{1}{2} \\ 0 &< \frac{10}{2n} + \frac{1}{2} < 1 \\ \lfloor \frac{10}{2n} + \frac{1}{2} \rfloor &= 0 \end{aligned}$$

Thus for every  $\epsilon > 0$  we can pick  $N = 10$  and it follows that

$$\lfloor \lfloor (10 + n)/2n \rfloor \rfloor < 0$$

Therefore  $(a_n) \rightarrow 0$ , as desired.

Reflecting on these examples, comment on the statement following Definition 2.2.3 that "the smaller the  $\epsilon$ -neighborhood, the larger  $N$  may need to be".

The key word here is "may". It may have to be larger, it may not. In some cases one value works for all of the  $\epsilon$ 's. I bet my money on the fact, that it can be rigorously proven, that this is the case only for the sequences, that are constant after some term, but I'll skip that

## 2.2.6

Suppose that for a particular  $\epsilon > 0$  we have found a suitable value of  $N$  that "works" for a given sequence in the sense of Definition 2.2.3.

(a) Then, any larger/smaller (pick one)  $N$  will also work for the same  $\epsilon > 0$ .

Larger. This fact follows from definition.

(b) Then, this same  $N$  will also work for any larger/smaller value of  $\epsilon$ .

Larger.  $x \in V_\epsilon \rightarrow x \in V_{\epsilon+s}$

## 2.2.7

*Informally speaking, the sequence  $\sqrt{n}$  "converges to infinity".*

*(a) Imitate the logical structure of Definition 2.2.3 to create a rigorous definition for the mathematical statement  $\lim x_n = \infty$ . Use this definition to prove  $\lim \sqrt{n} = \infty$ .*

**Definition of convergence to infinity** A sequence  $(a_n)$  converges for infinity if, for every  $\epsilon \in R$ , there exists an  $N \in \mathbf{N}$  such that whenever  $n \geq N$  it follows that  $a_n > \epsilon$ .

I relaxed a bit statement about the  $\epsilon$ , namely substituted  $\epsilon > 0$  to  $\epsilon \in R$ . In case you are wondering, Both cases are the equivalent.

Let  $\epsilon \in R$  be arbitrary. Then we can pick  $N \in \mathbf{N} : N > \epsilon^2$ . Then for  $n \in \mathbf{N} > N$ :

$$n > \epsilon^2$$

$$\sqrt{n} > \epsilon$$

Therefore the sequence converges to infinity

*(b) What does your definition in (a) say about the particular sequence  $(1, 0, 2, 0, 3, 0, 4, 0, 5, 0, \dots)$*

It says, that it doesn't converge to infinity (specifically for the case  $\epsilon > 0$ )

## 2.2.8

*Here are two useful definitions:*

*(i) A sequence  $(a_n)$  is eventually in a set  $A \subseteq R$  if there exists  $N \in \mathbf{N}$  such that  $a_n \in A$  for all  $n \geq N$ .*

*(ii) A sequence  $(a_n)$  is frequently in a set  $A \subseteq R$  if for every  $N \in \mathbf{N}$  there exists  $n \geq N$  such that  $a_n \in A$ .*

*(a) Is the sequence  $(-1)^n$  eventually or frequently in the set  $\{1\}$ ?*

Frequently

*(b) Which definition is stronger? Does frequently imply eventually or does eventually imply frequently?*

Eventually is stronger. Eventually implies frequently.

*(c) Give an alternative rephrasing of Definition 2.2.3B using either frequently or eventually. Which is the term we want?*

We want to use eventually

A sequence  $(a_n)$  converges to  $a$  if, given any  $\epsilon$ -neighborhood  $V_\epsilon(a)$  of  $a$ , sequence is eventually in  $V_\epsilon(a)$ .

*(d) Suppose an infinite number of terms of a sequence  $(x_n)$  are equal to 2. Is  $(x_n)$  necessarily eventually in the interval  $(1.9, 2.1)$ ? Is it frequently in  $(1.9, 2.1)$ ?*

It is not necessarily eventually in  $(1.9, 2.1)$ . Example:

$$(1, 2, 2, 2, 3, 2, 4, 2, 5, 2, 6, 2, \dots)$$

It is indeed frequently in (1.9, 2.1). This stems from the fact, that before  $a_n$ , there exist only a finite number of elements before it. Therefore 2 is bound to be met again at some time.

### 2.3.1

Show that the constant sequence  $(a, a, a, \dots)$  converges to  $a$ .

$$|a_n - a| = |a - a| = 0$$

therefore for every  $n \in \mathbf{N}$  it is true, that  $|a_n - a| < \epsilon$ . Therefore  $(a_n) \rightarrow a$ , as desired.

### 2.3.2

Let  $x_n \geq 0$  for all  $n \in \mathbf{N}$ .

(a) If  $(x_n) \rightarrow 0$ , show that  $(\sqrt{x_n}) \rightarrow 0$

Suppose that  $(x_n) \rightarrow 0$ . Then

$$\forall \epsilon \in \mathbf{R} > 0 : \exists N \in \mathbf{N} : \forall n \in \mathbf{N} : n \geq N : |x_n - 0| < \epsilon$$

.

$$|x_n - 0| < \epsilon$$

$$|x_n| < \epsilon$$

$$x_n < \epsilon$$

$$\sqrt{x_n} < \sqrt{\epsilon}$$

therefore

$$\forall \epsilon \in \mathbf{R} > 0 : \exists \epsilon' = \sqrt{\epsilon} > 0 : \exists N \in \mathbf{N} : \forall n \in \mathbf{N} : n \geq N : |\sqrt{x_n} - 0| < \epsilon'$$

. therefore

$$\forall \epsilon \in \mathbf{R} > 0 : \exists N \in \mathbf{N} : \forall n \in \mathbf{N} : n \geq N : |\sqrt{x_n} - 0| < \epsilon$$

. From which it follows that  $(\sqrt{x_n}) \rightarrow 0$  by definition of a limit.

(b) If  $(x_n) \rightarrow x$ , show that  $(\sqrt{x_n}) \rightarrow \sqrt{x}$

Suppose that  $(x_n) \rightarrow x$ . Then

$$\forall \epsilon \in \mathbf{R} > 0 : \exists N \in \mathbf{N} : \forall n \in \mathbf{N} : n \geq N : |x_n - x| < \epsilon$$

.

$$|x_n - x| < \epsilon$$

$$|(\sqrt{x_n} + \sqrt{x})(\sqrt{x_n} - \sqrt{x})| < \epsilon$$

$$|\sqrt{x_n} - \sqrt{x}| < \frac{\epsilon}{|\sqrt{x_n} + \sqrt{x}|}$$

$$|\sqrt{x_n} + \sqrt{x}| \leq |\sqrt{x_n}| + |\sqrt{x}| = \sqrt{x_n} + \sqrt{x} \rightarrow \frac{\epsilon}{|\sqrt{x_n} + \sqrt{x}|} < \frac{\epsilon}{\sqrt{x_n} + \sqrt{x}}$$

$$|\sqrt{x_n} - \sqrt{x}| < \frac{\epsilon}{\sqrt{x_n} + \sqrt{x}}$$

$(x_n)$  is convergent and therefore bounded. Therefore there exists  $M \in \mathbb{R} > 0 : x_n < M$ . Therefore  $\sqrt{x_n} < \sqrt{M}$ . Thus  $\sqrt{x_n} + \sqrt{x} < \sqrt{M} + \sqrt{x}$ . Therefore

$$|\sqrt{x_n} - \sqrt{x}| < \frac{\epsilon}{\sqrt{M} + \sqrt{x}}$$

therefore

$$\forall \epsilon \in \mathbb{R} > 0 : \exists \epsilon_1 = \frac{\epsilon}{\sqrt{M} + \sqrt{x}} > 0 : \exists N \in \mathbf{N} : \forall n \in \mathbb{N} : n > N : |\sqrt{x_n} - \sqrt{x}| < \epsilon$$

. therefore

$$\forall \epsilon \in \mathbb{R} > 0 : \exists N \in \mathbf{N} : \forall n \in \mathbb{N} : n > N : |\sqrt{x_n} - \sqrt{x}| < \epsilon$$

. From which it follows that  $(\sqrt{x_n}) \rightarrow x$  by definition of a limit.

### 2.3.3 (Squeeze Theorem).

Show that if  $x_n \leq y_n \leq z_n$  for all  $n \in \mathbb{N}$ , and if  $\lim x_n = \lim z_n = l$ , then  $\lim y_n = l$  as well.

$$\forall n \in \mathbb{N} : x_n \leq y_n \rightarrow \lim x_n \leq \lim y_n$$

$$\forall n \in \mathbb{N} : y_n \leq z_n \rightarrow \lim y_n \leq \lim z_n$$

therefore

$$\forall n \in \mathbb{N} : x_n \leq y_n \leq z_n \rightarrow \lim x_n \leq \lim y_n \leq \lim z_n$$

Therefore

$$\lim x_n = \lim z_n = l \rightarrow \lim y_n = l$$

as desired.

### 2.3.4

Show that limits, if they exist, must be unique. In other words, assume  $\lim a_n = l_1$  and  $\lim a_n = l_2$ , and prove that  $l_1 = l_2$ .

We will proceed with a proof by contradiction.

Suppose  $l_1 \neq l_2$ . Then

$$\forall \epsilon > 0 : \exists N \in \mathbb{N} : n \geq N \rightarrow |a_n - l_1| \leq \epsilon$$

$$\forall \epsilon > 0 : \exists N \in \mathbb{N} : n \geq N \rightarrow |a_n - l_2| \leq \epsilon$$

Let  $\epsilon = |l_1 - l_2|/2$ . Then  $\exists N_1 \in \mathbb{N}$  such that  $n_1 \geq N$  implies that

$$|a_{n_1} - l_1| < |l_1 - l_2|/2$$

also there exists  $N_2 \in \mathbb{N}$  such that  $n_2 \geq N_2$  implies that

$$|a_{n_2} - l_2| < |l_1 - l_2|$$

Let  $n = \max\{n_1, n_2\}$ . Then

$$|a_n - l_1| < |l_1 - l_2|/2$$

$$|a_n - l_2| < |l_1 - l_2|/2$$

$$|a_n - l_1| + |a_n - l_2| < |l_1 - l_2|$$

$$|a_n - l_1| + |l_2 - a_n| < |l_1 - l_2|$$

then by triangular inequality

$$|a_n - l_1 + l_2 - a_n| \leq |a_n - l_1| + |l_2 - a_n| < |l_1 - l_2|$$

$$|a_n - l_1 + l_2 - a_n| < |l_1 - l_2|$$

$$|-l_1 + l_2| < |l_1 - l_2|$$

$$|l_1 - l_2| < |l_1 - l_2|$$

which is a contradiction. Therefore

$$\lim a_n = l_1 \text{ and } \lim a_n \rightarrow l_2 \rightarrow l_1 = l_2$$

as desired.

### 2.3.5

Let  $(x_n)$  and  $(y_n)$  be given, and define  $(z_n)$  to be the "shuffled" sequence  $(x_1, y_1, x_2, y_2, x_3, y_3, \dots)$ . Prove that  $(z_n)$  is convergent if and only if  $(x_n)$  and  $(y_n)$  are both convergent with  $\lim x_n = \lim y_n$ .

**In the forward direction:**

Suppose that  $(z_n)$  is convergent to some  $l \in R$ . Then

$$\forall \epsilon > 0 : \exists N \in \mathbf{N} : n \geq N \rightarrow |z_n - l| < \epsilon$$

Because  $(z_n)$  is "shuffled" we can follow that

$$\forall \epsilon > 0 : \exists N \in \mathbf{N} : (n+1)/2 \geq N \rightarrow |x_{(n+1)/2} - l| < \epsilon$$

$$\forall \epsilon > 0 : \exists N \in \mathbf{N} : n/2 \geq N \rightarrow |y_{n/2} - l| < \epsilon$$

Therefore if we let  $m_1 = (n+1)/2$  and  $m_2 = n/2$

$$\forall \epsilon > 0 : \exists N \in \mathbf{N} : m_1 \geq N \rightarrow |x_{m_1} - l| < \epsilon$$

$$\forall \epsilon > 0 : \exists N \in \mathbf{N} : m_2 \geq N \rightarrow |y_{m_2} - l| < \epsilon$$

Therefore  $\lim x_n = \lim y_n = l$ , as desired.

**In the backward direction:**

Suppose that both  $(x_n)$  and  $(y_n)$  are convergent to some  $l$ . Then

$$\forall \epsilon > 0 : \exists N_1 \in \mathbf{N} : m_1 \geq N_1 \rightarrow |x_{m_1} - l| < \epsilon$$

$$\forall \epsilon > 0 : \exists N_2 \in \mathbf{N} : m_2 \geq N_2 \rightarrow |y_{m_2} - l| < \epsilon$$

If we pick  $N = (\max\{N_1, N_2\} + 1) * 2$ , then

$$\forall \epsilon > 0 : \exists N \in \mathbf{N} : n \geq N \rightarrow |x_{(n-1)/2} - l| < \epsilon$$

$$\forall \epsilon > 0 : \exists N \in \mathbf{N} : m_2 \geq N \rightarrow |y_{n/2} - l| < \epsilon$$

If  $n$  is odd then  $z_n = x_{(n-1)/2}$ , and if  $n$  is even then  $z_n = y_{n/2}$ . Therefore

$$\forall \epsilon > 0 : \exists N \in \mathbf{N} : n \geq N \rightarrow |z_n - l| < \epsilon$$

Thus  $(z_n) \rightarrow l$  as well, as desired.

### 2.3.6

Show that if  $(b_n) \rightarrow b$ , then the sequence of absolute values  $|b_n|$  converges to  $|b|$ .

As proven in 1.2.5

$$||a| - |b|| \leq |a - b|$$

Therefore  $||b_n| - |b|| \leq |b_n - b|$  Thus

$$\forall \epsilon > 0 : \exists N \in \mathbf{N} : n \geq N \rightarrow ||b_n| - |b|| < \epsilon$$

Thus  $(|b_n|) \rightarrow |b|$ , as desired.

(b) Is the converse of part (a) true? If we know that  $|b_n| \rightarrow |b|$ , can we deduce that  $(b_n) \rightarrow b$ ?

No. Glaring example is

$$a_n = (-1)^n$$

### 2.3.7

(a) Let  $(a_n)$  be a bounded (non necessarily convergent) sequence, and assume  $\lim b_n = 0$ . Show that  $\lim(a_nb_n) = 0$ . Why are not allowed to use Algebraic Limit Theorem to prove that?

Let  $M > |a_n|$  for all  $n \in N$  (it exists because of the boundness of  $a_n$ ). Then

$$|a_nb_n| = |a_n||b_n| \leq M|b_n|$$

$$|a_nb_n| \leq M|b_n|$$

$$-M|b_n| \leq a_nb_n \leq M|b_n|$$

Both  $(M|b_n|)$  and  $(-M|b_n|)$  converge to 0. Therefore  $(a_nb_n)$  converges to 0 as well by Squeeze theorem.

A little sidenote: this statement can be proven with the standart definition of limit as well, but that proof is longer and I wanted to use brand-new, self-proven theorem on this one.

We can't use Algebraic Limit Theorem on that one beacause it prerequisites that both sequences are convergent, and here it is not the case.

(b) Can we conclude anything about the convergence of  $(a_nb_n)$  if we assume that  $(b_n)$  converges to some nonzero limit  $b$ ?

We can probably conclude that if  $(a_n)$  is divergent, then  $(a_nb_n)$  is divergent as well, but we can't conclude nothing definitive about just a bounded sequence.

(c) Use (a) for prove Theoremt 2.3.3, part (iii), for the case when  $a = 0$ .

If  $(b_n)$  is convergent (possibly to zero), then it is bounded. Therefore, by (a), we can state that

$$(a_nb_n) \rightarrow ab = 0$$



But wait a second, we used something from the algebraic property in part (a)! Doesn't in mean, that we proved a theorem by assuming, that it is true? No. In part (a) we concluded, that  $M|b_n|$  is convergent to 0 by using part (i) of algebraic limit theorem. We didn't even use the case when  $M = 0$ , and therefore we can sleep well.

### 2.3.8

*Give an example of each of the following, or state that such a request is impossible by referencing the proper theorem(s):*

(a) *sequences  $(x_n)$  and  $(y_n)$ , which both diverge, but whose sum  $(x_n + y_n)$  converges*

$$(x_n) = n$$

$$(y_n) = -n$$

(b) *sequences  $(x_n)$  and  $(y_n)$ , where  $(x_n)$  converges,  $(y_n)$  diverges, and  $(x_n + y_n)$  converges*

Impossible.

I want to say, that the algebraic limit theorem prevents this statement to be true, but I don't know if we can apply it here.

Suppose that  $(x_n + y_n)$  converges to  $l$ . Let  $(ln) = l$  be a constant sequence. Then

$$(x_n + y_n - l) \rightarrow 0$$

Also, because both  $(x_n)$  and  $(l)$  converge,  $(l - x_n)$  converges as well. therefore

$$(x_n + y_n - l + l - x_n)$$

converges as well. Therefore  $(y)$  converges, which is a contradiction.

Turns out that yeah, it applies.

(c) *a convergent sequence  $(b_n)$  with  $b_n \neq 0$  for all  $n$  such that  $(1/b_n)$  diverges*

$$(b_n) = 1/n$$

(d) *an unbounded sequence  $(a_n)$  and a convergent sequence  $(b_n)$  with  $(a_n - b_n)$  bounded;*  
Impossible.

Same algebraic limit theorem with a bit of convergence to infinity

(e) *Two sequences  $(a_n)$  and  $(b_n)$ , where  $(a_n b_n)$  and  $(a_n)$  converge, but  $(b_n)$  diverge.*

$$(a_n) = 1/n$$

$$(b_n) = (-1)^n$$

### 2.3.9

Does Theorem 2.3.4 remain true, if all of the inequalities are assumed to be strict? If we assume, for instance, that a convergent sequence  $(x_n)$  satisfies  $x_n > 0$  for all  $n \in \mathbb{N}$ , what we may conclude about the limit?

If we swap all inequalities to strict inequalities, then the theorem is false. Example:  $(b_n) = 1/n \rightarrow 0$ , but  $b_n > 0$  for all  $n$ 's.

### 2.3.10

If  $(a_n) \rightarrow 0$  and  $|b_n - b| \leq a_n$ , then show that  $(b_n) \rightarrow b$ .

$(a_n) \rightarrow 0$  implies, that for every  $\epsilon \in \mathbb{R} > 0$  we can get  $|a_n| < \epsilon$ . Therefore

$$\forall \epsilon \in \mathbb{R} > 0 : \exists N \in \mathbb{N} : \forall n \geq N \rightarrow |b_n - b| < a_n \leq |a_n| < \epsilon$$

$$\forall \epsilon \in \mathbb{R} > 0 : \exists N \in \mathbb{N} : \forall n \geq N \rightarrow |b_n - b| < \epsilon$$

Therefore  $(b_n) \rightarrow b$ , as desired.

### 2.3.11 (Cesaro Means).

Show that if  $(x_n)$  is a convergent sequence, then the sequence given by the averages

$$y_n = \frac{x_1 + x_2 + \dots + x_n}{n}$$

also converges to the same limit.

The strategy for the proof for this theorem is to first show, that this statement is true for all of the sequences, that converge to 0. Then we'll show that this case is equivalent to the theorem in general.

First, let's prove that this is true for the case when  $(a_n) \rightarrow 0$

Strategy here is to show, that

Suppose that we have a given  $\epsilon \in \mathbb{R} > 0$ . Then there exists  $q \in \mathbb{N}$  such that  $|a_q| < \epsilon$ . Also, because of the fact, that  $(a_n) \rightarrow 0$  there exist  $k \in \mathbb{N} \geq q$  such that  $|a_k| \geq |a_q|$  (i.e. maximum element of this set) for all  $q_1 \in \mathbb{N} \geq q$ .

Then, let

$$N > \frac{|a_1| + |a_2| + \dots + |a_{q-1}| - q|a_k|}{\epsilon - |a_k|}$$

If, for some god forsaken reason,  $q \geq N$ , then we can set  $N$  to be more than  $q$ , and it wouldn't make the difference. Same applies to  $k$ , if it matters (it doesn't).

Therefore if we let  $n > N$ , then we can conclude that

$$n > \frac{|a_1| + |a_2| + \dots + |a_{q-1}| - q|a_k|}{\epsilon - |a_k|}$$

$$\begin{aligned}
& \frac{|a_1| + |a_2| + \dots + |a_{q-1}| - q|a_k|}{\epsilon - |a_k|} < n \\
& |a_1| + |a_2| + \dots + |a_{q-1}| - q|a_k| < \epsilon n - |a_k|n \\
& |a_1| + |a_2| + \dots + |a_{q-1}| + |a_k|n - q|a_k| < \epsilon n \\
& \frac{|a_1| + |a_2| + \dots + |a_{q-1}|}{n} + |a_k| - \frac{q|a_k|}{n} < \epsilon \\
& \frac{|a_1| + |a_2| + \dots + |a_{q-1}|}{n} + \frac{n|a_k|}{n} - \frac{q|a_k|}{n} < \epsilon \\
& \frac{|a_1| + |a_2| + \dots + |a_{q-1}|}{n} + \frac{n|a_k| - q|a_k|}{n} < \epsilon \\
& \frac{|a_1| + |a_2| + \dots + |a_{q-1}|}{n} + \frac{(n - q)|a_k|}{n} < \epsilon \\
& \frac{|a_1| + |a_2| + \dots + |a_{q-1}|}{n} + \frac{|a_k| + \dots + |a_k|}{n} < \epsilon
\end{aligned}$$

because  $|a_{q+}| \leq |a_k|$

$$\begin{aligned}
& \frac{|a_1| + |a_2| + \dots + |a_{q-1}|}{n} + \frac{|a_q| + \dots + |a_n|}{n} \leq \frac{|a_1| + |a_2| + \dots + |a_{q-1}|}{n} + \frac{|a_k| + \dots + |a_k|}{n} < \epsilon \\
& \frac{|a_1| + |a_2| + \dots + |a_{q-1}| + |a_q| + \dots + |a_n|}{n} < \epsilon \\
& \frac{|a_1| + |a_2| + \dots + |a_n|}{n} < \epsilon \\
& \left| \frac{a_1 + a_2 + \dots + a_n}{n} \right| = \frac{|a_1 + a_2 + \dots + a_n|}{n} \leq \frac{|a_1| + |a_2| + \dots + |a_n|}{n} < \epsilon \\
& \left| \frac{a_1 + a_2 + \dots + a_n}{n} \right| < \epsilon \\
& \left| \frac{a_1 + a_2 + \dots + a_n}{n} - 0 \right| < \epsilon
\end{aligned}$$

Therefore

$$\forall \epsilon \in R > 0 : \exists N \in \mathbf{N} : \forall n > N \rightarrow \left| \frac{a_1 + a_2 + \dots + a_n}{n} - 0 \right| < \epsilon$$

Therefore  $(a_n) \rightarrow 0$  implies that  $(\frac{a_1 + a_2 + \dots + a_n}{n}) \rightarrow 0$  as well.

Now, proud of our achievement, we can proceed with the initial argument. Suppose that  $(x_n) \rightarrow l$ . Then if we set  $l_n = l$ , then  $(l_n) \rightarrow l$ . Therefore  $(x_n - l_n) \rightarrow 0$ . And because of it, we can use our initial conclusion.

$$\left( \frac{x_1 - l_n + x_2 - l_n + \dots + x_n - l_n}{n} \right) \rightarrow 0$$

$$\begin{aligned}
& \left( \frac{x_1 + x_2 + \dots + x_n - l_n n}{n} \right) \rightarrow 0 \\
& \left( \frac{x_1 + x_2 + \dots + x_n}{n} - \frac{l_n n}{n} \right) \rightarrow 0 \\
& \left( \frac{x_1 + x_2 + \dots + x_n}{n} - l_n \right) \rightarrow 0
\end{aligned}$$

Therefore

$$\left( \frac{x_1 + x_2 + \dots + x_n}{n} \right) \rightarrow l$$

When push comes to shove, we can prove it axiomatically through

$$\begin{aligned}
& \left| \frac{x_1 - l_n + x_2 - l_n + \dots + x_n - l_n}{n} \right| < \epsilon \\
& \left| \frac{x_1 + x_2 + \dots + x_n - n l_n}{n} \right| < \epsilon \\
& \left| \frac{x_1 + x_2 + \dots + x_n}{n} - \frac{n l_n}{n} \right| < \epsilon \\
& \left| \frac{x_1 + x_2 + \dots + x_n}{n} - l_n \right| < \epsilon \\
& \left| \frac{x_1 + x_2 + \dots + x_n}{n} - l \right| < \epsilon
\end{aligned}$$

Which comes to the same conclusion. Therefore now we can affirmatively state that

$$(x_n) \rightarrow l \text{ implies } \frac{x_1 + x_2 + \dots + x_n}{n} \rightarrow l$$

as desired.

*Give an example to show that it is possible for the sequence  $(y_n)$  of averages to converge even if  $(x_n)$  does not.*

$$x_n = ([[n/2]])(-1)^n$$

Where  $[[\ ]]$  is a floor function. Therefore

$$x_n = (0, 0, 1, -1, 2, -2, \dots)$$

## 2.3.12

Consider the doubly indexed array  $a_{m,n} = m/(m+n)$ .

(a) Intuitively speaking, what should  $\lim_{m,n \rightarrow \infty} a_{m,n}$  represent? Compute the "iterated" limits

$$\lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} a_{m,n} \text{ and } \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} a_{m,n}$$

Intuitively speaking  $\lim_{m,n \rightarrow \infty} a_{m,n}$  should mean that the larger the  $m$  and  $n$  get, the closer  $a_{m,n}$  becomes.

$$\begin{aligned} \lim_{n \rightarrow \infty} a_{m,n} &= \lim_{n \rightarrow \infty} m/(m+n) = \lim_{n \rightarrow \infty} m/m + m/n = 1 \\ \lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} a_{m,n} &= \lim_{m \rightarrow \infty} 1 = 1 \end{aligned}$$

$$\begin{aligned} \lim_{m \rightarrow \infty} a_{m,n} &= \lim_{m \rightarrow \infty} m/(m+n) = \lim_{m \rightarrow \infty} m/m + m/n = \infty \\ \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} a_{m,n} &= \lim_{n \rightarrow \infty} \infty = \infty \end{aligned}$$

(b) Formulate a rigorous definition in the style of Definition 2.2.3 for the statement

$$\lim_{m,n \rightarrow \infty} a_{m,n} = l$$

A list is a set of length  $n$  in form

$$\{a_1, \{a_1, a_2\}, \{a_1, a_2, a_3\}, \dots, a_{n-1}, a_n\}\}\dots$$

It is denoted by

$$(a_1, a_2, a_3, \dots, a_n)$$

A double indexed sequence  $a_{m,n}$  is a function from the list  $(j, k)$ , where  $j, k \in \mathbf{N}$  to  $\mathbf{R}$ .

A sequence  $(a_{m,n})$  converges to a real number  $a$  if, for every positive number  $\epsilon$ , there exist  $N_1, N_2 \in \mathbf{N}$  such that whenever  $m > N_1$  and  $n > N_2$  it follows that  $|a_{m,n} - a| < \epsilon$ .

## 2.4.1

Complete the proof of Theorem 2.4.6 by showing that if the series  $\sum_{n=0}^{\infty} 2^n b_{2^n}$  diverges, then so does  $\sum_{n=1}^{\infty} b_n$ . Example 2.4.5 may be a useful reference.

First of all, let's state Theorem 2.4.6

**Theorem 2.4.6 (Cauchy condensation Test)** Suppose  $(b_n)$  is decreasing and satisfies  $b_n \geq 0$  for all  $n \in \mathbf{N}$ . Then, the series  $\sum_{n=1}^{\infty} b_n$  converges if and only if the series

$$\sum_{n=0}^{\infty} 2^n b_{2^n} = b_1 + 2b_2 + 4b_4 + 8b_8 + \dots$$

converges.

Suppose that  $\sum_{n=0}^{\infty} 2^n b_{2^n}$  diverges. Then a partial sum

$$s_{2^k} = b_1 + 2b_2 + 4b_4 + \dots = b_1 + b_2 + b_2 + b_4 + b_4 + b_4 + \dots \leq b_1 + b_2 + b_3 + b_4 + b_5 + b_6 + b_7 + \dots = s_n$$

Therefore, partial sum for  $\sum_{n=0}^{\infty} 2^n b_{2^n}$  is less than  $\sum_{n=1}^{\infty} b_n$ . Therefore  $\sum_{n=1}^{\infty} b_n$  diverges, as desired.

## 2.4.2

(a) Prove that the sequence defined by  $x_1 = 3$  and

$$x_{n+1} = \frac{1}{4 - x_n}$$

converges.

Firstly, let us write a couple of elements of this sequence to see what's going on:

$$(x_n) = 3, 1, 1/3, 3/11, \dots$$

Those elements of the sequence tell us, that the sequence is probably decreasing and also bounded below by some constant around 0.25, and therefore bounded below by 2. Also, the proof of the fact that the sequence is bounded above by 4 might come in handy.

To prove the proposition, that this sequence is convergent we will use property, that if a sequence is bounded and monotone, then it is convergent. Let's start with proposition that the this sequence is decreasing.

We'll use induction for this one, because it seems right;

Base:  $x_1 < 4$ .

Proposition:  $x_n < 4$

Step:

$$x_{n+1} = \frac{1}{4 - x_n} < 4$$

$$4 - x_n > \frac{1}{4}$$

$$-x_n > \frac{1}{4} - 4$$

$$x_n < 3\frac{1}{4}$$

Therefore  $x_n < 4$ . From this we can conclude that

$$x_n < 4$$

$$\begin{aligned}
4 - x_n &> 0 \\
\frac{1}{4 - x_n} &> 0 \\
x_{n+1} &> 0
\end{aligned}$$

Now let us prove, that the sequence is decreasing. We'll also do it with induction.

Base:  $x_1 - x_2 = 3 - 1 = 2 > 0$

Proposition:  $x_{n-1} - x_n > 0$ .

Step: We had proved already, that  $x_n > 0$ , and  $x_n < 4$  Therefore

$$\begin{aligned}
x_n - x_{n+1} &> 0 \\
x_n - \frac{1}{4 - x_n} &> 0 \\
(4 - x_n)x_n - 1 &> 0 \\
4x_n - x_n^2 - 1 &> 0 \\
x_n^2 - 4x_n + 1 &< 0 \\
(x_n - (2 - \sqrt{3}))(x_n - (2 + \sqrt{3})) &< 0
\end{aligned}$$

It can be shown, that our sequence bounded by  $2 - \sqrt{3}$  and  $2 + \sqrt{3}$ .

Thus the sequence is decreasing and bounded, and therefore convergent.

(b) Now that we know  $\lim x_n$  exists, explain why  $\lim x_{n+1}$  must also exist and equal the same value.

By the definition of limit of a sequence, for every  $\epsilon$  there exist an  $N$  such that for all  $n \geq N$   $|x_n - l| < \epsilon$ . Therefore, because every  $(x_{n+1}) \subset (x_n)$  we can conclude, that  $(x_{n+1})$  converges to the same value.

(c) Take the limit of each side of the recursive equation in part (a) of this exercise to explicitly compute  $\lim x_n$ .

$$\begin{aligned}
\lim x_{n+1} &= \frac{1}{4 - \lim x_n} \\
\lim x_n &= \frac{1}{4 - \lim x_n} \\
\lim x_n &= \frac{1}{4 - \lim x_n} \\
(4 - \lim x_n) \lim x_n &= 1 \\
4 \lim x_n - (\lim x_n)^2 - 1 &= 0 \\
(\lim x_n)^2 - 4 \lim x_n + 1 &= 0
\end{aligned}$$

From my 5th grade class I remember that.

$$\lim x_n = 2 - \sqrt{3} \text{ or } \lim x_n = 2 + \sqrt{3}$$

Given that  $2 + \sqrt{3} > 3$  and that the sequence is decreasing and first element of it is 3, we can state that  $x_n > 3$  for all  $n$ . Therefore

$$\lim x_n = 2 - \sqrt{3}$$

### 2.4.3

Following the model of Exercise 2.4.2, show that the sequence defined by  $y_1 = 1$  and  $y_{n+1} = 4 - 1/y_n$ .

Let's write down a few elements of this sequence:

$$y_1 = 1, 3, 11/3, 41/11, \dots$$

This one is probably increasing and bounded above by 4 and below by 1

Base:  $1 \leq y_1 = 1 \leq 4$

Proposition:  $1 \leq y_n \leq 4$

Step:

$$1 \leq y_n \leq 4$$

$$1 \geq 1/y_n \geq 1/4$$

$$-1 \leq -1/y_n \leq -1/4$$

$$3 \leq 4 - 1/y_n \leq 3\frac{3}{4}$$

$$1 \leq 4 - 1/y_n \leq 4$$

$$1 \leq y_{n+1} \leq 4$$

$$1 \leq y_n \leq 2 + \sqrt{3}$$

$$1 \geq 1/y_n \geq \frac{1}{2 + \sqrt{3}}$$

$$-1 \leq -1/y_n \leq -\frac{1}{2 + \sqrt{3}}$$

$$3 \leq 4 - 1/y_n \leq 4 - \frac{1}{2 + \sqrt{3}}$$



$$3 \leq 4 - 1/y_n \leq \frac{8 + 4\sqrt{3} - 1}{2 + \sqrt{3}}$$

$$3 \leq 4 - 1/y_n \leq \frac{7 + 4\sqrt{3}}{2 + \sqrt{3}}$$

$$3 \leq 4 - 1/y_n \leq \frac{(2 + \sqrt{3})^2}{2 + \sqrt{3}}$$

$$3 \leq 4 - 1/y_n \leq 2 + \sqrt{3}$$

as desired.

Therefore  $1 \leq y_n \leq 4$  for all  $n \in N$ .

Now, let us try to prove that the sequence is increasing

$$y_{n+1} \geq y_n$$

$$y_{n+1} - y_n \geq 0$$

$$4 - 1/y_n - y_n \geq 0$$

$$4y_n - 1 - y_n^2 \geq 0$$

$$y_n^2 - 4y_n + 1 \leq 0$$

$$(y - (2 + \sqrt{3}))(y - (2 - \sqrt{3})) \leq 0$$

Therefore if we prove that the sequence is bounded by  $2 + \sqrt{3}$  and  $2 - \sqrt{3}$ , then we'll have our proof

Base:  $2 - \sqrt{3} \leq y_1 = 1 \leq 2 + \sqrt{3}$

Proposition:  $2 - \sqrt{3} \leq y_n \leq 2 + \sqrt{3}$

Step:

$$2 - \sqrt{3} \leq y_n \leq 2 + \sqrt{3}$$

$$\frac{1}{2 - \sqrt{3}} \geq 1/y_n \geq \frac{1}{2 + \sqrt{3}}$$

$$-\frac{1}{2 - \sqrt{3}} \leq -1/y_n \leq -\frac{1}{2 + \sqrt{3}}$$

$$4 - \frac{1}{2 - \sqrt{3}} \leq 4 - 1/y_n \leq 4 - \frac{1}{2 + \sqrt{3}}$$

$$\frac{7 - 4\sqrt{3}}{2 - \sqrt{3}} \leq y_{n+1} \leq \frac{7 + 4\sqrt{3}}{2 + \sqrt{3}}$$

Given that  $7 + 4\sqrt{3} = (2 + \sqrt{3})^2$  and  $7 - 4\sqrt{3} = (2 - \sqrt{3})^2$

$$2 - \sqrt{3} \leq y_{n+1} \leq 2 + \sqrt{3}$$

Therefore all of elements of our sequence are bounded by  $2 - \sqrt{3}$  and  $2 + \sqrt{3}$ . Thus  $y_{n+1} \geq y_n$ , and therefore the sequence is increasing.

Now we have the proof that the sequence is both bounded and increasing, and thus converges to some number. After some calculation (and by using common sense) we can also conclude, that the sequence converges to  $2 + \sqrt{3}$ .

## 2.4.4

Show that

$$\sqrt{2}, \sqrt{2\sqrt{2}}, \sqrt{2\sqrt{2\sqrt{2}}}, \dots$$

converges and find the limit.

If we write the same expression as in exercise but using powers instead of radicals we'll get

$$2^{\frac{1}{2}}, 2^{\frac{3}{4}}, 2^{\frac{7}{8}}, \dots$$

Therefore the sequence can be written as

$$x_n = 2^{\sum_{m=1}^n 2^{-m}}$$

$2^m > 0$  for all  $m \in \mathbb{N}$ , therefore the sum is increasing. Thus, the power, in which we put 2 is also increasing, therefore the whole sequence is increasing.

Because each element of this sequence is a power of a positive number, we can conclude, that all of the elements are bound below by 0.

Therefore we just need to prove that

$$\sum_{m=1}^{\infty} 2^{-m} = 1$$

and we'll have our proof.

We'll do it by using the properties of this sum. We can see that

$$\sum_{m=1}^n 2^{-m} + 2^{-n} = 1$$

$2^{-n} > 0$ , and therefore  $1 - \sum_{m=1}^n 2^{-m} = 2^{-n} > 0$  for all  $m \in \mathbb{N}$ .

Thus, for every  $\epsilon > 0$  we can conclude that  $\exists M \in \mathbb{N}$  such that for all  $m \geq M$ .

$$1/m < \epsilon$$

therefore

$$\sum_{m=1}^{\infty} 2^{-m} = 1$$

and thus

$$(x_n) = 2^{\sum_{m=1}^n 2^{-m}} \rightarrow 2^1 = 2$$

## 2.4.5 (Calculating Square Roots)

Let  $x_1 = 2$ , and define

$$x_{n+1} = \frac{1}{2}\left(x_n + \frac{2}{x_n}\right)$$

a) Show that  $x_n^2$  is always greater than 2, and then use this to prove that  $x_n - x_{n+1} \geq 0$ .

Conclude that  $\lim x_n = \sqrt{2}$ .

We'll use induction to show that  $x_n^2$  is always greater than 2.

Base:  $2^2 = 4 > 2$ .

Proposition:  $x_n^2 > 2$

Step:

$$x_n^2 > 2$$

$$x_n^2 - 2 > 0$$

$$(x_n^2 - 2)^2 > 0$$

$$x_n^4 - 4x_n^2 + 4 > 0$$

$$x_n^2 - 4 + \frac{4}{x_n^2} > 0$$

$$\left(x_n^2 + 4 + \frac{4}{x_n^2}\right) > 8$$

$$\frac{1}{4}\left(x_n^2 + 4 + \frac{4}{x_n^2}\right) > 2$$

$$x_{n+1}^2 > 2$$

Thus  $x_n^2 > 2$  for all  $n \in \mathbf{N}$ .

Thus

$$x_n^2 > 2$$

$$x_n^2 - 2 > 0$$

$$\frac{1}{2}x_n - \frac{2}{2x_n} > 0$$

$$x_n - x_{n+1} = x_n - \frac{1}{2}\left(x_n + \frac{2}{x_n}\right) = \frac{1}{2}x_n - \frac{2}{2x_n} > 0$$

Therefore the sequence is decreasing. Now let us prove, that every element of the sequence is positive.

Base:  $x_1 > 0$

Step:

$$x_n > 0$$

$$x_n^2 > 0$$

$$\begin{aligned}
x_n^2 &> -2 \\
x_n^2 + 2 &> 0 \\
x_n + \frac{2}{x_n} &> 0 \\
(x_n + \frac{2}{x_n}) &> 0 \\
\frac{1}{2}(x_n + \frac{2}{x_n}) &> 0 \\
x_{n+1} &> 0
\end{aligned}$$

Thus every element of the sequence is positive and decreasing. Therefore the sequence is convergent (because it is bounded above by the first element and below by 0).

Thus this sequence converges to a positive limit. Let  $l = \lim x_n$ . Then also  $l = \lim x_{n+1}$ . Thus

$$\begin{aligned}
l &= \frac{1}{2}(l + 2/l) \\
2l &= l + 2/l \\
l &= 2/l \\
l^2 &= 2 \\
l &= \sqrt{2}
\end{aligned}$$

as desired.

(b) *Modify the sequence  $(x_n)$  so that it converges to  $\sqrt{c}$ .*

$$x_1 = c$$

$$x_n = \frac{1}{2}(x_n + \frac{c}{x_n})$$

The proof that this sequence works is obtained through the same logic, as in part (a).

## 2.4.6 (Limit Superior)

Let  $(a_n)$  be a bounded sequence.

(a) Prove that the sequence defined by  $y_n = \sup\{a_k : k \geq n\}$  converges

In order to prove that we need to show that this sequence is both bounded and monotone (in this case, decreasing).

Suppose that  $n_1 > n_2 \in \mathbf{N}$ . Then it follows that

$$\{a_k : k \geq n_1\} \subseteq \{a_k : k \geq n_2\}$$

Therefore

$$\forall l \in \{a_k : k \geq n_2\} \rightarrow l \leq \sup\{a_k : k \geq n_2\}$$

and thus

$$\forall l \in \{a_k : k \geq n_1\} \rightarrow l \leq \sup\{a_k : k \geq n_2\}$$

(i.e. if a number  $o$  is a supremum for  $\{a_k : k \geq n_1\}$ , then it is an upper bound for  $\{a_k : k \geq n_2\}$ ). And thus  $y_n \geq y_{n+1}$ . Therefore sequence  $(y_n)$  is decreasing.

Also, because the sequence is bounded,  $\exists M \in \mathbf{R} > 0$  such that  $|a_n| \leq M$  for all  $n \in \mathbf{N}$ . Thus,  $y_n$  is bounded below by  $-M$ . Therefore the sequence is decreasing and bounded. Therefore it converges.

(b) The limit superior of  $(a_n)$ , or  $\limsup a_n$ , is defined by

$$\limsup a_n = \lim y_n$$

where  $y_n$  is the sequence from part (a) of this exercise. Provide a reasonable definition for  $\liminf a_n$  and briefly explain why it always exists for any bounded sequence.

For a bounded sequence  $(a_n)$ ,  $\liminf a_n$  is a limit of a sequence, defined by

$$y_n = \inf\{a_k : k \geq n\}$$

Because for any  $n_1 > n_2 \in \mathbf{N}$  it follows that  $\{a_k : k \geq n_1\} \subseteq \{a_k : k \geq n_2\}$ , we will get that the sequence  $y_n$  is bounded, and because of the boundness of  $(a_n)$  we will get that the sequence is increasing.

(c) Prove that  $\liminf a_n \leq \limsup a_n$  for every bounded sequence, and give an example of a sequence, for which this inequality is strict.

We'll use a proof by contradiction on this one.

Suppose that  $\liminf a_n > \limsup a_n$ . Then let  $l = \liminf a_n - \limsup a_n$ .

Thus there exists  $n \in \mathbf{N}$  for which

$$|\sup\{a_k : k \geq n\} - \limsup a_n| < l/2$$

$$-l/2 < \sup\{a_k : k \geq n\} - \limsup a_n < l/2$$

$$\liminf a_n - l/2 < \sup\{a_k : k \geq n\} + \liminf a_n - \limsup a_n < \liminf a_n + l/2$$

$$\limsup a_n + l/2 < \sup\{a_k : k \geq n\} + l < \liminf a_n + l/2$$

$$\sup\{a_k : k \geq n\} + l/2 < \liminf a_n$$

$$\sup\{a_k : k \geq n\} < \liminf a_n - l/2$$

Therefore  $a_n$

At the same time there exists  $m \in M$  for which

$$|\inf\{a_k : k \geq m\} - \liminf a_n| < l/2$$

$$-l/2 < \inf\{a_k : k \geq m\} - \liminf a_n < l/2$$

$$\liminf a_n - l/2 < \inf\{a_k : k \geq m\} < l/2 + \liminf a_n$$

$$\liminf a_n - l/2 < \inf\{a_k : k \geq m\} < l/2 + \liminf a_n$$

Thus

$$\sup\{a_k : k \geq n\} < \inf\{a_k : k \geq m\}$$

Pick  $j = \max\{m, n\}$ . Thus

$$\{a_k : k \geq n\} \subseteq \{a_k : k \geq j\}$$

$$\{a_k : k \geq m\} \subseteq \{a_k : k \geq j\}$$

Therefore

$$\sup\{a_k : k \geq n\} \geq \sup\{a_k : k \geq j\}$$

$$\inf\{a_k : k \geq n\} \leq \inf\{a_k : k \geq j\}$$

Thus

$$\sup\{a_k : k \geq j\} \leq \sup\{a_k : k \geq n\} < \inf\{a_k : k \geq m\} \leq \inf\{a_k : k \geq j\}$$

$$\sup\{a_k : k \geq j\} < \inf\{a_k : k \geq j\}$$

Therefore supremum of the set is less than an infimum of a set, which is a contradiction.

Therefore  $\liminf a_n \leq \limsup a_n$ .

Example of a sequence, for which this inequality is strict is

$$a_n = -1^n$$

(d) Show that  $\liminf a_n = \limsup a_n$  if and only if  $\lim a_n$  exists. In this case, all three share the same value.

**In forward direction:**

Suppose that  $\liminf a_n = \limsup a_n = l$ . Then for every  $\epsilon > 0$  there exists  $n \in N$  such that

$$|\sup\{a_k : k \geq n\} - l| < \epsilon$$

Also there exists  $m \in N$  such that

$$|\inf\{a_k : k \geq m\} - l| < \epsilon$$

for the same  $\epsilon$ .

Let us set  $j = \max\{n, m\}$ . Then

$$-\epsilon < \sup\{a_k : k \geq j\} - l < \epsilon$$

$$l - \epsilon < \sup\{a_k : k \geq j\} < \epsilon + l$$

Also

$$-\epsilon < \inf\{a_k : k \geq j\} - l < \epsilon$$

$$l - \epsilon < \inf\{a_k : k \geq j\} < l + \epsilon$$

For any  $j \in N \geq k$

$$a_j \leq \sup\{a_k : k \geq j\} < \epsilon + l$$

$$l - \epsilon < \inf\{a_k : k \geq j\} \leq a_j$$

therefore

$$l - \epsilon < a_j < l + \epsilon$$

$$-\epsilon < a_j - l < \epsilon$$

$$|a_j - l| < \epsilon$$

Therefore

$$\forall \epsilon > 0 : \exists N \in \mathbf{N} : \forall n \geq N \rightarrow |a_n - l| < \epsilon$$

or in other words,  $\lim a_n = l = \liminf a_n = \limsup a_n$ , as desired.

**In other direction:**

Suppose that  $\lim a_n = l$ . From this we can state that

$$|a_n - l| < \epsilon$$

$$\epsilon < a_n - l < \epsilon$$

$$l - \epsilon < a_n < \epsilon + l$$

Thus both  $\sup\{a_k : k \geq n\}$  and  $\inf\{a_k : k \geq n\}$  are bounded by  $l - \epsilon$  and  $l + \epsilon$ . In other words

$$l - \epsilon \leq \inf\{a_k : k \geq n\} \leq \sup\{a_k : k \geq n\} \leq \epsilon + l$$

This is true for all  $\epsilon > 0$ . Thus if we set  $\epsilon = 1/j$  where  $j \in N$ , there exists appropriate  $i \in N$  for which

$$l - 1/j \leq \inf\{a_k : k \geq i\} \leq \sup\{a_k : k \geq i\} \leq l + 1/j$$

Therefore if we take a limit of all sides of this inequality we can state that

$$\lim(l - 1/j) \leq \liminf\{a_k : k \geq i\} \leq \limsup\{a_k : k \geq i\} \leq \lim(l + 1/j)$$

$$l \leq \liminf a_n \leq \limsup a_n \leq l$$

Here, the ability to take limits of all sides is justified by the fact, that  $\forall n \in \mathbf{N} > k : a_n \geq b_n \rightarrow \lim a_n \geq \lim b_n$ .

Thus  $\liminf a_n = \limsup a_n = l$ , as desired.

## 2.5.1

*Prove Theorem 2.5.2*

First of all, let us state the theorem itself.

**Theorem 2.5.2** Subsequences of a convergent sequence converge to the same limit.

Suppose that  $a_n$  is a convergent sequence,  $b_n$  is a subsequence of  $a_n$ , and  $l$  is a limit of  $a_n$ . Because  $a_n$  is a convergent sequence we can state that

$$\forall \epsilon \in R > 0 : \exists N \in \mathbf{N} : \forall n \geq N \rightarrow |a_n - l| < \epsilon$$

For each  $N \in \mathbf{N}$  there exists  $M \in \mathbf{N}$ , such that  $M \geq N$  and  $b_M = a_N$ . Also, for each  $m > M \in \mathbf{N}$  there exists  $n > N \in \mathbf{N}$  such that  $b_m = a_n$ . Thus we can state that

$$\forall \epsilon \in R > 0 : \exists N \in \mathbf{N} : \exists M \in \mathbf{N} > N : \forall m \geq M \rightarrow |b_m - l| < \epsilon$$

$$\forall \epsilon \in R > 0 : \exists M \in \mathbf{N} : \forall m \geq M \rightarrow |b_m - l| < \epsilon$$

or in other words,  $(b_n) \rightarrow l$ , as desired.

## 2.5.2

(a) *Prove that if an infinite series converges, then the associative property holds. Assume  $a_1 + a_2 + a_3 + a_4 + a_5 + \dots$  converges to a limit  $L$  (i.e. the sequence of partial sums  $(s_n) \rightarrow L$ ). Show that any regrouping of the terms*

$$(a_1 + a_2 + \dots + a_{n_1}) + (a_{n_1+1} + a_{n_1+2} + \dots + a_{n_2}) + (a_{n_2+1} + a_{n_2+2} + \dots + a_{n_3}) + \dots$$

*leads to a series that also converges to  $L$ .*

Suppose that we regrouped terms of the above-given sequence. Then each of the terms will be defined by sequence of partial sums

$$s_1 = (a_1 + a_2 + \dots + a_{n_1})$$

$$s_2 = (a_1 + a_2 + \dots + a_{n_1}) + (a_{n_1+1} + a_{n_1+2} + \dots + a_{n_2})$$



$$s_n = a_1 + \dots + a_{n_n}$$

Therefore,  $s_n$  is a subsequence of an original sequence. Therefore, because the original sequence converges to the same limit,  $s_n$  will be convergent to the same limit, as desired.

(b) Compare this result to the example discussed at the end of Section 2.1 where infinite addition was shown not to be associative. Why doesn't our proof in (a) apply to this example?

Our proof doesn't apply to any of the sequences, discussed in the 2.1 because each one of them was not convergent, which is a prerequisite for part (a).

### 2.5.3

Give an example of each of the following, or argue that such a request is impossible.

(a) A sequence that does not contain 0 or 1 as a term but contains subsequences, converging to each of those values

$$a_n = \begin{cases} 1/n & \text{if } n \text{ is even} \\ 1 - 1/n & \text{if } n \text{ is odd} \end{cases} \quad (9)$$

(b) A monotone sequence that diverges but has a convergent subsequence

Suppose that we have such a sequence. Then, because subsequence is convergent, it is bounded. Thus, there exist  $M > 0$  for which  $|a_{n_j}| \geq M \rightarrow -M \leq a_{n_j} \leq M$ . For each of elements of the original sequence, there exists an element of subsequence, that is "further down the line" (i.e.  $\forall n \in N : \exists m, j \in N : n < m$  and  $a_m = a_{n_j}$ ). Then

$$a_n \leq a_{m_j} \leq M$$

in case of the increasing sequence

$$a_n \geq a_{m_j} \geq -M$$

in case of the decreasing sequence.

Thus, the original sequence is bounded. Thus the original sequence is bounded and monotone. Therefore the original sequence is convergent. Therefore we have a contradiction.

(c) A sequence that contains subsequences converging to every point in the infinite set  $\{1, 1/2, 1/3, 1/4, 1/5, \dots\}$

Let

$$a_n = [1], [1, 1/2], [1, 1/2, 1/3], [1, 1/2, 1/3, 1/4], [1, 1/2, 1/3, 1/4, 1/5], \dots$$

where [ and ] are added just for the visual clue.

Therefore we have in it a subsequence, that converges to an arbitrary number  $1/n$  for  $n \in \mathbf{N}$ .

(d) *An unbounded sequence with a convergent subsequence.*

$$a_n = \begin{cases} n & \text{if } n \text{ is odd} \\ 0 & \text{if } n \text{ is even} \end{cases} \quad (10)$$

(e) *A sequence that has a subsequence that is bounded but contains no subsequence that converges*

Impossible. Because there exists bounded subsequence, which is in and of itself a sequence, there exists a convergent subsequence ( by Bolzano-Weierstrass Theorem). Subsequence of a subsequence is a subsequence of original sequence. Thus an original sequence contains a convergent subsequence.

## 2.5.4

*Assume  $(a_n)$  is a bounded sequence with the property that every convergent subsequence of  $(a_n)$  converges to the same limit  $a \in \mathbf{R}$ . Show that  $(a_n)$  must converge to  $a$ .*

Because  $(a_n)$  is bounded, there exists  $M > 0$  such that  $|a_n| < M$ .

Suppose that we have an  $\epsilon$ -neighborhood around  $a$ . Then  $[-M, M] \setminus V_\epsilon(a)$  could contain elements of  $(a_n)$ . Suppose that  $[-M, M] \setminus V_\epsilon(a)$  contains infinite amount of elements of  $(a_n)$ . Then there exists a subsequence of  $(a_n)$ , that it is outside of  $V_\epsilon$ . Thus, there exists a convergent subsequence of sequence, that converges to some number, other, than  $a$ , which is a contradiction. Therefore there exists only finite amount of elements in  $[-M, M] \setminus V_\epsilon(a)$ .

Therefore, for any  $\epsilon$ , there exists an  $N$  (which is a maximum index of a finite elements of elements outside the neighborhood, or 1 in case that there are no elements outside of the neighborhood), for which it is true, that  $n > N \in \mathbf{N} \rightarrow a_n \in V_\epsilon(a)$ . Therefore,  $(a_n) \rightarrow a$  by topological version of definition of convergence.

## 2.5.5

*Extend the result proved in Example 2.5.3 to the case  $|b| < 1$ . Show that  $\lim(b^n) = 0$  whenever  $-1 < b < 1$ .*

Let  $-1 < b < 1$ . Then we need to prove that

$$\forall \epsilon : \exists N \in \mathbf{N} : \forall n \geq N \rightarrow |b^n - 0| < \epsilon$$

$$\forall \epsilon : \exists N \in \mathbf{N} : \forall n \geq N \rightarrow |b^n| < \epsilon$$

We already know, that for  $0 < b_1 < 1$  it is true, that  $\lim(b_1^n) = 0$ . Thus, for  $b = 0$ , the case is trivial, and for  $b < 0$  it defaults to  $|b| = b_1$ . Therefore  $\lim(b^n) = 0$  for  $-1 < b < 1$ .

## 2.5.6

Let  $(a_n)$  be a bounded sequence, and define the set

$$S = \{x \in \mathbb{R} : x < a_n \text{ for indefinitely many terms } a_n\}$$

Show that there exists a subsequence  $(a_{n_k})$  converging to  $s = \sup S$ . (This is a direct proof of the Bolzano-Weierstrass Theorem using Axiom of Completeness.)

The wording of this exercise doesn't make it easy to understand what exactly are we trying to prove. After some pondering, I concluded, that what this particular exercise is trying to say, is that there exists a subsequence, that is convergent to a limit supremum of this sequence. If it does, then some rewording of 2.4.6 part(a) will give you the desired result. If it doesn't, then I don't know what do I need to prove, and therefore declare this particular exercise as unfinished.

This exercise already took too much time which was spent on anything but math, therefore instead of pondering further, I will rather spend time on other exercises from this book.

After some research on this topic, I found out that there exist another proof of Bolzano-Weierstrass Theorem, that doesn't use NIP, and my idea now is that this particular exercise tried to ask us to give that proof. That proof can be (kinda) derived from the limit superior theorem, by the fact, that if a limit superior exists, then there exists a subsequence in the original sequence, that converges to limit superior.

## 2.6.1

Give an example of each of the following, or argue that such a request is impossible

(a) A Cauchy sequence that is not monotone

$$a_n = (-0.5)^n$$

It is convergent (as proven in previous exercises), and is a Cauchy sequence (because it is convergent)

(b) A monotone sequence that is not Cauchy

$$a_n = n$$

It is not convergent, therefore not a Cauchy sequence.

(c) A Cauchy sequence with a divergent subsequence.

Impossible, because a Cauchy sequence is a convergent sequence, and all of the subsequences of convergent sequence are convergent to the same number.

(d) An unbounded sequence containing a subsequence that is Cauchy

$$a_n = \begin{cases} n & \text{if } n \text{ is odd} \\ 0 & \text{if } n \text{ is even} \end{cases} \quad (11)$$

## 2.6.2

Supply a proof for Theorem 2.6.2

Firstly, let us state the theorem itself

**Theorem 2.6.2** *Every convergent sequence is a Cauchy sequence*

Suppose that  $(x_n)$  converges to  $x$ . Thus,

$$\forall \epsilon/2 > 0 : \exists N \in \mathbf{N} : \forall n \geq N \rightarrow |x_n - x| < \epsilon/2$$

Let  $m > N$ . Then

$$|x_n - x| < \epsilon/2$$

$$|x_m - x| < \epsilon/2$$

thus

$$|x_n - x| + |x_m - x| < \epsilon$$

$$|x_n - x| + |x - x_m| < \epsilon$$

$$|x_n - x + x - x_m| \leq |x_n - x| + |x - x_m| < \epsilon$$

$$|x_n - x + x - x_m| < \epsilon$$

$$|x_n - x_m| < \epsilon$$

Thus any convergent sequence is a Cauchy sequence, as desired.

## 2.6.3

(a) Explain how the following pseudo-Cauchy property differs from the proper definition of a Cauchy sequence: A sequence  $(s_n)$  is pseudo-Cauchy if, for all  $\epsilon > 0$ , there exists an  $N$  such that if  $n \geq N$ , then  $|s_{n+1} - s_n| < \epsilon$ .

It obviously differs in the fact, that in given definition we are only considering the element, that goes after one element, instead of all of elements, that are after.

(b) If possible, give an example of a divergent sequence  $(s_n)$  that is pseudo-Cauchy  
Harmonic series.

## 2.6.4

Assume  $(a_n)$  and  $(b_n)$  are Cauchy sequences. Use a triangle inequality argument to prove  $c_n = |a_n - b_n|$  is Cauchy.

Therefore for each  $\epsilon$  there exists

$$\forall \epsilon/2 > 0 : \exists N_1 \in \mathbf{N} : \forall n, m \geq N_1 \rightarrow |a_n - a_m| < \epsilon/2$$

$$\forall \epsilon/2 > 0 : \exists N_2 \in \mathbf{N} : \forall n, m \geq N_2 \rightarrow |b_n - b_m| < \epsilon/2$$

Let us pick  $N = \max\{N_1, N_2\}$ . Then

$$|a_n - a_m| + |b_n - b_m| < \epsilon$$

$$\begin{aligned} |c_n - c_m| &= ||a_n - b_n| - |a_m - b_m|| \leq |a_n - b_n - a_m + b_m| \leq |a_n - a_m| + |b_n - b_m| < \epsilon \\ |c_n - c_m| &< \epsilon \end{aligned}$$

Thus  $(c_n)$  is also Cauchy, as desired.

## 2.6.5

If  $(x_n)$  and  $(y_n)$  are Cauchy sequences, then one easy way to prove that  $(x_n + y_n)$  is Cauchy is to use the Cauchy Criterion. By Theorem 2.6.4,  $(x_n)$  and  $(y_n)$  must be convergent, and the Algebraic Limit Theorem then implies  $(x_n + y_n)$  is convergent and hence Cauchy

(a) Give a direct argument that  $(x_n + y_n)$  is a Cauchy sequence that does not use the Cauchy Criterion or the Algebraic Limit Theorem.

Suppose that  $(x_n)$  and  $(y_n)$  are Cauchy sequences. Therefore

$$\forall \epsilon/2 > 0 : \exists N_1 \in \mathbf{N} : \forall m, n \geq N_1 \rightarrow |x_n - x_m| < \epsilon/2$$

and

$$\forall \epsilon > 0 : \exists N_2 \in \mathbf{N} : \forall m, n \geq N_2 \rightarrow |y_n - y_m| < \epsilon/2$$

Therefore let  $N = \max\{N_1, N_2\}$ . Then

$$\forall \epsilon/2 > 0 : \exists N \in \mathbf{N} : \forall m, n \geq N \rightarrow |x_n - x_m| < \epsilon/2$$

and

$$\forall \epsilon > 0 : \exists N \in \mathbf{N} : \forall m, n \geq N \rightarrow |y_n - y_m| < \epsilon/2$$

Thus

$$|x_n - x_m| + |y_n - y_m| < \epsilon$$

$$|x_n - x_m + y_n - y_m| = |x_n - x_m + y_n - y_m| \leq |x_n - x_m| + |y_n - y_m| < \epsilon$$

therefore  $(x_n + y_n)$  is a Cauchy sequence as well, as desired.

(b) Do the same for the product  $(x_n y_n)$ .

Suppose that  $(x_n)$  and  $(y_n)$  are Cauchy sequences. Therefore they are bounded by  $|x_n| < M_1$  and  $|y_n| < M_2$ . Let us pick the greatest  $M = \max\{M_1, M_2\}$ . Then

$$\forall \epsilon/2 > 0 : \exists N_1 \in \mathbf{N} : \forall m, n \geq N_1 \rightarrow |x_n - x_m| < \frac{\epsilon}{2M}$$

and

$$\forall \epsilon > 0 : \exists N_2 \in \mathbf{N} : \forall m, n \geq N_2 \rightarrow |y_n - y_m| < \frac{\epsilon}{2M}$$

Thus

$$\begin{aligned} |x_n y_n - x_m y_m| &= |x_n y_n - x_n y_m + x_n y_m - x_m y_m| = |x_n(y_n - y_m) + y_m(x_n - x_m)| \leq \\ &\leq |x_n||y_n - y_m| + |y_m||x_n - x_m| \leq M \frac{\epsilon}{2M} + M \frac{\epsilon}{2M} = \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \end{aligned}$$

Therefore  $(x_n y_n)$  is a Cauchy sequence as well.

## 2.6.6

(a) Assume the Nested Interval Property (Theorem 1.4.1) is true and use a technique similar to the one employed in the proof of the Bolzano-Weierstrass Theorem to give a proof for the Axiom of Completeness (The reverse implication was given in Chapter 1. This shows, that AoC is equivalent to NIP

Suppose that we have a bounded set  $A$ . Then, because it is bounded, there exists an interval  $I_1 = [a, b]$ , where  $a \in A$ , and  $b \in \mathbf{R} \setminus A$  - upper bound of  $A$ . Then, let us divide this set (and in general set  $I_n$ ) into two sets of equal length  $[a, l]$  and  $[l, b]$ . Then set  $I_{n+1}$  to the set,  $[l, b]$ , if it has any elements of  $A$ ; otherwise set it to  $[a, l]$ . Because of the NIP

$$\cap_{n=1}^{\infty} I_n = B \neq \emptyset$$

Suppose now, that there exist two elements  $c < d \in B$ . Then let us look at the interval  $[c, d]$  with length  $h$ . Because of the Archimedian property, there exists a  $n \in \mathbf{N}$ , such that  $1/n < |d - c|$ , and thus there must exist  $1/2^n < |d - c|$ . Therefore there exists a set  $I_j$ , which length is less than distance between  $c$  and  $d$ . Therefore we have a contradiction. (The absolute value here is redundant, but it doesn't hurt). Therefore there exists only one number in  $B$ . Let us henceforth call it  $l$ .

Suppose, that there exists  $q \in A$ , such that  $q > l$ . Then it follows, that there exists an interval  $[q, l]$ . By the same logic, as in the previous paragraph, we can conclude, that such element does not exist. Therefore, for all  $q \in A$  it is true, that  $q \leq l$ . Thus,  $l$  is an upper bound.

Suppose now, that  $r$  is an upper bound of  $A$  with the property, that  $r < l$ . Then  $[r, l]$  contains no elements of  $A$ . Therefore there exists  $I_g \subseteq [r, l]$  such that it contains no elements of  $A$ , which is a contradiction. Therefore, for any upper bound  $r$  it is true, that  $r \geq l$ .

Therefore  $l$  is an upper bound with the property, that any upper bound is either greater or equal to  $l$ . Thus,  $l$  is a lowest upper bound (or supremum) of  $A$ . Therefore, by existence of such a bound, any bounded set of  $\mathbf{R}$  has a least upper bound. In other words,

$$NIP \iff AoC$$

, as desired.

(b) Use the Monotone Convergence Theorem to give a proof of the Nested Interval Property. (This establishes the equivalence of AoC, NIP, and MCT)

Suppose, that we have nested intervals

$$I_1 \subseteq I_2 \subseteq I_3 \subseteq I_4 \dots \subseteq I_n \subseteq \dots$$

Then, let  $(a_n)$  = lower bound of  $I_n$ . Then it follows, that

$$a_1 \leq a_2 \leq a_3 \dots \leq a_n \leq \text{upper bound of any of } I_n$$

Therefore, this sequence is increasing and bounded above (and below by  $a_1$ ). Thus, it is convergent to some number  $l_1 \in \mathbb{R}$ .

Let us also define  $(b_n)$  = upper bound of  $I_n$

$$b_1 \geq b_2 \geq b_3 \dots \geq b_n \geq \text{lower bound of any of } I_n$$

Therefore, this sequence is decreasing and bounded below (and above by  $b_1$ ). Thus, it is convergent to some number  $l_2 \in \mathbb{R}$ .

Now look at  $l_1$ . Let us pick  $I_j$  with lower bound  $a_1$  and upper bound  $b_1$ . Suppose, that  $l_1 < a_1$ . Then let  $\epsilon = a_1 - l_1$ . Then it follows, that all of the elements of the sequence after  $g$  (i.e.  $a_n$  for  $n \geq g$ ) are not in  $V_\epsilon(l_1)$ . Thus, we have a contradiction. Thus,  $l \geq a_1$ . With the same logic and opposite sign we can prove that  $l \leq b_1$ . Thus,  $l \in I_g$  for any  $g \in \mathbb{N}$ . Thus,

$$l \in \bigcap_{n=1}^{\infty} I_n$$

Thus, the intersection of any number of nested sets is non-empty. Thus,

$$MCT \iff NIP$$

, as desired.

(c) This time, start with the Bolzano-Weierstrass Theorem and use it to construct a proof of the Nested Interval Property. (Thus, BW is equivalent to NIP, and hence to AoC and MCT as well.)

Suppose that we have nested intervals

$$I_1 \subseteq I_2 \subseteq I_3 \subseteq I_4 \dots \subseteq I_n \subseteq \dots$$

Now let us define a sequence, such that every member of a sequence is a number in a corresponding interval  $I_n$ .

$$(a_n) = \{x : x \in I_n\}$$

First of all, this sequence is going to be bounded by  $I_1$ . Therefore, by BW, it has a convergent subsequence. Let us call this subsequence  $(a_{n_j})$ , and call a limit of such a sequence  $l$ . Then by the same logic as in part (b),  $l \in I_g$  for any  $g \in \mathbb{N}$ , and therefore

$$l \in \bigcap_{n=1}^{\infty} I_n$$

Thus,

$$BW \iff NIP$$

(d) Finally, use the Cauchy Criterion to prove the Bolzano-Weierstrass Theorem. This is a final link in the equivalence of the five characterizations of completeness discussed at the end of Section 2.6

Suppose that we have a bounded sequence  $(a_n)$ . Then, it follows, that there exists  $M > 0$  such that  $a_n \in [-M, M]$  for any  $n \in \mathbb{N}$ . We are going to describe how to get from  $I_j$  to  $I_{j+1}$ , and by extension from  $a_{n_j}$  to  $a_{n_{j+1}}$ . For the complete construction let  $I_1 = [-M, M]$  and proceed from there. Now let's start constructing sequence.

Pick  $a_{n_j}$  from interval  $I_j = [a, b]$ . Then divide  $I_j$  into two equal-length intervals  $[a, l]$  and  $[l, b]$ . As discussed in previous exercises, one of them will have infinite amount of elements. Let  $I_{j+1}$  be interval with infinite amount of elements of  $(a_n)$ .

Therefore for each  $\epsilon > 0$  there will exist  $j \in \mathbb{N}$  such that it will be true, that  $I_j \subseteq V_\epsilon(a_j)$ . And thus will contain every element of subsequence  $\{a_{n_k} : k \geq j\}$  (the proof for validity of this statement was already discussed in original proof of BW). Thus,  $(a_{n_j})$  will be a Cauchy sequence, and therefore convergent to some value. Therefore

$$CC \iff BW$$

## 2.7.1

Proving the Alternating Series Test (Theorem 2.7.7) amounts to showing that the sequence of partial sums

$$s_n = a_1 - a_2 + a_3 - \dots \pm a_n$$

converges. (The opening example in Section 2.1 includes a typical illustration of  $(s_n)$ .) Different characterizations of completeness lead to different proofs.

Let us state the theorem itself first:

**Theorem 2.7.7 (Alternating Series Test)** Let  $(a_n)$  be a sequence satisfying

$$(i) \ a_1 \geq a_2 \geq a_3 \geq \dots \geq a_n \geq \dots \text{ and}$$

$$(ii) \ (a_n) \rightarrow 0$$

Then, the alternating series  $\sum_{n=1}^{\infty} (-1)^{n+1} a_n$  converges

Firstly, let us draw some conclusions from the theorem. If  $a_n < 0$  for some  $n \in \mathbb{N}$ , then all of the  $a_j$  for  $j > n$  are  $a_j \leq a_n < 0$ , and thus it does not converge to zero, which is a contradiction. Thus we can state that  $a_n \geq 0$  for all  $n \in \mathbb{N}$ .

(a) Prove the Alternating Series Test by showing that  $(s_n)$  is a Cauchy sequence.

Let  $\epsilon > 0$ . Then, because  $(a_n)$  converges, there exist  $N \in \mathbb{N}$  such that for all  $n > N$  it follows that  $|a_n| < \epsilon$ .



Let  $s_n = a_1 - a_2 + a_3 - \dots \pm a_n$ . Now let  $s_m = a_1 - a_2 + a_3 - \dots \pm a_m$  for  $m \geq n$ . Then

$$|s_m - s_n| = |a_{n+1} - a_{n+2} + \dots \pm a_m|$$

Let us examine interval  $[0, a_n]$  with regards to elements, that follow  $a_n$ :

$$a_n \geq a_{n+1} \rightarrow a_n - a_{n+1} \in [0, a_n]$$

Also, because  $a_n \geq a_{n+1} \geq a_{n+2}$  we can state that  $a_n - a_{n+1} + a_{n+2} \in [0, a_n]$ . Thus

$$[a_n - a_{n+1} + a_{n+2}, a_{n+2}] \subseteq [a_n - a_{n+1}, a_n] \subseteq [0, a_n]$$

Thus

$$a_n \geq a_{n+1} - a_{n+2} + a_{n+3} - \dots \pm a_m$$

for any  $m \geq n$ . Therefore

$$|s_m - s_n| = |a_{n+1} - a_{n+2} + \dots \pm a_m| \leq a_n < \epsilon$$

Therefore we can state that for any  $\epsilon > 0$  there exist  $N$ , such that for all  $n, m \geq N$  it follows that

$$|s_n - s_m| \leq a_n < \epsilon$$

Therefore  $(s_n)$  is a Cauchy sequence. Thus it converges, as desired.

(b) *Supply another proof for this result using the NIP (Theorem 1.4.1).*

Let us define interval  $[a_n - a_{n+1}, a_n]$  for some  $n \in N$ . We can see that  $a_n$  and  $a_n - a_{n+1}$  are in this interval. Because  $a_{n+2} \geq 0$  and  $a_{n+2} \leq a_{n+1}$  we can state that

$$a_n - a_{n+1} \leq a_{n+2} + a_n - a_{n+1} \leq a_n$$

It also doesn't hurt to state here that

$$a_{n+2} + a_n - a_{n+1} \leq a_n \rightarrow [a_n - a_{n+1}, a_{n+2} + a_n - a_{n+1}] \subseteq [a_n - a_{n+1}, a_n]$$

Now let us define

$$I_n = [a_{2n} - a_{2n+1}, a_{2n}]$$

from our discussion in previous paragraph we can state that

$$a_{2n+m} \in I_n$$

where  $m \in N \cup \{0\}$ . Also

$$I_n \subseteq I_{n+1}$$

For the future use let us also note the fact, that for every  $j \in N$  there exists an interval  $I_n$ , whose length is less or equal to  $a_j$ .

Thus we have a collection of nested intervals. NIP states that

$$\cap_{n=1}^{\infty} I_n = B \neq \emptyset$$

Suppose that we have  $a \neq b \in B$ . Then there exist  $\epsilon = |a - b|$ , for which by convergence of  $(a_n)$  there exists  $N$  for which it is true that

$$\forall n \geq N \rightarrow |a_n| \leq \epsilon$$

Therefore there exists an interval, whose length is less than the distance between  $a$  and  $b$ . Therefore both  $a$  and  $b$  cannot be at some  $I_n$  at the same time. Therefore we have a contradiction, and we can state that we cannot have two distinct elements in  $B$ . Thus, there exists only one element in  $B$ .

Note that for every partial sum  $s_n$  we can find corresponding  $I_m$  such that

$$s_n \in I_m$$

Thus we can state that

$$(s_n) \rightarrow l$$

or in other words,

$$\sum_{n=1}^{\infty} (-1)^{n+1} a_n = \lim s_n = l$$

where  $l$  is the sole element of  $B$ .

(c) Consider the subsequences  $(s_{2n})$  and  $(s_{2n+1})$ , and show how the Monotone Convergence Theorem leads to a third proof for the Alternating Series Test

Let us note that

$$s_{2n} = a_1 - a_2 + \dots + a_{2n-1} - a_{2n}$$

Thus

$$s_{2n} - s_{2(n+1)} = a_{2n+1} - a_{2n+2} \geq 0$$

thus series  $(s_{2n})$  is increasing. Also

$$s_{2n+1} - s_{2(n+1)+1} = -a_{2n+2} + a_{2n+3} \leq 0$$

thus series  $(s_{2n+1})$  is decreasing.

Because  $(s_{2n+1})$  is decreasing, there exists  $a = \sup(s_{2n+1})$ . Because  $(s_{2n})$  is increasing, there exists  $b = \inf(s_{2n})$ . Every  $s_j$  is either equal to some  $s_{2n}$  if  $j$  is even or to some  $s_{2n+1}$  if  $j$  is odd. Thus  $(s_n)$  is bounded above by  $a$  and bounded below by  $b$ .

By MCT we can state that

$$\lim s_{2n} = l_1$$

$$\lim s_{2n+1} = l_2$$

Now let note that for all  $n \in N$

$$s_{2n} - s_{2n+1} = a_{2n}$$

Because all subsequences of convergent sequence converge to the same number we can state that

$$\lim(s_{2n} - s_{2n+1}) = \lim(a_{2n}) = 0$$

thus

$$\lim s_{2n} = \lim s_{2n+1} = l_1 = l_2 = l$$

Because for every  $n$  there exists  $s_{2n} \leq s_j$  and  $s_{2n+1} \leq s_j$  we can conclude that by Squeeze theorem

$$\sum_{n=1}^{\infty} (-1)^{n+1} a_n = \lim s_n = l$$

as desired.

## 2.7.2

(a) Provide the details for the proof of the Comparison Test (Theorem 2.7.4) using the Cauchy Criterion for Series.

Let us firstly state the theorem itself

**Theorem 2.7.4 (Comparison Test)** Assume  $(a_k)$  and  $(b_k)$  are sequences satisfying  $0 \leq a_k \leq b_k$  for all  $k \in N$ .

(i) If  $\sum_{k=1}^{\infty} b_k$  converges, then  $\sum_{k=1}^{\infty} a_k$  converges

(ii) If  $\sum_{k=1}^{\infty} a_k$  diverges, then  $\sum_{k=1}^{\infty} b_k$  diverges

(i)

Let us pick some  $\epsilon > 0$ . Then, by Cauchy Criterion for sequences, there exists  $m > n \geq N \in \text{textbf{N}}$ . such that

$$|s_n - s_m| = |b_{m+1} + \dots + b_n| = b_{m+1} + \dots + b_n < \epsilon$$

for  $s_n$  - partial sum for  $\sum_{k=1}^{\infty} b_k$ .

Let  $l_n$  be a partial sum for  $\sum_{k=1}^{\infty} a_k$ . Then

$$|l_n - l_m| = |a_{m+1} + \dots + a_n| = a_{m+1} + \dots + a_n \leq b_{m+1} + \dots + b_n = |s_n - s_m| \leq \epsilon$$

Thus, by Cauchy Criterion for Series,  $\sum_{k=1}^{\infty} a_k$  converges.

(ii)

By Cauchy Criterion for sequences (or rather by negation of it), there exists  $\epsilon > 0$  such that for all  $m > n \in N$  it is true that

$$|l_n - l_m| = |a_{m+1} + \dots + a_n| = a_{m+1} + \dots + a_n > \epsilon$$

for  $l_n$  - partial sum for  $\sum_{k=1}^{\infty} a_k$ .

Let  $s_n$  be a partial sum for  $\sum_{k=1}^{\infty} b_k$ . Then

$$\epsilon < |l_n - l_m| = |a_{m+1} + \dots + a_n| = a_{m+1} + \dots + a_n \leq b_{m+1} + \dots + b_n = |s_n - s_m|$$

For all  $m > n \geq N$

Thus, by Cauchy Criterion for Series,  $\sum_{k=1}^{\infty} b_k$  diverges.

(b) Give another proof for the Comparison Test, this time using the Monotone Convergence Theorem

Firstly, let us look closer at  $\sum_{k=1}^{\infty} (b_k)$ : first thing that we could notice, is that every term of the sum is non-negative, and therefore we can state that the sequence of partial sums for it is increasing. Same can be said about  $\sum_{k=1}^{\infty} (a_k)$ .

(i)

Let us define  $s_k$  to be a partial sum for  $\sum_{k=1}^{\infty} (b_k)$  and  $l_k$  to be a partial sum for  $\sum_{k=1}^{\infty} (a_k)$ . Because  $s_k$  is increasing and convergent, it is true that

$$s_k \leq l$$

for any  $k \in N$  where  $l = \lim s_k$ . Thus

$$l_k \leq s_k \leq l$$

for any  $k \in N$ . Therefore  $l_k$  is bounded above by  $l$  as well. Bringing up to our attention again that  $a_k$  is non-negative, we can state that

$$0 \leq l_k \leq l$$

thus  $\sum_{k=1}^{\infty} (a_k)$  is bounded and monotone and therefore by MCT, convergent.

(ii)

Here we will not necessarily use MCT itself, but the fact, that any convergent function is bounded.

Partial sums for  $\sum_{k=1}^{\infty} (a_k)$  are not convergent and increasing, therefore we can state that they are unbounded, precisely that it is convergent to infinity. Because  $l_k \leq s_k$ , the same can be said about partial sums for  $\sum_{k=1}^{\infty} (b_k)$ .

### 2.7.3

Let  $\sum a_n$  be given. For each  $n \in N$ , let  $p_n = a_n$  if  $a_n$  is positive and assign  $p_n = 0$  if  $a_n$  is negative. In a similar manner, let  $q_n = a_n$  if  $a_n$  is negative and  $q_n = 0$  if  $a_n$  is positive.

(a) Argue that if  $\sum a_n$  diverges, then at least one of  $\sum p_n$  or  $\sum q_n$  diverges.

We are going to proceed with a proof by contradiction.

Suppose that both  $\sum p_n$  and  $\sum q_n$  converge. Let  $s_n$  be a partial sum for  $\sum a_n$ ,  $s1_n$  a partial sum for  $p_n$  and  $s2_n$  a partial sum for  $q_n$ . Then

$$s1_n + s2_n = p_1 + p_2 + \dots + p_n + q_1 + q_2 + \dots + q_n = a_1 + a_2 + a_3 + \dots + a_n = s_n$$

Thus  $\sum a_n$  converges, which is a contradiction. Therefore both sums  $\sum q_n$  and  $\sum p_n$  cannot be convergent. Therefore at least one of  $\sum q_n$  or  $\sum p_n$  must diverge.

(b) Show that if  $\sum a_n$  converges conditionally, then both  $\sum p_n$  and  $\sum q_n$  diverge.

We are going to proceed with a proof by contradiction as well. Suppose that  $p_n$  converges (we are only looking at the case with  $p_n$  here, but the same logic can be applied for  $q_n$  as well).

Then, we can state that  $\sum -p_n$  converges as well. Thus  $\sum a_n - p_n$  converges. Therefore  $\sum q_n$  also converges. Therefore  $\sum p_n - q_n$  converges as well. Therefore  $\sum a_n$  converges absolutely, which is a contradiction.

Therefore we can state, that in order to  $\sum a_n$  to converge absolutely, both  $\sum p_n$  and  $\sum q_n$  must diverge, as desired.

## 2.7.4

Give an example to show that it is possible for both  $\sum x_n$  and  $\sum y_n$  diverge, but for  $\sum x_n y_n$  converge.

$$x_n = y_n = 1/n \rightarrow x_n y_n = 1/n^2$$

## 2.7.5

(a) Show that if  $\sum a_n$  converges absolutely, then  $\sum a_n^2$  also converges absolutely. Does this proposition hold without absolute convergence?

Let  $s_n$  be a partial sum for  $\sum |a_n|$ . Then, because  $\sum |a_n|$  converges, there exists  $M > 0$  such that  $s_n \leq M$ . Therefore each  $|a_n| \leq M$ . Therefore

$$s_n^2 = |a_1|^2 + |a_2|^2 + \dots + |a_n|^2 \leq M|a_1| + M|a_2| + \dots + M|a_n| = M(|a_1| + |a_2| + \dots + |a_n|) = Ms_n$$

Therefore, because of the convergence  $\sum |a_n|$  we can state, that  $M \sum |a_n| = \sum M|a_n|$  converges as well. Thus,  $\sum |a_n|^2$  also converges. Thus  $\sum a_n^2$  converges absolutely, as desired.

If we substitute  $a_n$  with some conditionally convergent series, then this proof collapses on the moment, when we state that  $|a_n| \leq M$ . That doesn't necessarily disprove the proposition, only shows where does absolute convergence plays the role in our proof. In

order to disprove this proposition we need an example of a convergent sequence, that contradicts our proposition. One such sequence is

$$\sum a_n = \sum \frac{(-1)^{n+1}}{\sqrt{n}}$$

$$\sum a_n^2 = \sum \frac{1}{n}$$

where the first sum is convergent by Alternating Series Test, and the last is divergent, because it is a harmonic series.

(b) If  $\sum a_n$  converges and  $a_n \geq 0$ , can we conclude anything about  $\sum \sqrt{a_n}$ ?

Not really.

$a_n = \frac{1}{n^2}$  implies that  $\sum a_n$  converges, but  $\sum \sqrt{a_n}$  is a Harmonic Series. For  $a_n = \frac{1}{n^2}$  both  $\sum a_n$  and  $\sum \sqrt{a_n}$  converge (both conclusions are justified by the fact, that we had drawn some time ago that the series  $\sum 1/n^p$  converges if and only if  $p > 1$ ).

## 2.7.6

(a) Show that if  $\sum x_n$  converges absolutely, and the sequence  $(y_n)$  is bounded. then the sum  $\sum x_n y_n$  converges.

Let  $s_n$  be a partial sum for  $\sum |x_n|$  and  $l_n$  be a partial sum for  $\sum |y_n x_n|$ . Then, because  $(y_n)$  is bounded, there exists  $M > 0$  such that  $|y_n| \leq M$ . Therefore

$$l_n = |x_1 y_1| + |x_2 y_2| + \dots + |x_n y_n| \leq M|x_1| + M|x_2| + \dots + M|x_n| = M(|x_1| + |x_2| + \dots + |x_n|) = M s_n$$

Therefore  $\sum x_n$  being absolutely convergent implies that  $\sum M x_n$  is also absolutely convergent, which implies that  $\sum x_n y_n$  is also absolutely convergent.

(b) Find a counterexample that demonstrates that part (a) does not always hold if the convergence of  $\sum x_n$  is conditional

Same as in exercise 2.7.5

$$\sum x_n = \sum \frac{(-1)^{n+1}}{\sqrt{n}}$$

$$(y_n) = \frac{(-1)^{n+1}}{\sqrt{n}}$$

$$\sum y_n x_n = \sum \frac{1}{n}$$

## 2.7.7

Now that we have proved the basic facts about geometric series, supply a proof for Corollary 2.4.7

Corollary 2.4.7 states that the series

$$\sum_{n=1}^{\infty} 1/n^p$$

converges if and only if  $p > 1$

Let us employ Cauchy Condensation Test here. Let us first play with the term of the sequence for it.

$$2^n a_{2^n} = 2^n \frac{1}{(2^n)^p} = \frac{2^n}{2^{np}} = 2^{n-np} = 2^{n(1-p)} = (2^{1-p})^n$$

We know, that geometric series converges if and only if for its term  $ar^k$ ,  $|r| < 1$ . Thus, we can state, that by Cause Condensation Test, the original series converges if and only if

$$|2^{1-p}| < 1$$

$$2^{1-p} < 1$$

$$1 - p < \log_2(1)$$

$$1 - p < 0$$

$$p > 1$$

as desired.

## 2.7.8

*Prove Theorem 2.7.1 part (ii)*

Theorem 2.7.1 part (ii) states that

If  $\sum_{k=1}^{\infty} a_k = A$  and  $\sum_{k=1}^{\infty} b_k = B$  then  $\sum_{k=1}^{\infty} (a_k + b_k) = A + B$

Let  $s_n$  be a partial sum for the series  $\sum_{k=1}^{\infty} a_k = A$ ,  $t_n$  be a partial sum for  $\sum_{k=1}^{\infty} b_k = B$  and  $p_n$  be a partial sum for  $\sum_{k=1}^{\infty} (a_k + b_k)$ . Then

$$s_n + t_n = a_1 + a_2 + \dots + a_n + b_1 + b_2 + \dots + b_n = a_1 + b_1 + a_2 + b_2 + \dots + a_n + b_n = p_n$$

for every  $n \in \mathbb{N}$ .

Thus we can state that  $\lim(a_n + b_n) = A + B$  by Algebraic Limit Theorem. Thus

$$\lim(a_n + b_n) = \lim p_n = \sum (a_n + b_n) = A + B$$

as desired.

## 2.7.9 (Ratio Test)

Given a series  $\sum_{n=1}^{\infty} a_n$  with  $a_n \neq 0$ , the Ratio Test states that if  $(a_n)$  satisfies

$$\lim \left| \frac{a_{n+1}}{a_n} \right| = r < 1$$

then the series converges absolutely.

(a) Let  $r'$  satisfy  $r < r' < 1$ . (Why must such an  $r'$  exist?) Explain why there exists an  $N$  such that  $n \geq N$  implies  $|a_{n+1}| \leq |a_n|r'$

First of all,

$$\left| \frac{a_{n+1}}{a_n} \right| = |a_{n+1}| \frac{1}{|a_n|} = |a_{n+1}| \left| \frac{1}{a_n} \right|$$

If  $a_n \geq 0$ , then  $\left| \frac{1}{a_n} \right| = \frac{1}{a_n} = \frac{1}{|a_n|}$ . If  $a_n < 0$  then  $\left| \frac{1}{a_n} \right| = -\frac{1}{a_n} = \frac{1}{-a_n} = \frac{1}{|a_n|}$ . Therefore  $\left| \frac{1}{a_n} \right| = \frac{1}{|a_n|}$ . Thus

$$\left| \frac{a_{n+1}}{a_n} \right| = |a_{n+1}| \left| \frac{1}{a_n} \right| = \frac{|a_{n+1}|}{|a_n|}$$

Thus

$$\lim \left| \frac{a_{n+1}}{a_n} \right| = \lim \frac{|a_{n+1}|}{|a_n|} = r < 1$$

$r'$  exists, because for any  $\epsilon > 0$  there exists  $n \in \mathbf{N}$  such that  $1/n < \epsilon$ . Therefore if we set  $\epsilon = 1 - r$ , then there exists  $1/n_1 < \epsilon$ . Therefore

$$0 < 1/n_1 < \epsilon$$

$$0 < 1/n_1 < 1 - r$$

$$r < r + 1/n_1 < 1$$

Suppose that we set  $\epsilon = r' - r$ . Then, by definition of limit, there exists  $N \in \mathbf{N}$ , such that for any  $n \geq N$  it follows that

$$\left| \frac{|a_{n+1}|}{|a_n|} - r \right| < \epsilon$$

$$\left| \frac{|a_{n+1}|}{|a_n|} - r \right| < r' - r$$

$$-(r' - r) < \frac{|a_{n+1}|}{|a_n|} - r < r' - r$$

$$-r' + r < \frac{|a_{n+1}|}{|a_n|} - r < r' - r$$



$$-r' + 2r < \frac{|a_{n+1}|}{|a_n|} < r'$$

$$\frac{|a_{n+1}|}{|a_n|} < r'$$

$$|a_{n+1}| < r'|a_n|$$

as desired.

(b) Why does  $|a_N| \sum (r')^n$  necessarily converge?

Because we have  $r$  is a limit of non-negative terms, we can conclude, that  $r \geq 0$ . Thus,  $0 \leq r' < 1$ . Thus,  $|r'| < 1$ . Therefore,  $\sum (r')^n$  is a geometric series with  $|r| < 1$ . Therefore it converges. Therefore  $|a_N| \sum (r')^n$  is convergent. Also,  $\sum |a_N|(r')^n$  is convergent as well.

(c) Now, show that  $\sum |a_n|$  converges.

Let  $s_n$  be a partial sum for  $\sum a_n$  and let  $q_n$  be a partial sum for  $\sum |a_N|r'^n$ .

Let  $\epsilon$  be arbitrary. Then, there exists  $N_1 \in \mathbf{N}$  such that  $n \geq N_1 \rightarrow |a_{n+1}| \leq |a_n|r'$ .  $\sum a_n$ . There also exists  $N_2 \in \mathbf{N}$  such that  $m > n \geq N_2$  implies that  $|q_m - q_n| < \epsilon$  (this comes from Cauchy Criterion for Series). Then let  $N = \max\{N_1, N_2\}$ .

Then for  $m > n \geq N$ .

$$\begin{aligned} |s_m - s_n| &= ||a_{n+1}| + |a_{n+2}| + \dots |a_m|| = |a_{n+1}| + |a_{n+2}| + \dots |a_m| \leq \\ &\leq |a_n|r' + |a_{n+1}|r' + \dots + |a_{m-1}|r' \leq |a_{n-1}|r'^2 + |a_n|r'^2 + \dots + |a_{m-2}|r'^2 \leq \\ &\leq |a_N|r'^{N-n} + |a_N|r'^{N-n+1} + \dots + |a_N|r'^{N-m} = |q_m - q_n| < \epsilon \end{aligned}$$

Therefore there exists  $N \in \mathbf{N}$  such that for every  $m > n \geq N$

$$|s_m - s_n| < \epsilon$$

Thus, by Cauchy criterion for series, the series  $\sum |a_n|$  is convergent. Thus  $\sum a_n$  is absolutely convergent.

## 2.7.10

(a) Show that if  $a_n > 0$  and  $\lim(na_n) = l$  with  $l \neq 0$ , then the series  $\sum a_n$  diverges.

I think that for this one the strategy is to somehow compare element of a given sequence, and then conclude, that the partial sum of a desired series is more than a partial sum for harmonic series, and therefore is divergent.

First of all,  $n \in N \rightarrow n \geq 0$ , therefore  $a_n n \geq 0$ . Therefore we can state that  $l \geq 0$ .

Let  $(s_n)$  be a sequence of partial sums for  $\sum a_n$ . Then let  $\epsilon = l$ . Then there exists  $N \in \mathbf{N}$  such that for all  $n \in \mathbf{N} \geq N$  it follows that

$$|na_n - l| > \epsilon$$

$$na_n - l > \epsilon$$

$$na_n - l > l$$

$$na_n > 2l$$

$$a_n > \frac{2l}{n}$$

Let us now look at the Harmonic Series. Define  $q_n$  to be a partial sum for the Harmonic Series. We know, that it is divergent. Thus, there exists an  $\epsilon$ , such that

$$|q_m - q_n| \geq \epsilon$$

for any  $m > n \in \mathbf{N}$ . Thus

$$\left| \frac{1}{m} + \frac{1}{m-1} + \dots + \frac{1}{n+1} \right| \geq \epsilon$$

$$\frac{1}{m} + \frac{1}{m-1} + \dots + \frac{1}{n+1} \geq \epsilon$$

$$\frac{2l}{m} + \frac{2l}{m-1} + \dots + \frac{2l}{n+1} \geq 2l\epsilon$$

$$a_m + a_{m-1} + \dots + a_{n+1} \geq 2l\epsilon$$

$$|s_m - s_n| \geq 2l\epsilon$$

Thus, there exists  $2l\epsilon \in R$  such that for every  $m > n \in \mathbf{N}$  it follows that

$$|s_m - s_n| \geq 2l\epsilon$$

Thus, by Cauchy Criterion for series, the series  $\sum a_n$  is divergent.

(b) Assume  $a_n > 0$  and  $\lim(n^2 a_n)$  exists. Show that  $\sum a_n$  converges.

Suppose that all assumption in the exercise are true. Then, by definition of limit, for every  $\epsilon > 0$  there exists  $N \in \mathbf{N}$  such that if  $n \geq N$ , then it follows that

$$|n^2 a_n - l| < \epsilon$$

$$-\epsilon < n^2 a_n - l < \epsilon$$

$$-\epsilon + l < n^2 a_n < l + \epsilon$$

$$\frac{l - \epsilon}{n^2} < a_n < \frac{l + \epsilon}{n^2}$$

$$0 < a_n < \frac{l + \epsilon}{n^2}$$

We know, that the sum  $\sum 1/n^2$  is convergent. Therefore  $\sum \frac{l+\epsilon}{n^2}$  is convergent as well for any  $l + \epsilon \in R$ . Thus, by Comparison Test,  $\sum a_n$  is convergent as well, as desired.

## 2.7.11

Find examples of two series  $\sum a_n$  and  $\sum b_n$  both of which diverge, but for which  $\sum \min\{a_n, b_n\}$  converges. To make it more challenging, produce examples where  $(a_n)$  and  $(b_n)$  are positive and decreasing.

If the relax the exercise and try to make just normal versions of given things, then we'll get

$$a_x = \begin{cases} 1 & \text{if } n \text{ is even} \\ 1/n^2 & \text{if } n \text{ is odd} \end{cases} \quad (12)$$

and

$$b_x = \begin{cases} 1 & \text{if } n \text{ is odd} \\ 1/n^2 & \text{if } n \text{ is even} \end{cases} \quad (13)$$

Then the obvious minimum will converge, as desired. In this case we even have every element of every sequence to be positive.

The idea behind this one is probably to construct the desired case from "barely" divergent sequences and series (In this case we probably need something along the lines of Harmonic Series )

It (probably) can be shown, that the desired sequences are convergent to zero (or at the very least one of them).

On this particular exercise we are going to assume some things without any hope for proof or concrete base, and we are going to pull'em out of thin air. First of all: the required series are most likely to be "barely" divergent. Secondly, required sequences (not series) are probably convergent to 0. It's actually hard to believe, that such sequences exist.

I looked this exercise up on the internet for some hints, and I got some. Suppose that:

$a_n$  - equal to  $1/n^2$  for "odd" turns (i.e. for [1], [3, 4], [7, 8, 9], ...), and maximum of itself for other terms  $b_n$  - equal to  $1/n^2$  for "even" turns (i.e. for [2], [5, 6], [10, 11, 12], ...), and maximum of itself for other terms

Then I think, that both of them diverge, but cannot prove it. There exists a more robust example, but I think, that it is pretty cursed anyways. I'm done with this exercise anyways, not gonna spend more time on it.

## 2.7.12 (Summation by Parts)

Let  $(x_n)$  and  $(y_n)$  be sequences, and let  $s_n = x_1 + x_2 + \dots + x_n$ . Use the observation that  $x_j = s_j - s_{j-1}$  to verify formula

$$\sum_{j=m+1}^n x_j y_j = s_n y_{n+1} - s_m y_{m+1} + \sum_{j=m+1}^n s_j (y_j - y_{j+1})$$

$$\begin{aligned}
& s_n y_{n+1} - s_m y_{m+1} + \sum_{j=m+1}^n s_j (y_j - y_{j+1}) = \\
& = s_n y_{n+1} - s_m y_{m+1} + s_{m+1} (y_{m+1} - y_{m+2}) + s_{m+2} (y_{m+2} - y_{m+3}) + \dots + s_n (y_n - y_{n+1}) = \\
& = s_n y_{n+1} - s_m y_{m+1} + s_{m+1} y_{m+1} - s_{m+1} y_{m+2} + s_{m+2} y_{m+2} - s_{m+2} y_{m+3} + \dots + s_n y_n - s_n y_{n+1} = \\
& = -s_m y_{m+1} + s_{m+1} y_{m+1} - s_{m+1} y_{m+2} + s_{m+2} y_{m+2} - s_{m+2} y_{m+3} + \dots + s_{n-1} y_{n-1} - s_{n-1} y_n + s_n y_n = \\
& = y_{m+1} (s_{m+1} - s_m) + y_{m+2} (s_{m+2} - s_{m+1}) + \dots + y_n (s_n - s_{n-1}) = \\
& = y_{m+1} x_{m+1} + y_{m+2} x_{m+2} + \dots + y_n x_n = \sum_{j=m+1}^n x_j y_j
\end{aligned}$$

as desired.

### 2.7.13 (Dirichlet's Test)

*Dirichlet's Test for convergence states that if the partial sums of  $\sum_{n=1}^{\infty} x_n$  are bounded (but not necessarily convergent), and if  $(y_n)$  is a sequence satisfying  $y_1 \geq y_2 \geq y_3 \geq \dots \geq 0$  with  $\lim y_n = 0$ , then the series  $\sum_{n=1}^{\infty} x_n y_n$  converges.*

(a) Let  $M > 0$  be an upper bound for the partial sums of  $\sum_{n=1}^{\infty} x_n$ . Use Exercise 2.7.12 to show that

$$\left| \sum_{j=m+1}^n x_j y_j \right| \leq 2M |y_{m+1}|$$

Suppose that everything in exercise is true. Then it follows that  $|s_j| < M$ . Then

$$\begin{aligned}
\left| \sum_{j=m+1}^n x_j y_j \right| &= \left| s_n y_{n+1} - s_m y_{m+1} + \sum_{j=m+1}^n s_j (y_j - y_{j+1}) \right| \leq \\
&\leq |s_n y_{n+1}| + |s_m y_{m+1}| + \left| \sum_{j=m+1}^n s_j (y_j - y_{j+1}) \right| = \\
&\leq |s_n| |y_{n+1}| + |s_m| |y_{m+1}| + |M| \left| \sum_{j=m+1}^n (y_j - y_{j+1}) \right| \leq \\
&\leq M |y_{n+1}| + M |y_{m+1}| + M \left| \sum_{j=m+1}^n (y_j - y_{j+1}) \right| =
\end{aligned}$$

$$\begin{aligned}
&= M \left( |y_{n+1}| + |y_{m+1}| + \left| \sum_{j=m+1}^n (y_j - y_{j+1}) \right| \right) \leq \\
&\leq M (|y_{n+1}| + |y_{m+1}| + |y_{m+1} - y_{m+2} + y_{m+2} - y_{m+3} + \dots + y_n - y_{n+1}|) = \\
&= M (|y_{n+1}| + |y_{m+1}| + |y_{m+1} - y_{n+1}|) = \\
&= M (y_{n+1} + y_{m+1} + y_{m+1} - y_{n+1}) = 2M|y_{m+1}|
\end{aligned}$$

Therefore

$$\left| \sum_{j=m+1}^n x_j y_j \right| \leq 2M|y_{m+1}|$$

as desired

(b) *Prove Dirichlet's Test just stated.*

Suppose that we are given some  $\epsilon > 0$ . By the convergence of  $(y_n)$ , there exists  $N$ , such that for  $\epsilon_1 = \epsilon/2M$  and  $m+1 \geq N$  it follows that

$$\begin{aligned}
|y_{m+1}| &< \epsilon_1 \\
|y_{m+1}| &< \epsilon/2M \\
2M|y_{m+1}| &< \epsilon \\
\left| \sum_{j=m+1}^n x_j y_j \right| &\leq 2M|y_{m+1}| < \epsilon \\
\left| \sum_{j=m+1}^n x_j y_j \right| &< \epsilon
\end{aligned}$$

Thus, by Cauchy Criterion for Series, it follows that  $\sum_{n=1}^{\infty} x_j y_j$  converges, as desired.

(c) *Show how the Alternating Series Test (Theorem 2.7.7) can be derived as a special case of Dirichlet's Test.*

Firstly, 2.7.7 states that

**Theorem 2.7.7 (Alternating Series Test)** *Let  $(a_n)$  be a sequence satisfying*

$$(i) \ a_1 \geq a_2 \geq a_3 \geq \dots \geq a_n \geq \dots \text{ and}$$

$$(ii) \ (a_n) \rightarrow 0$$

*Then, the alternating series  $\sum_{n=1}^{\infty} (-1)^{n+1} a_n$  converges*

We can set  $x_n = (-1)^{n+1}$ , which is gonna be bounded by  $[-1, 1]$ , and  $y_n = a_n$ , which converges to 0 and satisfies all of the necessary prerequisites for Dirichlet's Test to be applied.

## 2.7.14 (Abel's Test).

Abel's Test for convergence states that if the series  $\sum_{n=1}^{\infty} x_n$  converges, and if  $y_n$  is a sequence satisfying

$$y_1 \geq y_2 \geq y_3 \geq \dots \geq 0$$

then the series  $\sum_{n=1}^{\infty} x_n y_n$  converges.

(a) Carefully point out how the hypothesis of Abel's Test differs from that of Dirichlet's Test in Exercise 2.7.13.

First of all, Abel's Test requires  $\sum x_n$  to be not only bounded, but convergent as well. Secondly,  $(y_n)$  is not required to be convergent.

(b) Assume that  $\sum a_n$  has partial sums, that are bounded by a constant  $A > 0$ , and assume  $b_1 \geq b_2 \geq b_3 \geq \dots \geq 0$ . Use Exercise 2.7.12 to show that

$$\left| \sum_{j=1}^n a_j b_j \right| \leq 2Ab_1$$

let  $s_n$  be a sequence of partial sums for  $\sum a_n$ . Then

$$\begin{aligned} \left| \sum_{j=1}^n a_j b_j \right| &= \left| s_n b_{n+1} - s_1 b_2 + \sum_{j=1}^n s_j (b_j - b_{j+1}) \right| \leq \\ &\leq |s_n b_{n+1}| + |s_1 b_2| + \left| \sum_{j=1}^n s_j (b_j - b_{j+1}) \right| \\ &\leq |s_n b_{n+1}| + |s_1 b_2| + \sum_{j=1}^n |s_j (b_j - b_{j+1})| \\ &\leq |s_n b_{n+1}| + |s_1 b_2| + \sum_{j=1}^n |s_j (b_j - b_{j+1})| \\ &\leq |s_n| |b_{n+1}| + |s_1| |b_2| + \sum_{j=1}^n |s_j| |b_j - b_{j+1}| \\ &\leq A |b_{n+1}| + A |b_2| + \sum_{j=1}^n A |b_j - b_{j+1}| \\ &\leq A (|b_{n+1}| + |b_2| + \sum_{j=1}^n |b_j - b_{j+1}|) \end{aligned}$$

$$\begin{aligned}
&\leq A(b_{n+1} + b_1 + \sum_{j=1}^n (b_j - b_{j+1})) \\
&\leq A(b_{n+1} + b_1 + b_1 - b_2 + b_2 - b_3 + \dots + b_n - b_{n+1}) \\
&\leq A(b_1 + b_1) = 2Ab_1
\end{aligned}$$

as desired.

(c) *Prove Abel's Test via the following strategy. For a fixed  $m \in \mathbf{N}$ , apply part (b) to  $\sum_{j=m+1}^n x_j y_j$  by setting  $a_n = x_{m+n}$  and  $b_n = y_{m+n}$ . (Argue that an upper bound on the partial sums of  $\sum_{n=1}^{\infty} a_n$  can be made arbitrarily small by taking  $m$  to be large.)*

Let us fix  $m \in \mathbf{N}$ . Then set  $a_n = x_{m+n}$  and  $b_n = y_{m+n}$ . Then  $\sum_{j=1}^n a_j$  is equal to  $\sum_{m+1}^{m+n} x_n$ . Thus, for every  $\epsilon > 0$  we can find  $m \in \mathbf{N}$ , such that  $\sum_{j=1}^n a_j < \epsilon$  by convergence of  $\sum x_j$  and by using Cauchy Criterion. Therefore we always have  $A = \epsilon$ . Thus

$$\left| \sum_{j=m+1}^n x_j y_j \right| = \left| \sum_{j=1}^n a_n b_n \right| < 2Ab_1$$

Therefore by Cauchy Criterion for Series,  $\sum x_j y_j$  is convergent.

### 2.8.1

*Using the particular array  $(a_{ij})$  from Section 2.1, compute  $\lim_{n \rightarrow \infty} s_{nn}$ . How does this value compare to the two iterated values for the sum already computed?*

Firstly, let us state the requested array:  $\{a_{ij} : i, j \in \mathbf{N}\}$ , where  $a_{ij} = 1/2^{j-i}$  if  $j > i$ ,  $a_{ij} = -1$  if  $j = i$ , and  $a_{ij} = 0$  if  $j < i$  (I'd gladly portraint the grid here if I knew how to do it).

I don't think, that we need concrete proofs and/or some other rigorous stuff, so it'll suffice to show that

$$s_{1,1} = -1$$

$$s_{2,2} = -3/2$$

$$s_{3,3} = -7/4$$

Or in general,  $(s_{n,n}) \rightarrow -2$ , or in other words,

$$\lim_{n \rightarrow \infty} s_{n,n} = -2$$

as desired.

The resulting value is equal to one of the earlier computed results.

## 2.8.2

Show that if the iterated series

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} |a_{ij}|$$

converges (meaning that for each fixed  $i \in \mathbf{N}$  the series  $\sum_{j=1}^{\infty} |a_{ij}|$  converges to some real number  $b_i$ , and the series  $\sum_{i=1}^{\infty} b_i$  converges as well), then the iterated series

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} a_{ij}$$

converges.

Suppose that

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} |a_{ij}|$$

converges. This implies that for every  $i \in \mathbf{N}$  it is true that  $\sum_{j=1}^{\infty} |a_{ij}|$  converges. This implies that  $\sum_{j=1}^{\infty} a_{ij}$  converges. It is also true, that  $\sum_{i=1}^{\infty} b_i$  converges. Triangular inequality implies that

$$b_i = \sum_{j=1}^{\infty} |a_{ij}| \geq \left| \sum_{j=1}^{\infty} a_{ij} \right|$$

$$0 \leq \left| \sum_{j=1}^{\infty} a_{ij} \right| \leq b_i$$

Thus, by Comparison Test

$$\sum_{i=1}^{\infty} \left| \sum_{j=1}^{\infty} a_{ij} \right|$$

converges. Thus, by Absolute Convergence Test

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} a_{ij}$$

converges too, as desired.



### 2.8.3

**Theorem 2.8.1.** Let  $\{a_{ij} : i, j \in \mathbf{N}\}$  be a doubly indexed array of real numbers. If

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} |a_{ij}|$$

converges, then both  $\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} a_{ij}$  and  $\sum_{j=1}^{\infty} \sum_{i=1}^{\infty} a_{ij}$  converge to the same value. Moreover,

$$\lim_{n \rightarrow \infty} s_{nn} = \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} a_{ij} = \sum_{j=1}^{\infty} \sum_{i=1}^{\infty} a_{ij}$$

where  $s_{nn} = \sum_{i=1}^n \sum_{j=1}^n a_{ij}$

(a) Prove that the set  $\{t_{mn} : m, n \in \mathbf{N}\}$  is bounded above, and use this fact to conclude that the sequence  $(t_{nn})$  converges.

Because

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} |a_{ij}| = l$$

is convergent, we can state that  $t_{mn}$  is less than  $l$ . Also,  $t_{m,m} - t_{n,n} \geq 0$  for  $m \geq n$ . Thus, the sequence is bounded and monotone, and therefore convergent.

(b) Now, use the fact that  $(t_{nn})$  is a Cauchy sequence to argue that  $(s_{nn})$  is a Cauchy sequence and hence converges.

Let  $\epsilon > 0$ . Then there exists  $N \in \mathbf{N}$  such that for  $m > n \in \mathbf{N}$  it follows that

$$|t_{mm} - t_{nn}| < \epsilon$$

$$||a_{m,m}| + |a_{m,m-1}| + \dots + |a_{n+1,n+1}|| < \epsilon$$

$$|a_{m,m}| + |a_{m,m-1}| + \dots + |a_{n+1,n+1}| < \epsilon$$

$$|a_{m,m} + a_{m,m-1} + \dots + a_{n+1,n+1}| \leq |a_{m,m}| + |a_{m,m-1}| + \dots + |a_{n+1,n+1}| < \epsilon$$

$$|a_{m,m} + a_{m,m-1} + \dots + a_{n+1,n+1}| < \epsilon$$

$$|s_{mm} - s_{nn}| < \epsilon$$

Therefore  $(s_{nn})$  is a Cauchy sequence, and therefore it is convergent, as desired.

## 2.8.4

(a) Argue that there exists an  $N_1 \in \mathbf{N}$  such that  $m, n \geq N_1$  implies  $B - \frac{\epsilon}{2} < t_{mn} \leq B$ .

There exist  $m_0, n_0 \in \mathbf{N}$  such that

$$B - \frac{\epsilon}{2} < t_{m_0, n_0} \leq B$$

We know, that  $t_{m_1, n_0} - t_{m_0, n_0} \geq 0$  for  $m_1 \geq m_0$ . Thus, if we pick  $N_1 = \max\{m_0, n_0\}$ , then  $m, n \geq N_1$  implies that

$$B - \frac{\epsilon}{2} < t_{m, n} \leq B$$

as desired

(b) Now, show that there exists an  $N$  such that

$$|s_{mn} - S| < \epsilon$$

for all  $m, n \geq N$ .

We know, that  $(t_{n, n})$  is a Cauchy sequence, and thus, for the given  $\epsilon$  there exists  $N$ , such that  $m > n \geq N$  implies that

$$|t_{m, m} - t_{n, n}| < \epsilon/2$$

$$||a_{m, m}| + |a_{m-1, m}| + \dots + |a_{n+1, n+1}|| < \epsilon/2$$

$$|a_{m, m}| + |a_{m-1, m}| + \dots + |a_{n+1, n+1}| < \epsilon/2$$

$$|a_{m, m}| + |a_{m-1, m}| + \dots + |a_{n+1, m}| < \epsilon/2$$

$$|a_{m, m} + a_{m-1, m} + \dots + a_{n+1, m}| < \epsilon/2$$

$$|a_{m, m} + a_{m-1, m} + \dots + a_{n+1, m}| < \epsilon/2$$

$$|s_{nn} - S| < \epsilon/2$$

$$|s_{nn} - S| + |a_{m, m} + a_{m-1, m} + \dots + a_{n+1, m}| < \epsilon$$

$$|s_{nn} + a_{m, n} + \dots + a_{n+1, n} - S| \leq |s_{nn} - S| + |a_{m, m} + a_{m-1, m} + \dots + a_{n+1, m}| < \epsilon$$

$$|s_{nn} + a_{m, n} + \dots + a_{n+1, n} - S| < \epsilon$$

$$|s_{mn} - S| < \epsilon$$

Thus, there exists  $N \in \mathbf{N}$  such that  $m, n \geq N$  implies

$$|s_{mn} - S| < \epsilon$$

as desired

## 2.8.5

Use the Algebraic Limit Theorem (Theorem 2.3.3) and the Order Limit (Theorem 2.3.4) to show that for all  $m \geq N$

$$|(r_1 + r_2 + \dots + r_m) - S| \leq \epsilon$$

Conclude that the iterated sum  $\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} a_{ij}$  converges to  $S$ .

We know, that for every  $\epsilon > 0$  there exists  $N \in \mathbf{N}$  such that for  $m, n \geq N$  it follows that

$$|s_{mn} - S| < \epsilon$$

We also should state, that because the sum of absolute values converge to some number, then it follows, that  $\sum r_m$  also converges to some number. Therefore we can use both Order Limit Theorem and Algebraic Limit Theorem.

$$\begin{aligned} & \left| \left( \sum_{j=1}^{\infty} a_{1,j} + \sum_{j=1}^{\infty} a_{2,j} + \dots + \sum_{j=1}^{\infty} a_{m,j} \right) - S \right| \leq \epsilon \\ & \left| \sum r_n - \sum s_{mn} \right| = \\ & = \left| \left( \sum_{j=1}^{\infty} a_{1,j} + \sum_{j=1}^{\infty} a_{2,j} + \sum_{j=1}^{\infty} a_{m,j} \right) - \left( \sum_{j=1}^n a_{1,j} + \sum_{j=1}^n a_{2,j} + \dots + \sum_{j=1}^n a_{m,j} \right) \right| = \\ & \left| \sum_{j=n+1}^{\infty} a_{1,j} + \sum_{j=n+1}^{\infty} a_{2,j} + \dots + \sum_{j=n+1}^{\infty} a_{m,j} \right| \leq \\ & \leq \sum_{j=n+1}^{\infty} |a_{1,j}| + \sum_{j=n+1}^{\infty} |a_{2,j}| + \dots + \sum_{j=n+1}^{\infty} |a_{m,j}| \rightarrow 0 \end{aligned}$$

By convergence of  $\sum \sum |a_{m,n}|$  we know, that every  $\sum_{j=1}^{\infty} |a_{m,n}|$  converges to some number. Suppose that it converges to  $r_m$ . Thus we can state that

$$\sum_{j=n+1}^{\infty} |a_{m,j}| = \lim_{j \rightarrow \infty} \left( \sum_{j=1}^{\infty} |a_{m,j}| - \sum_{j=1}^n |a_{m,j}| \right) = \lim_{j \rightarrow \infty} (r_m - r_m) = 0$$

Thus,  $q_n = |\sum r_n - \sum s_{mn}|$  is a sequence, each element of which is less than a sum of elements of sequences, convergent to 0, and thus, by Algebraic Limit Theorem, it itself is also convergent to 0.

Thus,  $\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} a_{ij} = S$ , as desired.

## 2.8.6

Finish the proof by showing that the other iterated sum,  $\sum_{j=1}^{\infty} \sum_{i=1}^{\infty} a_{ij}$ , converges to  $S$  as well. Notice that the same argument can be used once it is established that, for each fixed column  $j$ , the sum  $\sum_{i=1}^{\infty} a_{ij}$  converges to some real number  $c_j$ .

We can make an argument here, that we don't really need two different proofs on the account that it doesn't matter in which particular direction (rows or columns) we construct  $r$ 's.

To make the argument more concrete, substitute  $i$  for  $j$  in the previous section of the proof and you'll get the desired result.

Therefore if  $\sum_{j=1}^{\infty} \sum_{i=1}^{\infty} |a_{ij}|$  exists, then it follows that

$$\lim_{n \rightarrow \infty} s_{nn} = \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} a_{ij} = \sum_{j=1}^{\infty} \sum_{i=1}^{\infty} a_{ij}$$

## 2.8.7

(a) Assuming the hypothesis – and hence the conclusion of Theorem 2.8.1, show that  $\sum_{k=2}^{\infty} d_k$  converges absolutely.

This follows from the fact that  $\sum_{k=2}^{\infty} |d_k|$  is increasing, and bounded above by  $\sum \sum |a_{ij}|$ , and therefore convergent absolutely.

(b) Imitate the strategy in the proof of Theorem 2.8.1 to show  $\sum_{k=2}^{\infty} d_k$  converges to  $S = \lim_{n \rightarrow \infty} s_{nn}$

Let  $n \in \mathbf{N}$  and let  $m = 2n$ . Then it follows, that every  $a_{ij}$ , that comprises  $s_{nn}$  is in  $d_m$ . Also, every  $a_{ij}$  in  $d_m$  is contained in  $s_{mm}$ . Thus, let  $q_k$  be a sequence, such that  $q_k = d_m - s_{nn}$ . We know that  $t_{nn}$  (which is sequence of partial sums for the absolute sum) converges. Thus, if we define  $l_k$  to be an absolute sum of evements of  $q_k$ , then it follows, that

$$|t_m m - t_n n| < \epsilon$$

$$t_m m - t_n n < \epsilon$$

$$t_m m - t_n n < \epsilon$$

$$l_k < \epsilon$$

$$|q_k| < \epsilon$$

Thus,  $(q_k) \rightarrow 0$ . Therefore we can state that

$$\lim d_m = \lim q_m + \lim s_{nn}$$

Thus,  $\lim d_m = S$ , as desired.

## 2.8.8

Assume that  $\sum_{i=1}^{\infty} a_i$  converges to  $A$ , and  $\sum_{i=1}^{\infty} b_i$  converges absolutely to  $B$ .

(a) Show that the set

$$\left\{ \sum_{i=1}^m \sum_{j=1}^n |a_i b_j| : m, n \in \mathbf{N} \right\}$$

is bounded. Use this to show that the iterated sum  $\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} |a_i b_j|$  converges so that we may apply Theorem 2.8.1

First of all, because sum of both sequences converge, we can state that  $(a_n) \rightarrow 0$  and  $(b_n) \rightarrow 0$ . Thus, both of them are bounded. Thus let us take a look at a partial sum of  $\sum_{i=1}^m \sum_{j=1}^n |a_i b_j|$

$$\begin{aligned} s_n &= |a_1 b_1| + |a_2 b_2| + \dots + |a_n b_n| = |a_1| |b_1| + |a_2| |b_2| + \dots + |a_n| |b_n| \leq \\ &\leq |M| |b_1| + |M| |b_2| + \dots + |M| |b_n| = |M| (b_1 + b_2 + \dots + b_n) \end{aligned}$$

where  $M$  is a number such that  $|a_n| < M$ .

Thus we can state that every element of a sequence of partial sums for the original sequence is less or equal to an element of a sequence of partial sums for  $\sum |M b_n|$ . Thus, the set is bounded above by —MB— (by Algebraic Limit Theorem for Series). Because the sum it is a sum of non-negative terms we can conclude, that is is also bounded below by 0 (to be more rigorous, every element in sequence of partial sums is greater, than an element of a sequence of 0's, and therefore is non-negative).

By the fact that this sum is a sum of non-negative terms, we can state that the sequence of partial sums is increasing, and therefore, by MCT, the sum is convergent.

(b) Let  $s_{nn} = \sum_{i=1}^n \sum_{j=1}^n a_i b_j$ , and use the Algebraic Limit Theorem to show that  $\lim_{n \rightarrow \infty} s_{nn} = AB$ . Conclude that

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} a_j b_i = \sum_{j=1}^{\infty} \sum_{i=1}^{\infty} a_j b_i = \sum_{k=2}^{\infty} d_k = AB$$

$$\begin{aligned} s_n &= a_1 b_1 + a_1 b_2 + a_1 b_3 + \dots + a_n b_n = \\ &= a_1 (b_1 + b_2 + \dots + b_n) + a_2 (b_1 + b_2 + \dots + b_n) + \dots + a_n (b_1 + b_2 + \dots + b_n) = \\ &= (a_1 + a_2 + \dots + a_n) (b_1 + b_2 + \dots + b_n) = p_n q_n \end{aligned}$$

where  $p_n$  and  $q_n$  are sequences of partial sums for  $\sum a_n$  and  $b_n$  respectively. Thus we can state that

$$\lim_{n \rightarrow \infty} s_{nn} = \lim_{n \rightarrow \infty} \left( \sum_{i=1}^n a_i \sum_{i=1}^n b_i \right) = AB$$

Therefore, by Theorem 2.8.1 and its consequences

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} a_j b_i = \sum_{j=1}^{\infty} \sum_{i=1}^{\infty} a_j b_i = \sum_{k=2}^{\infty} d_k = AB$$

as desired.

### 3.2.1

(a) Where in the proof of Theorem 3.2.3 part (ii) does the assumption that the collection of open sets be finite get used

Theorem 3.2.3 states that

(ii) The intersection of a finite collection of open sets is open.

The assumption of the finality of the set is used in the fact, that we need the minimum of the epsilons.

(b) Give an example of an infinite collection of nested open sets

$$O_1 \supseteq O_2 \supseteq O_3 \supseteq O_4 \supseteq \dots$$

whose intersection  $\cap_{n=1}^{\infty} O_n$  is closed and nonempty.

First of all, we should state that open is not an opposite of closed in this context. We can get  $O_n = (-\infty; \infty)$ . Then this definition (technically) fits into the requirement of qs

Let  $O_n = (1 - 1/n, 2 + 1/n)$ . Let us then define

$$A = \cap_{n=1}^{\infty} O_n$$

Then it follows that  $x \in A \rightarrow 1 \leq x \leq 2$ . Thus,  $A = [1, 2]$ , which is closed, as desired.

### 3.2.2

Let

$$B = \left\{ \frac{(-1)^n n}{n+1} : n = 1, 2, 3, \dots \right\}$$

(a) Find the limit points of B

1 and -1. This can be seen from the fact that  $\lim_{n \rightarrow \infty} \frac{n}{n+1} = 1$  and  $\lim_{n \rightarrow \infty} \frac{-n}{n+1} = -1$ , where both of the expressions in the limit are subsequences of the original sequence (numbers were not derived by using some process, just pure intuition).

$x \in B$  implies that  $x = \frac{n}{n+1}$  or  $x = \frac{-n}{n+1}$  for some  $n \in \mathbb{N}$ . Thus it can be shown, that -1 and 1 are the only limit points of B.

(b) Is B a closed set?

No, it isn't, because it doesn't contain its limit points. This stems from the fact that

$$\frac{(-1)^n n}{n+1} = 1$$

$$(-1)^n n = n + 1$$

and no such number exists. The same can be shown for  $-1$ .

(c) *Is  $B$  an open set?*

No, because any neighborhood around any element will contain some numbers, that are not in  $B$  (e.g. irrational numbers).

(d) *Does  $B$  contain any isolated points*

Given that none of the limit points are in the  $B$ , we can conclude that the set  $B$  consists exclusively of isolated points.

(e) *Find  $\bar{B}$*

$$\bar{B} = B \cup \{-1, 1\}$$

### 3.2.3

*Decide whether the following sets are open, closed, or neither. If a set is not open, find a point in the set for which there is no  $\epsilon$ -neighborhood contained in the set. If a set is not closed, find a limit point that is not contained in the set.*

(a)  $\mathbf{Q}$

Neither closed, nor open. Any number will suffice for both tests. (proof - look at the irrational numbers).

(b)  $\mathbf{N}$

Not open - any number will suffice. Closed - does not have limit points, and thus contains the set of limit points (i.e. empty set).

(c)  $\{x \in \mathbf{R} : x > 0\}$

Open. Is not closed, does not contain 0.

(d)  $(0, 1]$

Not open, 1. Not closed, 0.

(e)  $\{1 + 1/4 + 1/9 + \dots + 1/n^2 : n \in \mathbf{N}\}$

Not open, number 1 is an example. Not closed, does not contain its supremum.

### 3.2.4

*Prove the converse of Theorem 3.2.5 by showing that if  $x = \lim a_n$  for some sequence  $(a_n)$  contained in  $A$  satisfying  $a_n \neq x$ , then  $x$  is a limit point of  $A$ .*

Firstly, let us state the theorem itself

**Theorem 3.2.5** *A point  $x$  is a limit point of a set  $A$  if and only if  $x = \lim a_n$  for some sequence  $(a_n)$  contained in  $A$  satisfying  $a_n \neq x$  for all  $n \in \mathbf{N}$ .*

Let  $\epsilon > 0$ . Then, by definition of convergence, there exists  $N \in \mathbf{N}$ , such that  $n \geq N \rightarrow a_n \in V_\epsilon(x)$  (don't forget, that  $x = \lim a_n$ ). Therefore there exists  $a_n \in V_\epsilon(x) \neq x$ . Also, because the sequence is contained in  $A$ ,  $a_n \in V_\epsilon(x) \cap A$ . Therefore every  $\epsilon$ -neighborhood of  $x$  intersects  $A$  in some point other than  $x$ . Therefore  $x$  is a limit point, as desired.

### 3.2.5

*Let  $a \in A$ . Prove that  $a$  is an isolated point of  $A$  if and only if there exists an  $\epsilon$ -neighborhood  $V_\epsilon(a)$  such that  $V_\epsilon(a) \cap A = \{a\}$*

**In one direction:**

Suppose that  $a$  is an isolated point of  $A$ . It follows, that it is not a limit point. Thus, by definition of a limit point, there does not exist an  $V_\epsilon(a)$  such that it intersects  $A$  at some other point than  $a$ . In other words,  $V_\epsilon(a) \cap A = \{a\}$ , as desired.

**In other direction:**

We'll prove this one by contradiction. Suppose that  $V_\epsilon(a) \cap A = \{a\}$  and  $a$  is a limit point. Then, there exists a sequence  $(a_n)$  fully contained in  $A$ , that converges to  $a$ . Therefore, there exists  $N \in \mathbf{N}$  such that  $n \geq N \rightarrow a_n \in V_\epsilon(a)$ . Therefore,  $\{a, a_n\} \subseteq V_\epsilon(a) \cap A$ , which is a contradiction. Thus,  $a$  is not a limit point and therefore is an isolated point, as desired.

### 3.2.6

*Prove Theorem 3.2.8*

Firstly, let us state the theorem

**Theorem 3.2.8** *A set  $F \subseteq R$  is closed if and only if every Cauchy sequence contained in  $F$  has a limit that is also an element of  $F$ .*

**In one direction:**

Let  $F \subseteq R$  be closed. We are going to proceed with proof by contradiction.

Suppose that there exists a Cauchy sequence  $(s_n)$ , that is contained in  $F$ , that has a limit outside of  $F$ . Then  $x$  is a limit point of  $F$ . Then it is contained in  $F$ , because  $F$  is closed, which is a contradiction.

**In other direction:**

Assume that every Cauchy sequence contained in  $F$  has a limit that is also an element of  $F$ .

We are going to use a proof by contradiction on this one as well. Suppose that  $F$  is not closed. Then, there exists  $x$ , that is a limit point of  $F$ , that is not in  $F$ . Therefore, there exists a sequence, that is contained in  $F$ , that converges to  $x$ . Then, this sequence is Cauchy. Thus, we have a Cauchy sequence, that is contained in  $F$ , which converges to a point outside of  $F$ , which is a contradiction.



### 3.2.7

Let  $x \in O$  where  $O$  is an open set. If  $(x_n)$  is a sequence converging to  $x$ , prove that all but a finite number of the terms of  $(x_n)$  must be contained in  $O$ .

Because  $x$  is in open set, there exists an  $\epsilon$  such that the neighborhood  $V_\epsilon(x) \subseteq O$ . Thus, by the convergence of  $(x_n)$  to  $x$ , there exists  $N \in \mathbf{N}$  such that  $n \geq N \rightarrow x_n \in V_\epsilon$ . Therefore for this sequence only  $N$  or less elements are outside of  $O$ . Therefore for any convergent sequence  $(x_n)$  it is true, that all but a finite number of terms of  $(x_n)$  are outside of  $O$ , as desired.

### 3.2.8

Given  $A \subseteq \mathbf{R}$ , let  $L$  be the set of all limit points of  $A$ .

(a) Show that the set  $L$  is closed.

Suppose that  $L$  is not closed. Thus, there exists  $x \notin L$ , that is a limit point of  $L$ . Therefore there exists a sequence  $(s_n)$ , that is convergent to  $x$ , that is fully contained in  $L$ . Therefore for every  $\epsilon$  there exists  $N \in \mathbf{N}$  such that  $n \geq N$  implies that  $s_n \in V_\epsilon(x)$ . Because  $V_\epsilon(x)$  is an open set, there exists a neighborhood  $V_{\epsilon_1}(s_n) \subseteq V_\epsilon(x)$ . Because  $(s_n)$  is contained in  $L$ ,  $s_n$  is a limit point of  $A$ . Therefore,  $V_{\epsilon_1}(s_n)$  contains some element of  $A$  other than  $s_n$ . Therefore every neighborhood of  $V_\epsilon(x)$  contains an element of  $A$ . Therefore  $x$  is a limit point of  $A$ . Therefore  $x \in L$ , which is a contradiction. Therefore  $L$  is closed, as desired.

(b) Argue that if  $x$  is a limit point of  $A \cup L$ , then  $x$  is a limit point of  $A$ . Use this observation to furnish a proof for Theorem 3.2.12

Suppose that  $x$  is a limit point of  $A \cup L$ . Then it follows that every  $\epsilon$ -neighborhood of  $x$  intersects  $A \cup L$  at some point other than  $x$ . Let us look at particular  $V_\epsilon(x)$ . We can follow that there exists a point in the neighborhood, that it is in  $A \cup L$  and is distinct from  $x$ . Let us call this point  $a$ . From the fact, that  $a \in A \cup L$  we can state that  $a$  is either in  $A$ ,  $L$ , or both. If the point is in  $L$ , then it follows, that it is a limit point of  $A$ . Because  $a \in V_\epsilon(x)$  and the fact, that  $V_\epsilon(x)$  is an open set, we can state that there exists a neighborhood  $V_{\epsilon_1}(a) \subseteq V_\epsilon(x)$ . Therefore, because  $a$  is a point in  $L$ , is a limit point of  $A$ . Thus  $V_{\epsilon_1}(a)$  intersects  $A$  at some point. Thus, there exists a point  $q \in A$  and  $q \in V_{\epsilon_1}(a)$ . Therefore  $q \in V_\epsilon(x)$ . Therefore any neighborhood of  $x$  contains a point in  $A$ . Therefore it is a limit point of  $A$ , as desired.

Now let us state Theorem 3.2.12

**Theorem 3.2.12** For any  $A \subseteq \mathbf{R}$ , the closure  $\bar{A}$  is a closed set and is the smallest set containing  $A$ .

From what we've argued we can follow that any limit point of  $\bar{A} = A \cup L$  is a limit point of  $A$ , and therefore is an element of  $L$  and therefore an element of  $\bar{A}$ . Thus all limit points of  $\bar{A}$  is in  $\bar{A}$ . Therefore it is a closed set.

Any closed set containing  $A$  must contain  $L$  as well. This shows, that  $\bar{A}$  is the smallest closed set containing  $A$ .

### 3.2.9

(a) If  $y$  is a limit point of  $A \cup B$ , show that  $y$  is either a limit point of  $A$  or a limit point of  $B$  (or both).

We are going to proceed with a proof by contradiction. Suppose that there exists a limit point  $x$  of  $A \cup B$ , that is not a limit point of  $A$  or  $B$ . Then it follows, that there exists a sequence  $(s_n)$ , that is contained in  $A \cup B$ , converges to  $x$  and such that  $s_n \neq x$  for every  $n \in \mathbb{N}$ . Then this sequence contains infinite amount of elements in either  $A$ ,  $B$  or both (rigorous conclusion can be drawn from the argument, that there could not be finite amount of elements of the sequence in both  $A$  and  $B$ ). Therefore there exists a subsequence in either  $A$  or  $B$ , that is convergent to  $x$ . Therefore  $x$  is a limit point for either  $A$  or  $B$ .

(b) Prove that  $\overline{A \cup B} = \bar{A} \cup \bar{B}$

We already concluded, that if  $y$  is a limit point of  $A \cup B$ , then  $y$  is a limit point of  $A$ , or a limit point of  $B$ . In other words,

$$y \in L_{A \cup B} \rightarrow y \in L_A \text{ or } y \in L_B$$

Then it follows that

$$y \in \overline{A \cup B} \rightarrow y \in L_A \text{ or } y \in L_B \text{ or } y \in A \cup B$$

$$y \in \overline{A \cup B} \rightarrow y \in L_A \text{ or } y \in L_B \text{ or } y \in A \text{ or } y \in B$$

$$y \in \overline{A \cup B} \rightarrow y \in A \text{ or } y \in L_A \text{ or } y \in B \text{ or } y \in L_B$$

$$y \in \overline{A \cup B} \rightarrow y \in \bar{A} \text{ or } y \in \bar{B}$$

$$y \in \overline{A \cup B} \rightarrow y \in \bar{A} \cup y \in \bar{B}$$

The usage of the word "or" is a little liberal here, but the idea is the same - both  $\overline{A \cup B}$  and  $\bar{A} \cup \bar{B}$  exhaustively decompose to the same 4 sets, and therefore are equal.

(c) Does the result about closures in (b) extend to infinite

The current argument extends at least to the finite amount of element by the virtue of induction. Something tells me, that this particular thing does not extend though to the infinite amount of element.

One possible example on why this proposition might be false can be drawn from the fact, that closure of any rational number is the number itself, therefore the union of closures of all rational numbers is the set of rational numbers, but the closure of the set of all rational numbers is the set of real numbers.

I'm gonna roll with this particular counter-example as the proof of the fact, that given proposition is false, because I don't see anything that is wrong with it.

Another example is the union of sets, where each set contains one element -  $1/n$  for some  $n \in N$ . Same logic applies, closure of any given set is the set itself, but the closure of union of sets includes zero.

### 3.2.10 (De Morgan's Laws)

A proof for the De Morgan's Laws in the case of two sets is outlined in Exercise 1.2.3. The general argument is similar.

(a) Given a collection of sets  $\{E_\lambda : \lambda \in \Lambda\}$ , show that

$$(\cup_{\lambda \in \Lambda} E_\lambda)^c = \cap_{\lambda \in \Lambda} E_\lambda^c$$

and

$$(\cap_{\lambda \in \Lambda} E_\lambda)^c = \cup_{\lambda \in \Lambda} E_\lambda^c$$

Suppose that  $x \in (\cup_{\lambda \in \Lambda} E_\lambda)^c$ . It follows, that  $x$  is not in the union of given sets. Therefore there is no set  $E_n$  such that  $x \in E_n$  (because if there would be such a set, then  $x$  wouldn't be in  $(\cup_{\lambda \in \Lambda} E_\lambda)^c$ ). Therefore  $x \in \cap_{\lambda \in \Lambda} E_\lambda^c$ . Therefore

$$(\cup_{\lambda \in \Lambda} E_\lambda)^c \subseteq \cap_{\lambda \in \Lambda} E_\lambda^c$$

The proof of reverse inclusion is the same as with the forward, but in reverse order.

$x \in (\cap_{\lambda \in \Lambda} E_\lambda)^c$  implies that  $x$  is not in every  $E_n$ . Therefore there exists  $x \in E_n^c$  for some  $E_n$ . therefore it is in  $\cup_{\lambda \in \Lambda} E_\lambda^c$ . The proof of reverse inclusion uses the same argument, but in other direction.

(b) Now, provide the details for the proof of Theorem 3.2.14

Let us state the theorem first.

**Theorem 3.2.14** (i) The union of a finite collection of closed sets is closed.

(ii) The intersection of an arbitrary collection of closed sets is closed

Let  $\{E_\lambda : \lambda \in \Lambda\}$  be a finite collection of closed sets. Therefore  $\{E_\lambda^c : \lambda \in \Lambda\}$  is a finite collection of open sets. Therefore intersection

$$\cap_{\lambda \in \Lambda} E_\lambda^c = (\cup_{\lambda \in \Lambda} E_\lambda)^c$$

is open. Therefore  $\cup_{\lambda \in \Lambda} E_\lambda$  is closed, as desired.

Let  $\{E_\lambda : \lambda \in \Lambda\}$  be a arbitrary collection of closed sets. Therefore  $\{E_\lambda^c : \lambda \in \Lambda\}$  is a arbitrary collection of open sets. Therefore union

$$\cup_{\lambda \in \Lambda} E_\lambda^c = (\cap_{\lambda \in \Lambda} E_\lambda)^c$$

is open. Therefore  $\cap_{\lambda \in \Lambda} E_\lambda$  is closed, as desired.

### 3.2.11

Let  $A$  be bounded above so that  $s = \sup A$  exists. Show that  $s \in \overline{A}$

For any  $\epsilon > 0$  there exists an element  $a \in A$  such that

$$a > s - \epsilon$$

$$a - s > -\epsilon$$

$$s - a < \epsilon$$

$s$  is a supremum of  $A$ , therefore  $s \geq a$ , therefore  $s - a \geq 0$ . Thus  $s - a = |s - a|$ . Therefore

$$|s - a| < \epsilon$$

Therefore  $a \in V_\epsilon(s)$  for any  $\epsilon > 0$ . Thus,  $s$  is a limit point of  $A$ . Therefore  $s \in \overline{A}$ , as desired.

### 3.2.12

Decide whether the following statements are true or false. Provide counterexamples for those that are false, and supply proofs for those that are true.

(a) For any set  $A \subseteq \mathbf{R}$ ,  $\overline{A}^c$  is open.

True. Closure of any set is closed, therefore complement of closure is open.

(b) If a set  $A$  has an isolated point, it cannot be an open set

True. We're gonna proceed with a proof by contradiction. Suppose that  $A$  is an open set and  $x \in A$  is an isolated point. Then there exists a neighborhood  $V_\epsilon(x) \subseteq A$ . Therefore any neighborhood of  $x$  intersects  $A$  at some point (this can be proven rigorously, but it's tedious and redundant). Therefore  $x$  is a limit point of  $A$ , which is a contradiction. Therefore it is not isolated, therefore we have a contradiction.

(c) A set  $A$  is closed if and only if  $\overline{A} = A$ .

True

In one direction: if  $A$  is closed, then  $L \subseteq A$ , therefore  $\overline{A} = A \cup L = A$ .

In other direction:  $\overline{A} = A \cup L = A$  implies that  $L \subseteq A$ . Therefore  $A$  is closed. (By the way,  $L$  denotes a set of limit points of  $A$ )

(d) If  $A$  is a bounded set, then  $s = \sup A$  is a limit point of  $A$

False. Any finite set is counterexample.

(e) Every finite set is closed.

True. Finite sets consist of isolated points, therefore they don't have limit points, therefore they contain their limit points.

(f) An open set that contains every rational number must necessarily be all of  $\mathbf{R}$ .

False.  $\mathbf{R} \setminus \{\sqrt{2}\}$ .

### 3.2.13

*Prove that the only sets that are both open and closed are  $R$  and the empty set  $\emptyset$*

Suppose that there exists a set  $B \neq R \neq \emptyset$ , that is both closed and open. Then let  $x \in B^c$ . It follows, that  $x$  cannot be a limit point of  $B$  (because if it was, then it would be in  $B$ , because  $B$  is closed). Therefore we have an isolated point in an open set  $B^c$ , which is impossible (as proven in the previous exercise). Therefore we have a point which is neither a limit point, nor a isolated point, which is a contradiction.

Same argument doesn't work on  $R$  and  $\emptyset$ , because of the fact, that  $\emptyset$  doesn't contain any points.

### 3.2.14

*A set  $A$  is called  $F_\sigma$  set if it can be written as the countable union of closed sets. A set  $B$  is called  $G_\delta$  set if it can be written as the countable intersection of open sets*

*(a) Show that a closed interval  $[a, b]$  is a  $G_\delta$  set.*

Let there be a collection of open sets, defined by

$$\{(-1/n + a, b + 1/n) : n \in N\}$$

Then its intersection is equal to  $[a, b]$  (This can be shown by using the fact, that  $[a, b]$  is in every set of the collection, but any number outside of  $[a, b]$  isn't). Therefore it is a  $G_\delta$  set.

*(b) Show that the half-open interval  $(a, b]$  is both a  $G_\delta$  and an  $F_\sigma$  set.*

The intersection of

$$\{(a, b + 1/n) : n \in N\}$$

is equal to  $(a, b]$ , therefore it is a  $G_\delta$  set.

The union of

$$\{[a + 1/n, b] : n \in N\}$$

is equal to  $(a, b]$  (if  $a + 1/n < b$ , then let the set corresponding be an empty set), therefore it is a  $F_\sigma$  set.

*(c) Show that  $Q$  is an  $F_\sigma$ , and set of irrationals  $I$  forms a  $G_\sigma$  set.*

We know, that  $Q$  is a countable set. Also, singleton  $\{m\}$  is a closed set. Therefore union of  $\{m : m \in Q\}$  is a countable union of closed sets, that is equal to  $Q$ . Therefore  $Q$  is a  $F_\sigma$  set.

Also, let  $\{R \setminus m : m \in Q\}$  be a collection of open sets. Then its intersection is equal to  $I$ . Therefore  $I$  is a  $G_\delta$  set.

### 3.3.1

*Show that if  $K$  is compact, then  $\sup K$  and  $\inf K$  both exist and are elements of  $K$*

If  $K$  is compact, then it is bounded (i.e. bounded above and below), and thus have supremum and infimum.

Now let us take into account the fact, that for every  $\epsilon > 0$  it is true, that there exist elements  $k_1$  and  $k_2$  (with a possibility that  $k_1 = k_2$ ) such that

$$k_1 > \sup K - \epsilon$$

$$k_2 < \inf K + \epsilon$$

thus

$$\sup K - k_1 < \epsilon$$

$$k_2 - \inf K < \epsilon$$

because of the properties of supremum and infimum  $\sup K - k_1 \geq 0$  and  $k_2 - \inf K \geq 0$ . Thus

$$|\sup K - k_1| < \epsilon$$

$$|k_2 - \inf K| < \epsilon$$

Therefore we can state, that every neighborhood around supremum and infimum has a member of  $K$  in it. Thus  $\sup K$  and  $\inf K$  are limit points of  $K$ . Thus, because  $K$  is closed, we can conclude that  $\sup K \in K$  and  $\inf K \in K$ , as desired.

### 3.3.2

*Prove the converse of Theorem 3.3.4 by showing that if a set  $K \subseteq \mathbb{R}$  is closed and bounded, then it is compact.*

Firstly, let us state the theorem itself:

**Theorem 3.3.4** *A set  $K \subseteq \mathbb{R}$  is compact if and only if it is closed and bounded.*

Let  $(s_n)$  be a sequence, that is contained in  $K$ . Thus, because  $K$  is bounded,  $(s_n)$  is bounded as well. Thus, by BW, it has a convergent subsequence  $(s_{n_k})$ . If this subsequence has its limit as an element of subsequence (i.e.  $s_{n_k} = c$  for some  $c \in \mathbb{R}$  and some of  $n_k \in \mathbb{N}$ ), then its limit is in  $K$  by the virtue of the fact, that  $s_n \in K$  for all  $n \in \mathbb{N}$ . If subsequence doesn't have its limit as its element, then we can state, that its limit is a limit point of  $K$ , and therefore in  $K$  by the fact, that  $K$  is closed.

### 3.3.3

*Show that the cantor set defined in Section 3.1 is compact set.*

I'll not provide a definition here, because it is too lengthy, but you can look it up online. The Cantor set is bounded by definition, therefore we don't need to worry about that. Now by the same definition, the complement of Cantor set is a infinite union of open sets, therefore the complement of Cantor set is an open set itself. Thus, we can state that a complement of a complement (i.e. the set itself) is closed. Thus, Cantor set is closed and bounded, and therefore compact, as desired.

### 3.3.4

*Show that if  $K$  is compact and  $F$  is closed, then  $K \cap F$  is compact.*

The intersection of arbitrary collection of closed sets is closed, therefore  $K \cap F$  is closed as well.  $x \in K \cap F \rightarrow x \in K \rightarrow |x| \leq M$ , where  $M$  is a bound of  $K$ . Therefore  $K \cap F$  is bounded. Therefore  $K \cap F$  is compact, as desired.

### 3.3.5

*Decide which of the following sets are compact. For those that are not compact, show how Definition 3.3.1 breaks down. In other words, give an example of a sequence contained in the given set that does not possess a subsequence converging to a limit in the set.*

(a)  $\mathbf{Q}$

Not compact.  $(s_n) = \{1, 1.4, 1.41, 1.414, 1.4142, \dots\}$  - sequence, that converges to  $\sqrt{2}$ .

(b)  $\mathbf{Q} \cap [0, 1]$

Not compact.  $(s_n) = \{0.7, 0.7, 0.707, 0.7071, \dots\}$  - sequence, that converges to  $\sqrt{2}/2$ .

(c)  $\mathbf{R}$

Not compact  $(x_n) = n$  - doesn't converge, and none of its subsequences converge as well.

(d)  $\mathbf{R} \cap [0, 1]$

Compact.

(e)  $\{1, 1/2, 1/3, 1/4, \dots\}$

Not compact.  $(s_n) = 1/n$  converges to 0, but 0 is not in the set.

(d)  $\{1, 1/2, 2/3, 3/4, 4/5, \dots\}$

Compact.

### 3.3.6

*As some more evidence of the surprising nature of the Cantor set, follow these steps to show that the sum  $C + C = \{x + y : x, y \in C\}$  is equal to the closed interval  $[0, 2]$ . (Keep*

in mind that  $C$  has zero length and contains no intervals. )

The observation that  $\{x + y : x, y \in C\} \subseteq [0, 2]$  passes for obvious, so we only need to prove the reverse inclusion  $[0, 2] \subseteq \{x + y : x, y \in C\}$ . Thus, given  $s \in [0, 2]$ , we must find two elements  $x, y \in C$  satisfying  $x + y = s$ .

(a) Show that there exist  $x_1, y_1 \in C_1$  for which  $x_1 + y_1 = s$ . Show in general that, for an arbitrary  $n \in \mathbf{N}$ , we can always find  $x_n, y_n \in C_n$  for which  $x_n + y_n = s$ .

If  $s \in C_1$ , then we can set  $x_n = s$ .

If  $s \in (1/3, 2/3)$  pick  $x_n = 1/3$ .

If  $s \in (1, 4/3]$  pick  $x_n = 1$

If  $s \in (4/3, 2]$  pick  $x_n = 2/3$ .

We are going to proceed with induction on this one. The base case is taken care of ( $C_1$ ).

Now proposition is that  $x_{n-1} + y_{n-1} \in C_{n-1}$ ,  $x_{n-1} \in C_{n-1}$  and  $y_{n-1} \in C_{n-1}$ .

Let the length of closed interval, that composes  $C_n$  be equal to  $l$  (i.e.  $1/3$  for  $C_1$ ,  $1/9$  for  $C_2$  and so on; rigorously it'll be equal to  $3^{-n}$ ). Then pick interval  $I_x \subseteq C_n$  such  $I_x$  is the "largest" interval, that composed  $C_n$  with the property that  $\exists i \in I_x : i \leq x_{n-1}$  (largest in this context rigorously means that  $\forall n \in \mathbf{N} : u \in I_n, o \in I_x \rightarrow u < o$ ; in a more lay terms pick the one, that is furthest to the right, but which either contains or less than  $x_{n-1}$ ). Define in the same style  $I_y$  for  $y_{n-1}$ . Then pick  $q_x$  and  $q_y$  be the lowest numbers in corresponding intervals.

Let us now look closer at  $q_x$  and  $q_y$ . Because  $q_x$  is a lower bound for interval in  $C_n$ , we can state, that  $[q_x, q_x + l]$  is in  $C_n$ . If  $x_{n-1} \in C_n$ , then it is the end of the story and the only can set new  $x_n$  to the  $[q_x + l]$ . Otherwise, we also get interval  $[q_x + 2l, q_x + 3l]$  in our disposal.

$x_{n-1} \in C_{n-1}$  implies that  $x_{n-1} - q_x \leq 2l$ . Same applies to  $y_{n-1}$ . Thus it follows, that  $x_{n-1} + y_{n-1} - (q_x + q_y) \leq 4l$ , where  $l$  is the length of an interval in  $C_n$ .

Now let  $k = x_{n-1} + y_{n-1} - (q_x + q_y) \leq l$ .

If  $k \leq l$  then set  $x_n = q_x$  and  $y_n = q_y + k$ .

If  $l < k \leq 2l$  then set  $x_n = q_x + l$  and  $y_n = (k - l)$ .

If  $2l < k \leq 3l$  then set  $x_n = q_x + 2l$  and  $y_n = (k - 2l)$  (given that  $x_{n-1} \notin C_n$ ; If  $x_{n-1} \in C_n$ , set  $x_n = q_x(k - 2l)$  and  $y = q_y + 2l$ )

If  $k \leq 4l$  then set  $x_n = q_x + 3l$  and  $y_n = (k - 3l)$ .

Thus we can state, that  $\exists x_n, y_n \in C_n$  for all  $n \in \mathbf{N}$ .

(b) Keeping in mind that the sequences  $(x_n)$  and  $(y_n)$  do not necessarily converge, show how they can nevertheless be used to produce the desired  $x$  and  $y$  in  $C$  satisfying  $x + y = s$

Because sequences  $(x_n)$  and  $(y_n)$  are contained in  $C$ , we can conclude, that they are bounded. Therefore they have a convergent subsequence  $(x_{n_k}) \rightarrow x$  and  $(y_{n_k}) \rightarrow y$ . Therefore  $(x_{n_k} + y_{n_k}) \rightarrow x + y$ . But we know that  $x_n + y_n = s$  for every  $n \in \mathbf{N}$  (by the definition of the sequence). Thus, sequence  $(x_n + y_n) \rightarrow s$ . We know, that subsequences of convergent sequence converge to the same limit. Thus,  $(x_{n_k} + y_{n_k}) \rightarrow s$ . Because same sequence



cannot converge to two different limits we can conclude that  $s = x + y$ , as desired.

### 3.3.7

*Decide whether the following propositions are true or false. If the claim is valid, supply a short proof, and if the claim is false, provide a counterexample.*

(a) *An arbitrary intersection of compact sets is compact.*

I want to note here firstly, that by our definitions,  $\emptyset$  is a compact set

Every element of our arbitrary intersection is contained in the first set. Thus, the intersection is bounded. We know, that arbitrary intersection of closed sets is closed. Thus, the intersection is closed as well. Thus it is closed and bounded. Thus it is compact.

(b) *Let  $A \subseteq \mathbf{R}$  be arbitrary, and let  $K \subseteq \mathbf{R}$  be compact. Then, the intersection  $A \cap K$  is compact*

False.  $[0, 1] \cap (0, 1) = (0, 1)$ .

(c) *If  $F_1 \supseteq F_2 \supseteq F_3 \dots$  is a nested sequence of nonempty closed sets, then the intersection  $\bigcap_{n=1}^{\infty} F_n \neq \emptyset$*

False. Let  $F_n = [n, \infty)$ .

(d) *A finite set is always compact.*

True. It contains only isolated points and therefore is closed. Also, we can find the maximum and minimum of it, therefore it is bounded.

(e) *A countable set is always compact*

False.  $\mathbf{N}$  is not bounded,  $\mathbf{Q}$  is not closed.

### 3.3.8

*Follow these steps to prove the final implication in Theorem 3.3.8.*

*Assume  $K$  satisfies (i) and (ii), and let  $\{O_\lambda : \lambda \in \Lambda\}$  be an open cover for  $K$ . For contradiction, let's assume that no finite subcover exists. Let  $I_0$  be a closed interval containing  $K$ , and bisect  $I_0$  into two closed intervals  $A_1$  and  $B_1$ .*

(a) *Why must either  $A_1 \cap K$  or  $B_1 \cap K$  (or both) have no finite subcover consisting of sets from  $\{O_\lambda : \lambda \in \Lambda\}$*

Suppose that both of  $A_1 \cap K$  and  $B_1 \cap K$  have finite subcover. Then if we take a union of those subcovers it will cover whole  $K$  and we'll have a finite subcover for  $K$ , which is a contradiction.

(b) *Show that there exists a nested sequence of closed intervals  $I_0 \supseteq I_1 \supseteq I_2 \dots$  with the property that, for each  $n$ ,  $I_n \cap K$  cannot be finitely covered and  $\lim |I_n| = 0$ .*

Let us have  $I_n$  ( $I_0$  for the base case) such that  $I_n \cap K$  does not contain finite subcovers. Divide this set into two equally sized closed intervals  $A_n$  and  $B_n$ . Then at least one of  $A_n \cap K$  and  $B_n \cap K$  will not be finitely covered (because if both of them were finitely covered, then  $I_n$  would be finitely covered). Therefore set  $I_{n+1}$  to either  $A_n$  or  $B_n$ , whichever is not

finitely covered by original set. Because they are equally sized, the sizes of intervals are convergent to 0 (size of each of them is  $2^{-n}$  times the size of  $I_0$ , and  $I_0$  is a closed interval (not closed set) and therefore is bounded, therefore has a finite length).

(c) Show that there exists an  $x \in K$  such that  $x \in I_n$  for all  $n$ .

Each  $I_n \cap K$  is an intersection of two compact sets (closed interval and a compact set), and therefore compact. Therefore we have a series of nested compact sets, therefore its intersection is non-empty. Therefore there exists  $x \in K$  such that  $x \in I_n$  for all  $n \in N$ .

(d) Because  $x \in K$ , there must exist an open set  $O_{\lambda_0}$  from the original collection that contains  $x$  as an element. Argue that there must be an  $n_0$  large enough to guarantee that  $I_{n_0} \subseteq O_{\lambda_0}$ . Explain why this furnished us with the desired contradiction.

Because  $O_{\lambda_0}$  is an open set and  $x \in O_{\lambda_0}$ , there exists neighborhood  $V_\epsilon(x)$  such that  $V_\epsilon(x) \subseteq O_{\lambda_0}$ . Also, there exists  $I_{n_0}$  with length  $\epsilon/2$  (because lengths of  $I_n$ 's are convergent to 0). Thus,

$$\forall q \in R : -\epsilon/2 \leq q - x \leq \epsilon \rightarrow -\epsilon < q - x < \epsilon \rightarrow |q - x| < \epsilon$$

Therefore

$$I_{n_0} \subseteq V_\epsilon(x) \rightarrow I_{n_0} \subseteq O_{\lambda_0}$$

Therefore  $I_{n_0}$  is finitely covered, which is a contradiction. Therefore there does not exist an open cover for compact set, for which there is no finite subcover, as desired.

### 3.3.9

Consider each of the sets listed in Exercise 3.3.5. For each one that is not compact, find an open cover for which there is no finite subcover.

For

$$Q$$

it is  $(-n, n)$

For

$$Q \cup [0, 1]$$

it is  $\{(-1, 0.7) \cup (0.8, 2), (-1, 0.707) \cup (0.708, 2), (-1, 0.7071) \cup (0.7072, 2), \dots\}$  where boundaries of corresponding intervals converge to  $\sqrt{2}/2$ .

For

$$R$$

it is  $(-n, n)$

For

$$\{1, 1/2, 1/3, 1/4, 1/5\}$$

it is  $(1/n, 2)$ .

### 3.3.10

Let's call a set clomcompact if it has the property that every closed cover (i.e., a cover consisting of closed sets) admits a finite subcover. Describe all of the clomcompact subsets of  $R$ .

First of all, no open set can be a clomcompact set, because for any number in the open set there exists a neighborhood, and we add to a collection of sets a collection  $[\epsilon - 1/n + x, x + \epsilon + 1/n]$ , that will prevent us from creating a finite subcover.

Same applies to any set, that has an interval in it.

Finite sets are clomcompact (empty set included).

Cantor set is not clomcompact ( $\{[0, 0], [0, 1 - 1/n], 1\}$  will prevent us from constructing a finite subcover.)

Any compact set cannot be a clomcompact.

Basically, any isolated point is a closed set in and of itself, therefore infinite sets cannot be clomcompact.

Therefore the only clomcompact sets are the finite ones.

If a set is finite, then it'll certainly have in its clomcompact cover a set, where each one of the points are located, therefore finite sets are clomcompact.

Therefore set is clomcompact if and only if it is finite.

### 3.4.1

If  $P$  is a perfect set and  $K$  is compact, is the intersection  $P \cap K$  always compact? Always perfect?

$x \in P \cap K \rightarrow x \in K$ , thus  $P \cap K$  is bounded. Intersection of arbitrary collection of closed sets are closed, therefore  $P \cap K$  is closed as well. Therefore  $P \cap K$  is always compact.

The intersection is not always perfect -  $[1, 2] \cap [2, 3] = \{2\}$ .

### 3.4.2

Does there exist a perfect set consisting of only rational numbers?

If we count empty set as a set, that consists of only rational numbers (i.e. if we translate the text of this exercise to  $x \in B \rightarrow x \in Q$ ), then yes.

If not, then no. This stems from the fact, that the set of rationals is countable, and non-empty perfect sets are uncountable.

### 3.4.3

Review the portion of the proof given for Theorem 3.4.2 and follow these steps to complete the argument

First of all, Theorem 3.4.2 states that Cantor set is perfect.

(a) Because  $x \in C_1$  argue that there exists an  $x_1 \in C \cap C_1$  with  $x_1 \neq x$  satisfying  $|x - x_1| \leq 1/3$

Here we can even go as far as to say, that there exists  $x \in C$  such that  $|x - x_1| \leq 1/6$ . The idea here is that the  $x_1 \in C_1$  implies, that there exists a closed interval  $I_1 \subseteq C_1$ , which length is  $1/3$  such that  $x \in I_1$ . Now the fact, that  $x \in I_1$  and  $|I_1| = 1/3$  implies that the minimal distance between the boundary of  $I_1$  and  $x$  is the half the length of the interval. Boundaries of closed intervals, that constitute  $C_1$  (or  $C_n$  in general) are inside of  $C$ . Therefore there exists  $x \in C$  (and  $C \cap C_1$ , because  $C \cap C_n = C$  for every  $n \in N$ ) such that  $|x - x_1| \leq 1/6 \leq 1/3$ .

(b) Finish the proof by showing that for each  $n \in N$  there exists  $x_n \in C \cap C_n$ , different from  $x$ , satisfying  $|x - x_n| \leq 1/3^n$

Length of the interval, that constitutes  $C_n$  is  $3^{-n}$ , therefore  $x \in C \rightarrow x \in C_n \rightarrow \exists x_n : x_n \in C, |x_n - x| < 1/3^n$ . Therefore for any  $\epsilon$ -neighborhood of  $x$  we will have  $n \in N$  such that  $1/3^n < \epsilon$  (by convergence of  $1/3^n$  to 0) and therefore there will exist  $x_n \in C$ , that is also in desired neighborhood. Therefore any  $x \in C$  is a limit point of  $C$ . Therefore  $C$  is a closed set, that does not contain any isolated points. Therefore it is perfect (how nice).

### 3.4.4

Repeat the Cantor construction from Section 3.1 starting with the open interval  $[0, 1]$ . This time, however, remove the open middle fourth from each component.

Is the resulting set compact? Perfect?

The set will surely be bounded. It will also be equal to the compliment of union of arbitrary collection of open sets, and therefore will be closed.

By the same logic, as in previous exercise (but perhaps this time with a slightly lower length of an interval) it will be perfect.

(b) Using the algorithms from Section 3.1, compute the length and dimension of this Cantor-like set.

The process will be the same as in 3.1, because I didn't get which exact formula they used in order to get to the desired conclusion.

Let us call the desired set  $E$ . Then it follows, that after each iteration we remove  $1/4$  of  $E_n$  to get to  $E_{n+1}$ .

Therefore, the length of a  $E$  will be defined as a limit of a sequence

$$a_1 = 1$$

$$a_{n+1} = a_n - a_n/4 = 3/4a_n$$

Python shows, that it'll be equal to 0, but we'll need to use a more rigorous approach.

First of all, let us show by induction that  $a_n$  is more or equal to 0 for any  $n \in N$ . Base case is already taken care of ( $a_1 = 1 \geq 0$ ). Proposition is that  $a_n \geq 0$ . Therefore the step

is trivial:

$$a_n \geq 0 \rightarrow 3/4 a_n \geq 0 \rightarrow a_{n+1} \geq 0$$

as desired.

$$a_{n+1} = a_n * 3/4 \rightarrow a_{n+1} - a_n = 1/4 a_n \geq 0$$

therefore the series is decreasing. Therefore it is bounded below and decreasing, therefore it is convergent. Also it shows, that it'll be bounded above by 1.

Thus, we can conclude that

$$a_n = a_{n-1} * 3/4 = a_{n-2} * 9/16 = \dots = a_1 * (3/4^n) = 3/4^n$$

We know, that  $|b| < 1$  implies that  $(b^n) \rightarrow 0$ .  $|E_n| = 3/4^{n-1}$ . Therefore the length of  $E$  is equal to  $\lim 3/4^n = 0$ .

Probably the half of this proof is redundant, but I'll roll with it anyways.

When we remove the first quarter from  $[0, 1]$  we'll get  $[0, 3/8] \cup [5/8, 1]$ . Therefore if we magnify the set by  $8/3$ , we'll get  $[0, 1] \cup [5/3, 8/3]$ , or two copies of  $E$ . Thus, the dimension of  $E$  (with processes and definitions from 3.1) will be

$$2 = (8/3)^x$$

$$\ln(2)/\ln(8/3) = x$$

$$x \approx 0.70669505\dots$$

Idk why do I need to know this, but here's the answer.

### 3.4.5

Let  $A$  and  $B$  be subsets of  $\mathbf{R}$ . Show that if there exist disjoint open sets  $U$  and  $V$  with  $A \subseteq U$  and  $B \subseteq V$ , then  $A$  and  $B$  are separated.

Suppose that  $A$  and  $B$  are not separated and let  $x \in \overline{A} \cap B$ . (If  $\overline{A} \cap B = \emptyset$ , then swap names for  $A$  with  $B$  and  $U$  with  $V$ ).

It can be shown, that if  $l$  is a limit point of  $A$ , then it is also a limit point of  $U \supseteq A$  (I won't rigorously prove it here because it was earlier proven rigorously somewhere in this chapter's exercises; it is also somewhat obvious and redundant). Thus,  $\overline{A} \subseteq \overline{U}$ . Therefore,

$$l \in \overline{A} \cap B \rightarrow l \in \overline{U} \cap V \rightarrow l \in V$$

Because  $l$  is in  $V$  and  $V$  is open, it implies that there exists neighborhood  $V_\epsilon(l) \subseteq V$ . Also, because  $l$  is a limit point of  $U$  we can state, that every neighborhood of  $l$  has a member of  $U$  in it. Thus,  $\exists x \in U : x \in V_\epsilon(l) \rightarrow \exists x \in U : x \in V$ . Therefore  $U$  and  $V$  are not disjoint, which is a contradiction.

### 3.4.6

*Prove Theorem 3.4.6*

Firstly, let us state the theorem itself:

**Theorem 3.4.6** *A set  $E \subseteq \mathbf{R}$  is connected if and only if, for all nonempty disjoint sets  $A$  and  $B$  satisfying  $E = A \cup B$ , there always exists a convergent sequence  $(x_n) \rightarrow x$  with  $(x_n)$  contained in one of  $A$  or  $B$ , and  $x$  an element of the other.*

Suppose that we divided  $E$  into disjoint  $A$  and  $B$ .

**In one direction:**

$E$  being connected implies that  $\overline{A} \cap B \neq \emptyset$  (If it is not the case, then swap names for  $A$  and  $B$ ). Let  $l \in \overline{A} \cap B$ . Then it follows, that  $l$  is a limit point for  $\overline{A}$ . Thus, there exists a sequence, that is contained in  $A$ , that is convergent to  $l \in \overline{A} \cap B \rightarrow l \in B$ , as desired.

**In other direction:**

Let us divide  $E$  into disjoint  $A$  and  $B$ . Then it follows, that there exists a sequence  $(x_n) \rightarrow x$  that is contained in  $A$ , that converges to a limit in  $B$  (once again, if it is not the case, then swap names). Also, because  $A$  and  $B$  are disjoint, it follows that  $x_n \neq x$  for all  $n \in \mathbf{N}$ . Then it follows, that  $x$  is a limit point of  $A$ . Thus  $x \in \overline{A} \cap B \neq \emptyset$ . Therefore  $A$  and  $B$  are not separated. Therefore  $E$  is connected, as desired.

### 3.4.7

(a) *Find an example of disconnected set whose closure is connected*

$Q$

(b) *If  $A$  is connected, is  $\overline{A}$  necessarily connected? If  $A$  is perfect, is  $\overline{A}$  necessarily perfect?*

If  $A$  is empty or a singleton, then  $A = \overline{A}$ , therefore  $\overline{A}$  is connected as well. Therefore we need only to handle the case, when  $A$  is nonempty and not a singleton.

Suppose that  $A$  is connected and  $\overline{A}$  is disconnected. Thus we can find two nonempty  $Q$  and  $W$  such that  $Q \cup W = \overline{A}$ . Now let  $A_1 = Q \cap A$ ,  $A_2 = W \cap A$ ,  $L_1 = Q \cap L$  and  $L_2 = W \cap L$ .

$\overline{Q} \cap W = \emptyset$  implies that any limit points of  $A_1$  are not in  $W$  and therefore not in  $L_2$ . Also, by the same logic, any limit points of  $A_2$  are not in  $L_1$ .

Let  $A_1$  be empty. It follows, that  $A \subseteq W$ . Therefore  $A_2 = A$ . Therefore  $Q$  consists of some of limit points of  $A$ . But those limit points cannot be a limit for any  $A_2 = A$ . Therefore  $Q$  is empty, which is a contradiction. (If  $A_2$  is empty, then swap names and the same logic will apply; both sets cannot be empty, because they have to contain at least  $A$ , which is nonempty)

Therefore there exist two sets, such that  $A_1 \cup A_2 = A$ , and  $\overline{A_1} \cap A_2 = \emptyset$  and  $\overline{A_2} \cap A_1 = \emptyset$ . Therefore we can write  $A_1$  and  $A_2$  are separated and nonempty. Therefore  $A$  is disconnected, which is a contradiction.

If  $A$  is perfect, then it is closed, therefore  $A = \overline{A}$ . Therefore  $\overline{A}$  is perfect as well.

### 3.4.8

A set  $E$  is totally disconnected if, given two points  $x, y \in E$ , there exist separated  $A$  and  $B$  with  $x \in A$ ,  $y \in B$ , and  $E = A \cup B$

I probably should add here, that we are also assuming that  $x \neq y$ , otherwise this definition is meaningless.

(a) Show that  $\mathbb{Q}$  is totally disconnected

We know, that between any two real numbers there exists an irrational number. Thus for  $x < y \in \mathbb{Q}$  (swap names if it is not the case) there exists an irrational number  $i$  such that  $x < i < y$ . Thus, for those two points there would exist two intervals  $A = \mathbb{Q} \cap (-\infty, i)$  and  $B = \mathbb{Q} \cap (i, \infty)$ , which are separated.

(b) Is the set of irrational numbers totally disconnected.

Yes. It is also true that between two real numbers there exists a rational number. Thus, by the same logic as in (a), the set of irrationals is totally disconnected.

### 3.4.9

Follow these steps to show that the Cantor set  $C = \bigcap_{n=0}^{\infty} C_n$  described in Section 3.1 is totally disconnected in the sense described in Exercise 3.4.8

(a) Given  $x, y \in C$ , with  $x < y$ , set  $\epsilon = y - x$ . For each  $n = 0, 1, 2, \dots$ , the set  $C_n$  consists of a finite number of closed intervals. Explain why there must exist an  $N$  large enough so that it is impossible for  $x$  and  $y$  both to belong to same closed interval  $C_N$ .

The length of an interval, that constitutes  $C_n$  is  $3^{-n+1}$ . The sequence of lengths of those intervals converges to 0, therefore there exists a  $C_N$  such that maximum length of an interval in it is less than  $\epsilon$ .

(b) Argue that there exists a point  $z \notin C$  such that  $x < z < y$ . Explain how this proves that there can be no interval of the form  $(a, b)$  with  $a < b$  contained in  $C$ .

Because for every  $x, y \in C$  there exists  $C_N$ , which intervals cannot hold both of  $x$  and  $y$ , there exists a point  $z \notin C_N$  such that  $x < z < y$  (otherwise they would be in the same interval).

Thus for any two points  $a < b \in C$  there exists  $z \notin C$  such that  $a < z < b$ .

(c) Show that  $C$  is totally disconnected.

We can take two intervals  $A = (-\infty, z) \cap C$  and  $B = (z, \infty) \cap C$ , which would satisfy the constraint from the definition.

### 3.4.10

Let  $\{r_1, r_2, r_3, \dots\}$  be an enumeration of the rational numbers, and for each  $n \in \mathbb{N}$  set  $\epsilon_n = 1/2^n$ . Define  $O = \bigcup_{n=1}^{\infty} V_{\epsilon_n}(r_n)$ , and let  $F = O^c$ .

(a) Argue that  $F$  is a closed, nonempty set consisting only of irrational numbers.

$O$  is defined to be a union of arbitrary collection of open sets, therefore it is open itself. Thus, its compliment,  $F$ , must be closed.

The non-emptiness of this set might stem from the fact, that even if every neighborhood is disjoint, then length of  $O$  is equal to  $2 \sum 1/2^n = 2$ .

If  $F$  is empty, then  $O = \mathbf{R}$ . Therefore  $O$  is closed and perfect.

Let  $\{[0, 3] \setminus \cup_{i=1}^n r_i\}$  be a collection of compact sets. Then each one of them will be non-empty. Therefore their intersection is nonempty. Therefore  $F$ , which is a superset of this set, is nonempty.

(b) *Does  $F$  contain any nonempty open intervals? Is  $F$  totally disconnected?*

If  $(a, b)$  is an interval in  $F$ , then it follows, that it contains a real number, which is not in  $F$ . Therefore there are no open sets in  $F$ .

Therefore by this logic,  $F$  is totally disconnected (pick a rational number between  $a$  and  $b$ , call it  $r$ , and produce sets  $(-\infty, r) \cup F$  and  $(r, \infty) \cup F$ )

(c) *Is it possible to know whether  $F$  is perfect? If not, can we modify this construction to produce a nonempty perfect set of irrational numbers?*

It depends on how exactly do we define  $r_1, r_2, \dots$  whether  $F$  will be perfect or not. We can define those neighborhoods to converge around some irrational number, (i.e. make it so first one has an upper bound of 3, then define the next one to be bound below by 3.2, then upper to 3.14, then lower to 3.142 and so on to  $\pi$ ; Maybe we'll need to increase the speed with which we converge in order to accomodate for the fact, that epsilons are getting smaller pretty fast)

I don't think that we can somehow put a constraint on the epsilons, if we don't define how to produce the  $r$ 's

I don't consider this particular exercise to be even close to being correctly done, maybe i'll return to it later

### 3.5.1

*Argue that a set  $A$  is a  $G_\delta$  set if and only if its complement is an  $F_\sigma$  set.*

**In one direction:** Suppose that  $G_\delta$  set is composed by  $\cap_{n=1}^\infty s_n$ , where  $s_n$  is a closed set. Then by De Morgan rule

$$G_\delta^c = (\cap_{n=1}^\infty s_n)^c = \cup_{n=1}^\infty s_n^c$$

Because  $s_n$  is a closed set, then  $s_n^c$  is an open set. Therefore

$$G_\delta^c = \cup_{n=1}^\infty s_n^c$$

is a  $F_\sigma$  set.

*In other direction:* Suppose that  $F_\sigma$  set is composed by  $\cup_{n=1}^\infty s_n$ , where  $s_n$  is an open set. Then by De Morgan rule

$$F_\sigma^c = (\cup_{n=1}^\infty s_n)^c = \cap_{n=1}^\infty s_n^c$$



Because  $s_n$  is an open set, then  $s_n^c$  is a closed set. Therefore

$$F_\sigma^c = \bigcup_{n=1}^{\infty} s_n^c$$

is a  $G_\delta$  set.

### 3.5.2

Replace each \_\_\_\_\_ with the word *finite* or *countable*, depending of which is more appropriate

- (a) The countable union of  $F_\sigma$  sets is a  $F_\sigma$  set
- (b) The finite intersection of  $F_\sigma$  sets is an  $F_\sigma$  set.
- (c) The finite union of  $G_\delta$  sets is a  $G_\delta$  set
- (d) The countable intersection of  $G_\delta$  sets is an  $G_\delta$  set.

### 3.5.3

(This exercise has already appeared as Exercise 3.2.14.)

### 3.5.4

**Theorem 3.5.2** If  $\{G_1, G_2, G_3, \dots\}$  is a countable collections of dense, open sets, then the intersection  $\bigcap_{n=1}^{\infty} G_n$  is not empty.

(a) Starting with  $n = 1$ , inductively construct a nested sequence of closed intervals  $I_1 \supseteq I_2 \supseteq I_3 \supseteq \dots$  satisfying  $I_n \subseteq G_n$ . Give special attention to the issue of the endpoints of each  $I_n$

Let us pick any  $x_1 \in G_1$ . Then, because it is open, there exists  $V_{\epsilon_1}(x) \subseteq G_1$ . Let  $I_1 = [x_1 - \epsilon/2, x_1 + \epsilon/2]$ . Then it follows, that there exist  $x_2 \in G_2$  such that  $x_1 - \epsilon/2 < x_2 < x_1 + \epsilon/2$ . Then it follows that  $x_2 \in G_1 \cap G_2$ .

In general let  $I_n = [x_n - \epsilon/2, x_n + \epsilon/2]$ . Then it follows, that there exist  $x_{n+1} \in G_{n+1}$  such that  $x_n - \epsilon/2 < x_{n+1} < x_n + \epsilon/2$ . Then it follows that  $x_{n+1} \in G_n \cap G_{n+1}$ .

(b) Now, use Theorem 3.3.5 or the NIP to furnish the proof

Therefore we have got the collection of nonempty sets, which satisfies

$$I_n \subseteq G_n$$

and

$$I_{n+1} \subseteq I_n$$

Therefore  $\bigcap I_n \neq \emptyset$ . Therefore

$$\bigcap_{n=1}^{\infty} I_n \subseteq \bigcap_{n=1}^{\infty} G_n \neq \emptyset$$

### 3.5.5

Show that it is impossible to write

$$\mathbf{R} = \bigcup_{n=1}^{\infty} F_n$$

where for each  $n \in \mathbf{N}$ ,  $F_n$  is a closed set containing no nonempty open intervals.

Suppose that  $\mathbf{R} = \bigcup_{n=1}^{\infty} F_n$  where  $F_n$  are closed sets. Then

$$\mathbf{R}^c = \emptyset = (\bigcup_{n=1}^{\infty} F_n)^c = \bigcap_{n=1}^{\infty} (F_n)^c$$

Then  $\bigcap_{n=1}^{\infty} F_n^c = \emptyset$ , where  $F_n$  is a closed set and therefore  $F_n^c$  is an open set.

Let  $x, y \in \mathbf{R}$  such that  $x < y$ . Then suppose that there is no  $z \in F_n^c$  such that  $x < z < y$ . Then it follows, that  $(x, y) \not\subseteq F_n^c$ . Therefore  $(x, y) \subseteq F_n$ , which is a contradiction. Therefore  $F_n^c$  is dense in  $\mathbf{R}$ .

Therefore  $\bigcap_{n=1}^{\infty} F_n^c$  is a countable intersection of dense, open sets (countable because we sum from  $n$  to  $\infty$ , therefore on  $\mathbf{N}$ ). Therefore it cannot be empty. Therefore  $\bigcup_{n=1}^{\infty} F_n \neq \mathbf{R}$ , which is a contradiction.

### 3.5.6

Show how the previous exercise implies that the set  $I$  of irrationals cannot be a  $F_\sigma$  set, and  $Q$  cannot be a  $G_\delta$  set.

We had already proven that  $Q$  is a  $F_\sigma$  set and  $I$  is a  $G_\delta$  set.

Suppose that  $Q$  is a  $G_\delta$  set. Then it follows that  $Q^c = I$  is a  $F_\sigma$  set. Therefore  $Q \cup Q^c = \mathbf{R} = I \cup I^c$  is a  $F_\sigma$  set.

Because  $Q$  and  $I$  is a  $F_\sigma$  there exists collection  $F_n$  and  $S_n$  such that every  $F_n$  and  $S_n$  is a closed set. Also, for every  $x < y \in F_n$  it follows that  $x, y \in Q$ . Therefore, there exists  $z \in Q^c = I$  such that  $x < z < y$ . Therefore there does not exist a nonempty open interval  $(x, y) \subseteq F_n$ . Same can be said about  $S_n$ . Thus  $Q \cup I = \mathbf{R}$  can be written as the union of sets  $J_n$  where for each  $n \in \mathbf{N}$ ,  $J_n$  is a closed set containing no nonempty open intervals, which is a contradiction.

Thus  $Q$  cannot be a  $G_\delta$  set. Because  $I$  being  $F_\sigma$  set implies that  $Q$  is a  $G_\delta$  set,  $I$  cannot be a  $F_\sigma$  set.

### 3.5.7

Using Exercise 3.5.6 and versions of the statements in Exercise 3.5.2, construct a set that is neither in  $F_\sigma$  nor a  $G_\delta$ .

First of all, it is necessary to state that  $\emptyset$  is in and of itself a closed set. Therefore  $\bigcup_{n=1}^{\infty} \emptyset = \emptyset$ . Thus  $\emptyset$  is a  $F_\sigma$  set. Also,  $\emptyset$  is an open set. Thus  $\bigcap_{n=1}^{\infty} \emptyset = \emptyset$  is a  $G_\delta$  set.

If  $A$  is a closed set, then  $A = \bigcup_{n=1}^{\infty} A$  is a  $F_\sigma$  set. If  $A$  is an open set, then  $A = \bigcap_{n=1}^{\infty} A$  is a  $G_\delta$  set.

Let  $W$  be a desired set. Then it follows that  $W$  is not a  $F_\sigma$ , and is not a  $G_\delta$  set. Also,  $W^c$  has the same properties.

Any finite set is closed and therefore is a  $F_\sigma$  set. Their compliment is a  $G_\delta$  set.

Therefore the set that we are looking for is neither closed nor open.

Cantor set won't do, because it is closed.

Set  $\{1, 1/2, 1/3, \dots\}$  won't do, because it can be written as a countable union of closed singletons, and therefore it is  $F_\sigma$  set. Its compliment won't do for obvious reasons.

$Q$  is an  $F_\sigma$  set. Then  $Q \cap (0, 1)$  is not closed, because it doesn't contain 0 and 1. It can be still written as the countable union of closed singletons, therefore it is still  $F_\sigma$ .

I decided to look up a hint in the internet saw the answer.  $((-\infty, 0) \cap I) \cup ((0, \infty) \cap Q)$  is neither  $F_\sigma$  nor  $G_\delta$ . What a shameful waste of an exercise.

### 3.5.8

*Show that a set  $E$  is nowhere-dense in  $R$  if and only if the compliment of  $\overline{E}$  is dense in  $R$*

**In one direction:** Suppose that  $E$  is nowhere-dense. Then it follows, that  $\overline{E}$  does not contain any nonempty intervals. Thus, given any  $x, y \in R$  it follows that  $(x, y) \not\subseteq \overline{E}$ . Thus, there exists  $k \in (x, y)$  such that  $k \notin \overline{E}$ . Therefore  $k \in (\overline{E})^c$ . Thus, given two real numbers, there exists  $k \in (\overline{E})^c$  such that  $x < k < y$ . Therefore  $(\overline{E})^c$  is dense in  $R$ , as desired.

**In other direction:** Suppose that  $(\overline{E})^c$  is dense in  $R$ . Then, for any  $x < y \in R$  there exists  $z \in (\overline{E})^c$  such that  $x < z < y$ . Therefore there does not exist  $x_1, y_1 \in R$  such that  $(x, y) \subseteq \overline{E}$ . Therefore  $E$  is nowhere-dense.

### 3.5.9

*Decide whether the following sets are dense in  $R$ , nowhere-dense in  $R$ , or somewhere in between*

(a)  $A = \mathbf{Q} \cap [0, 5]$

$\overline{A} = [0, 5]$ , therefore the set is not nowhere-dense. It isn't dense in  $R$  though, because for  $-1$  and  $-1/2$  there is no number in the  $A$ , that is between them.

(b)  $B = \{1/n : n \in \mathbf{N}\}$

Nowhere-dense

(c) *the set of irrationals*

Is dense in  $R$ .

(d) *the Cantor set*

Cantor set is closed, and therefore  $C = \overline{C}$ . Also, it is totally disconnected, and therefore  $x < y \in C \rightarrow \exists z \in C^c : x < z < y$ . Therefore it does not contain any open intervals. Therefore it is nowhere-dense.

### 3.5.10

Finish the proof by finding a contradiction to the results in this section

**Theorem 3.5.4 (Baire's Theorem)** The set of real numbers  $\mathbf{R}$  cannot be written as the countable union of nowhere-dense sets.

For contradiction, assume that  $E_1, E_2, E_3, \dots$  are each nowhere-dense and satisfy  $\mathbf{R} = \bigcup_{n=1}^{\infty} E_n$ . For each  $E_n$  we can state that  $E_n \subseteq \overline{E_n}$ . Thus,  $\bigcup_{n=1}^{\infty} E_n \subseteq \bigcup_{n=1}^{\infty} \overline{E_n}$ . Therefore  $R \subseteq \bigcup_{n=1}^{\infty} \overline{E_n}$ . Therefore  $\bigcup_{n=1}^{\infty} \overline{E_n} = R$ .  $E_n$  is nowhere-dense, therefore  $\overline{E_n}$  contains no nonempty intervals. Therefore  $R$  can be written as countable union of closed sets, where each of those sets contain no nonempty open intervals, which is a contradiction.

### 4.2.1

Use Definition 4.2.1 to supply a proof for the following limit statements.

- (a)  $\lim_{x \rightarrow 2} (2x + 4) = 8$ .
- (b)  $\lim_{x \rightarrow 0} x^3 = 0$ .
- (c)  $\lim_{x \rightarrow 2} x^3 = 8$ .
- (d)  $\lim_{x \rightarrow \pi} \lfloor x \rfloor = 3$ , where  $\lfloor x \rfloor$  denotes the greatest integer less than or equal to  $x$ .

Let's first state Definition 4.2.1

**Definition 4.2.1.** Let  $f : A \rightarrow \mathbf{R}$ , and let  $c$  be a limit point of the domain  $A$ . We say that  $\lim_{x \rightarrow c} f(x) = L$  provided that, for all  $\epsilon > 0$ , there exists a  $\delta > 0$  such that whenever  $0 < |x - c| < \delta$  (and  $x \in A$ ) it follows that  $|f(x) - L| < \epsilon$ .

(a):

$$|f(x) - L| = |2x + 4 - 8| = |2x - 4| = 2|x - 2| < \epsilon$$

$$|x - 2| < \frac{\epsilon}{2}$$

$$\delta = \frac{\epsilon}{2} \rightarrow |2x + 4 - 8| < \epsilon$$

as desired.

(b):

$$|f(x) - L| = |x^3 - 0| = |x^3| = |x|^3 < \epsilon$$

$$|x| < \sqrt[3]{\epsilon}$$

$$\delta = \sqrt[3]{\epsilon} \rightarrow |x^3| < \epsilon$$

as desired.

(c):

$$|f(x) - L| = |x^3 - 8| = |(x - 2)(x^2 + 2x + 4)| = |x - 2||x^2 + 2x + 4| < \epsilon$$

$$|x - 2| < \frac{\epsilon}{|x^2 + 2x + 4|}$$

Suppose that we set the maximum delta at 1; then upper bound for  $|x^2 + 2x + 4|$  is:

$$\begin{aligned} |x^2 + 2x + 4| &\leq |x^2| + |2x| + 4 = |x|^2 + 2|x| + 4 \leq (|c| + 1)^2 + 2(|c| + 1) + 4 = \\ &= (2 + 1)^2 + 2(2 + 1) + 4 = 9 + 6 + 4 = 19 \end{aligned}$$

Therefore

$$\delta = \min\{1, \epsilon/19\} \rightarrow |x^3 - 8| = |x - 2||x^2 + 2x + 4| < \frac{\epsilon}{19} * 19 = \epsilon$$

as desired.

(d):

$$|[[x]] - 3| = |[0.1415926...]| = 0 < \epsilon$$

Suppose that we pick  $\delta = 0.1$ , then any  $x \in V_\delta$  will satisfy  $|[[x]] - 3| = 0 < \epsilon$  for any  $\epsilon > 0$  as desired.

## 4.2.2

*Assume a particular  $\delta > 0$  has been constructed as a suitable response to a particular  $\epsilon$  challenge. Then, any larger/smaller (pick one)  $\delta$  will also suffice.*

Smaller. This follows from the fact, that

$$\delta_1 < \delta_2 \rightarrow V_{\delta_1} \subset V_{\delta_2}$$

## 4.2.3

*Use Corollary 4.2.5 to show that each of the following limits does not exist.*

(a)  $\lim_{x \rightarrow 0} |x|/x$

(b)  $\lim_{x \rightarrow 1} g(x)$  where  $g$  is Dirichlet's function from Section 4.1.

I'll not state corollary 4.2.5 function here, because it's tedious, but it'll be obvious which corollary I'm talking about by the context.

(a): let

$$(x_n) = 1/n$$

$$(y_n) = -1/n$$

then

$$(x_n) \rightarrow 0; (y_n) \rightarrow 0$$

but

$$|x_n|/x_n = 1$$

$$|y_n|/y_n = -1$$

therefore the limit does not exist.

(b):

The Dirichlet function is

$$D(x) = \begin{cases} 1 & \text{if } x \in \mathbf{Q} \\ 0 & \text{if } x \notin \mathbf{Q} \end{cases} \quad (14)$$

let

$$(x_n) = 2/n + 1$$

$$(y_n) = \sqrt{2}/n + 1$$

then

$$(x_n) \rightarrow 1; (y_n) \rightarrow 1$$

but

$$(x_n) = 2/n + 1 \in \mathbf{Q}$$

$$(y_n) = \sqrt{2}/n + 1 \notin \mathbf{Q}$$

therefore

$$D(x_n) = 1$$

$$D(y_n) = 0$$

thus the function is not continuous at 1

#### 4.2.4

*Review the definition of Thomae's function  $t(x)$  from Section 4.1.*

*(a) Construct three different sequences  $(x_n)$ ,  $(y_n)$ , and  $(z_n)$ , each of which converges to 1 without using the number 1 as a term in the sequence.*

*(b) Now, compute  $\lim t(x_n)$ ,  $\lim t(y_n)$ , and  $\lim t(z_n)$ .*

*(c) Make an educated conjecture for  $\lim_{x \rightarrow 1} t(x)$ , and use Definition 4.2.1B to verify the claim. Given  $\epsilon > 0$ , consider the set of points  $\{x \in \mathbf{R} : t(x)\epsilon\}$ . Argue that all the points in this set are isolated.*

The definition of Thomae function is

$$t(x) = \begin{cases} 1 & \text{if } x = 0 \\ 1/n & \text{if } x = m/n \in \mathbf{Q} \setminus \{0\} \\ 0 & \text{if } x \notin \mathbf{Q} \end{cases} \quad (15)$$

(a): Let our three sequences be

$$(x_n) = n/(n+1)$$

$$(y_n) = (n+1)/n$$

$$(z_n) = \sum_{i=1}^n \frac{1}{2^i}$$

(b):

$$t(x_n) = \{1/2, 1/3, 1/4, 1/5, 1/6, 1/7, \dots\}$$

$$t(y_n) = \{1, 1/2, 1/3, 1/4, 1/5, 1/6, \dots\}$$

$$t(z_n) = \{1/2, 1/4, 1/8, 1/16, \dots\}$$

(c): The educated conjecture here is that  $\lim_{x \rightarrow 1} t(x) = 0$

In order to prove that conjecture author suggests, that we use  $\epsilon - \delta$  definition. Let's try it;

$$|t(x)| < \epsilon$$

For all  $\epsilon \in \mathbf{R} > 0$

Therefore by archimedes property there exists a number  $n \in \mathbf{N}$  s.t.  $\frac{1}{n} < \epsilon$ . Thus suppose that we have  $\delta = 1/n$ . Then our proposition is that

$$\forall b \in (1 - 1/n; 1 + 1/n) \rightarrow |t(b)| < \epsilon$$

If  $b \notin \mathbf{Q}$  then  $t(b) = 0$  and therefore  $|t(b)| < \epsilon$ ; therefore we need to prove, that any number  $b = m_1/n_1 \in (1 - 1/n; 1 + 1/n) \cap \mathbf{Q}$  is such, that  $|t(b)| = 1/n_1 < 1/n$ . Also suppose  $m_1 = n_1 + k$  (it's worth noting that in this case  $k \in \mathbf{Z}$ ); then

$$1 - \frac{1}{n} < \frac{m_1}{n_1} < 1 + \frac{1}{n}$$

$$1 - \frac{1}{n} < \frac{n_1 + k}{n_1} < 1 + \frac{1}{n}$$

$$1 - \frac{1}{n} < 1 + \frac{k}{n_1} < 1 + \frac{1}{n}$$

$$-\frac{1}{n} < \frac{k}{n_1} < \frac{1}{n}$$

$$\left| \frac{k}{n_1} \right| < \frac{1}{n}$$

$$|k| \left| \frac{1}{n_1} \right| = |k| |t(\frac{1}{n_1})| < \frac{1}{n}$$

therefore because  $k \in \mathbf{Z}$

$$|t(\frac{1}{n_1})| = |\frac{1}{n_1}| < \frac{1}{n|k|} < \frac{1}{n}$$

thus for each  $\epsilon > 0$  we can find a corresponding  $\delta > 0$  as desired.

## 4.2.5

Suppose that  $\lim_{x \rightarrow c} f(x) = L$  and  $\lim_{x \rightarrow c} g(x) = M$

(ii)  $\lim_{x \rightarrow c} [f(x) + g(x)] = L + M$

(iii)  $\lim_{x \rightarrow c} [f(x)g(x)] = LM$

(a) *Supply the details for how Corollary 4.2.4 part (ii) follows from the sequential criterion for functional limits in Theorem 4.2.3 and the Algebraic Limit Theorem for sequences proved in Chapter 2.*

From the algebraic limit theorem we know, that if  $(a_n) \rightarrow a$  and  $(b_n) \rightarrow b$  then

$$(a_n) + (b_n) = a + b$$

We also know, that for any sequence  $(c_n) \rightarrow c$  it is true, that  $f(c_n) \rightarrow L$  and  $g(c_n) \rightarrow M$ ; therefore by the algebraic limit theorem

$$f(c_n) + g(c_n) = L + M$$

for any sequence  $(c_n) \rightarrow c$ . Therefore we can state that

$$\lim_{x \rightarrow c} (f(x) + g(x)) = L + M$$

as desired

(b) *Now, write another proof of Corollary 4.2.4 part (ii) directly from Definition 4.2.1 without using the sequential criterion in Theorem 4.2.3.*

$\lim_{x \rightarrow c} f(x) = L$  and  $\lim_{x \rightarrow c} g(x) = M$ ; therefore for any  $\epsilon_1 > 0$  we can find  $\delta_1 > 0$  s.t.

$$|x - c| < \delta_1 \rightarrow |f(x) - L| < \epsilon_1$$

Also for the same  $\epsilon_1$  there exist  $\delta_2 > 0$  s.t.

$$|x - c| < \delta_2 \rightarrow |g(x) - M| < \epsilon_1$$

let  $\delta_3 = \min\{\delta_1, \delta_2\}$ ; then it is true that

$$|x - c| < \delta_3 \rightarrow |f(x) - L| < \epsilon_1$$

$$|x - c| < \delta_3 \rightarrow |g(x) - M| < \epsilon_1$$



because  $V_{\delta_1} \subseteq V_{\delta_3}$  and  $V_{\delta_2} \subseteq V_{\delta_3}$   
therefore

$$|f(x) - L| + |g(x) - M| < 2\epsilon_1$$

Therefore

$$|f(x) + g(x) - L - M| = |f(x) - L + g(x) - M| \leq |f(x) - L| + |g(x) - M| < 2\epsilon_1$$

Thus for any  $\epsilon > 0$  there exist corresponding  $\epsilon_1 = \frac{\epsilon}{2}$  for which there exist corresponding  $\delta = \min\{\delta_1, \delta_2\}$  (where  $\delta_1$  is a delta for  $f(x)$  and  $\delta_2$  is a delta for  $g(x)$ ) which satisfies

$$|x - c| < \delta \rightarrow |f(x) + g(x) - (L + M)| < \epsilon$$

therefore  $\lim_{x \rightarrow c} (f(x) + g(x)) = L + M$  as desired.

(c) Repeat (a) and (b) for Corollary 4.2.4 part (iii).

(a):

From the algebraic limit theorem we know, that if  $(a_n) \rightarrow a$  and  $(b_n) \rightarrow b$  then

$$(a_n)(b_n) = ab$$

We also know, that for any sequence  $(c_n) \rightarrow c$  it is true, that  $f(c_n) \rightarrow L$  and  $g(c_n) \rightarrow M$ ;  
therefore by the algebraic limit theorem

$$f(c_n)g(c_n) = LM$$

for any sequence  $(c_n) \rightarrow c$ . Therefore we can state that

$$\lim_{x \rightarrow c} (f(x)g(x)) = LM$$

as desired

(b):

$\lim_{x \rightarrow c} f(x) = L$  and  $\lim_{x \rightarrow c} g(x) = M$ ;

In order to prove the needed limit let's first use some algebra

$$|f(x)g(x) - LM| =$$

$$|f(x)g(x) + f(x)M - f(x)M - LM| =$$

$$|f(x)(g(x) - M) + M(f(x) - L)| \leq |f(x)(g(x) - M)| + |M(f(x) - L)| =$$

$$|f(x)||g(x) - M| + |M||f(x) - L|$$

our strategy is to show that both elements of the last sum are less or equal to  $\epsilon/2$   
Let  $\epsilon > 0$ .

$$|M||f(x) - L| < \frac{\epsilon}{2}$$

If  $M = 0$  then the abovementioned inequality always holds and we are free to choose any  $\delta_1$ ;

Otherwise let us pick  $\delta_1$  such that inequality

$$|f(x) - L| < \frac{\epsilon}{2|M|}$$

holds.

The next step is a little bit more complicated because we need to work with  $f(x)$ ; let us pick  $\epsilon = 1$ ; then because  $\lim_{x \rightarrow c} f(x) = L$  we know that there exists  $\delta_2$  s.t.  $|x - c| < \delta_2 \rightarrow |f(x) - L| < 1$ .

Therefore

$$|f(x) - L| < 1$$

Little sidenote: let's prove that

$$|a - b| < c \rightarrow |a| < |b| + c$$

Firstly some preliminary stuff

$$|a - b| \geq 0 \rightarrow c > |a - b| > 0 \rightarrow c > 0$$

$$|a - b| < c \rightarrow -c < a - b < c$$

$$b - c < a < b + c$$

Now let's see all the cases for  $a, b \in \mathbf{R}$

if  $a \geq 0$  and  $b \geq 0$  then

$$a < b + c$$

$$|a| < |b| + c$$

if  $a < 0$  and  $b \geq 0$  then

$$b + c \geq 0 > a$$

$$a < b + c$$

$$|a| < |b| + c$$

if  $a \geq 0$  and  $b < 0$  then

$$b - c < a < b + c$$

$$-b + c > -a > -b - c$$

$$|b| + c > -a > |b| - c$$

$$-|b| - c < a < c - |b|$$

$$|a| < c - |b| \leq c + |b|$$

$$|a| < c + |b|$$

if  $a < 0$  and  $b < 0$  then

$$b - c < a < b + c$$

$$-b + c > -a > -b - c$$

$$|b| + c > |a| > |b| - c$$

$$|b| + c > |a|$$

$$|a| < |b| + c$$

Therefore  $\forall a, b \in \mathbf{R}$

$$|a - b| < c \rightarrow |a| < |b| + c$$

as desired.

Back to our exercise:

$$|f(x) - L| < 1$$

$$|f(x)| < |L| + 1$$

Therefore we can state that upper bound for our  $|f(x)|$  with  $\epsilon = 1$  is  $|L| + 1$

Thus if we pick  $\delta_2$  sufficient for

$$|g(x) - M| < \frac{\epsilon}{2(|L| + 1)}$$

therefore if we pick  $\delta = \min\{\delta_1, \delta_2\}$  then

$$|x - c| < \delta \rightarrow$$

$$|f(x)g(x) - LM| \leq |f(x)||g(x) - M| + |M||f(x) - L| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$$

therefore  $\lim_{x \rightarrow c} [f(x)g(x)] = LM$  as desired

## 4.2.6

Let  $g : A \rightarrow \mathbf{R}$  and assume that  $f$  is bounded function on  $A \subseteq \mathbf{R}$  (i.e. there exists  $M > 0$  satisfying  $|f(x)| \leq M$  for all  $x \in A$ ). Show that if  $\lim_{x \rightarrow c} g(x) = 0$ , then  $\lim_{x \rightarrow c} g(x)f(x) = 0$  as well.

Here we can't use an intuitive approach of just using algebraic limit theorem because  $f(x)$  may not have limit at  $c$ . Anyways we proceed by  $\epsilon - \delta$  approach.

Therefore we need to show that

$$\exists \delta : |f(x)g(x)| < \epsilon$$

First of all,

$$|f(x)g(x)| = |f(x)||g(x)|$$

Then we notice, that because  $f(x)$  is bounded

$$\exists M \in \mathbf{R} > 0 : |f(x)| \leq M$$

therefore

$$|f(x)||g(x)| < |M||g(x)| = M|g(x)|$$

therefore if we pick  $\delta$  sufficient for  $|g(x)| < \frac{\epsilon}{M}$  then it follows that

$$|f(x)g(x)| \leq M|g(x)| < \epsilon$$

therefore

$$\forall \epsilon \in \mathbf{R} \exists \delta : |x - c| < \delta \rightarrow |f(x)g(x)| < \epsilon$$

therefore

$$\lim_{x \rightarrow c} [f(x)g(x)] = 0$$

as desired.

## 4.2.7

(a) The statement  $\lim_{x \rightarrow 0} 1/x^2 = \infty$  certainly makes intuitive sense. Construct a rigorous definition in the "challenge-response" style of Definition 4.2.1 for a limit statement of the form  $\lim_{x \rightarrow c} f(x) = \infty$  and use it to prove the previous statement

**Definition of limit to infinity** Let  $f : A \rightarrow \mathbf{R}$ , and let  $c$  be a limit point of the domain  $A$ . We say that  $\lim_{x \rightarrow c} f(x) = \infty$  provided that, for all  $\epsilon \in \mathbf{R}$ , there exists a  $\delta > 0$  such that whenever  $0 < |x - c| < \delta$  (and  $x \in A$ ) it follows that  $f(x) > \epsilon$ .

Now we need to show that for  $f(x) = 1/x^2$

$$\lim_{x \rightarrow 0} f(x) = \infty$$

First

$$f(x) > \epsilon$$

$$\frac{1}{x^2} > \epsilon$$

$$x^2 < \frac{1}{\epsilon}$$

$$x < \sqrt{\frac{1}{\epsilon}}$$

therefore if we pick  $\delta = \sqrt{\frac{1}{\epsilon}}$ , then it follows that

$$f(x) > \epsilon$$

as desired.

Quick (and insufficient) test in Python seems to corroborate this statement

(b) Now construct a definition for the statement  $\lim_{x \rightarrow \infty} f(x) = L$ . Show  $\lim_{x \rightarrow \infty} 1/x = 0$

**Definition of infinite limit** Let  $f : A \rightarrow \mathbf{R}$ , and let  $c$  be a limit point of the domain  $A$ . We say that  $\lim_{x \rightarrow \infty} f(x) = L$  provided that, for all  $\epsilon \in \mathbf{R} > 0$ , there exists a  $\delta$  such that whenever  $x > \delta$  (and  $x \in A$ ) it follows that  $|f(x) - c| < \epsilon$ .

We start as usual at the  $\epsilon$

$$|f(x) - 0| < \epsilon$$

$$|1/x| < \epsilon$$

Given that we can pick any  $\delta$  as we want, we can pick it at the very least at 0 to get rid of the absolute value

$$1/x < \epsilon$$

$$x > 1/\epsilon$$

therefore  $\delta = 1/\epsilon$  then it follows that

$$|f(x) - 0| < \epsilon$$

as desired.

(c) What would a rigorous definition for  $\lim_{x \rightarrow \infty} f(x) = \infty$  would look like? Give an example of such a limit

**Definition of infinite limit to infinity** Let  $f : A \rightarrow \mathbf{R}$ , and let  $c$  be a limit point of the domain  $A$ . We say that  $\lim_{x \rightarrow \infty} f(x) = \infty$  provided that, for all  $\epsilon \in \mathbf{R}$ , there exists a  $\delta$  such that whenever  $x > \delta$  (and  $x \in A$ ) it follows that  $f(x) > \epsilon$ .

The corresponding example of such a limit is  $f(x) = x$ .

### 4.2.8

Assume  $f(x) \geq g(x)$  for all  $x$  in some set  $A$  on which  $f$  and  $g$  are defined. Show that for any limit point  $c$  of  $A$  we must have

$$\lim_{x \rightarrow c} f(x) \geq \lim_{x \rightarrow c} g(x)$$

I'm gonna do it by using contradiction; suppose that  $f(x)$  and  $g(x)$  are defined as in the exercise, but

$$\lim_{x \rightarrow c} f(x) < \lim_{x \rightarrow c} g(x)$$

then it follows that there exist a sequence  $(a_n) \rightarrow c$  such that  $f(a_n) \geq g(a_n)$  for all  $n \in \mathbf{N}$ ; Therefore  $\lim(f(a_n)) \geq \lim(g(a_n))$  and but it contradicts our initial assumption.

### 4.2.9 (Squeeze Theorem)

Let  $f, g$  and  $h$  satisfy  $f(x) \geq g(x) \geq h(x)$  for all  $x$  in some common domain  $A$ . If  $\lim_{x \rightarrow c} f(x) = L$  and  $\lim_{x \rightarrow c} h(x) = L$  at some limit point  $c$  of  $A$ , show  $\lim_{x \rightarrow c} g(x) = L$  as well

As proven in the previous exercise

$$\forall x \in A : f(x) > g(x) \rightarrow \lim_{x \rightarrow c} f(x) \geq \lim_{x \rightarrow c} g(x)$$

therefore

$$\lim_{x \rightarrow c} f(x) = L \geq \lim_{x \rightarrow c} g(x)$$

and

$$\lim_{x \rightarrow c} g(x) \geq \lim_{x \rightarrow c} h(x) = L$$

Thus

$$L \geq \lim_{x \rightarrow c} g(x) \geq L$$

therefore

$$\lim_{x \rightarrow c} g(x) = L$$

as desired.

### 4.3.1

Let  $g(x) = \sqrt[3]{x}$ .

(a) Prove that  $g$  is continuous at  $c = 0$

We're gonna use  $\epsilon - \delta$  definition. First of all, let's state that  $g(0) = 0$ . Therefore

$$|f(x) - f(c)| = |\sqrt[3]{x} - 0| < \epsilon$$

$$|\sqrt[3]{x}| < \epsilon$$

Here I would like to prove that  $\forall x \in \mathbf{R} : |\sqrt[3]{x}| = \sqrt[3]{|x|}$ : if  $x \geq 0$  then  $|\sqrt[3]{x}| = \sqrt[3]{x} = \sqrt[3]{|x|}$ ; if  $x < 0$  then  $|\sqrt[3]{x}| = \sqrt[3]{-x} = \sqrt[3]{|x|}$ . Therefore

$$|\sqrt[3]{x}| = \sqrt[3]{|x|} < \epsilon$$

is justified.

Therefore we can state that

$$|x| < \epsilon^3$$

Thus if we pick  $\delta = \epsilon^3$  then

$$|x - c| = |x| < \delta \rightarrow |f(x) - f(c)| = |\sqrt[3]{x} - 0| = |\sqrt[3]{x}| = \sqrt[3]{|x|} < \sqrt[3]{\epsilon^3} = \epsilon$$

Therefore  $g$  is continuous at 0

(b) Prove that  $g$  is continuous at a point  $c \neq 0$ . (The identity  $a^3 - b^3 = (a - b)(a^2 + ab + b^2)$  will be helpful)

We're gonna use  $\epsilon - \delta$  definition once again.

$$|f(x) - f(c)| = |\sqrt[3]{x} - \sqrt[3]{c}| < \epsilon$$

First, let's use a little algebra

$$|\sqrt[3]{x} - \sqrt[3]{c}| = |\sqrt[3]{x} - \sqrt[3]{c}| * 1 = |\sqrt[3]{x} - \sqrt[3]{c}| \frac{(\sqrt[3]{x}^2 + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c}^2)}{(\sqrt[3]{x}^2 + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c}^2)} = \frac{|\sqrt[3]{x} - \sqrt[3]{c}|(\sqrt[3]{x}^2 + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c}^2)}{(\sqrt[3]{x}^2 + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c}^2)}$$

Let's look now at the sum  $\sqrt[3]{x}^2 + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c}^2$ :  $\sqrt[3]{x}^2 \geq 0$  because it is a square. For  $\sqrt[3]{x} + \sqrt[3]{x}^2$  we need to be able to articulate  $\delta$  so that both  $x$  and  $c$  are the same sign; if we do that then it becomes nonnegative.  $\sqrt[3]{c}^2 \geq 0$  because it is a square

Therefore if right now we pinky-promise that we will account for unusual delta in the future, then we are able to say that

$$\sqrt[3]{x}^2 + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c}^2 \geq 0$$

And therefore

$$\sqrt[3]{x}^2 + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c}^2 = |\sqrt[3]{x}^2 + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c}^2|$$

Continuing with our initial algebra

$$\frac{|\sqrt[3]{x} - \sqrt[3]{c}|(\sqrt[3]{x^2} + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c^2})}{(\sqrt[3]{x^2} + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c^2})} = \frac{|\sqrt[3]{x} - \sqrt[3]{c}||\sqrt[3]{x^2} + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c^2}|}{(\sqrt[3]{x^2} + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c^2})} = \frac{|x - c|}{(\sqrt[3]{x^2} + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c^2})} =$$

$$|x - c| \frac{1}{(\sqrt[3]{x^2} + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c^2})} < \epsilon$$

As we disussed earlier  $(\sqrt[3]{x^2} + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c^2}) \geq 0$  and therefore

$$|x - c| < \epsilon(\sqrt[3]{x^2} + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c^2})$$

Thus, if we pick  $\delta = \min\{\epsilon(\sqrt[3]{x^2} + \sqrt[3]{x}\sqrt[3]{c} + \sqrt[3]{c^2}), |x - 0|\}$  (we need the second value because we need the sum to be equal to its absolute value; ) then

$$|x - c| < \delta \rightarrow |f(x) - f(c)| < \epsilon$$

Therefore  $f(x) = \sqrt[3]{x}$  is continous on  $\mathbf{R}$ .

### 4.3.2

(a) Supply a proof for Theorem 4.3.9 using the  $\epsilon - \delta$  characterization of continuity.

First, let's state the theorem

**Theorem 4.3.9 (Composition of Continuous Functions).** *Given  $f : A \rightarrow \mathbf{R}$  and  $g : B \rightarrow \mathbf{R}$ , assume that the range  $f(A) = \{f(x) : x \in A\}$  is contained in the domain  $B$  so that the composition  $g \circ f = g(f(x))$  is well-defined on  $A$ .*

*If  $f$  is continous ac  $c \in A$ , and if  $g$  is continous at  $f(c) \in B$ , then  $g \circ f$  is continous at  $c$ .*

Firstly, the fact that both  $f$  and  $g$  are continous tells that

$$\forall \epsilon_1 \in \mathbf{R} : \exists \delta_1 : |x - c| < \delta_1 \rightarrow |f(x) - f(c)| < \epsilon_1$$

$$\forall \epsilon_2 \in \mathbf{R} : \exists \delta_2 : |x - c| < \delta_2 \rightarrow |g(x) - g(c)| < \epsilon_2$$

And we need to prove that

$$\forall \epsilon \in \mathbf{R} : \exists \delta : |x - c| < \delta \rightarrow |g(f(x)) - g(f(c))| < \epsilon$$

The main strategy for this one is to plug some delta into some epsilon (or vice versa), and get some use out of it.

Firstly, let's get some things out of the way: let us fix particular  $c \in A$  and  $\epsilon \in \mathbf{R} > 0$ . Then, let's plug this  $\epsilon$  at  $f(c)$  into the continuity of  $g(x)$  so we can get a  $\delta_g > 0$ . Therefore we will have

$$\forall \epsilon \in \mathbf{R} : \exists \delta_g : |x - f(c)| < \delta_g \rightarrow |g(x) - g(f(c))| < \epsilon$$



which is kinda close to the thing, that we're trying to prove.

We also know that

$$\forall \epsilon_f \in \mathbf{R} : \exists \delta_f : |x - c| < \delta_f \rightarrow |f(x) - f(c)| < \epsilon_f$$

therefore it is true that

$$\begin{aligned} \forall \epsilon \in \mathbf{R} : \exists \delta_g : |y - f(c)| < \delta_g \rightarrow |g(y) - g(f(c))| < \epsilon \\ \exists \delta_f : |x - c| < \delta_f \rightarrow |f(x) - f(c)| < \delta_g \end{aligned}$$

From this we can state that

$$\forall \epsilon \in \mathbf{R} > 0 : \exists \delta_f : |x - c| < \delta_f \rightarrow |f(x) - f(c)| < \delta_g \rightarrow |g(f(x)) - g(f(c))| < \epsilon$$

This doesn't sound too persuasive for me, so I probably need to explore it a little but more.

Suppose that with all the present assumptions, we get the given  $\epsilon$ . If we plug it into definition of continuity for  $g(x)$  at  $g(f(c))$ , then we'll get the necessary  $\delta_g$ . If we plug  $\delta_g$  as an  $\epsilon$  for the definition of continuity of  $f(x)$ , then we'll get  $\delta_f$ .

We can probably prove it with a little bit more clarity. We need to prove that

$$\forall \epsilon \in \mathbf{R} : \exists \delta : |x - f(c)| < \delta \rightarrow |g(f(x)) - g(f(c))| < \epsilon$$

Firstly, definition of continuity of  $g(x)$  gives us the fact, that

$$\forall \epsilon \in \mathbf{R} : \exists \delta_g : |x - f(c)| < \delta_g \rightarrow |g(x) - g(f(c))| < \epsilon$$

then if  $x \in f(A)$  then  $\exists y \in A$  s.t.  $f(y) = x$  therefore

$$\forall \epsilon \in \mathbf{R} : \exists \delta_g : |f(y) - f(c)| < \delta_g \rightarrow |g(f(y)) - g(f(c))| < \epsilon$$

From the definition of continuity of  $f$  we know that

$$\forall \epsilon_f \in \mathbf{R} : \exists \delta_f : |x - c| < \delta_f \rightarrow |f(x) - f(c)| < \epsilon_f$$

Therefore

$$\forall \epsilon \in \mathbf{R} : \exists \delta : |x - f(c)| < \delta \rightarrow |g(f(x)) - g(f(c))| < \epsilon$$

as desired.

(b) Give another proof of this theorem using the sequential characterization of continuity (from Theorem 4.3.2 (iv))

Theorem 4.3.2 (iv) states that if  $(x_n) \rightarrow c$  (with  $x_n \in A$ ), then  $f(x_n) \rightarrow f(c)$ .

Because  $f(x)$  is continuous we can state that for every sequence  $(x_n) \rightarrow c$  it is true that  $f(x_n) \rightarrow f(c)$ . Therefore because  $f(x_n)$  is a sequence itself, we can state that  $g(f(x_n)) \rightarrow g(f(c))$ . Therefore it is true, that for every sequence  $(x_n) \rightarrow c$  it follows, that  $g(f(x_n)) \rightarrow g(f(c))$ . Therefore  $g(f(x))$  is continuous, as desired.

### 4.3.3

Using the  $\epsilon - \delta$  characteriation of continuity (and tus using no previous results anbout the sequences), show that the linear function  $f(x) = ax + b$  is continous at every poinnt of  $\mathbf{R}$ .

Let's start with our usual stuff

$$\begin{aligned} |f(x) - f(c)| &< \epsilon \\ |ax + b - (ac + b)| &= |ax + b - ac - b| = |a||x - c| < \epsilon \\ |x - c| &< \epsilon/a \end{aligned}$$

Therefore if we pick  $\delta = \epsilon/a$  then it follows that  $|f(x) - f(c)| < \epsilon$ , as desired.

### 4.3.4

(a) Show using Definition 4.3.1 that any function  $f$  with domain  $\mathbf{Z}$  with necessarily be continous at every point in its domain.

Suppose that  $f : Z \rightarrow R$ . We need to prove that

$$\forall \epsilon : \exists \delta : |x - c| < \delta \rightarrow |f(x) - f(c)| < \epsilon$$

Suppose that we pick  $\delta = 0.1$  (or any other value, such that the only one of the domain values will be in the needed neighborhood). Then there will be only one number in the domain neighborhood, and because of that we can state that

$$|f(x) - f(c)| = |f(c) - f(c)| = 0 < \epsilon$$

Therefore the fucntion is continous, as desired.

(b) Show in general that if  $c$  is an isolated point of  $A \subseteq \mathbf{R}$ , then  $f : A \rightarrow \mathbf{R}$  is continous at  $c$ .

Because  $c$  is an isolated point, we can follow, that it is not a limit point. Therefore there exists  $\delta$  such that neighborhood  $V_\delta(c) \cap A = \{c\}$ . Then it follows that for all  $\epsilon > 0$  we have  $\delta > 0$  such that

$$x \in V_\delta(c) \rightarrow f(x) = f(c) \in V_\epsilon(c)$$

as desired.

### 4.3.5

In Theorem 4.3.4, statement (iv) says that  $f(x)/g(x)$  is continous at  $c$  if both  $f$  and  $g$  are, provided that the quotent is defined. Show that if  $g$  is continous at  $c$  and  $g(c) \neq 0$ , then there exists an open interval containing  $c$  on which  $f(x)/g(x)$  is always defined.

Suppose that  $f(x)/g(x)$  is continuous at  $c$ . Then it follows, that for any  $\epsilon > 0$  there exists  $\delta > 0$  such that

$$x \in V_\delta \rightarrow f(x)/g(x) \in V_\epsilon$$

Thus,  $f(x)/g(x)$  is defined on open interval  $V_\delta$ .

### 4.3.6

(a) Referring to the proper theorems, give a formal argument that Dirichlet's function from Section 4.1 is nowhere-continuous at  $R$ .

Let  $c \in R$  and  $\epsilon = 0.5$

Suppose now that  $\delta > 0$ . Then it follows, that there exists both  $z_1 \in I$  and  $z_2 \in Q$  such that

$$c - \delta < z_1 < c + \delta$$

$$c - \delta < z_2 < c + \delta$$

therefore exists  $f(z_1) = 0$  and  $f(z_2) = 1$ . Therefore for any  $c \in R$  there does not exist a  $\delta$ , such that

$$x \in V_\delta(c) \rightarrow f(x) \in V_\epsilon(f(c))$$

. Therefore Dirichlet's function is not continuous at any point in  $R$ .

(b) Review the definition of Thomae's function in Section 4.1 and demonstrate that it fails to be continuous at every rational point.

Thomae's function is defined as

$$t(x) = \begin{cases} 1 & \text{if } x = 0 \\ 1/n & \text{if } x = m/n \in \mathbf{Q} \setminus \{0\} \\ 0 & \text{if } x \notin \mathbf{Q} \end{cases} \quad (16)$$

Suppose now that  $c \in Q$ . Then it follows that if we set  $\epsilon = f(c)$ . For every  $\delta > 0$  there exists  $z \in I$  such that  $z \in V_\delta$ . Therefore, for  $\epsilon = f(c)$   $f(z) = 0$ . Therefore

$$|f(c) - f(x)| < \epsilon$$

$$-\epsilon < f(x) - f(c) < \epsilon$$

$$-\epsilon < f(x) - f(c) < f(c)$$

$$-\epsilon < f(x) - f(c) < f(c)$$

$$f(x) - f(c) < f(c)$$

$$f(x) < 0$$

$$0 < 0$$

Which is false. Therefore for every  $c \in Q$  there exist a  $\epsilon$ , such that there is no  $\delta > 0$  for which it is true that

$$x \in V_\delta \rightarrow f(x) \in V_\delta(f(c))$$

(c) Use the characterization of continuity in Theorem 4.3.2 (iii) to show that Thomae's function is continuous at every irrational point in  $\mathbf{R}$ . (Given  $\epsilon > 0$ , consider the set of points  $\{x \in \mathbf{R} : t(x) \geq \epsilon\}$ . Argue that all the points in this set are isolated )

Let  $\epsilon > 0$  and  $z \in I$ . Therefore we can state, that there exists  $N \in \mathbf{N}$  such that  $1/n < \epsilon$ . Now let us look at  $V_{1/n}(z)$ . There exists a finite amount of natural numbers, that are less than  $N$ . If there exist natural numbers  $n_1 < n$ , such that there exists  $m_1 \in \mathbf{Z}$  for which  $m_1/n_1 \in V_{1/n}(z)$ , then pick one, for which  $|m_1/n_1 - z|$  is the lowest and let  $\delta = |m_1/n_1 - z|$ . If such numbers do not exist, pick  $\delta = 1/n$ . That insures, that any rational number in  $V_\delta(z)$  has the property, that if we put it as  $m_1/n_1$ , then  $n_1 > n$ .

Then it follows, that for any  $\epsilon > 0$  there exists  $\delta > 0$  such that  $x \in V_\delta \rightarrow f(x) \in V_\epsilon(t(z))$  (if  $x \in I$ , then  $x = 0 < \epsilon$ ; if  $x \in Q$ , then  $f(x) < 1/n < \epsilon$  because of how had we constructed  $\delta$  in the previous paragraph). Therefore Thomae's function is continuous at any irrational point.

In the text of the exercise we were asked to consider a specific set and all that other stuff, and that method would certainly work, but I consider this particular method to be a little more straightforward.

### 4.3.7

Assume  $h : R \rightarrow R$  is continuous on  $R$  and let  $K = \{x : h(x) = 0\}$ . Show that  $K$  is a closed set.

Suppose that  $K$  is not closed. Then it follows, that there exists a limit point of  $K$ , that is not in  $K$ . Let  $y \in R$  be a limit point of  $K$ , that is not in  $K$ . It follows, that there exists a sequence  $(h(x_n)) \rightarrow y$ , such that  $h(x_n) \in K$  and  $h(x_n) \neq y$ .