SECOND-LARGEST AND MINIMUM DISTANCE MULTIPLICITIES IN PLANAR POINT SETS: AN UPPER BOUND OF 15/11, AND RELATED RESULTS ON ERDŐS PROBLEM #132

WINGATE JONES

ABSTRACT. For a finite set $X \subset \mathbb{R}^2$ let Δ_2 and δ be the second-largest and the smallest distance determined by X, and let $\mu(d)$ be the number of pairs of X at distance d. Clemen, Dumitrescu and Liu asked for $L = \limsup_{n\to\infty} \max_{|X|=n} \min\{\mu(\Delta_2), \mu(\delta)\}/n$; the known bounds were $9/7 \le L \le 3/2$, the upper bound coming from Vesztergombi's inequality $\mu(\Delta_2) \le 3n/2$. We prove that $\min\{\mu(\Delta_2), \mu(\delta)\} \le \frac{15}{11}n + C_0$ for an absolute constant C_0, so $L \le 15/11 < 3/2$. In one of the regimes of the proof ("Region II", defined by $\Delta \le 1.94\,\Delta_2$ and $\tau(X) \ge \sqrt{3}/2 + 0.03$, where $\tau(X)$ is an explicit scale-invariant measure of the excess of Δ over Δ_2 in units of δ) the bound improves to $\frac{4}{3}n + C_0$.

We also give three results on Erdős Problem #132, which asks whether some distance other than the diameter must occur at most n times. First, if no vertex of the convex hull has four points of X at distance Δ_2 (for instance, if no four points of X are concyclic), then $\mu(\Delta_2) \le \frac{3}{2}|L_1| + |L_2|$, where L_1, L_2 are the first two convex layers; so, under this hypothesis, the answer is yes whenever at least half as many points lie at depth ≥ 3 as on the hull. This rests on a layer bound whose inputs from Vesztergombi's structure theory we prove directly. Second, for n points in convex position there are at least three distances occurring between 1 and n times when n is even and at least 6, and when $6 \le n \le 18$. The cases $n = 11, 13, 15, 17$ are computer-assisted: they reduce to the unsatisfiability of finite ordinal problems, certified by DRAT (and for $n \le 13$ LRAT) proof checkers, and for $n = 7, 11, 13$ also by a Lean 4 formalisation. Third, if all but w of the n points in convex position lie on one circle and $n + w \ge 91$, $n \ge 3w$, then at least $(n - 3w + 1)/4$ distances are rare; the proof combines Kneser's theorem with a bound of Beukers and Smyth on torsion points of plane curves.

AI DISCLOSURE

The author used an AI assistant (Claude, Anthropic) for exploration, computation, formalisation and drafting. The checks performed on the results are described in Appendix A.

1. INTRODUCTION

Let $X \subset \mathbb{R}^2$ be a set of n points. Write $\Delta = \Delta(X)$ for its diameter, Δ_2 for its second-largest distance and δ for its minimum distance, and write $\mu(d) = \mu_X(d)$ for the number of unordered pairs of points of X at distance d. Hopf and Pannwitz [16] proved that $\mu(\Delta) \le n$.

Erdős Problem #132. Erdős and Pach asked whether there must be a *second* distance, besides the diameter, that occurs at least once and at most n times, and whether the number of such distances must tend to infinity (perhaps being at least $n^{1-o(1)}$). This is Problem #132 on the Erdős problems website [6], where it is recorded as open and where Erdős's offer of \$100 for "any nontrivial result" is recorded (reference [Er97e] there). The statement fails for $n = 4$ (two equilateral triangles glued along an edge) and holds for $n = 5, 6$ by Erdős and Fishburn [11]. We call a distance d *rare* if $1 \le \mu(d) \le n$.

Date: Draft of 24 September 2026.

Clemen, Dumitrescu and Liu [7] proved that the answer to the first question is yes for sets in convex position (their Theorem 1.2), and that $\mu(\Delta_2) \le n$ whenever

$$\min\Big\{\tfrac{3}{2}(|L_1|+|L_2|),\ \tfrac{4}{3}|L_1|+2|L_2|,\ 2|L_1|+|L_2|\Big\} \le n, \tag{1}$$

where L_1, L_2 are the first two convex layers (their Theorem 1.3). They showed that neither Δ_2 nor δ is always rare: they constructed sets with $\min\{\mu(\Delta_2), \mu(\delta)\} \ge 9n/8 - o(n)$ (their Proposition 1.5). They asked (their Problem 1.6) for the value of

$$L := \limsup_{n\to\infty} \max_{|X|=n} \frac{\min\{\mu(\Delta_2), \mu(\delta)\}}{n}.$$

Vesztergombi [23] proved that $\mu(\Delta_2) \le \frac{3}{2}n$ for every n-point planar set, which gives $L \le 3/2$. On the Erdős problems forum, the user ienjoymath [17] gave a construction showing $L \ge 9/7$ and conjectured $L = 9/7$. As far as we know, $9/7 \le L \le 3/2$ were the best bounds before this paper.

Results. Our main result is an upper bound strictly below 3/2. For a set with at least two distances put

$$\tau(X) := \frac{\Delta^2 - \Delta_2^2 - \delta^2}{2\,\Delta_2\,\delta},$$

a scale-invariant measure of how much the diameter exceeds Δ_2 (in the scaling $\delta = 1$ used in the proof, $\tau = t - (1-t^2)/(2\Delta_2)$ with $t = \Delta - \Delta_2$; see Section 2).

Theorem 1. *There is an absolute constant C_0 such that every n-point set $X \subset \mathbb{R}^2$ satisfies*

(a) $\min\{\mu(\Delta_2), \mu(\delta)\} \le \frac{15}{11}\,n + C_0$;

(b) *if moreover $\Delta \le 1.94\,\Delta_2$ and $\tau(X) \ge \sqrt{3}/2 + 3/100$, then* $\min\{\mu(\Delta_2), \mu(\delta)\} \le \frac{4}{3}\,n + C_0$.

Consequently $L \le 15/11 \approx 1.3636$.

(Here X is assumed to determine at least two distances; otherwise $n \le 3$.)

The constant C_0 is explicit; the proof gives $C_0 = 3N_0 + 2\cdot 10^8$ with $N_0 = 1.51\cdot 10^9$ (Section 8). We have made no attempt to optimise it. A weaker bound, $\frac{54}{37}n + C_0$ with $C_0 = 3N_0 + 10^8$, obtained by a different argument in Regions II and III and with the trivial bound $\mu(\delta) \le e(S) + 6|R|$ in place of Section 6, is formalised in Lean 4; that argument is written out in Appendix B, and the formalisation is described in Section 10.2. The bound $\frac{15}{11}$ is not formalised.

Theorem 9.6 and Corollary 9.7 give a new class of sets for which the answer to the first question of Problem #132 is yes. Let $m_3 := n - |L_1| - |L_2|$ be the number of points of depth at least 3. Condition (1) contains the case $2|L_1| + |L_2| \le n$, i.e. $|L_1| \le m_3$. We show that $|L_1| \le 2m_3$ suffices under a genericity hypothesis that holds, for instance, when no four points of X are concyclic. The inputs we need from Vesztergombi's structure theory [24] are proved here from scratch (Proposition 9.2 and Theorem 9.5).

Section 10 treats the second question of Problem #132 in convex position. Theorem 10.1 states that a set of n points in convex position has at least three rare distances for every n with $6 \le n \le 18$. Some of these cases follow from the literature, some follow from a parity-sharp version of the counting argument of [7], and the cases $n = 11, 13, 15, 17$ are new and computer-assisted. Theorem 10.14 shows that a set in convex position with all but w points on one circle has at least $(n - 3w + 1)/4$ rare distances once $n + w \ge 91$ (and $n \ge 3w$); so the second question of Problem #132 has a positive answer for such sets when $w = o(n)$. In Section 11 we list open problems.

Idea of the proof of Theorem 1. Let S be the set of points that have a partner at distance Δ_2, let $Q \subseteq S$ be the set of points with exactly one such partner, and put $s = |S|$, $p = |Q|$, $r = n - s$. Vesztergombi's bound applied to $S \setminus Q$ gives $\mu(\Delta_2) \le \frac{3}{2}s - \frac{1}{2}p$. On the δ side we use the exact identity $\mu(\delta) = e(S) + \sum_{y \in R} \frac{1}{2}(\deg y + m_y)$, where $e(S)$ is the number of δ-pairs inside S, $R := X \setminus S$, $\deg y$ is the number of δ-neighbours of y and m_y the number of those in S. We prove two estimates: $e(S) \le s + p + O(1)$, and $\deg y + m_y \le 10$ for all but $O(1)$ points $y \in R$ (and ≤ 8 in Region II). The combination with weights $\frac{8}{11}$ and $\frac{3}{11}$ gives $\frac{15}{11}n + O(1)$ (Lemma 3.5).

For the first estimate: almost all points of S lie on the boundary of a convex body, either the "ball polygon" $K = \bigcap_{w \in X} \mathrm{B}(w, \Delta_2)$ or the convex hull, and each has a well-defined inward direction. All but $O(1)$ of the δ-edges inside S join points with almost equal inward directions. Such edges are either tangential edges, of which there are at most s, or *rungs* joining the two boundaries, whose geometry is fixed by an exact identity (Lemma 2.2(c)). A corner lemma shows that the inner end of a rung has only one Δ_2-partner, up to $O(1)$ exceptions, unless a second partner sits in one exactly determined reflected position; in that case the rung forces a missing tangential edge elsewhere. So every rung is paid for, either on the Δ_2 side (through p) or by a missing tangential edge. For the second estimate: a point of R that is not a diametral endpoint lies in the interior of K, while its neighbours in S do not; where ∂K is nearly flat on a long stretch, this confines those neighbours to an open arc of less than $240°$, so there are at most 4 of them, and the points y near which ∂K is not flat are $O(1)$ in number. The regime where the diameter is much larger than Δ_2 is handled separately by an incidence argument.

Notation. $\mathrm{B}(c, r)$ and $\mathrm{C}(c, r)$ are the closed disc and the circle with centre c and radius r. For unit vectors u, u', $\angle(u, u') \in [0, \pi]$ is the angle between them; recall $|u - u'| \le \angle(u, u')$. Angles are in radians unless marked with a degree sign.

2. Set-up for the proof of Theorem 1

We scale so that $\delta = 1$. Throughout Sections 2 to 8 and Appendix B, X is an n-point set with $n \ge 4$ and at least two distinct distances, so $\Delta > \Delta_2 \ge \delta = 1$. We put

$$t := \Delta - \Delta_2 > 0, \qquad \tau := t - \frac{1 - t^2}{2\Delta_2}, \qquad \beta := \frac{1}{100}.$$

Then $\tau = (\Delta^2 - \Delta_2^2 - 1)/(2\Delta_2)$, which is the quantity $\tau(X)$ of the introduction, and $\tau - t = (t^2 - 1)/(2\Delta_2)$. Hence $\tau \le t$ when $t \le 1$, and $\tau > 1$ if and only if $t > 1$.[C2] A crucial feature of X is that it has *no distance in the open interval* (Δ_2, Δ).

We use the following objects.

- D is the set of endpoints of diametral pairs. Each point of D is a vertex of $P := \operatorname{conv} X$.
- S is the set of points of X that have a partner at distance Δ_2; $R := X \setminus S$ and $r := |R|$, $s := |S|$.
- $Q \subseteq S$ is the set of points with exactly one partner at distance Δ_2 (Δ_2-*degree* 1), and $p := |Q|$.
- $K := \bigcap_{w \in X} \mathrm{B}(w, \Delta_2)$, a compact convex set (a "ball polygon").
- $e(Y)$ is the number of δ-pairs inside $Y \subseteq X$. For $y \in X$, $\deg y$ is the number of δ-neighbours of y, and for $y \in R$, m_y is the number of δ-neighbours of y that lie in S.

The *degenerate regime* is $\Delta > 1.94\,\Delta_2$ (Section 7); the *nondegenerate regime* is $\Delta \le 1.94\,\Delta_2$ (Sections 3 to 6).

Lemma 2.1 (ball polygon). *(a)* $X \setminus D \subseteq K$.
(b) $S \setminus D \subseteq \partial K$.
(c) $\operatorname{dist}(z, K) \ge t$ *for every* $z \in D$.

Proof. (a) A point $y \notin D$ has all its distances at most Δ_2, since there are none in (Δ_2, Δ). (b) If $v \in S \setminus D$ and $|vw| = \Delta_2$, then $v \in K \cap \mathrm{C}(w, \Delta_2) \subseteq \partial K$. (c) If $z \in D$ and $|zx| = \Delta$, then $K \subseteq \mathrm{B}(x, \Delta_2)$, while $|zx| = \Delta_2 + t$. □

Direction field. For each $p \in S$ we fix a unit vector u_p.

- If $p \in S \setminus D$, choose a Δ_2-partner w_p and put $u_p := (w_p - p)/\Delta_2$. Since $K \subseteq \mathrm{B}(w_p, \Delta_2)$, u_p is an inward normal of K at p.
- If $p \in S \cap D$, choose a diametral partner x_p and put $u_p := (x_p - p)/\Delta$. This is an inward normal of P at p.

Lemma 2.2 (exact identities)**.** *Let $y \in X$, $y \neq p$.*

(a) If $p \in S \cap D$, then $(y - p) \cdot u_p \geq |y - p|^2/(2\Delta) > 0$.
(b) If $p \in S \setminus D$, then either $(y - p) \cdot u_p \geq |y - p|^2/(2\Delta_2)$ (we say y is normal *for p), or $|y - w_p| = \Delta$, so that $y \in D$ (we say y is an* exception *for p).*
(c) In the exception case with $|y - p| = 1$ we have exactly $(y - p) \cdot u_p = -\tau$. Hence $y - p = -\tau u_p \pm \sigma u_p^{\perp}$ with $\sigma := \sqrt{1 - \tau^2}$. In particular exceptions at distance 1 exist only when $\tau \leq 1$.

Proof. (a) $|y - x_p|^2 = |y - p|^2 - 2\Delta(y - p) \cdot u_p + \Delta^2 \leq \Delta^2$. (b) $|y - w_p|^2 = |y - p|^2 - 2\Delta_2(y - p) \cdot u_p + \Delta_2^2$, and $|y - w_p| \in [0, \Delta_2] \cup \{\Delta\}$. (c) With $|y - p| = 1$ and $|y - w_p| = \Delta = \Delta_2 + t$ the same identity gives $(y - p) \cdot u_p = (1 + \Delta_2^2 - \Delta^2)/(2\Delta_2) = (1 - 2\Delta_2 t - t^2)/(2\Delta_2) = -\tau$.[C2] □

The proof uses only $|p - x_p| = \Delta$, resp. $|p - w_p| = \Delta_2$. So (a) holds for every $p \in D$ (also $p \in R \cap D$) and every diametral partner x of p in place of x_p, and (b), (c) hold for every Δ_2-partner w of p in place of w_p, with u_p replaced by $(w - p)/\Delta_2$. We use these variants below.

Lemma 2.3 (no common normal)**.** *If $p \neq q$ both lie in $S \setminus D$, then $u_p \neq u_q$.*

Proof. Neither point lies in D, so each is normal for the other: $(q - p) \cdot u_p > 0$ and $(p - q) \cdot u_q > 0$. These cannot both hold if $u_p = u_q$. □

Good edges, T-edges and rungs. A δ-edge pq inside S is *good* if $\angle(u_p, u_q) \leq \beta$, and *bad* otherwise. For a good edge put $e := q - p$, $a := e \cdot u_p$ and $b := -e \cdot u_q$. Then $|a + b| = |e \cdot (u_p - u_q)| \leq \beta$.

Lemma 2.4 (dichotomy)**.** *Assume $\tau > 2\beta$. Each good edge is of exactly one of the following two kinds.*

- T-edge: $a, b \in (0, \beta)$.
- Rung: *one end $v \in S \setminus D$ sees the other end $z \in S \cap D$ as an exception. Then $(z - v) \cdot u_v = -\tau$ and $(v - z) \cdot u_z \in [\tau - \beta, \tau + \beta]$.*

Proof. By Lemma 2.2, from each end the other end is either normal (then the corresponding one of a, b is positive) or an exception (then it equals $-\tau$). An exception target lies in D, and points of D see everything as normal, so both ends cannot be exceptions. If neither is, then $a, b > 0$ and $a + b \leq \beta$. If one is, say $a = -\tau$, then $b \in [\tau - \beta, \tau + \beta]$, and $b > 0$ shows that b is not an exception value. □

We say a T-neighbour q of p is on the *right* (resp. *left*) of p if $(q - p) \cdot u_p^{\perp} > 0$ (resp. < 0), where $u^{\perp}$ denotes u rotated by $-90°$.

(T1) Each p has at most one T-neighbour on each side. Indeed, two T-directions on one side both make an angle in $(0, \arcsin \beta)$ with $\pm u_p^{\perp}$, so they are less than $60°$ apart, whereas two points at distance 1 from p and at distance ≥ 1 from each other are at least $60°$ apart as seen from p.

(T2) If q is the right T-neighbour of p then p is the left T-neighbour of q, since $\angle(u_p, u_q) \le \beta$.

Hence the T-edges form vertex-disjoint directed paths and cycles, and every cycle has length at least 3 (a vertex cannot be both the left and the right T-neighbour of another). We say each point of S has a *left slot* and a *right slot*; a T-edge from p to its right T-neighbour q uses the right slot of p and the left slot of q. So the number e_T of T-edges is half the number of used slots, and $e_T \le s$. T-edges with both ends in $S \setminus D$ or both in $S \cap D$ are *pure*; the others are *mixed*. Note that (T1) and (T2) apply to mixed T-edges too.

Turning. Let γ be a closed arc of the boundary of a compact convex set with nonempty interior. Its *turning* $\operatorname{turn}(\gamma)$ is the length of the set $N(\gamma) \subseteq S^1$, the union of the normal cones at *all* points of γ, including the full normal cones at its endpoints.

Lemma 2.5 (multiplicity)**.** *Let $\gamma_1, \dots, \gamma_m$ be closed boundary arcs of one convex body, and suppose every boundary point lies in at most M of them. Then $\sum_i \operatorname{turn}(\gamma_i) \le 2\pi M$.*

Proof. For all but countably many directions $\nu \in S^1$ the set of boundary points with outer (or inner) normal ν is a single point y_ν, and $\nu \in N(\gamma_i)$ if and only if $y_\nu \in \gamma_i$. Integrate over ν. □

Points of $S \setminus D$ lie on ∂K and points of $S \cap D$ lie on ∂P; in each case distinct points are at Euclidean, hence arc-length, distance at least 1. So a boundary arc of length ℓ contains at most $\ell + 1$ of them, and a boundary of perimeter ℓ contains at most ℓ of them.

3. Reductions

Lemma 3.1 ((VS))**.** *For every $S' \subseteq S$, the number of Δ_2-pairs inside S' is at most $\frac{3}{2}|S'|$. In particular $\mu(\Delta_2) \le \frac{3}{2}s$.*

Proof. Every Δ_2-pair of X lies in S, and every distance within S' is a distance of X. If S' contains no Δ_2-pair there is nothing to prove. If S' contains a diametral pair of X, then $\operatorname{diam} S' = \Delta$ and, because X has no distances in (Δ_2, Δ), Δ_2 is the second-largest distance of S'; Vesztergombi's theorem [23] gives at most $\frac{3}{2}|S'|$ such pairs. Otherwise every distance in S' is at most Δ_2, so $\Delta_2 = \operatorname{diam} S'$ and Hopf–Pannwitz [16] gives at most $|S'|$. □

We use Vesztergombi's theorem in exactly the form of its main theorem: for any set of m points in the plane, the second-largest distance occurs at most $\frac{3}{2}m$ times [23, Theorem]. There is no threshold on m and no additive constant.

Lemma 3.2 (weighted Δ_2 bound)**.** $\mu(\Delta_2) \le p + \frac{3}{2}(s - p) = \frac{3}{2}s - \frac{1}{2}p$.

Proof. At most p Δ_2-pairs touch Q, since each point of Q lies in exactly one Δ_2-pair. Every other Δ_2-pair lies inside $S \setminus Q$, and there are at most $\frac{3}{2}(s - p)$ of them by Lemma 3.1. □

Lemma 3.3 (exact identity)**.** $\mu(\delta) = e(S) + \sum_{y \in R} \frac{1}{2}(\deg y + m_y)$. *Moreover* $\deg y \le 6$ *and* $m_y \le \deg y$*, so* $\deg y + m_y \le 12$ *for every* $y \in R$.

Proof. The δ-pairs inside S number $e(S)$. The pairs with one end in S and one in R number $\sum_{y \in R} m_y$. The pairs inside R are counted twice in $\sum_{y \in R}(\deg y - m_y)$. Adding, $\mu(\delta) = e(S) + \sum_y m_y + \frac{1}{2}\sum_y(\deg y - m_y)$, which is the identity. Each point has at most 6 points at distance δ, since these are pairwise at least δ apart. □

Corollary 3.4 ((E))**.** $\mu(\delta) \le e(S) + 6r$. □

Lemma 3.5 (linear combination)**.** *Suppose $e(S) \le s + p + C$, and that $\deg y + m_y \le 10$ for all but at most C' points $y \in R$. Then*

$$\min\{\mu(\Delta_2), \mu(\delta)\} \le \tfrac{15}{11}\, n + C + C'.$$

If instead $\deg y + m_y \le 8$ *for all but at most* C' *points* $y \in R$, *then* $\min\{\mu(\Delta_2), \mu(\delta)\} \le \frac{4}{3}n + C + C'$.

Proof. For an exceptional y we still have $\deg y + m_y \le 12$ (Lemma 3.3). So in the first case $\mu(\delta) \le s + p + 5r + C + C'$, and with Lemma 3.2

$$\min\{\mu(\Delta_2), \mu(\delta)\} \le \tfrac{8}{11}(\tfrac{3}{2}s - \tfrac{1}{2}p) + \tfrac{3}{11}(s + p + 5r + C + C') = \tfrac{15}{11}(s + r) - \tfrac{1}{11}p + \tfrac{3}{11}(C + C').$$

In the second case $\mu(\delta) \le s + p + 4r + C + 2C'$, and the weights $\frac{2}{3}, \frac{1}{3}$ give $\frac{4}{3}(s + r) + \frac{1}{3}(C + 2C')$.[LP] □

Remark 3.6. If $\deg y + m_y \le 2c$ for all but $O(1)$ points of R, the best weights give $\max\{\frac{4}{3}, \frac{3c}{2c+1}\}\, n + O(1)$ under the hypothesis $e(S) \le s + p + O(1)$, and $\frac{3c}{2c+1}\, n + O(1)$ under the stronger hypothesis $e(S) \le s + O(1)$; the weight on $\mu(\Delta_2)$ must be at least $\frac{2}{3}$ to cancel the p-term.[LP] For $2c = 12, 10, 8, 6$ these values are $\frac{18}{13}, \frac{15}{11}, \frac{4}{3}, \frac{4}{3}$, respectively $\frac{18}{13}, \frac{15}{11}, \frac{4}{3}, \frac{9}{7}$. In particular the trivial bound (E) together with $e(S) \le s + p + O(1)$ already gives $\frac{18}{13}n + O(1)$, with weights $\frac{10}{13}, \frac{3}{13}$.

So in the nondegenerate regime it suffices to show $e(S) \le s + p + O(1)$ in every τ-regime (Section 5), and $\deg y + m_y \le 10$ (resp. ≤ 8 in Region II) for all but $O(1)$ points $y \in R$ (Section 6).

Small cases. Suppose we are in the nondegenerate regime and $\Delta_2 < 10^4$. Then $\Delta < 1.94 \cdot 10^4$, so X lies in a disc of radius Δ about any of its points, and since the discs of radius $\frac{1}{2}$ about points of X are disjoint, $n \le (2\Delta + 1)^2 < N_0 := 1.51 \cdot 10^9$.[C11] Since $\mu(\delta) \le 3n$, Theorem 1 holds for such X once $C_0 \ge 3N_0$. *From now on, in the nondegenerate regime,* $\Delta_2 \ge 10^4$; in particular $2/\Delta_2 \le \beta/2$.[C3]

4. Bad edges are $O(1)$ in the nondegenerate regime

4.1. A lens lemma.

Lemma 4.1 (lens)**.** *Let* $\Lambda = \mathrm{B}(c_1, \rho) \cap \mathrm{B}(c_2, \rho)$ *with* $0 < |c_1 c_2| = d < 2\rho$, *and put* $\cos\varphi_0 := d/(2\rho)$. *Assume*

$$\rho \;\ge\; \frac{2}{\cos(\varphi_0/2) - \cos\varphi_0}. \tag{2}$$

Let p *lie on the arc of* $\partial\Lambda$ *belonging to* $\mathrm{C}(c_1, \rho)$, *let* q *lie on the arc belonging to* $\mathrm{C}(c_2, \rho)$, *and suppose* $|pq| \le 1$. *Then there is a corner* c *of* Λ *such that the boundary arcs from* p *to* c *and from* c *to* q *each have length at most* $\rho_0 := 1/\sin(\varphi_0/2)$.

Proof. Take $c_1 = (-d/2, 0)$, $c_2 = (d/2, 0)$, $p = c_1 + \rho(\cos\varphi, \sin\varphi)$ and $q = c_2 + \rho(-\cos\psi, \sin\psi)$ with $|\varphi|, |\psi| \le \varphi_0$. *Step 1.* $p_x - q_x = \rho(\cos\varphi - \cos\varphi_0) + \rho(\cos\psi - \cos\varphi_0)$ with both terms ≥ 0; since $|p_x - q_x| \le 1$, each term is at most 1. By (2), $\cos\varphi < \cos(\varphi_0/2)$, i.e. $|\varphi| > \varphi_0/2$; likewise $|\psi| > \varphi_0/2$. *Step 2.* By the mean value theorem, $\rho(\varphi_0 - |\varphi|) = \rho(\cos\varphi - \cos\varphi_0)/\sin\xi \le 1/\sin\xi$ for some $\xi \in (\varphi_0/2, \varphi_0) \subseteq (0, \pi/2)$, so $\rho(\varphi_0 - |\varphi|) \le \rho_0$; likewise for ψ. *Step 3.* If φ, ψ had opposite signs, then $|p_y - q_y| \ge 2\rho\sin(\varphi_0/2)$. Now (2) gives $\rho \ge 2/(1 - \cos\varphi_0) = 1/\sin^2(\varphi_0/2)$, so $2\rho\sin(\varphi_0/2) \ge 2/\sin(\varphi_0/2) \ge 2 > 1$, a contradiction. *Step 4.* So the corner $c = (0, \mathrm{sign}(\varphi)\rho\sin\varphi_0)$ works. □

Lemma 4.2 (lens-corner arc length)**.** *In the situation of Lemma 4.1, suppose moreover that* $p, q \in \partial K'$ *for a compact convex* $K' \subseteq \Lambda$, *and* $|pq| = 1$. *Let* H *be the closed half-plane bounded by the line* pq *that contains* c, *and let* $\gamma := \partial K' \cap H$ *be the boundary arc of* K' *from* p *to* q *on* c*'s side. Then* $\mathrm{length}(\gamma) \le 2\rho_0$.

Proof. $\partial\Lambda$ contains no segment, so the line pq meets it only in p and q, and $\partial\Lambda \cap H$ is exactly the arc $p \to c \to q$, of length at most $2\rho_0$. Since $K' \cap H \subseteq \Lambda \cap H$ are convex, monotonicity of perimeter gives $\text{length}(\gamma) + 1 \le 2\rho_0 + 1$. (If $K' \cap H$ is the segment $[p, q]$, then γ is that segment, of length $1 \le 2\rho_0$.) □

Constants.[C7] For *KK lenses* ($\rho = \Delta_2$, $d \le \Delta \le 1.94\Delta_2$) we have $\cos\varphi_0 \le 0.97$, so $\rho_0^2 = 2/(1-\cos\varphi_0) \le 66.7 < 8.2^2$. The function $c \mapsto \cos(\varphi_0/2) - c = \sqrt{(1+c)/2} - c$ is decreasing on $[0, 1)$, so (2) holds as soon as $\rho \ge 89.01$, in particular for $\rho = \Delta_2 \ge 10^4$. For *DD lenses* ($\rho = \Delta$, $d \le \Delta$) we have $\cos\varphi_0 \le \frac{1}{2}$, $\rho_0 \le 2$, and (2) needs only $\rho \ge 5.5$.

4.2. **The count.** We split bad edges pq by type: *KK* (both ends in $S \setminus D$), *DD* (both in $S \cap D$) and *cross* (one end in each).

If the perimeter of K is below 1202, then $|S \setminus D| < 1202$ and at most $6 \cdot 1202$ edges touch $S \setminus D$; similarly for P and $S \cap D$. We may therefore assume both perimeters are at least 1202 when counting the bad edges touching the corresponding row; these small cases contribute a constant below $1.5 \cdot 10^4$ in total.

KK edges. Let $p, q \in S \setminus D$ form a bad edge. If $w_p = w_q$, then $\angle(u_p, u_q) \le 2\arcsin(1/(2\Delta_2)) < \beta$,[C3] so the edge would be good; hence $w_p \ne w_q$. Then $K \subseteq \Lambda := \mathrm{B}(w_p, \Delta_2) \cap \mathrm{B}(w_q, \Delta_2)$, p lies on the $\mathrm{C}(w_p, \Delta_2)$-arc and q on the $\mathrm{C}(w_q, \Delta_2)$-arc of $\partial\Lambda$, and $d = |w_p w_q| \le \Delta \le 1.94\Delta_2$. Lemmas 4.1 and 4.2 give an arc $\gamma_{pq} \subseteq \partial K$ from p to q of length at most $2\rho_0$. The union of the normal cones along the closed arc γ_{pq} is connected and contains u_p and u_q, so $\text{turn}(\gamma_{pq}) \ge \angle(u_p, u_q) > \beta$. If $y \in \gamma_{pq}$ then p lies within arc distance $2\rho_0$ of y; there are at most $4\rho_0 + 1$ such p, each of degree at most 6, so every point lies on at most $M = 6(4\rho_0 + 1)$ of the arcs. By Lemma 2.5,

$$\#\text{KK-bad} \le 12\pi(4\rho_0 + 1)/\beta < 1.28 \cdot 10^5.\text{[C9]}$$

DD edges. The same argument, on ∂P with the lens $\mathrm{B}(x_p, \Delta) \cap \mathrm{B}(x_q, \Delta)$ and $\rho_0 \le 2$, gives $\#\text{DD-bad} \le 12\pi \cdot 9/\beta < 3.4 \cdot 10^4$. (If $x_p = x_q$ the angle is less than $2/\Delta < \beta$ and the edge is good.)

Cross edges. Let zv be a bad edge with $z \in S \cap D$ and $v \in S \setminus D$. Since $z \in X$, $|zw_v| \in [0, \Delta_2] \cup \{\Delta\}$.

(c1) $|zw_v| = \Delta$. By Lemma 2.2(a), $(w_v - z)/\Delta$ is a second inward normal of P at z. Its angle to u_v is the angle at w_v of the triangle zvw_v; that angle is opposite the side of length 1, hence acute, and by the law of sines it is at most $\arcsin(1/\Delta) < 2/\Delta_2 \le \beta/2$.[C3] As $\angle(u_z, u_v) > \beta$, the normal cone of P at z has angle greater than $\beta/2$. Normal cones at distinct vertices have disjoint interiors, so fewer than $4\pi/\beta$ points z qualify, each of degree at most 6: $\#(\text{c1}) < 24\pi/\beta < 7600$.

(c2) $|zw_v| \le \Delta_2$. Let g be the unit vector from v towards x_z, and $\alpha := \angle(u_v, g)$.

(i) *Depth.* $v \notin D$ gives $|vx_z| \le \Delta_2$, while $|zx_z| = \Delta$ and $|zv| = 1$ give $|vx_z| \ge \Delta - 1$. So $v \in \mathrm{B}(x_z, \Delta_2)$ at depth $d_0 := \Delta_2 - |vx_z| \in [0, 1-t]$; in particular this case needs $t \le 1$. Also $K \subseteq \mathrm{B}(x_z, \Delta_2)$.

(ii) *α is not small.* $\angle(u_z, g)$ is the angle at x_z of the triangle zvx_z, which is at most $\arcsin(1/\Delta) < \beta/2$; hence $\alpha \ge \angle(u_v, u_z) - \angle(u_z, g) > \beta/2$.

(iii) *α is not close to π.* Since $w_v, x_z \in X$, $|w_v x_z| \le \Delta \le 1.94\Delta_2$. The law of cosines in the triangle vw_vx_z (sides Δ_2 and $\Delta_2 - d_0$ at v, enclosing the angle α) gives $\cos\alpha \ge (1 + \varrho^2 - 1.94^2)/(2\varrho)$, where $\varrho := (\Delta_2 - d_0)/\Delta_2 \in [1 - 10^{-4}, 1]$. The right side is increasing in ϱ, so $\cos\alpha \ge -0.8821$, i.e. $\alpha \le 151.9°$. By concavity of sin on $[0, \pi]$, $\sin\alpha \ge \min\{\sin(\beta/2), 0.471\} = \sin(\beta/2)$.[C8]

(iv) *The walk.* Let $e_0 \perp u_v$ be the unit vector with $e_0 \cdot g = -\sin\alpha < 0$. Follow ∂K from v in the direction whose forward one-sided tangent is closest to e_0, parametrised by arc length, $\gamma(0) = v$. Suppose the turning of $\gamma([0,\sigma])$ is at most $\beta/4$. This turning includes the whole normal cone $N_K(v)$. Then every normal along the arc is within $\beta/4$ of u_v, so every forward tangent γ' is within $\beta/4$ of e_0 (at a corner v, the forward tangent is perpendicular to an extreme normal of $N_K(v)$, which is also within $\beta/4$ of u_v). Hence $\gamma'\cdot g \le -\sin\alpha + 2\sin(\beta/8) \le -\sin(\beta/2) + 2\sin(\beta/8) < -\beta/5$.[C8] Integrating, $(\gamma(\sigma)-v)\cdot g \le -\sigma\beta/5$, and so

$$|\gamma(\sigma) - x_z|^2 \;\ge\; (\Delta_2 - d_0)^2 + \tfrac{2}{5}(\Delta_2 - d_0)\,\sigma\beta.$$

For $\sigma = L := 6/\beta = 600$ the right side exceeds Δ_2^2,[C8] which puts $\gamma(L)$ outside $\mathrm{B}(x_z, \Delta_2) \supseteq K$, a contradiction. Hence the closed arc J_v of length L from v in the chosen direction has turning greater than $\beta/4$.

(v) *Count.* A point y lies in J_v only if v is within arc distance L of y, which allows at most $2L+1$ points v. By Lemma 2.5, $\#\{v\} \le 2\pi(2L+1)/(\beta/4)$. Each such v has degree at most 6, so $\#(\mathrm{c2}) \le 48\pi(2L+1)/\beta < 1.82 \cdot 10^7$.[C9]

Proposition 4.3. *In the nondegenerate regime the number C_{bad} of bad edges satisfies $C_{\mathrm{bad}} < 1.83 \cdot 10^7$.[C9]*

5. The nondegenerate regime: bounding $e(S)$

In this section we prove $e(S) \le s + p + C$ with an absolute constant C in every τ-regime of the nondegenerate regime. Rungs are counted with the bad edges excluded: $e(S) \le e_T + \rho + C_{\mathrm{bad}}$, where ρ is the number of good rungs. (When $\tau \le 2\beta$, Lemma 2.4 is not used.)

5.1. **Region I: $\tau \le \sqrt{3}/2 - 2\beta$.** Let $p \in S \setminus D$ and let q be a good neighbour of p; the *elevation* of $q - p$ is $\arcsin((q-p)\cdot u_p)$. If q is normal for p, then $b > 0$ (by Lemma 2.2(a) if $q \in D$, by (b) if $q \notin D$), so $a < \beta$. Otherwise q is an exception and $a = -\tau$.

- If $\tau \ge 0$, every good neighbour on one side of $\mathbb{R}u_p$ has elevation in $[-\arcsin\tau, \arcsin\beta)$, a window of width less than $\arcsin(\sqrt{3}/2 - 2\beta) + \arcsin\beta \approx 58.35^\circ < 60^\circ$.[C4]
- If $\tau < 0$, then $|\tau| \le 1/(2\Delta_2)$ and all elevations lie in $(0, \arcsin\beta) \cup \{\arcsin|\tau|\}$; the window is again less than 60°.[C4]

Let $p \in S \cap D$. Then $a > 0$ always, and either $a < \beta$, or p is the exception of q and $a \in [\tau - \beta, \tau + \beta]$; the window is at most $\arcsin(\sqrt{3}/2 - \beta) \approx 58.87^\circ < 60^\circ$.[C4]

Since two neighbours of p are at least 60° apart, p has at most one good neighbour on each side. Hence

$$e(S) \le s + C_{\mathrm{bad}} \qquad \text{in Region I.}$$

5.2. **Common facts for $\tau > \sqrt{3}/2 - 2\beta$.** Here $\tau > 2\beta$,[C5,C6] so Lemma 2.4 applies: every good edge is a T-edge or a rung. *Region II* is $\tau \ge \sqrt{3}/2 + 3\beta$; *Region III* is $\sqrt{3}/2 - 2\beta < \tau < \sqrt{3}/2 + 3\beta$. If $\tau > 1$ then $t > 1$, and by Lemma 2.2(c) there are no rungs; so $e(S) \le e_T + C_{\mathrm{bad}} \le s + C_{\mathrm{bad}}$ by (T1). *From now on in this section assume* $\tau \le 1$; then $t \le 1$ and $t \ge \tau$ (Section 2). Put

$$\theta_0 := \arccos\tau \in [0, \pi/2), \qquad \sigma = \sqrt{1-\tau^2} = \sin\theta_0, \qquad o_\pm(u) := -\tau u \pm \sigma u^\perp.$$

By Lemma 2.2(c), a rung (v, z) with $v \in S \setminus D$ has $z = v + o_+(u_v)$ or $z = v + o_-(u_v)$; in particular *a K-vertex has at most two rungs.* The map $u \mapsto o_\pm(u)$ is a rotation, so $|o_\pm(u) - o_\pm(u')| = |u - u'| \le \angle(u, u')$, and $|o_+(u) - o_-(u)| = 2\sigma$. The *K-row* is $S \setminus D$ and the *D-row* is $S \cap D$; rungs join the two rows.

T-steps. If q is the right T-neighbour of p, then $q - p = \cos\varepsilon\, u_p^\perp + \sin\varepsilon\, u_p$ with $\varepsilon \in (0, \arcsin\beta)$, since $a = (q-p)\cdot u_p \in (0, \beta)$.

Lemma 5.1 (mixed T-edges)**.** *Assume $\sqrt{3}/2 - 2\beta < \tau \le 1$. Let $v \in S \setminus D$ and $z \in S \cap D$ be joined by a T-edge. Then the closed arc of ∂K of length 2 starting at v on z's side has turning at least $t - \arcsin\beta$. Consequently there are fewer than $2\pi \cdot 10/(t - \arcsin\beta)$ mixed T-edges: fewer than 71 in Region II and fewer than 76 in Region III.*

Proof. Here $t \ge \tau > \sqrt{3}/2 - 2\beta > 0.846$. Suppose the turning is $\Theta < t - \arcsin\beta$; note $t - \arcsin\beta < 1 < \pi/3$. Let e_0 be the unit tangent direction at v perpendicular to u_v pointing to z's side. Every forward tangent along the arc is within Θ of e_0, so $|\gamma(\sigma') - v| \ge \sigma' \cos\Theta \ge \sigma'/2$ for the arc-length parametrisation γ, and some $\sigma^* \le 2$ has $|\gamma(\sigma^*) - v| = 1$. The unit vector $\gamma(\sigma^*) - v$ lies within Θ of e_0, and the unit vector $z - v$ lies within $\arcsin\beta$ of e_0. Hence $|\gamma(\sigma^*) - z| \le \Theta + \arcsin\beta < t \le \operatorname{dist}(z, K)$ by Lemma 2.1(c). But $\gamma(\sigma^*) \in K$, a contradiction.

For the count: each v has at most two T-edges, and v must lie within arc distance 2 of a point y of its arc, so each boundary point lies on at most $M \le 10$ of these arcs. By Lemma 2.5 the number of mixed T-edges is less than $2\pi \cdot 10/(t - \arcsin\beta)$. In Region II, $t - \arcsin\beta \ge 0.886$ and the bound is below 71;[C5] in Region III, $t - \arcsin\beta > 0.836$ and the bound is below 76.[III] □

Lemma 5.2 (corner lemma)**.** *Let $v \in S \setminus D$ have a rung partner z and a second Δ_2-partner $w' \ne w_v$, and put $u' := (w' - v)/\Delta_2$ and $c := z - v$. Then exactly one of the following holds.*

(i) *$|zw'| \le \Delta_2$. Then $c \cdot u' \ge 1/(2\Delta_2) > 0$ and $\angle(u_v, u') > \pi/2 - \theta_0$.*
(ii) *$|zw'| = \Delta$. Then $u' = u^*$, the reflection of u_v in the line $\mathbb{R}c$, and $\angle(u_v, u') = 2\theta_0 > 0$.*

In both cases u_v and u' are inward normals of K at v, and the normal cone $N_K(v)$ has angle at least $\angle(u_v, u')$.

Proof. Since $z \in X$, $|zw'| \in [0, \Delta_2] \cup \{\Delta\}$, so exactly one case occurs. We have $c \cdot u_v = -\tau$ and $|c| = 1$, so $\angle(c, u_v) = \pi - \theta_0$. (i) By Lemma 2.2(b) applied with w' in place of w_v, z is normal for v with respect to w': $c \cdot u' \ge |c|^2/(2\Delta_2) > 0$. So $\angle(c, u') < \pi/2$ and $\angle(u_v, u') \ge \angle(c, u_v) - \angle(c, u') > \pi/2 - \theta_0$. (ii) By Lemma 2.2(c) applied with w', $c \cdot u' = -\tau$. The unit vectors u with $c \cdot u = -\tau$ are the two vectors at angle θ_0 from $-c$, namely u_v and its reflection u^* in $\mathbb{R}c$. Since $w' \ne w_v$ and both are at distance Δ_2 from v, $u' \ne u_v$; so $u' = u^*$, $\theta_0 > 0$, and $\angle(u_v, u') = 2\theta_0$. Both u_v and u' are inward normals of K at v, because $K \subseteq \mathrm{B}(w_v, \Delta_2) \cap \mathrm{B}(w', \Delta_2)$. Moreover $\angle(u_v, u') < \pi$, because $|w_v w'| \le \Delta < 2\Delta_2$; so the (convex) normal cone contains the shorter arc between them. □

Normal cones of K at distinct boundary points have disjoint interiors. We say that v is *of type* (i) if some second Δ_2-partner of v is as in (i), and *of type* (ii) if v has Δ_2-degree at least 2 and is not of type (i). A rung K-end of type (ii) has Δ_2-degree exactly 2, since only the directions u_v and u^* are available.

5.3. **Region II: $\tau \ge \sqrt{3}/2 + 3\beta$.** We keep $\tau \le 1$. Then $\theta_0 \le \arccos(\sqrt{3}/2 + 3\beta) < 26.4^\circ < 30^\circ$ and $\sigma \le \sigma_{\max} \approx 0.4440$.[C5]

(R1) *Each vertex has at most one rung.* At $v \in S \setminus D$: the two candidate positions $v + o_\pm(u_v)$ are $2\sigma < 1$ apart. At $z \in S \cap D$: if z had rungs to $v \ne v'$, both rungs are good, so $\angle(u_v, u_{v'}) \le 2\beta$ and $|v - v'| = |o_\pm(u_v) - o_{\pm'}(u_{v'})| \le 2\sigma + 2\beta \approx 0.908 < 1$,[C5] a contradiction.

So rungs form a matching. By Lemma 5.2, a type-(i) vertex has a normal cone of angle greater than $\pi/2 - \theta_0 > \pi/3$, so *there are at most* 5 *type-*(i) *vertices.* Every other rung K-end either lies in Q or is of type (ii).

Lemma 5.3 (far partners persist)**.** *Let $z \in D$ and $w \in X$ with $|zw| = \Delta$, put $\hat{w} := (w - z)/\Delta$, and let $y \in X$ with $|zy| = 1$ and $\eta := (y - z) \cdot \hat{w}$. Then $|yw|^2 = \Delta^2 + 1 - 2\Delta\eta$. If*

$$\eta < \frac{\Delta^2 - \Delta_2^2 + 1}{2\Delta} = t + \frac{1 - t^2}{2\Delta},$$

then $|yw| > \Delta_2$, hence $|yw| = \Delta$. When $t \le 1$ the threshold is at least t.

Proof. The identity is $|y-w|^2 = |y-z|^2 - 2(y-z)\cdot(w-z) + |w-z|^2$. The inequality $\Delta^2 + 1 - 2\Delta\eta > \Delta_2^2$ is the displayed condition, and $\Delta^2 - \Delta_2^2 = 2\Delta t - t^2$ gives the second form of the threshold. There are no distances in (Δ_2, Δ).[II] □

Lemma 5.4. *Let (v, z) be a rung whose K-end v is of type* (ii)*. Then z has no T-neighbour at all, pure or mixed.*

Proof. Let $w := w_v$ and let w^* be the second Δ_2-partner of v, in direction u^* from v. Since the rung is an exception for v with respect to both partners (Lemma 5.2(ii)), $|zw| = |zw^*| = \Delta$. Put $\hat{w} := (w-z)/\Delta$ and $\hat{w}^* := (w^*-z)/\Delta$. The angle between $\hat{w}$ and u_v is the angle at w of the triangle vzw; it is opposite the side vz of length 1, hence acute, and by the law of sines it is at most $\arcsin(1/\Delta)$. Likewise $\angle(\hat{w}^*, u^*) \le \arcsin(1/\Delta)$. Also $\angle(u^*, u_v) = 2\theta_0$, and $\angle(u_v, u_z) \le \beta$ because the rung is good.

Let z_1 be a T-neighbour of z and $e := z_1 - z$, a unit vector with $e \cdot u_z \in (0, \beta)$, so $\angle(e, u_z) \ge \pi/2 - \arcsin\beta$. Put $\varepsilon := \beta + \arcsin\beta + \arcsin(1/\Delta)$. By the triangle inequality for angles, $\angle(e, \hat{w}) \ge \pi/2 - \varepsilon$ and $\angle(e, \hat{w}^*) \ge \pi/2 - (2\theta_0 + \varepsilon)$, so

$$\eta := e \cdot \hat{w} \le \sin\varepsilon, \qquad \eta^* := e \cdot \hat{w}^* \le \sin(2\theta_0 + \varepsilon).$$

Both are less than t: since $t \ge \tau = \cos\theta_0 = \sin(\pi/2 - \theta_0)$ and $0 \le \varepsilon \le 2\theta_0 + \varepsilon < \pi/2 - \theta_0$, it suffices that $3\theta_0 + \varepsilon < \pi/2$, and indeed $\theta_0 < 26.4^\circ$ and $\varepsilon < 1.2^\circ$ give $3\theta_0 + \varepsilon < 80.4^\circ$. By Lemma 5.3 (with $t \le 1$), $|z_1 w| = |z_1 w^*| = \Delta$. Now $w \ne w^*$, and z, z_1 are two distinct points of $\mathrm{C}(w, \Delta) \cap \mathrm{C}(w^*, \Delta)$. These two points are $2\sqrt{\Delta^2 - |ww^*|^2/4} \ge \sqrt{3}\,\Delta > 1$ apart, because $|ww^*| \le \Delta$. This contradicts $|zz_1| = 1$. □

The count in Region II. By (T1) every point of S has T-degree at most 2. By Lemma 5.4 the D-ends of rungs with a type-(ii) K-end have T-degree 0, and they are distinct by (R1). Let d be the number of these rungs. Summing T-degrees, $2e_T \le 2(s-d)$. Every other rung has a K-end in Q or of type (i), and each K-vertex has at most one rung (R1), so $\rho \le d + p + 5$. Hence

$$e(S) \le e_T + \rho + C_{\mathrm{bad}} \le (s-d) + (d+p+5) + C_{\mathrm{bad}} = s + p + 5 + C_{\mathrm{bad}} \qquad \text{in Region II.}$$

Mixed T-edges need no separate count here.

5.4. **Region III: $\sqrt{3}/2 - 2\beta < \tau < \sqrt{3}/2 + 3\beta$.** Here $\theta_0 \in (26.36^\circ, 32.22^\circ)$ and $t \ge \tau > 0.846$. By Lemma 5.2, a rung K-vertex of Δ_2-degree at least 2 has a normal cone of angle at least $\min\{2\theta_0, \pi/2 - \theta_0\} > 2\pi/7$, because $\tau < \sqrt{3}/2 + 3\beta < \cos(\pi/7)$ and $\tau > \sqrt{3}/2 - 2\beta > \cos(3\pi/14)$;[C6] so there are at most 6 such vertices.

Let q_1 and q_2 be the numbers of K-vertices with exactly one and exactly two good rungs. Then $\rho \le q_1 + 2q_2$, and $p \ge q_1 + q_2 - 6$. It therefore suffices to prove

$$e_T \le s - q_2 + C, \tag{3}$$

since then $e(S) \le e_T + \rho + C_{\mathrm{bad}} \le s + q_1 + q_2 + C + C_{\mathrm{bad}} \le s + p + C + C_{\mathrm{bad}} + 6$. A K-vertex with two rungs is called a *two-rung vertex*; its rungs are $z = v + o_+(u_v)$ and $z' = v + o_-(u_v)$, and $z - z' = 2\sigma u_v^\perp$.

(a) The case $\tau > \sqrt{3}/2$. The two positions $v + o_\pm(u_v)$ are $2\sigma < 1$ apart, so $q_2 = 0$, and (3) holds with $C = 0$ by (T1).

(b) The case $\tau < \sqrt{3}/2$: slot killing. Here $2\sigma > 1$; more precisely $0 < 2\sigma - 1 < 0.067$ and $\sigma < 0.5332$.[III]

Lemma 5.5. *Let v be a two-rung vertex with rungs $z = v + o_+(u_v)$ and $z' = v + o_-(u_v)$. Then z' has no T-neighbour (pure or mixed) on its right, and z has no T-neighbour on its left.*

Proof. Let y be the right T-neighbour of z'. Then $y - z' = \cos\varepsilon\, u_{z'}^{\perp} + \sin\varepsilon\, u_{z'}$ with $\varepsilon \in (0, \arcsin\beta)$, and $\angle(u_v, u_{z'}) \le \beta$ (good rung). So the angle γ' between $y - z'$ and $z - z' = 2\sigma u_v^{\perp}$ is at most $\gamma := \beta + \arcsin\beta < 0.0201$. Hence

$$|y - z|^2 = 1 + 4\sigma^2 - 4\sigma\cos\gamma' \le (2\sigma - 1)^2 + 4\sigma(1 - \cos\gamma) < 0.005 < 1.^{[III]}$$

Also $y \ne z$, since $|yz'| = 1 \ne 2\sigma = |zz'|$. This contradicts $\delta = 1$. The statement for z is symmetric. □

Injectivity. Suppose $v \ne \bar v$ are two-rung vertices with the same $z' = v + o_-(u_v) = \bar v + o_-(u_{\bar v})$. Put $\bar z := \bar v + o_+(u_{\bar v}) = z' + 2\sigma u_{\bar v}^{\perp}$. Since $\angle(u_v, u_{\bar v}) \le 2\beta$ (both within β of $u_{z'}$), $|\bar z - z| \le 2\sigma \cdot 2\beta < 1$, so $\bar z = z$.[III] Then $v, \bar v$ are the two points of $\mathrm{C}(z, 1) \cap \mathrm{C}(z', 1)$, mirror images in the line zz', which has direction $u_v^{\perp}$. As $v - z' = \tau u_v + \sigma u_v^{\perp}$, we get $\bar v - z' = -\tau u_v + \sigma u_v^{\perp}$, and $(\bar v - z') \cdot u_{z'} \le -\tau\cos\beta + \sigma\sin\beta < 0$,[III] contradicting Lemma 2.2(a) at $z' \in D$. The same holds for the z-ends. So the right slots killed at the z'-ends are pairwise distinct, and so are the left slots killed at the z-ends: Lemma 5.5 kills $2q_2$ distinct slots. As e_T is half the number of used slots, $e_T \le \frac{1}{2}(2s - 2q_2) = s - q_2$, which is (3).

(c) The case $\tau = \sqrt{3}/2$. Now $\sigma = \frac{1}{2}$ and $\theta_0 = 30°$. For a two-rung vertex v with $w := w_v$, $z - z' = u_v^{\perp}$ has length 1, so $vz'z$ is an equilateral triangle of side 1, the *triangle at v*; its *base* $z'z$ is a δ-edge inside $S \cap D$. By definition of rungs, $|zw| = |z'w| = \Delta$. (Property (R1) at D-vertices can fail here, since $2\sigma + 2\beta > 1$; the argument below does not use it.) We prove (3) with $C = C_{\mathrm{bad}} + 75$.

Base determines apex. The other point of $\mathrm{C}(z, 1) \cap \mathrm{C}(z', 1)$ is $\bar v = z' - \tau u_v + \frac{1}{2}u_v^{\perp}$, and $(\bar v - z') \cdot u_{z'} \le -\tau\cos\beta + \frac{1}{2}\sin\beta < 0$, contradicting Lemma 2.2(a). So a δ-edge is the base of at most one triangle, and at most C_{bad} triangles have a bad base. A good base joins two points of D, so by Lemma 2.4 it is a pure D-row T-edge; since $z - z' = u_v^{\perp}$ is within β of $u_{z'}^{\perp}$, z is the right T-neighbour of z'.

Small rows. If the K-row has at most 2 points then $q_2 \le 2$; if the D-row has at most 2 points, then there is at most one base and $q_2 \le 1$. In both cases (3) holds with $C = 2$ since $e_T \le s$. So assume both rows have at least 3 points.

Lemma 5.6 (rows are cyclic orders)**.** *If v' is the right pure T-neighbour of v in the K-row, then v' is the successor of v in the counterclockwise cyclic order of $S \setminus D$ on ∂K. The same holds in the D-row, on ∂P.*

Proof. Take coordinates with $v = (0, 0)$, $v' = (1, 0)$. By the T-step form, $u_v = (\sin\varepsilon, \cos\varepsilon)$ and $u_{v'} = (-\sin\varepsilon', \cos\varepsilon')$ with $\varepsilon, \varepsilon' \in (0, \arcsin\beta)$. These are inward normals of K at v and v', so the part of K with $y \le 0$ lies in the triangle $\{y \le 0,\ y \ge -x\tan\varepsilon,\ y \ge (x - 1)\tan\varepsilon'\}$. That triangle has $x \in [0, 1]$ and depth at most $\tan(\arcsin\beta)$, so each of its points is within $\sqrt{1/4 + \tan^2(\arcsin\beta)} < 1$ of v or v'.[III] Hence the boundary arc of K from v to v' in $\{y \le 0\}$ contains no other point of the row. Traversed from v to v' with K on the side of u_v (above), this arc runs counterclockwise, because $u^{\perp}$ is u rotated by $-90°$. For the D-row, u_z is an inward normal of P by Lemma 2.2(a), and the same argument applies on ∂P. □

Let $N_K := |S \setminus D|$ and $N_D := |S \cap D|$, and let B_K (resp. B_D) be the number of pairs $(x, \operatorname{succ} x)$ of cyclically consecutive points of the K-row (resp. D-row) that are not joined by a

pure T-edge. By Lemma 5.6 the pure T-edges of a row are among its N consecutive pairs, so there are $N_K - B_K$ and $N_D - B_D$ of them. With Lemma 5.1,

$$e_T \le s - B_K - B_D + 75. \tag{4}$$

If a point x has no pure right T-neighbour, then $(x, \operatorname{succ} x)$ is counted in B_K (resp. B_D): otherwise x would be the right T-neighbour of $\operatorname{succ} x$, and then $x = \operatorname{succ}(\operatorname{succ} x)$ by Lemma 5.6, so the row would have only 2 points.

Rotational sense. In both rows "right" means counterclockwise along the boundary (Lemma 5.6). Moreover, if $z_k \in S \cap D$ and its right T-neighbour z_{k+1} both lie on $\mathrm{C}(w, \Delta)$, then z_{k+1} is counterclockwise from z_k about w. Indeed, put $\hat{w}_k := (w - z_k)/\Delta$ and $e := z_{k+1} - z_k$. Then $e \cdot \hat{w}_k = 1/(2\Delta)$, so $e = \frac{1}{2\Delta}\hat{w}_k \pm \sqrt{1 - \frac{1}{4\Delta^2}}\,\hat{w}_k^{\perp}$. Since x_{z_k} and w are both diametral partners of z_k and $|x_{z_k} w| \le \Delta$, we have $\angle(u_{z_k}, \hat{w}_k) \le \pi/3$; and $\angle(e, u_{z_k}^{\perp}) < \arcsin\beta$ by the T-step form. Hence $\angle(e, \hat{w}_k^{\perp}) < \pi/3 + \arcsin\beta < \pi/2$, the sign is $+$, and moving along $\hat{w}_k^{\perp}$ ($\hat{w}_k$ rotated by $-90°$) is moving counterclockwise about w.

Lemma 5.7 (a D-row component lies on one circle)**.** *Let Z be the component of the graph of pure D-row T-edges that contains a good base $z'z$ of the triangle at v, and let $w := w_v$. Then $Z \subseteq \mathrm{C}(w, \Delta)$, consecutive points of Z are at angle exactly $\alpha_1 := 2\arcsin(1/(2\Delta))$ about w, all in the counterclockwise sense, and Z spans an angle at most $\pi/3$ about w. In particular Z is a path.*

Proof. We know $|zw| = |z'w| = \Delta$. Let $z_k \in Z$ have both a left T-neighbour z_{k-1} and a right T-neighbour z_{k+1}, and put $u := u_{z_k}$. By the T-step form, $z_{k+1} - z_k = \cos\varepsilon_1 u^{\perp} + \sin\varepsilon_1 u$ with $\varepsilon_1 \in (0, \arcsin\beta)$, and $z_{k-1} - z_k = -\cos\varepsilon_2 u^{\perp} + \sin\varepsilon_2 u$ where $\sin\varepsilon_2 = (z_{k-1} - z_k) \cdot u_{z_k}$ is the value b of the incoming T-edge, so $\varepsilon_2 \in (0, \arcsin\beta)$. Let $\nu = \cos\varphi\, u + \sin\varphi\, u^{\perp}$ be any inward normal of P at z_k. Then $\nu \cdot (z_{k+1} - z_k) = \sin(\varepsilon_1 + \varphi) \ge 0$ and $\nu \cdot (z_{k-1} - z_k) = \sin(\varepsilon_2 - \varphi) \ge 0$, so $\varphi \in [-\varepsilon_1, \varepsilon_2]$.

Suppose $|z_k w| = \Delta$. Then w is a diametral partner of z_k, and $\hat{w}_k := (w - z_k)/\Delta$ is an inward normal of P at z_k (Lemma 2.2(a) and the remark after it). So $\eta := (z_{k+1} - z_k) \cdot \hat{w}_k = \sin(\varepsilon_1 + \varphi) \in [0, \sin(2\arcsin\beta)]$, and $\sin(2\arcsin\beta) < 0.0201 < t$.[III] By Lemma 5.3, $|z_{k+1} w| = \Delta$. In the same way $(z_{k-1} - z_k) \cdot \hat{w}_k = \sin(\varepsilon_2 - \varphi) \in [0, 0.0201)$ and $|z_{k-1} w| = \Delta$.

Now induct outwards from the base, with no wrap-around: going right from z, each vertex reached has a left T-neighbour, so whenever it has a right T-neighbour it is an interior vertex and the step applies; going left from z' is symmetric. So every vertex of Z lies on $\mathrm{C}(w, \Delta)$. Two points of $\mathrm{C}(w, \Delta)$ at distance 1 are at angle α_1 about w, and by the rotational-sense remark each step to the right is counterclockwise about w. Points of X are at most Δ apart, and a chord $2\Delta\sin(\theta/2) \le \Delta$ has $\theta \le \pi/3$; so Z spans at most $\pi/3$, and monotone steps cannot close up into a cycle. □

Lemma 5.8 (one centre per component)**.** *If A and B are triangles, at v_A and v_B, whose good bases lie in the same component Z, then $w_{v_A} = w_{v_B}$.*

Proof. Write $w_A := w_{v_A}$, $w_B := w_{v_B}$, and let $z'_B z_B$ be the base of B. By Lemma 5.7 for A, $z'_B, z_B \in \mathrm{C}(w_A, \Delta)$; by the definition of rungs, $z'_B, z_B \in \mathrm{C}(w_B, \Delta)$. Suppose $w_A \ne w_B$; then w_A is the mirror image of w_B in the line $z'_B z_B$. Put $a := (w_B - z_B)/\Delta$ and $e := z'_B - z_B$. From $|z'_B - w_B| = \Delta$ we get $a \cdot e = 1/(2\Delta)$ exactly. The mirror image of a in the line $\mathbb{R}e$ is $a^* = 2(a \cdot e)e - a$, and $w_A - z_B = \Delta a^*$. So $(w_A - z_B) \cdot a = \Delta(2(a \cdot e)^2 - 1) < 0$. But w_B is a diametral partner of z_B and $w_A \in X$, so $(w_A - z_B) \cdot a \ge 0$ by Lemma 2.2(a). Contradiction. □

So all triangles with good base in Z share one centre w. The apex v_A of such a triangle lies on $\mathrm{C}(w, \Delta_2)$ and on the perpendicular bisector of its base (both v_A and w are equidistant from the

base endpoints), on the same side as the base; so, seen from w, it lies at the angle of the base midpoint. Order the triangles with good base in Z along Z. Let A, B be consecutive, B to the right, with bases (z_i, z_{i+1}) and (z_{i+k}, z_{i+k+1}). Then the apexes are at angle $k\alpha_1$ about w, with $k\alpha_1 \le \pi/3 - \alpha_1 < \pi/3$ by Lemma 5.7, and $k\alpha_1 \ge \alpha_2 := 2\arcsin(1/(2\Delta_2))$ because $|v_A v_B| \ge 1$.

The arc between consecutive apexes. Let γ_{AB} be the shorter arc of $\mathrm{C}(w, \Delta_2)$ from v_A to v_B. For $x \in X \setminus \{w\}$, $\mathrm{B}(x, \Delta_2) \cap \mathrm{C}(w, \Delta_2)$ is the arc $\{\theta : \cos(\theta - \theta_x) \ge |wx|/(2\Delta_2)\}$ of angular width at most π. It contains $v_A, v_B \in K$, which are less than π apart about w, so it contains γ_{AB}. Hence $\gamma_{AB} \subseteq K \subseteq \mathrm{B}(w, \Delta_2)$, and therefore $\gamma_{AB} \subseteq \partial K$.

Lemma 5.9 (the K-path is forced along the arc)**.** *Let $v_0 := v_A$ and follow pure right T-edges in the K-row: $v_0, v_1, \ldots$. As long as $v_i \in \gamma_{AB} \setminus \{v_B\}$ and v_i has a pure right T-neighbour, that neighbour is the point v^* of γ_{AB} at angle α_2 counterclockwise from v_i about w.*

Proof. Normal fact. $u_{v_i} = \hat{w}_i := (w - v_i)/\Delta_2$. For $i = 0$ this holds because $w = w_{v_A}$. For v_i in the relative interior of γ_{AB}, K contains the arc on both sides of v_i, and $K \subseteq \mathrm{B}(w, \Delta_2)$; so every inward normal ν at v_i satisfies $(y - v_i) \cdot \nu \ge 0$ for y on the arc on both sides, which forces ν to be a positive multiple of $\hat{w}_i$. Since u_{v_i} is an inward normal of K at v_i, $u_{v_i} = \hat{w}_i$.

v^* lies on γ_{AB}. Otherwise v_B would be at angle in $(0, \alpha_2)$ from v_i, and $|v_i v_B| < 1$.

The step. Use coordinates with second axis $u := \hat{w}_i$ and first axis $u^\perp$ (counterclockwise tangent). Then $v_{i+1} - v_i = (\cos\psi, \sin\psi)$ with $\psi \in (0, \arcsin\beta)$ (T-step form, with $u_{v_i} = \hat{w}_i$). From $v_{i+1} \in K \subseteq \mathrm{B}(w, \Delta_2)$, $|v_{i+1} - w|^2 = \Delta_2^2 + 1 - 2\Delta_2 \sin\psi \le \Delta_2^2$, so $\psi \ge \psi^* := \arcsin(1/(2\Delta_2))$, the elevation of v^*. Suppose $\psi > \psi^*$. Write $u_{v_{i+1}} = (\sin\varphi, \cos\varphi)$; $|\varphi| \le \beta$ because the T-edge is good. From $v_i \in K$, $(v_i - v_{i+1}) \cdot u_{v_{i+1}} = -\sin(\psi + \varphi) \ge 0$, so $\varphi \le -\psi < 0$. From $v^* \in K$ (as $v^* \in \gamma_{AB}$),

$$(v^* - v_{i+1}) \cdot u_{v_{i+1}} = (\cos\psi^* - \cos\psi)\sin\varphi + (\sin\psi^* - \sin\psi)\cos\varphi \ge 0.$$

But both terms are negative. So $\psi = \psi^*$ and $v_{i+1} = v^*$. □

So the path moves along γ_{AB} in steps of exactly α_2 about w, and it cannot pass v_B. It reaches v_B only if $j\alpha_2 = k\alpha_1$ for some integer $j \ge 1$; otherwise it stops at a point $b_{AB} \in \gamma_{AB} \setminus \{v_B\}$ of the K-row that has no pure right T-neighbour. (Its right slot is unused or used by a mixed T-edge.) In particular a pure right T-neighbour of a point of γ_{AB} never lies on another part of ∂K.

The path cannot land on v_B. Since $\alpha_2 > \alpha_1$, $j\alpha_2 = k\alpha_1$ forces $k > j$, so $j(\alpha_2 - \alpha_1) = (k - j)\alpha_1 \ge \alpha_1$ and

$$k\alpha_1 = j\alpha_2 \ge \frac{\alpha_1\alpha_2}{\alpha_2 - \alpha_1} = \frac{1}{1/\alpha_1 - 1/\alpha_2}.$$

For $\rho \ge 10^4$ put $f(\rho) := 1/(2\arcsin(1/(2\rho)))$. From $x \le \arcsin x \le x + x^3/(6(1 - x^2))$ (the Taylor coefficients of arcsin beyond the first are at most $\frac{1}{6}$) we get $\rho - 1/(23\rho) \le f(\rho) \le \rho$. Hence $1/\alpha_1 - 1/\alpha_2 = f(\Delta) - f(\Delta_2) \le t + 1/(23\Delta_2)$. At $\tau = \sqrt{3}/2$ we have $t \le 1$ and $t - \tau = (1 - t^2)/(2\Delta_2) \le 1/(2\Delta_2)$, so $t < 0.8661$. Therefore $k\alpha_1 \ge 1/(t + 1/(23\Delta_2)) > 1/0.8662 > 1.154 > \pi/3$,[III] contradicting $k\alpha_1 < \pi/3$. So every pair (A, B) of consecutive triangles in the same Z yields a point b_{AB} of the K-row in the half-open arc $[v_A, v_B)$ with no pure right T-neighbour, and hence a pair $(b_{AB}, \operatorname{succ} b_{AB})$ counted in B_K.

Counting. Distinct points b give distinct pairs $(b, \operatorname{succ} b)$. The half-open arcs $[v_A, v_B) \subseteq \partial K$ are pairwise disjoint:

- *Same Z:* they are consecutive intervals on one circle.
- *Different components $Z \ne Z'$ with the same centre w:* their angular ranges about w are disjoint, since a point of one strictly inside the range of the other would be at angle less

than α_1, hence at distance less than 1, from one of its points. The apex arcs lie in these ranges.

- *Different centres $w \neq w'$:* let $\gamma \subseteq \mathrm{C}(w, \Delta_2)$ and $\gamma' \subseteq \mathrm{C}(w', \Delta_2)$ be nondegenerate arcs of ∂K sharing a point x. If x were in the relative interior of γ, then near x the boundary ∂K coincides with γ on both sides, and γ' would contain a nondegenerate piece of ∂K lying on both circles, forcing $w = w'$. So x is an endpoint of both arcs; for the half-open arcs this means $x = v_A = v_{A'}$. The triangle at a K-vertex is unique, so $A = A'$.

Hence $B_K \geq \sum_Z (m_Z - 1)$, where m_Z is the number of triangles with good base in Z and the sum is over components with $m_Z \geq 1$. Each such Z is a path (Lemma 5.7); its right end has no pure right T-neighbour and so gives a pair counted in B_D, and distinct Z give distinct pairs. So $B_D \geq \#\{Z : m_Z \geq 1\}$. Adding, $B_K + B_D \geq \sum_Z m_Z \geq q_2 - C_{\mathrm{bad}}$, and (4) gives

$$e_T \leq s - q_2 + C_{\mathrm{bad}} + 75,$$

which is (3).

Conclusion for Region III. In all three cases, $e(S) \leq s + p + 2C_{\mathrm{bad}} + 81$.

Remark 5.10. In Regions II and III, the argument of Appendix B (ownership and pairing, rhombi, and the weighted bound with $2p$) is no longer needed. What replaces it is the observation that a rung beyond the first-order count is always paid for, either through p on the Δ_2 side or by a missing T-edge.

6. Points without a Δ_2-partner

In this section we are in the nondegenerate regime with $\Delta_2 \geq 10^4$; no assumption on τ is made. We bound $\deg y + m_y$ for $y \in R$.

Theorem 6.1. *All but fewer than $7.1 \cdot 10^7$ points $y \in R$ satisfy $m_y \leq 4$, and hence $\deg y + m_y \leq 10$.*

Theorem 6.2. *Let $0 < \varepsilon_0 \leq 1/50$, put $\varepsilon := \tan \varepsilon_0$, and suppose $t > \sqrt{3}/2 + \varepsilon$. Then all but at most $2\pi \cdot 468^2/\varepsilon_0 + 5106 + 465^2$ points $y \in R$ satisfy $m_y \leq 3$, and $m_y \leq 2$ if $\deg y = 6$; hence $\deg y + m_y \leq 8$. In Region II this holds with $\varepsilon_0 = 1/50$, and then the exceptional set has fewer than $7.1 \cdot 10^7$ points.*

In Region II, $t \geq \tau \geq \sqrt{3}/2 + 3\beta$ if $t \leq 1$, and $t > 1$ otherwise; since $\tan(1/50) < 0.0201 < 3\beta$, the hypothesis of Theorem 6.2 holds with $\varepsilon_0 = 1/50$.

6.1. Structure of R.

(a) *Points of $R \cap D$.* If $y \in R \cap D$, then by Lemma 2.2(a) (which holds for every point of D) all other points of X lie in an open half-plane bounded by a line through y. The δ-neighbours of y are pairwise at least 60° apart as seen from y, so $\deg y \leq 3$ and $\deg y + m_y \leq 6$.

(b) $\partial K \subseteq \bigcup_{w \in X} \mathrm{C}(w, \Delta_2)$. A point $x \in K$ with $|x - w| < \Delta_2$ for all (finitely many) $w \in X$ is an interior point of K. Hence a point of $X \setminus D$ on ∂K has a Δ_2-partner, i.e. lies in S. With Lemma 2.1(a): $R \setminus D \subseteq \operatorname{int} K$ and $S \setminus D = (X \setminus D) \cap \partial K$. So every δ-neighbour of a point of X lies in $\operatorname{int} K$ (and then in $R \setminus D$), in $S \setminus D \subseteq \partial K$, or in D, which is disjoint from K.

(c) *Depth.* For $y \in R \setminus D$ put $h := \operatorname{dist}(y, \partial K) > 0$. Every neighbour q of y in S is outside $\operatorname{int} K$: either $q \in S \setminus D \subseteq \partial K$, or $q \in D$ with $\operatorname{dist}(q, K) \geq t$ (Lemma 2.1(c)). The segment $[y, q]$ leaves K at a point $b \in \partial K$, so $h \leq |yb| \leq 1$, and $h \leq 1 - t$ if $q \in D$. *Consequently, if $h > 1$ then $m_y = 0$ and $\deg y + m_y \leq 6$.*

From now on let $y \in R \setminus D$ with $h \le 1$. Let $p_0 \in \partial K$ be a nearest boundary point, $n_0 := (p_0 - y)/h$, which is an outer normal of K at p_0, $H_0 := \{x : (x - p_0) \cdot n_0 \le 0\} \supseteq K$, and $\ell_0 := \partial H_0$.

Small K. If the perimeter of K is less than 464, then $\operatorname{diam} K < 232$ and, since the discs of radius $\frac{1}{2}$ about points of $X \setminus D \subseteq K$ are disjoint, $|X \setminus D| \le (2 \cdot 232 + 1)^2 = 465^2$. We may therefore assume that the perimeter of K is at least 464, and we define

$$\gamma_y := \text{the closed arc of } \partial K \text{ of points at arc-length distance at most 232 from } p_0.$$

6.2. **Thin K.**

Lemma 6.3 (thin K)**.** *If some point $b \in \partial K \cap \mathrm{B}(y, 1)$ does not lie on γ_y, then K lies in a rectangle of length $L < 400$ and width $W < 9$. Hence $|X \setminus D| < 5106$.*

Proof. Let ν_b be an outer normal of K at b, $H_b := \{x : (x - b) \cdot \nu_b \le 0\} \supseteq K$ and $\ell_b := \partial H_b$. Then $\operatorname{dist}(y, \ell_0) = h \le 1$ and $\operatorname{dist}(y, \ell_b) \le |yb| \le 1$. Both arcs of ∂K from p_0 to b have length greater than 232.

Case $\nu_b = n_0$. Then $\ell_0 = \ell_b$: otherwise one of H_0, H_b is strictly contained in the other, and the point $p_0 \in \ell_0$ or $b \in \ell_b$ lies outside the smaller half-plane, although both lie in K. So $[p_0, b] \subseteq K \cap \ell_0 \subseteq \partial K$ is a boundary arc of length $|p_0 b| \le 2 < 232$, a contradiction.

Case $\nu_b \ne \pm n_0$. The lines ℓ_0, ℓ_b meet at a point A, and $Z := H_0 \cap H_b$ is a closed cone with apex A and opening $\theta := \pi - \angle(n_0, \nu_b) \in (0, \pi)$. The point $y \in Z$ has distance at most 1 from both bounding lines; the set of such points of Z is a rhombus whose far vertex is on the bisector at distance $1/\sin(\theta/2)$ from A, so $|yA| \le 1/\sin(\theta/2)$. The point p_0 lies on the boundary ray of Z in ℓ_0 and b on the other boundary ray, and $|Ap_0| \le |Ay| + |yp_0| \le 1 + 1/\sin(\theta/2)$, and likewise for $|Ab|$. If $[p_0, b] \subseteq \partial K$ we conclude as before. Otherwise the line $p_0 b$ meets ∂K only in p_0, b; let H^+ be the closed half-plane bounded by it that contains A. Then $\partial K \cap H^+$ is one of the two arcs from p_0 to b, and $K \cap H^+$ is a convex subset of the triangle $Ap_0 b$. By monotonicity of perimeter, the length of this arc is at most $|Ap_0| + |Ab| \le 2 + 2/\sin(\theta/2)$. If $\theta \ge 1^\circ$ this is less than $231.2 < 232$,[N] a contradiction. So $\theta < 1^\circ$.

Fatness step ($\theta < 1^\circ$, or $\nu_b = -n_0$, in which case Z is a strip and we put $\theta := 0$). Let e be the direction of the bisector of Z (the direction of the strip if $\theta = 0$), oriented so that the cone opens towards increasing values of the *level* $x \cdot e$, and let $e^\perp$ be a unit normal; call $x \cdot e^\perp$ the *height*. The cross-section of Z at level λ has length $w(\lambda) = w(\lambda_y) + 2\tan(\theta/2)(\lambda - \lambda_y)$, where λ_y is the level of y and $w(\lambda_y) = (d_1 + d_2)/\cos(\theta/2) \le 2/\cos(\theta/2) =: w_y$, with $d_1, d_2 \le 1$ the distances from y to the two lines. Let $[E_-, E_+] \ni \lambda_y$ be the range of levels of K, $L := E_+ - E_-$, which is positive since K has interior points, and let W be the range of heights of K. The cross-sections of Z are nested, so $W \le w_y + 2\tan(\theta/2)L$; also $L \le \operatorname{diam} K \le 2\Delta_2$.

Pick $a, a' \in K$ at levels E_-, E_+, and let $\lambda_m := (E_- + E_+)/2$. The section of K at level λ_m is a segment of positive length, with top p_{top} and bottom p_{bot}. Let $n_{\mathrm{top}} = \cos\varphi\, e^\perp + \sin\varphi\, e$ be an outer normal of K at p_{top}. Writing $a - p_{\mathrm{top}} = -\frac{L}{2}e + v_a e^\perp$ and $a' - p_{\mathrm{top}} = \frac{L}{2}e + v_{a'} e^\perp$ with $|v_a|, |v_{a'}| \le W$, we have

$$-\tfrac{L}{2}\sin\varphi + v_a \cos\varphi \le 0, \qquad \tfrac{L}{2}\sin\varphi + v_{a'}\cos\varphi \le 0.$$

If $\cos\varphi = 0$, these two inequalities contradict each other, since $L > 0$. If $\cos\varphi < 0$, adding them gives $v_a + v_{a'} \ge 0$; but the midpoint of aa' lies in K at level λ_m, so its height is at most that of p_{top}, i.e. $v_a + v_{a'} \le 0$. Hence $v_a + v_{a'} = 0$, the midpoint of aa' is p_{top}, and then both inequalities are equalities, so a, a' lie on the supporting line at p_{top}; now $p_{\mathrm{bot}} = p_{\mathrm{top}} - \mu e^\perp$ with $\mu > 0$ has $n_{\mathrm{top}} \cdot (p_{\mathrm{bot}} - p_{\mathrm{top}}) = -\mu\cos\varphi > 0$, so it violates the supporting half-plane. Therefore $\cos\varphi > 0$, and the inequalities give $|\tan\varphi| \le 2W/L$. The same holds at p_{bot} with $-e^\perp$ in place of $e^\perp$.

By (b) above, every boundary point x of K lies on some $\mathrm{C}(w,\Delta_2)$ with $w\in X$, and then $(x-w)/\Delta_2$ is an outer normal of K at x (as $K\subseteq \mathrm{B}(w,\Delta_2)$). Choose such w_{top} for p_{top} and w_{bot} for p_{bot}, and use these normals above. Then, with $\tan\varphi^* := 2W/L$,

$$(w_{\mathrm{bot}}-w_{\mathrm{top}})\cdot e^\perp = (p_{\mathrm{bot}}-p_{\mathrm{top}})\cdot e^\perp + \Delta_2(n_{\mathrm{top}}\cdot e^\perp - n_{\mathrm{bot}}\cdot e^\perp) \geq -W + 2\Delta_2\cos\varphi^*,$$

while $|w_{\mathrm{bot}}-w_{\mathrm{top}}|\leq \Delta \leq 1.94\,\Delta_2$. Suppose $L\geq 400$. Then $\tan\varphi^*\leq 2w_y/400 + 4\tan(\theta/2)$ and $W\leq w_y + 4\Delta_2\tan(\theta/2)$, so

$$\frac{2\Delta_2\cos\varphi^* - W}{\Delta_2} \geq 2\Big(1-\frac{\tan^2\varphi^*}{2}\Big) - 4\tan(\theta/2) - \frac{w_y}{10^4} > 1.96 > 1.94,$$ [N]

a contradiction. Hence $L<400$ and $W\leq w_y + 800\tan(0.5^\circ) < 9$. So K lies in a rectangle of size less than 400×9, and the disjoint discs of radius $\frac12$ about the points of $X\setminus D$ lie in a rectangle of area less than $401\cdot 10$; so $|X\setminus D| < 4010/(\pi/4) < 5106$. □

The argument uses only boundary points of K and centres in X, so corners of K and boundary points that are not in X are covered.

6.3. **Big turning.**

Lemma 6.4. *Let $\varepsilon_0>0$. The number of $y\in R\setminus D$ with $h\leq 1$ and $\mathrm{turn}(\gamma_y)>\varepsilon_0$ is less than $2\pi M/\varepsilon_0$, where $M=468^2$. For $\varepsilon_0=1/50$ this is less than $6.89\cdot 10^7$.*

Proof. If $b\in\gamma_y$ then $|b-p_0|\leq 232$ (a chord is at most the arc), so $|b-y|\leq 233$. The points y are at least 1 apart, so for fixed b at most $(2\cdot 233.5)^2\leq 468^2$ of them qualify (disjoint discs of radius $\frac12$ in $\mathrm{B}(b,233.5)$). Apply Lemma 2.5 to the arcs γ_y, indexed by y.[N] □

6.4. **The flat case.**

Lemma 6.5 (flat case)**.** *Suppose $\partial K\cap \mathrm{B}(y,1)\subseteq\gamma_y$ and $\mathrm{turn}(\gamma_y)\leq\varepsilon_0$ with $0<\varepsilon_0\leq 1/50$, and put $\varepsilon := \tan\varepsilon_0$. Use coordinates with origin p_0, first axis a unit tangent $t_0\perp n_0$ and second axis n_0; so $y=(0,-h)$. Then there is a concave function f on $[-231,231]$ with $f(0)=0$, $f\leq 0$ and $|f(x)|\leq\varepsilon|x|$, whose graph is contained in $\gamma_y\subseteq\partial K$, and $U := \mathrm{B}(y,1)\cap\{(x,z): z<f(x)\}\subseteq \operatorname{int}K$. Write a neighbour q of y as $q=y+(\sin\psi,\cos\psi)$, with ψ measured from n_0. Then:*

- (F1) *every neighbour $q\notin\operatorname{int}K$ (in particular every neighbour in S, and every neighbour in D) has $\cos\psi\geq h+f(\sin\psi)\geq h-\varepsilon|\sin\psi|$; hence $\cos\psi>-\varepsilon|\sin\psi|$, i.e. $|\psi|<\pi/2+\varepsilon_0$;*
- (F2) *every neighbour $q\in S\setminus D$ has $\cos\psi = h+f(\sin\psi)\in[h-\varepsilon,h]$;*
- (F3) *every neighbour $q\in D$ (in S or in R) has $\cos\psi\geq h+t+f(\sin\psi)\geq h+t-\varepsilon$;*
- (F4) *every neighbour $q\in\operatorname{int}K$ has $\cos\psi<h$.*

Proof. The normal set $N(\gamma_y)$ is an arc of S^1 of length at most ε_0 containing n_0. Parametrise γ_y by arc length $\lambda\in[-232,232]$ with $\gamma(0)=p_0$. Every one-sided tangent is perpendicular to a normal in $N(\gamma_y)$, hence within ε_0 of $\pm t_0$, and after orienting, $\gamma'(\lambda)\cdot t_0\geq\cos\varepsilon_0>0$. So $x(\lambda):=\gamma(\lambda)\cdot t_0$ is strictly increasing with $|x(\pm 232)|\geq 232\cos\varepsilon_0>231$, and γ_y contains the graph of a function f on $[-231,231]$ with $|f'|\leq\varepsilon$ almost everywhere; as $f(0)=0$, $|f(x)|\leq\varepsilon|x|$. Also $f\leq 0$ because $K\subseteq H_0$. At each graph point the outer normals have n_0-component at least $\cos\varepsilon_0>0$, and K lies in the corresponding supporting half-plane; so K lies below every supporting line of the graph, and f is concave. The set U is convex (a disc intersected with the strict hypograph of a concave function; the x-range of $\mathrm{B}(y,1)$ is $[-1,1]$), it contains y (as $-h<0=f(0)$), and $U\cap\partial K=\emptyset$, since $\partial K\cap\mathrm{B}(y,1)\subseteq\gamma_y$ and, x being monotone along γ_y, the points of $\gamma_y\cap\mathrm{B}(y,1)$ lie on the graph, where $z=f(x)$. A connected set meeting $\operatorname{int}K$ and missing ∂K lies in $\operatorname{int}K$; so $U\subseteq\operatorname{int}K$.

(F1) If $q\notin\operatorname{int}K$ then $q\notin U$, so $q_z\geq f(q_x)$, i.e. $\cos\psi-h\geq f(\sin\psi)$; and $h>0$. The inequality $\cos\psi+\tan\varepsilon_0|\sin\psi|>0$ is equivalent to $\cos(|\psi|-\varepsilon_0)>0$, i.e. $|\psi|<\pi/2+\varepsilon_0$. (F2) A neighbour

in $S \setminus D$ lies on $\partial K \cap \mathrm{B}(y,1)$, hence on the graph. (F3) The graph point $g = (\sin\psi, f(\sin\psi))$ lies in K and, by (F1), directly below q; so $t \le \operatorname{dist}(q,K) \le |q-g| = \cos\psi - h - f(\sin\psi)$. (F4) A neighbour in $\operatorname{int} K$ is not on ∂K. If it had $\cos\psi \ge h$, it would satisfy $z \ge 0 \ge f(x)$; points strictly above a graph point are outside K (which lies below the supporting lines), and points on the graph are on ∂K. So $\cos\psi - h < f(\sin\psi) \le 0$. □

By (b) above, every neighbour of y lies in $\operatorname{int} K$, in $S \setminus D$, or in D.

6.5. **Proofs of Theorems 6.1 and 6.2.**

Proof of Theorem 6.1. Take $\varepsilon_0 = 1/50$. Points of $R \cap D$ have $\deg y + m_y \le 6$, and points of $R \setminus D$ with $h > 1$ have $m_y = 0$. Let $y \in R \setminus D$ with $h \le 1$. If the perimeter of K is less than 464, there are at most 465^2 such points in total. If some point of $\partial K \cap \mathrm{B}(y,1)$ is not on γ_y, then Lemma 6.3 applies, and there are fewer than 5106 such points in total. If $\operatorname{turn}(\gamma_y) > \varepsilon_0$, then y is one of fewer than $6.89 \cdot 10^7$ points (Lemma 6.4). Otherwise Lemma 6.5 applies: by (F1) every neighbour of y in S lies in the open arc $|\psi| < \pi/2 + \varepsilon_0$, of length less than $4\pi/3$. Distinct neighbours are at least $60°$ apart as seen from y, and an open arc of length less than $240°$ contains at most 4 such points. So $m_y \le 4$. The exceptional points number fewer than $6.89 \cdot 10^7 + 5106 + 465^2 < 7.1 \cdot 10^7$. □

Proof of Theorem 6.2. The exceptional points are as in the proof of Theorem 6.1; so let y be in the flat case of Lemma 6.5, with $h \le 1$. Note $t > \sqrt{3}/2 + \varepsilon > \frac{1}{2} + 2\varepsilon$.

(i) *At most one neighbour in D.* By (F3) a neighbour in D has $\cos\psi \ge t - \varepsilon > \sqrt{3}/2$, i.e. $|\psi| < 30°$; two neighbours at angular distance at least $60°$ cannot both lie in this open arc.

(ii) *At most one neighbour in $S \setminus D$ with $\psi \in [0,\pi)$, and at most one with $\psi \in (-\pi, 0]$.* If $0 \le \psi_1 < \psi_2$ with $\psi_2 - \psi_1 \ge 60°$ are both in $S \setminus D$, then $\psi_2 < 90° + \varepsilon_0 < 91.2°$ by (F1), so $(\psi_1 + \psi_2)/2 \in [30°, 61.2°)$ and $(\psi_2 - \psi_1)/2 \in [30°, 45.6°)$. Hence $\cos\psi_1 - \cos\psi_2 = 2\sin\frac{\psi_1+\psi_2}{2}\sin\frac{\psi_2-\psi_1}{2} \ge \frac{1}{2} > \varepsilon$, contradicting (F2), by which both cosines lie in $[h-\varepsilon, h]$.

Every neighbour in S lies in $S \setminus D$ or in D, so $m_y \le 3$; and if $m_y = 3$, the neighbours in S are one point of D at some ψ_D with $|\psi_D| < 30°$ and two points $K_\pm$ of $S \setminus D$.

(iii) *$\deg y = 6$ and $m_y = 3$ is impossible.* Six neighbours pairwise at least $60°$ apart are exactly $60°$ apart, at $\psi_D + 60°k$. So $K_\pm$ occupy two of the five directions $\psi_D \pm 60°$, $\psi_D \pm 120°$, $\psi_D + 180°$, and by (F2) their cosines differ by at most ε. We go through the ten pairs.

- A pair containing $\psi_D + 180°$ (four pairs): $\cos(\psi_D + 180°) = -\cos\psi_D < -\sqrt{3}/2 < -\varepsilon$, contradicting (F1).
- $\{\psi_D + 60°, \psi_D - 60°\}$: $|\cos(\psi_D + 60°) - \cos(\psi_D - 60°)| = \sqrt{3}|\sin\psi_D| \le \varepsilon$, so $\cos(\psi_D \pm 60°) \ge \frac{1}{2}\cos\psi_D - \frac{\sqrt{3}}{2}|\sin\psi_D| \ge \frac{1}{2}(1 - \varepsilon^2/3) - \frac{\varepsilon}{2} \ge \frac{1}{2} - \varepsilon$. By (F2), $h \ge \frac{1}{2} - \varepsilon$, and then (F3) gives $\cos\psi_D \ge h + t - \varepsilon \ge \frac{1}{2} + t - 2\varepsilon > 1$, a contradiction.
- $\{\psi_D + 120°, \psi_D - 60°\}$ (and symmetrically $\{\psi_D - 120°, \psi_D + 60°\}$): the two cosines are x and $-x$ with $x = \cos(\psi_D + 120°)$, and both lie in $[h-\varepsilon, h]$, so $|x| \le \varepsilon/2$ and $h \le \varepsilon$. As $\psi_D + 120° \in (90°, 150°)$, this forces $\psi_D + 120° \le 90° + \arcsin(\varepsilon/2)$, so $\psi_D + 60° \in (30°, 31.2°]$. The neighbour at $\psi_D + 60°$ is not in S (as $m_y = 3$), and $\cos(\psi_D + 60°) > \frac{1}{2} > h$, so by (F4) it is not in $\operatorname{int} K$; hence it lies in D, and by (F3) it has $|\psi| < 30°$, a contradiction.
- $\{\psi_D + 120°, \psi_D - 120°\}$: both directions have $|\psi| > 90°$, so their cosines are negative and $h < \varepsilon$ by (F2). The neighbours at $\psi_D \pm 60°$ are not in S, and one of them has $30° < |\psi| \le 60°$, so $\cos\psi \ge \frac{1}{2} > h$; by (F4) it is not in $\operatorname{int} K$, so it lies in D, and (F3) gives $|\psi| < 30°$, a contradiction.
- The two pairs on one side, $\{\psi_D + 60°, \psi_D + 120°\}$ and $\{\psi_D - 60°, \psi_D - 120°\}$, are excluded by (ii).

Hence $\deg y = 6$ implies $m_y \le 2$, and $\deg y \le 5$ implies $\deg y + m_y \le 8$. For $\varepsilon_0 = 1/50$ the exceptional set has at most $6.89 \cdot 10^7 + 5106 + 465^2 < 7.1 \cdot 10^7$ points. □

Remark 6.6 (regimes). Theorem 6.1 uses nothing about rungs or τ, so it holds in Regions I, II, III, including $\tau = \sqrt{3}/2$, $\tau \le 0$ and $\tau > 1$. Theorem 6.2 needs $t > \sqrt{3}/2 + \varepsilon$ with $\varepsilon > 0$ fixed; in Region III with $\tau \ge \sqrt{3}/2 + \eta$, $0 < \eta \le 1/50$, it holds with $\varepsilon_0 = \arctan(\eta/2)$, but then the exceptional count grows like $1/\eta$ and is not uniform as $\eta \to 0$. For $\tau \le \sqrt{3}/2$ we only have Theorem 6.1.

Remark 6.7 (local sharpness of Theorem 6.1 for $\tau < \sqrt{3}/2$). Consider a single disc, $K = \mathrm{B}(w, \Delta_2)$ near y, with $|y-w| = \Delta_2 - h$. A neighbour at angle ψ from the outward radial direction satisfies $|q-w|^2 = (\Delta_2-h)^2+1+2(\Delta_2-h)\cos\psi$. So q lies on ∂K if and only if $\cos\psi = c_K(h) := h+(h^2-1)/(2(\Delta_2-h))$, and $\operatorname{dist}(q,K) \ge t$ if and only if $\cos\psi \ge c_D(h) := h+t+((h+t)^2-1)/(2(\Delta_2-h))$; moreover $c_D(0) = \tau$ and c_D is increasing in h.[N] Two such D-feasible neighbours 60° apart need $c_D(h) \le \cos 30^\circ$, so this local picture requires $\tau < \sqrt{3}/2$. For $\tau < \sqrt{3}/2$ it occurs: with $w = 0$ and $\Delta_2 = 10^4$, take on $\mathrm{C}(w, \Delta_2)$ chords $[p_-, p_+]$ of length 2 (every other chord position), let y be the chord midpoint, $e := (p_+ - p_-)/2$, $\nu := y/|y|$, $z_\pm := y \pm e/2 + (\sqrt{3}/2)\nu$ and $a_\pm := y \pm e/2 - (\sqrt{3}/2)\nu$. In 60-digit decimal arithmetic (script `r4_nb_exact.py`) one checks that $|yp_\pm| = |yz_\pm| = |ya_\pm| = 1$, all pairwise distances are at least 1, the $z_\pm$ lie on $\mathrm{C}(w, \Delta_2 + t)$ with $t = 0.86598790\ldots$ and $\tau = \sqrt{3}/2 - 5.0 \cdot 10^{-5}$, $\operatorname{dist}(z_\pm, \mathrm{B}(w,\Delta_2)) = t$, and $a_\pm \in \operatorname{int}\mathrm{B}(w,\Delta_2)$. So every such y has degree 6, two neighbours on ∂K and two D-feasible neighbours, i.e. $\deg y + m_y = 10$ as soon as $z_\pm \in S$. In this one-centre configuration the $z_\pm$ have no Δ_2-partner; whether $\Theta(n)$ such points y with $z_\pm \in S$ can occur in a global configuration is open. So local arguments of the kind above cannot improve the constant 10 when $\tau < \sqrt{3}/2$.

7. The degenerate regime $\Delta > 1.94\Delta_2$

Proposition 7.1. *If $\Delta > 1.94\,\Delta_2$ and $n \ge 4$, then $\mu(\Delta_2) \le n + 1$.*

Put $\kappa := (\Delta_2/\Delta)^2 < 1/1.94^2$. Note that this section does not use δ.

Step 1: no two disjoint diametral pairs. Suppose (x, x') and (z, z') are disjoint diametral pairs with midpoints m, m'. The four cross distances satisfy $\sum |ab|^2 = 2\Delta^2 + 4|m - m'|^2 \ge 2\Delta^2$, and each is either at most Δ_2 or equal to Δ. If at most one equals Δ, the sum is at most $\Delta^2(1 + 3\kappa) < 2\Delta^2$,[C10] a contradiction. So at least two cross distances equal Δ.

- If two disjoint cross pairs equal Δ, say $|xz| = |x'z'| = \Delta$, then z and x' are the two points of $\mathrm{C}(x,\Delta) \cap \mathrm{C}(z',\Delta)$, so x, z, z', x' form a rhombus of side Δ whose diagonals satisfy $d_1^2 + d_2^2 = 4\Delta^2$. One diagonal is at least $\sqrt{2}\Delta$, a contradiction.
- If two cross pairs sharing a point equal Δ, say $|xz| = |xz'| = \Delta$, then xzz' is an equilateral triangle of side Δ, and $x' \in \mathrm{C}(x,\Delta)$ with $|x'z|, |x'z'| \le \Delta$ lies on the 60° arc of $\mathrm{C}(x,\Delta)$ between z and z', strictly inside it. One of $|x'z|, |x'z'|$ is at least $2\Delta\sin 15^\circ$, and $(2\sin 15^\circ)^2 = 2 - \sqrt{3} > \kappa$,[C10] so that distance lies in (Δ_2, Δ), which is impossible.

Step 2: the diameter graph is a star. A graph with no two disjoint edges is a star or a triangle. A triangle is impossible for $n \ge 4$: a fourth point is not diametral, so it is within Δ_2 of all three vertices, but $\Delta_2^2 < \Delta^2/3$ is less than the squared circumradius.[C10] So $D = \{y\} \cup A$ with A the set of diametral partners of y; put $B := X \setminus D \subseteq K$ and $W := \{y\} \cup A$.

Step 3: where the Δ_2-pairs are.

- *Within B: none.* For $a \in A$, $K \subseteq \mathrm{B}(y,\Delta_2) \cap \mathrm{B}(a,\Delta_2)$, a lens whose bounding box has squared diagonal less than $0.24\,\Delta_2^2$.[C10] So $\operatorname{diam} K < \Delta_2$.
- *Within A: at most one.* $A \subseteq \mathrm{C}(y,\Delta)$, and its pairwise distances are at most Δ_2 (a Δ-pair inside A would create a triangle in the diameter graph). A chord of $\mathrm{C}(y,\Delta)$ of length at

most $\Delta_2 < \Delta/1.94$ subtends an angle less than 30° at y. So A lies in an arc of angular width less than 60°, on which chord length is increasing in angular separation; a Δ_2-pair in A must realise the largest pairwise distance, hence be the two extreme points of A.

- *Between y and A: none*, as those distances equal Δ.

Hence $\mu(\Delta_2) \le 1 + I$, where $I := \#\{(b, w) : b \in B,\ w \in W,\ |bw| = \Delta_2\}$.

Step 4: $I \le |B| + |W|$. We may assume K has nonempty interior: an open chord between two points of K lies in the interior of every disc $\mathrm{B}(w, \Delta_2)$, so otherwise $|B| \le |K \cap X| \le 1$ and the bound is trivial. For $w \in X$ let $J_w := \mathrm{C}(w, \Delta_2) \cap K$.

- *Each J_w is a single closed arc (possibly a point).* For $x \in X \setminus \{w\}$, $\mathrm{C}(w, \Delta_2) \cap \mathrm{B}(x, \Delta_2)$ is the closed arc $\{\cos(\theta - \theta_x) \ge |wx|/(2\Delta_2)\}$, of angular length at most π. For $p, q \in J_w$ we have $|pq| < \Delta_2 < 2\Delta_2$, so p, q are not antipodal and the minor arc of $\mathrm{C}(w, \Delta_2)$ from p to q lies in every $\mathrm{B}(x, \Delta_2)$, hence in K. Since $\operatorname{diam} K < \Delta_2$, the angular spread of J_w is less than 60°, so J_w is a single closed arc.
- *No point is interior to two arcs.* Suppose b is a relative-interior point of a nondegenerate arc J_w and $b \in \mathrm{C}(w', \Delta_2)$ with $w' \in X \setminus \{w\}$. The function $f(\theta) = |w + \Delta_2 e(\theta) - w'|^2 - \Delta_2^2 = |w - w'|^2 + 2\Delta_2\, e(\theta) \cdot (w - w')$ is ≤ 0 on J_w (as $K \subseteq \mathrm{B}(w', \Delta_2)$) and vanishes at b, so it has an interior local maximum with value 0. But every local maximum of this shifted cosine is its global maximum $|w - w'|^2 + 2\Delta_2|w - w'| > 0$, a contradiction.
- *Consequences.* Each nondegenerate J_w is a subarc of ∂K, and distinct nondegenerate arcs have disjoint relative interiors. So a point $b \in B$ lies on at most two nondegenerate W-arcs (one on each side); any further incidences of b come from degenerate arcs $J_w = \{b\}$.
- *Counting.* Let I_{nd} be the number of incidences from nondegenerate arcs. Then $I_{\mathrm{nd}} \le |B| + \#\{b : \text{two } W\text{-arcs meet at } b\}$. Mapping each such b to the W-arc that starts at b in the clockwise direction is injective, so $I_{\mathrm{nd}} \le |B| + \#\{w : J_w \text{ nondegenerate}\}$. Each degenerate arc contributes at most one incidence. Hence $I \le |B| + |W|$.

Therefore $\mu(\Delta_2) \le 1 + |B| + |A| + 1 = n + 1$, proving Proposition 7.1.

8. Assembly of the proof of Theorem 1

regime ($\delta = 1$, $\beta = 1/100$)	$e(S) \le$	$\deg y + m_y \le$	$\min\{\mu(\Delta_2), \mu(\delta)\} \le$
degenerate: $\Delta > 1.94\Delta_2$, $n \ge 4$	—	—	$n + 1$ (Proposition 7.1)
nondegenerate, $\Delta_2 < 10^4$	—	—	$3n < 3N_0$ (Section 3)
I: $\tau \le \sqrt{3}/2 - 2\beta$ (incl. $\tau \le 0$)	$s + C$	10 (Theorem 6.1)	$\frac{15}{11}n + C$
III: $\sqrt{3}/2 - 2\beta < \tau < \sqrt{3}/2 + 3\beta$	$s + p + C$	10 (Theorem 6.1)	$\frac{15}{11}n + C$
II: $\sqrt{3}/2 + 3\beta \le \tau \le 1$	$s + p + C$	8 (Theorem 6.2)	$\frac{4}{3}n + C$
II: $\tau > 1$ ($\Leftrightarrow t > 1$)	$s + C$	8 (Theorem 6.2)	$\frac{4}{3}n + C$

Table 1. The regimes in the proof of Theorem 1. In the last column $C < 2 \cdot 10^8$; the R-bounds hold for all but fewer than $7.1 \cdot 10^7$ points of R.

The τ-intervals partition $\mathbb{R}$, the split at $\Delta = 1.94\,\Delta_2$ is exhaustive, and the cases $n \le 3$ (or X with a single distance) are trivial.

Constants. By Proposition 4.3, $C_{\mathrm{bad}} < 1.83 \cdot 10^7$. By Section 5, in every τ-regime of the nondegenerate regime with $\Delta_2 \ge 10^4$,

$$e(S) \le s + p + C_e, \qquad C_e := 2C_{\mathrm{bad}} + 81 < 3.7 \cdot 10^7$$

(Region I and $\tau > 1$: C_{bad}; Region II: $C_{\text{bad}}+5$; Region III: $C_{\text{bad}}+6$ for $\tau \neq \sqrt{3}/2$ and $2C_{\text{bad}}+81$ for $\tau = \sqrt{3}/2$). By Theorem 6.1, $\deg y + m_y \leq 10$ for all but $C' < 7.1 \cdot 10^7$ points $y \in R$, and in Region II, by Theorem 6.2, $\deg y + m_y \leq 8$ for all but $C' < 7.1 \cdot 10^7$ points. Lemma 3.5 gives

$$\min\{\mu(\Delta_2), \mu(\delta)\} \leq \tfrac{15}{11}n + C_e + C' \quad \text{in general, and}$$
$$\min\{\mu(\Delta_2), \mu(\delta)\} \leq \tfrac{4}{3}n + C_e + C' \quad \text{in Region II,}$$

and $C_e + C' < 1.1 \cdot 10^8 < 2 \cdot 10^8$. The degenerate regime gives $n + 1 \leq \frac{4}{3}n + 1$, and the small cases give at most $3N_0$. So Theorem 1 holds with $C_0 = 3N_0 + 2 \cdot 10^8$. Part (b) is the Region II row: its hypotheses are $\Delta \leq 1.94\,\Delta_2$ and $\tau \geq \sqrt{3}/2 + 3\beta$, and if $\Delta_2 < 10^4$ the small-case bound applies. □

Remark 8.1. (a) *The bottleneck is the R-term.* The estimate $e(S) \leq s + p + O(1)$ is sharp enough for $\frac{4}{3}$ (Remark 3.6); the value $\frac{15}{11}$ comes only from the bound $\deg y + m_y \leq 10$ of Theorem 6.1 in Regions I and III. By Remark 6.7, this bound cannot be improved by local arguments when $\tau < \sqrt{3}/2$, because the exact local configuration there has $\deg y + m_y = 10$; whether it can occur for $\Theta(n)$ points in a global configuration is open.

(b) *What remains for* 9/7. If $\deg y + m_y \leq 6$ held for all but $o(n)$ points of R, Region I would give $\frac{9}{7}$, but Regions II and III would still give only $\frac{4}{3}$ because of the p-term (Remark 3.6). So reaching $\frac{9}{7}$ with this method would also require sharpening the Δ_2 side in Regions II and III.

(c) *Consistency with the lower bound.* The construction of ienjoymath [17] for $L \geq 9/7$ lies in Region I ($t/\delta \to 0$). In it $e(S) = |S|$, every point of S has Δ_2-degree 2 or 4 (so $Q = \emptyset$ and $\mu(\Delta_2) = \frac{3}{2}|S|$), and every point of R has $m_y = 0$ and $\deg y + m_y \leq 6$; we checked the last statement numerically for $n = 60, 120, 250$ (Appendix A). This is consistent with Theorem 6.1, which is only an upper bound; in Region I the gap between $\frac{9}{7}$ and $\frac{15}{11}$ is exactly the gap between 6 and 10 in the R-term.

9. A layer bound and a new class of sets

In this section X is an arbitrary finite planar set with at least two distinct distances, $\Delta > \Delta_2$ are its two largest distances, and we scale so that $\Delta_2 = 1$. Every distance of X is at most 1 or equal to Δ. L_1 is the set of extreme points of X (vertices of $\operatorname{conv} X$), L_2 is the set of extreme points of $X \setminus L_1$, and $m_3 := n - |L_1| - |L_2|$. Let G be the graph on $L_1 \cup L_2$ whose edges are the Δ_2-pairs, let G' be its 2-core (repeatedly delete vertices of degree less than 2), and write $L_i' := L_i \cap V(G')$ and $\deg' = \deg_{G'}$. For $p \in X$ let $h(p)$ be the number of points of X at distance Δ_2 from p.

Lemma 9.1 (location)**.** *(i) Both endpoints of every Δ-pair lie in L_1.*
(ii) Every Δ_2-pair has an endpoint in L_1 and both endpoints in $L_1 \cup L_2$.
In particular G contains every Δ_2-pair of X.

Proof. (i) If $|pq| = \Delta$ then $X \subseteq \mathrm{B}(q, \Delta)$ and p lies on its boundary circle, so p is an extreme point of X. (ii) Let $|pq| = 1$. If $p \notin L_1$, then p has no Δ-partner by (i), so $X \subseteq \mathrm{B}(p, 1)$ and q, lying on the boundary circle, is in L_1. So some endpoint, say p, lies in L_1. Suppose $q \notin L_1$. Every point x with $|px| > 1$ has $|px| = \Delta$ and so lies in L_1. Hence $X \setminus L_1 \subseteq \mathrm{B}(p, 1)$ with q on the boundary circle, so q is extreme in $X \setminus L_1$, i.e. $q \in L_2$. □

Proposition 9.2 (core degree of L_2)**.** *Every $q \in L_2'$ has $\deg'(q) = 2$.*

Proof. Put $q = 0$. Since q is not extreme it has no Δ-partner (Lemma 9.1(i)), so $X \subseteq \mathrm{B}(0, 1)$ and the G-neighbours of q lie on the unit circle. Since $q \in V(G')$, $\deg'(q) \geq 2$. Suppose v_1, v_2, v_3 are three distinct G'-neighbours of q. Each $|v_i v_j|$ is either at most 1 or equal to Δ. The argument

follows the three cases of Vesztergombi's inner-point analysis [23, pp. 192–194], with one change: we delete no points, and the contradiction in Cases 1 and 2 is with membership in the 2-core, via $\deg_G \geq \deg'$.

Case 3: at most one pair v_iv_j at distance Δ. Label so that $|v_1v_2|, |v_2v_3| \leq 1$. Then v_2 lies on the shorter arc between v_1 and v_3, and in polar angle we may take v_2 at 0, v_1 at a and v_3 at $-b$ with $0 < a, b \leq 60°$. Since q is not extreme, the closed half-plane $\{x : x \cdot v_2 \leq 0\}$ contains a point $t \in X$, $t \neq q$. For $t \neq 0$ with $t \cdot v_2 \leq 0$ we get $|t - v_2|^2 = |t|^2 - 2t \cdot v_2 + 1 > 1$, so $|tv_2| = \Delta$. By symmetry assume the polar angle φ of t lies in $[90°, 180°]$. Then the angle between t and v_3 is $\varphi + b \in (90°, 240°]$, so $t \cdot v_3 < 0$ and $|tv_3| = \Delta$. Hence t lies on the perpendicular bisector of v_2v_3, which passes through 0, so $\varphi = 180° - b/2$. The angle between t and v_1 is then $180° - b/2 - a \geq 90°$, so $|tv_1|^2 \geq |t|^2 + 1 > 1$ and $|tv_1| = \Delta$. Now v_1, v_2, v_3 lie on both $\mathrm{C}(0, 1)$ and $\mathrm{C}(t, \Delta)$, two distinct circles, which meet in at most two points. Contradiction. (This case uses only that q is not extreme.)

Case 2: exactly two pairs at Δ. Say $|v_1v_2| = |v_1v_3| = \Delta$ and $|v_2v_3| \leq 1$. Then v_1 lies on the perpendicular bisector of v_2v_3, and we may take $v_1 = (-1, 0)$, $v_{2,3} = (\cos a, \pm \sin a)$ with $0 < a \leq 30°$ and $\Delta = 2\cos(a/2)$. We claim v_1 has no Δ_2-neighbour $r \neq q$; then $\deg_G(v_1) = 1$, contradicting $v_1 \in V(G')$. Such an $r = (x, y)$ satisfies $|r - v_1| = 1$ (so $|r|^2 = -2x$), $|r| \leq 1$, and $|r - v_i| \in [0, 1] \cup \{\Delta\}$ for $i = 2, 3$; by symmetry $y \geq 0$. Write $c = \cos a$, $s_a = \sin a$.

- *Both $|r - v_2|, |r - v_3| \leq 1$.* Impossible: $|r - v_3|^2 - 1 = -2x(1 + c) + 2ys_a \geq 0$, with equality only if $r = q$.
- $|r - v_2| = \Delta$ (the case $|r - v_3| = \Delta$ is symmetric). Using $|r|^2 = -2x$, this condition is the linear equation $-2x(1 + c) - 2s_ay = 1 + 2c$. Intersecting with the circle $|r - v_1| = 1$ gives $x = (\pm s_a\sqrt{8c + 7} - 3(1 + c))/(4(1 + c))$, so $|r|^2 = (3(1 + c) \mp s_a\sqrt{8c + 7})/(2(1 + c))$. The root with $+s_a\sqrt{8c + 7}$ in $|r|^2$ has $|r|^2 \geq 3/2$, outside the disc. For the other root, $|r|^2 \leq 1 \iff 1 + c \leq s_a\sqrt{8c + 7} \iff 1 + c \leq (1 - c)(8c + 7) \iff c^2 \leq \frac{3}{4} \iff a \geq 30°$. So only $a = 30°$ remains, and there $|r - v_3|^2 = 3 + c - s_a\sqrt{8c + 7} = 2$, which lies strictly between 1 and $\Delta^2 = 2 + \sqrt{3}$. Forbidden.
- *Both equal Δ.* Then r lies on the x-axis, so $r \in \{(0, 0), (-2, 0)\}$, and $(-2, 0)$ is outside the disc.

Case 1: all three pairs at Δ. Then v_1, v_2, v_3 form an equilateral triangle with circumcentre q, and $\Delta = \sqrt{3}$. Let $P \neq q$ be a further Δ_2-neighbour of some v_i, so $|P - v_i| = 1$ and $|P| \leq 1$. It cannot have both other distances at most 1: $|P - v_i| = 1$ gives $|P|^2 = 2P \cdot v_i$, the two conditions give $P \cdot (v_i - v_j) \leq 0$ for $j \neq i$, and adding and using $\sum_j v_j = 0$ gives $3P \cdot v_i \leq 0$, so $P = q$. Hence $|P - v_j| = \sqrt{3}$ for some $j \neq i$. Solving exactly, there are six candidate points, two for each v_i, each with $|P|^2 = (9 - \sqrt{33})/6$ and at squared distance $(7 - \sqrt{33})/2 \approx 0.63$ from the third vertex, so each is a Δ_2-neighbour of exactly one v_i. Their pairwise squared distances are $(29 - 5\sqrt{33})/6 \approx 0.046$ for three disjoint pairs (each joining candidates of different vertices), and otherwise $4/3$, $(9 - \sqrt{33})/2 \approx 1.628$ or $(17 - \sqrt{33})/6 \approx 1.876$, all strictly inside the forbidden interval $(1, 3)$. So a set of candidates that can coexist in X has at most two elements, at most two of the v_i have a second Δ_2-neighbour, and some v_i has $\deg_G(v_i) = 1$, contradicting $v_i \in V(G')$.

With four or more neighbours apply the argument to any three. Hence $\deg'(q) \leq 2$, so $\deg'(q) = 2$. □

Lemma 9.3 (opposite-side lemma)**.** *Let $|pa| = |pb| = |br| = 1$ with $r \neq p$. Suppose a lies strictly on one side of the line pb and r does not lie strictly on that side. Then $|ar| > 1$, so $|ar| = \Delta$.*

Proof. Put $p = 0$, $b = (1, 0)$, $a = (\cos A, -\sin A)$ with $0 < A < \pi$, and $r = b + (\cos F, \sin F)$ with $0 \leq F \leq \pi$; $F \neq \pi$ as $r \neq p$. Then $|r - a|^2 - 1 = 2(1 + \cos F)(1 - \cos A) + 2\sin F \sin A$, where the first term is positive and the second nonnegative. □

Lemma 9.4. *Let p be an extreme point of X with Δ_2-neighbours $w_1, \dots, w_k$ in angular order. Then $\deg_G(w_i) = 1$ for every $3 \le i \le k-2$.*

Proof. Since p is extreme, some line through p has $X \setminus \{p\}$ strictly on one side. So the directions of the w_j from p are distinct, lie in an open half-plane, and any two differ by less than $180°$; for $j < i < l$, w_j and w_l lie strictly on opposite sides of the line pw_i. Suppose $r \neq p$ and $|rw_i| = 1$. Put $p = 0$, $w_i = (1, 0)$. Then w_{i-2}, w_{i-1} lie strictly below the x-axis and w_{i+1}, w_{i+2} strictly above; by symmetry assume r is not strictly below. Write $w_{i-2} = e(-\alpha_1)$, $w_{i-1} = e(-\alpha_2)$ with $0 < \alpha_2 < \alpha_1 < 180°$. By Lemma 9.3, $|rw_{i-2}| = |rw_{i-1}| = \Delta$, so r lies on the perpendicular bisector of $w_{i-2}w_{i-1}$, the line through 0 in direction $-\gamma$ with $\gamma = (\alpha_1 + \alpha_2)/2 \in (0°, 180°)$. As $r \neq 0$ is not below the axis, $r = \lambda e(180° - \gamma)$ with $\lambda > 0$. From $|r - w_i| = 1$ we get $\lambda(\lambda + 2\cos\gamma) = 0$, so $\lambda = -2\cos\gamma$, whence $\gamma > 90°$ and $\alpha_1 > 90°$. Then $|w_{i-2}w_i| = 2\sin(\alpha_1/2) > \sqrt{2} > 1$, so $|w_{i-2}w_i| = \Delta$. Let $\theta > 0$ be the direction of w_{i+1}; then $\alpha_1 + \theta < 180°$, and since the chord length $2\sin(x/2)$ is strictly increasing on $[0°, 180°]$, $|w_{i-2}w_{i+1}| = 2\sin((\alpha_1 + \theta)/2) > 2\sin(\alpha_1/2) = \Delta$, contradicting the maximality of Δ. □

Theorem 9.5 (degree at most 4)**.** *Every $p \in L_1'$ has $\deg'(p) \le 4$.*

Proof. Suppose $p \in L_1'$ has five G'-neighbours $u_1, \dots, u_5$ in angular order. Among the G-neighbours of p, u_3 has at least two on each side, so $u_3 = w_i$ with $3 \le i \le k-2$, and $\deg_G(u_3) = 1$ by Lemma 9.4. But $u_3 \in V(G')$ forces $\deg_G(u_3) \ge 2$. □

Theorem 9.6 (layer bound)**.** *For every finite $X \subset \mathbb{R}^2$ with at least two distinct distances,*

$$\mu(\Delta_2) \;\le\; |L_1| + |L_2| + \tfrac{1}{2} \sum_{p \in L_1'} (\deg'(p) - 2).$$

Proof. By Lemma 9.1, $\mu(\Delta_2) = e(G)$. Each deletion made while pruning G to its 2-core removes at most one edge, so $e(G) \le e(G') + |V(G) \setminus V(G')|$. By Proposition 9.2, $e(G') = \frac{1}{2}\sum_{v \in V(G')} \deg'(v) = |V(G')| + \frac{1}{2}\sum_{p \in L_1'}(\deg'(p) - 2)$. Add the two estimates and use $|V(G)| = |L_1| + |L_2|$. □

Corollary 9.7. *Assume* (H3)*: no point of L_1 has four or more points of X at distance Δ_2. (This holds, for instance, if no four points of X are concyclic.) Then*

$$\mu(\Delta_2) \le |L_1| + |L_2| + \tfrac{1}{2}\#\{p \in L_1' : \deg'(p) = 3\} \le \tfrac{3}{2}|L_1| + |L_2|.$$

Consequently, under (H3)*, if $|L_1| \le 2m_3$ then Δ and Δ_2 are two distinct distances each occurring at least once and at most n times.*

Proof. Under (H3), $2 \le \deg'(p) \le h(p) \le 3$ for $p \in L_1'$. The condition $\frac{3}{2}|L_1| + |L_2| \le n$ is equivalent to $|L_1| \le 2m_3$, and $\mu(\Delta) \le n$ by [16]. □

Corollary 9.8 (unconditional form)**.** *For every finite X with at least two distances, $\mu(\Delta_2) \le |L_1| + |L_2| + \frac{1}{2}\sum_{p \in L_1}(\min\{h(p), 4\} - 2)^+$. Hence $\mu(\Delta_2) \le n$ whenever $\sum_{p \in L_1}(\min\{h(p), 4\} - 2)^+ \le 2m_3$.*

Proof. Combine Theorem 9.6 with $2 \le \deg'(p) \le \min\{h(p), 4\}$ on L_1' (Theorem 9.5). □

Remark 9.9. (a) Corollary 9.7 covers sets not covered by (1). For example, if $|L_1| = |L_2| = 0.4n$ and $m_3 = 0.2n$, the three quantities in (1) are $1.2n$, $1.33n$ and $1.2n$, while Corollary 9.7 gives $\mu(\Delta_2) \le n$ (under (H3)). Of course (1) needs no hypothesis like (H3).

(b) Using $\deg' \le 4$, Theorem 9.6 recovers the bound $2|L_1| + |L_2|$ of [7, Theorem 1.3]. Proposition 9.2 and Theorem 9.5 are the inputs from [24] (Propositions 3 and 5–8 there, as used in [7]) needed for this; we prove them directly in the 2-core, without deleting points. We have not checked the other inputs used in [7] for the first two bounds in (1).

(c) In Corollary 9.7, the bound $\frac{3}{2}|L_1| + |L_2|$ is attained under (H3) alone by the following example, which we have checked numerically for $m = 8, 12, 16$ (floating point): a regular m-gon with m even, with a point at distance Δ_2 from both endpoints of every other edge (CDL's "servers"). These examples contain concyclic quadruples; for sets with no four concyclic points see Question 11.1 below.

Proposition 9.10 (distinct middle neighbours)**.** *Let $p \neq p'$ be points of X with a common Δ_2-neighbour w, and suppose that each of p, p' has Δ_2-neighbours strictly on both sides of its line to w. Then X contains four concyclic points. In particular, if no four points of X are concyclic, then distinct vertices of* conv X *with three Δ_2-neighbours have distinct middle Δ_2-neighbours.*

Proof. The point p' is not strictly on both sides of the line pw, so choose a Δ_2-neighbour a of p strictly on the side of pw on which p' does not lie strictly. Lemma 9.3 (with $b = w$, $r = p'$) gives $|ap'| = \Delta$. Symmetrically there is a Δ_2-neighbour b of p' with $|bp| = \Delta$. Now $|pb| = |ap'| = \Delta$ and $|pa| = |bp'| = \Delta_2$, so p, b, a, p' are concyclic by the observation in Section 11 (the four points are distinct: $|pa| = \Delta_2 \neq \Delta = |pb|$, and $a \neq p'$, $b \neq p$ since those distances are Δ). For the last sentence: if $p \in L_1$ has exactly three Δ_2-neighbours, they lie in an open half-plane at p, and the middle one in angular order has neighbours of p strictly on both sides of its line to p. □

10. Sets in convex position

A set X is in *convex position* if every point of X is a vertex of conv X; in particular no three points of X are collinear. Throughout this section $|X| = n$, $m := \lfloor n/2 \rfloor$, $N := \binom{n}{2}$, $D = D(X)$ is the number of distinct distances, and $k = k(X)$ is the number of rare distances ($1 \leq \mu(d) \leq n$). A distance is *frequent* if $\mu(d) \geq n + 1$. R_M denotes the vertex set of a regular M-gon. By Hopf–Pannwitz, Δ is always rare; Clemen, Dumitrescu and Liu proved $k \geq 2$ for $n \geq 5$ [7, Theorem 1.2]. For $n = 5$ this is best possible: R_5 has $k = 2$.

Theorem 10.1. *Let X be a set of n points in convex position with $6 \leq n \leq 18$. Then at least three distinct distances each occur at least once and at most n times. The same holds for every even $n \geq 6$.*

The cases have different status:

- $n = 6, 7, 8$: a corollary of published results (Proposition 10.3).
- even $n \geq 10$: Theorem 10.4, a parity-sharp version of the proof of [7, Theorem 1.2];
- $n = 9$: Theorem 10.5, from the classification of Erdős and Fishburn [12];
- $n = 11, 13, 15, 17$: Theorems 10.8 and 10.9, new and computer-assisted. The $n = 7$ case is re-proved the same way, without the literature.

In Section 10.3 we show that sets in convex position with all but w points on one circle have at least $(n - 3w + 1)/4$ rare distances once $n + w \geq 91$.

10.1. Counting and the literature.

Lemma 10.2 (parity-sharp counting)**.** $(D - k)(n + 1) + k \leq N$. *Consequently $k \geq D - m + 1$ whenever $D \geq 1$. More precisely, if $n = 2m + 1$ and $D = m + K$ with $1 \leq K \leq m + 1$, then $k \geq K + 1$; if $n = 2m$ and $D = m + K$ with $1 \leq K < 2m$, then $k \geq K + 2$.*

Proof. The first inequality holds because frequent distances contribute at least $n + 1$ pairs each and rare ones at least 1. It gives $k \geq (D(n + 1) - N)/n = D - \frac{n-1}{2} + \frac{D}{n} > D - m$, hence $k \geq D - m + 1$. For $n = 2m + 1$ the right side is $K + (m + K)/(2m + 1)$, with $0 < (m + K)/(2m + 1) \leq 1$. For $n = 2m$ it is $K + 1 + K/(2m)$, with $0 < K/(2m) < 1$. □

We use the following results from the literature.

(Alt) Altman [1, 2]: $D \geq m$; for odd n, $D = m$ only for R_n (see also [15, Theorem 1]).
(Fi) Fishburn [15]: for even $n \geq 8$, $D = n/2$ if and only if X is R_n or n vertices of R_{n+1}. For $n = 6$ the hexagons with $D = 3$ are R_6, R_7 minus a vertex, and one further hexagon [15, Theorem 2], [10], with multiplicity profiles $(6, 6, 3)$, $(5, 5, 5)$ and $(6, 6, 3)$ respectively (as recorded in [7, §2]; we re-checked the profiles in exact arithmetic).
(EF96) Erdős–Fishburn [12]: a convex nonagon has exactly 5 distances if and only if its vertices are 9 vertices of R_{10} or of R_{11}.
(r_2) Erdős–Fishburn [10, §3], quoting Fishburn's technical report [14]: the largest possible second-largest multiplicity in a convex n-gon is $r_2^{(n)} = 1, 3, 5, 6, 7, 8$ for $n = 3, \ldots, 8$.

A set on a circle has every distance class of maximum degree at most 2, hence of size at most n; so for cocircular X all distances are rare and $k = D$.

Proposition 10.3 (from the literature)**.** *For $n = 6, 7, 8$ in convex position, $k \geq 3$.*

Proof. By (r_2), $r_2^{(n)} \leq n$ for $n \leq 8$, so at most one distance is frequent and $k \geq D - 1$. By (Alt), $D \geq 3$ for $n = 6, 7$ and $D \geq 4$ for $n = 8$. So $k \geq 3$ unless $n \in \{6, 7\}$ and $D = 3$. For $n = 7$, $D = 3$ forces R_7 (profile $(7, 7, 7)$, $k = 3$); for $n = 6$, (Fi) gives the profiles $(6, 6, 3)$ and $(5, 5, 5)$, so $k = 3$. □

The value $r_2^{(7)} = 7$, and the other values of $r_2^{(n)}$, rest on an unrefereed technical report [14] quoted in a refereed paper [10]; for $n = 7$ we give an independent proof in Theorem 10.8.

Theorem 10.4 (even n)**.** *Let X be in convex position with $n = 2m \geq 6$. Then $k \geq 3$. If $n \geq 8$ and $k \leq 3$, then $D = m + 1$.*

Proof. By (Alt), $D \geq m$. If $D \geq m + 1$, Lemma 10.2 gives $k \geq 3$, and $k \geq 4$ if $D \geq m + 2$. If $D = m$ and $n = 6$, (Fi) gives $k = 3$. If $D = m$ and $n \geq 8$, then X is R_n, with profile $(n, \ldots, n, n/2)$, or n vertices of R_{n+1}, with profile $(n-1, \ldots, n-1)$; in both cases $k = m \geq 4$. □

Theorem 10.4 is essentially the argument of [7, Theorem 1.2], with the counting done with parity. It adds to the literature only for even $n \geq 10$.

Theorem 10.5 ($n = 9$)**.** *Every set of 9 points in convex position has $k \geq 3$.*

Proof. Here $m = 4$. If $D = 4$, then $X = R_9$ by (Alt) and $k = 4$. If $D = 5$, then X is cocircular by (EF96) and $k = D = 5$. If $D \geq 6$, Lemma 10.2 gives $k \geq 3$. □

10.2. $n = 7, 11, 13, 15, 17$**: an ordinal reduction and SAT certificates.** Label $X = (p_0, \ldots, p_{n-1})$ in convex cyclic order, let $d_1 < \cdots < d_D$ be the distinct distances, and colour each pair $\{i, j\}$ with $c(ij) = t$ where $|p_i p_j| = d_t$.

Lemma 10.6 (quadrilateral inequality)**.** *For $i < j < k < l$ (in cyclic order), $|p_i p_k| + |p_j p_l| > |p_i p_j| + |p_k p_l|$ and $|p_i p_k| + |p_j p_l| > |p_j p_k| + |p_l p_i|$.*

Proof. The diagonals of the convex quadrilateral $p_i p_j p_k p_l$ meet at a point O that lies on neither of the sides $p_i p_j$, $p_k p_l$. So $|p_i p_k| + |p_j p_l| = (|p_i O| + |O p_j|) + (|p_k O| + |O p_l|) > |p_i p_j| + |p_k p_l|$, by the strict triangle inequality. The other inequality is the same. □

Ordinal form. Let $x_1 \leq x_2$ be the two diagonal colours, and $y_1 \leq y_2$ the colours of one pair of opposite sides. Then it is *not* the case that $x_1 \leq y_1$ and $x_2 \leq y_2$, since otherwise $|p_i p_k| + |p_j p_l| \leq$ the sum of the two sides. We call a colouring of the pairs of an n-cycle's vertex set satisfying this for all 4-tuples and both pairs of opposite sides a *Q-colouring*. This ordinal use of the quadrilateral inequality goes back to Altman [1]. Abstracting a convex distance problem to a finite search in this way is the method of Morić and Pritchard [20], whose Fact 3.1 is Lemma 10.6.

Some of our certificates also use a second valid constraint (*Lemma T*): if $w \neq w'$ lie on the same open boundary arc between p_i and p_j, then $(c(iw), c(jw)) \neq (c(iw'), c(jw'))$, because a point is determined by its distances to p_i, p_j and a side of the line $p_i p_j$.

Lemma 10.7 (reduction to profiles)**.** *If $k \leq 2$, then at least $D-2$ colour classes have size at least $n+1$, and $D \leq D_{\max}(n)$, the largest D with $(D-2)(n+1)+2 \leq N$. For $n = 7, 11, 13, 15, 17$, $D_{\max} = 4, 6, 7, 8, 9$.*

Call a Q-colouring using exactly D colours, all of them, with at least F classes of size at least $n+1$, an (n, D, F) *ordinal counterexample*. Since Lemma 10.6 (and Lemma T) are necessary for geometric realisability, the following suffices to prove $k \geq 3$ for a given n: *for every $2 \leq D \leq D_{\max}(n)$ there is no $(n, D, D-2)$ ordinal counterexample.* (We check the smaller D even where (Alt) would exclude them, so that no citation is needed; for $D \leq 4$ we check the stronger statement with $F = 0$, i.e. no Q-colouring with D colours at all.) No realisation step and no floating-point arithmetic enters.

Theorem 10.8. *Every set of n points in convex position with $n \in \{7, 11, 13\}$ has at least three rare distances. For these n, the ordinal quadrilateral inequality (Lemma 10.6) alone implies the statement.*

Proof (computer-assisted). By Lemma 10.7 it suffices that the following instances have no ordinal counterexample:

- $n = 7$: $(7, 2, 0)$, $(7, 3, 1)$, $(7, 4, 2)$;
- $n = 11$: $(11, 2, 0)$, $(11, 3, 0)$, $(11, 4, 0)$, $(11, 5, 3)$, $(11, 6, 4)$;
- $n = 13$: $(13, 2, 0)$, $(13, 3, 0)$, $(13, 4, 0)$, $(13, 5, 0)$, $(13, 6, 4)$, $(13, 7, 5)$.

Each is encoded in CNF and shown unsatisfiable, with the refutation checked as described below and in Appendix A. □

Encoding. Variables $x_{e,c}$ say that pair e has colour c; there are exactly-one constraints per pair and at-least-one constraints per colour. In the *direct* encoding of Lemma 10.6, each dominated 4-tuple of colours is forbidden. In the *order* encoding, auxiliary literals $y_{e,t} \Leftrightarrow c(e) \geq t$ are introduced and "$c(a) \leq t \leq c(b)$ and $c(a') \leq u \leq c(b')$" is forbidden for both diagonal/side matchings; this is equivalent, since sorted domination holds if and only if some matching dominates. Selector variables f_c imply "class c has at least $n+1$ pairs" through sequential counters, and $\sum_c f_c \geq F$.

Certification.

- *DRAT/LRAT.* For every instance with $n = 11$ and $n = 13$ listed above, and for $(7, 4, 2)$, a DRUP proof produced by Glucose 4 [3] was verified by `drat-trim` [25], which also emitted an LRAT proof that was re-verified by `lrat-check` [8]. The largest certificate, for $(13, 7, 5)$, has 446,686 lemmas (a 37 MB DRUP proof and an 82 MB LRAT proof). For $n = 13$ all instances were certified in the Q-only encoding. For $n = 11$, $(11, 6, 4)$ was certified in the Q-only encoding (35,510 lemmas), and $(11, 2..5, \cdot)$ in the encoding with Q and Lemma T (Q-only is covered by the Lean proof below). A truncated proof is correctly rejected. On this route, the trusted components are the CNF encoder (a short Python program, cross-validated as in Appendix A) and the proof checkers.
- *$n = 7$ without SAT.* A plain backtracking enumeration of all colourings, with Q checked as soon as its pairs are assigned, finds no ordinal counterexample for $D = 2, 3, 4$. For $D = 4$ it does so with Lemma T switched off.
- *Lean 4.* The statements for $n = 7$, 11 and 13 are formalised in Lean 4 [9] with Mathlib [22]; see below.

$n = 15$ and $n = 17$.

Theorem 10.9. *Every set of n points in convex position with $n \in \{15, 17\}$ has at least three rare distances. For these n, too, the ordinal quadrilateral inequality (Lemma 10.6) alone implies the statement.*

Proof (computer-assisted). Here $m = 7, 8$ and $D_{\max} = m+1$ (Lemma 10.7). It suffices to refute, for $n = 15$, the instances $(15, D, 0)$ for $2 \le D \le 6$ and $(15, D, D-2)$ for $D = 7, 8$; and for $n = 17$, the instances $(17, D, 0)$ for $2 \le D \le 7$ and $(17, D, D-2)$ for $D = 8, 9$. Only Lemma 10.6 is encoded (order encoding, without Lemma T). The instances with $F = D - 2$ are split by the pair $\{r_1, r_2\}$ of colours that are *not* required to be frequent: in the case $\{r_1, r_2\}$ every other colour class is required to have at least $n + 1$ pairs, the colours r_1, r_2 are required to be used, and the implied bound $|r_1| + |r_2| \le N - (D-2)(n+1)$ is added as a redundant constraint. Every ordinal counterexample falls under at least one of the $\binom{D}{2}$ cases. All $5 + 21 + 28$ cases for $n = 15$ and all $6 + 28 + 36$ cases for $n = 17$ are unsatisfiable; each refutation was produced by the solver Kissat [5] as a DRAT proof and verified by `drat-trim` [25]. □

Symmetry breaking. For $D \ge m$ most case encodings contain symmetry-breaking constraints, justified as follows. Q-colourings, and the case conditions, are invariant under the dihedral group acting on $\{0, \dots, n-1\}$: reversing the cyclic order maps diagonals to diagonals and pairs of opposite sides to pairs of opposite sides. (Reversing the order of the colours is not a symmetry, and is not used.) Let L^* be the largest cyclic length $\min(j-i,\, n-j+i)$ of a pair of the largest colour D. After a rotation we may assume that $\{0, L^*\}$ has colour D; this is encoded by selector variables s_L with $s_L \Rightarrow c(0, L) = D$ and "no pair of colour D has cyclic length $> L$", and $\bigvee_L s_L$. The reflection $i \mapsto L - i \pmod n$ fixes $\{0, L\}$, preserves cyclic lengths and exchanges the pairs $\{0, L+1\}$ and $\{L, n-1\}$, so we may also require $s_L \Rightarrow c(0, L+1) \le c(L, n-1)$. As a sanity check, all solutions of two satisfiable instances with lowered thresholds were enumerated and grouped into dihedral orbits (392 solutions in 42 orbits for $n = 7$, $D = 4$; 21,348 solutions in 1,280 orbits for $n = 9$, $D = 5$); every orbit meets the symmetry-breaking constraints.

Cost and trust. The hard cases are those whose non-frequent pair contains the diameter colour D: for $n = 17$, $D = 9$ these took 3 to 12 minutes each, and the largest DRAT proof has about 124 MB. The other cases took seconds. Three cases ($(15, 8, \{1, 2\})$, $(15, 7, \{3, 7\})$, $(17, 9, \{2, 3\})$) were re-solved with CaDiCaL [5] and re-verified with `drat-trim`. The same pipeline re-certifies the cases $D = m + 1$ for $n = 7, 9, 11, 13$. On this route the trusted components are the CNF encoder (whose Q part is the order encoding validated in Appendix A), the symmetry-breaking argument above, and `drat-trim`; the solver is not trusted. The proofs are not stored (they are regenerated deterministically by the scripts), and Theorem 10.9 is *not* part of the Lean formalisation.

The Lean formalisation ($n = 7$, 11, 13). The file `Erdos/Erdos/E132Convex.lean` proves, without `sorry`,

```
theorem erdos132_convex7 (p : Fin 7 → EuclideanSpace ℝ (Fin 2)) (hp :
    StrictConvexPolygon p) : 3 ≤ (rareDists p).card
```

and the analogous `erdos132_convex11` and `erdos132_convex13` (file `E132Convex13.lean`) for `Fin 11` and `Fin 13`. Here `StrictConvexPolygon p` means that *every* index-increasing triple (p_i, p_j, p_k), $i < j < k$, is positively oriented; that is, the vertices are given as a strictly convex polygon listed counter-clockwise. `rareDists p` is the set of distances realised by pairs $i < j$ with multiplicity at most n. The proof formalises Lemma 10.6 (via the intersection point of the diagonals and strict convexity of the Euclidean norm), the passage to ranks and Q-colourings, Lemma 10.7, and the correctness of a bit-vector encoding of the finite statement "every Q-colouring with colours below $D_{\max}$ has at least three classes of size in $[1, n]$". This covers

all $D \le D_{\max}$ at once, with unused colours allowed. The finite statement is closed by Lean's `bv_decide`, which bit-blasts the formula with a verified bit-blaster, calls the bundled SAT solver CaDiCaL, and checks the resulting LRAT certificate with a checker that is proved correct in Lean.

Trust base. `#print axioms` reports, for each theorem, the standard axioms `propext`, `Classical.choice`, `Quot.sound`, plus one axiom generated by `bv_decide` (`..._native.bv_decide.ax_1_14`). That axiom asserts that the verified LRAT checker `verifyBVExpr` returns `true` on the concrete formula and certificate. The value `true` is obtained by running compiled code, not by kernel reduction, so the trust base is comparable to that of `native_decide`: it includes the Lean compiler and runtime. Neither our Python encoder, nor any external SAT solver, nor `drat-trim` is trusted on this route. An attempt to check the $n = 7$ axiom statement by kernel reduction ran for more than 20 minutes without finishing and was stopped.

Unordered point sets. The file `E132ConvexFinset.lean` proves (using only the three standard axioms) that a finite set S in convex position, in the sense of Mathlib's `ConvexIndependent`, can be listed as a `StrictConvexPolygon`. The proof sorts $S \setminus \{p_0\}$ by orientation about a point $p_0 \in S$; a negatively oriented increasing triple would put a point of S inside a triangle spanned by others. It also proves that the rare distances of S (unordered pairs) coincide with `rareDists p` for any such listing. This gives `erdos132_convex{7,11,13}_finset`: for `S : Finset` with `S.card` $= n$ and `ConvexIndependent`, $3 \le$ `(rareDistsS S).card`.

The layer bound. The results of Section 9 are formalised in `E132Layer.lean` without `sorry`, and `#print axioms` reports only `propext`, `Classical.choice` and `Quot.sound` (no `bv_decide` or `native_decide`). This covers the localisation of Δ_2-pairs to $L_1 \cup L_2$, the degree-2 property of L_2-vertices in the 2-core, the bound deg ≤ 4 on hull vertices, Theorem A, Corollary A1 and its unconditional form. There L_1 is the set of extreme points of X, shown equal to the vertex set of conv X, and the 2-core is the largest subset with minimum degree at least 2.

The main theorem. The $\frac{54}{37}$ bound for Problem 1.6 of [7], proved in Appendix B, is formalised in full in `E132Main*.lean`, `E132ConvexCurve.lean` and `E132Ve87.lean` (about 7,000 lines):

```
theorem erdos132_main : ∃ C₀ : ℝ, ∀ X : Finset Pt, 2 ≤ X.card → (min (mult
        X (dist2 X)) (mult X (minDist X)) : ℝ) ≤ 54/37 * X.card + C₀
```

with `mult X d` the number of unordered pairs of distinct points of X at distance d, and `dist2`, `minDist` the second-largest and smallest distance. So the formal theorem `erdos132_main` is the $\frac{54}{37}$ bound; the stronger bound $\frac{15}{11}$ of Theorem 1 (and the bound $\frac{4}{3}$ in Region II) is *not* formalised. The formal proof follows the argument of Appendix B: Sections 2, 4 and 7 and Region I as in the text, the ownership and pairing argument in Region II, and the weighted Lemma B.2 in Region III. There is no `sorry`, and `#print axioms` reports only `propext`, `Classical.choice` and `Quot.sound`. The formalisation also proves Vesztergombi's bound $\mu(\Delta_2) \le \frac{3}{2}n$ and the Hopf–Pannwitz bound, so no result is used as a black box. In three places it takes a different route from the text: (a) Vesztergombi's bound is proved in the 2-core (every degree-4 core vertex has two degree-2 core neighbours, then double counting), with no point deletion; (b) the bad-edge bounds of Section 4 are obtained by counting over directions of outward normals instead of arc length, so Lemma 4.2 (lens-corner arc length) is not needed; (c) the degenerate-regime incidence bound uses a local charging argument, and the mixed-T bound of Lemma 5.1 is proved in a weaker form (at most 1200 instead of 71), which only changes the constant C_0.

Limitations of the formalisation. (i) The `bv_decide` axiom described above remains. (ii) Only the $n = 7$ LRAT certificate (1.6 MB) is stored and replayed with `bv_check`; for $n = 11, 13$ the certificates (23 MB and 97 MB) are not stored, so compilation re-runs CaDiCaL (about 40 s and 4 min respectively). (iii) Lemma 4.2 is not formalised; the formal proof of the $\frac{54}{37}$ bound

does not need it (see above). (iv) None of Sections 5.2 to 5.4 and 6, beyond what Appendix B uses, is formalised.

Remark 10.10. (a) The method is tight at "three": $(11, 6, 3)$ is satisfiable (e.g. with class sizes $(12, 9, 12, 12, 6, 4)$), as is $(9, 5, 2)$.
(b) The ordinal statement fails for $n = 5, 6$: $(6, 3, 1)$ has an ordinal solution with profile $(9, 3, 3)$, so at $n = 6$ geometry beyond Lemma 10.6 is needed.
(c) Without the case split by the non-frequent pair, the instance $(15, 8, 6)$ was unresolved after more than 20 minutes of CaDiCaL; the split into $\binom{8}{2}$ cases, together with the symmetry breaking, is what makes $n = 15, 17$ feasible. For $n = 19$ the analogous computation looks feasible but expensive (hours per hard case), and it has not been done.
(d) Theorem 10.8 for $n = 7$ also follows from the literature (Proposition 10.3).

Remark 10.11 (related work). Independently, a Lean project on the Lean Pool [19] proves that for point sets with coordinates in certain restricted fields (e.g. $\mathbb{Q}(\sqrt{2})$), $n \le 2^r$ where r is the number of rare distances. This gives at least three rare distances for $n \ge 8$ in those coordinate fields; it neither implies nor is implied by Theorem 10.1. Other contributions on the forum [13] concern the first question of Problem #132 (at least two rare distances) for small n and general sets.

10.3. **Almost cocircular sets.** For sets on one circle all distances are rare. We show that a bounded number of points off the circle cannot destroy this, up to a linear loss. The two ingredients are Kneser's theorem, which recognises cocircular sets with few distances as subsets of regular polygons, and a uniform bound for the number of vertices of a regular polygon that a point off its circle can see at chord distances, which follows from a theorem of Beukers and Smyth on torsion points of plane curves.

Lemma 10.12 (cocircular sets with few distances)**.** *Let A be a set of $q \ge 3$ points on a circle Γ with $D(A) \le q/2 + K'$, where $K' \ge 0$ and $q > 4K' + 2$. Then A is contained in the vertex set of a regular M-gon inscribed in Γ, for some M with $q \le M \le q + 2K' + 1$.*

Proof. Identify Γ with $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$ by the polar angle. The distance between points with angles a, b determines $a - b$ up to sign, so the difference set $A - A \subseteq \mathbb{T}$ has $|A - A| \le 2D(A) + 1 \le q + 2K' + 1$. By Kneser's theorem [18] (see also [21]), applied in the abelian group $\mathbb{T}$ to the finite sets A and $-A$, with H the stabiliser of $A - A$ (a finite subgroup, hence cyclic of some order M), we have $|A - A| \ge |A + H| + |-A + H| - |H| = (2t - 1)M$, where $A + H$ is a union of t cosets of H. Since $q \le tM$: if $t \ge 2$ then $(2t - 1)M \ge \frac{2t-1}{t} q \ge \frac{3}{2} q$, so $\frac{3}{2} q \le q + 2K' + 1$, i.e. $q \le 4K' + 2$, a contradiction. Hence $t = 1$, so A lies in one coset of H, i.e. in the vertex set of a regular M-gon on Γ, and $q \le M \le |A - A| \le q + 2K' + 1$. □

Scale Γ to the unit circle in $\mathbb{C}$, let $\zeta := e^{2\pi i/M}$ and $c_j := |1 - \zeta^j|$. We say that a point $u \in \mathbb{C}$ *hits* the vertex ζ^a if $|u - \zeta^a| = c_j$ for some j.

Lemma 10.13 (hit lemma)**.** *If $u \ne 0$ and $|u| \ne 1$, then u hits at most 23 vertices of the regular M-gon $\{\zeta^a\}$, for every M.*

Proof. Put $\lambda := \bar{u} \ne 0$ and $\tau := |u|^2 - 1 \ne 0$, and consider the Laurent polynomial

$$F(x, y) := \lambda x + \bar{\lambda} x^{-1} - y - y^{-1} - \tau.$$

For $|x| = |y| = 1$ we have $|u - x|^2 - |1 - y|^2 = -F(x, y)$. So if u hits ζ^a with $|u - \zeta^a| = c_j$, then (ζ^a, ζ^j) and (ζ^a, ζ^{-j}) are *torsion points* (both coordinates roots of unity) of the curve $F = 0$.

F is irreducible. Multiply by xy: $G := xyF = -x\,y^2 + (\lambda x^2 - \tau x + \bar{\lambda})\,y - x$. As a polynomial in y over $\mathbb{C}[x]$ it is primitive ($\gcd(x, \bar{\lambda}) = 1$), and it is not divisible by x or y. So a nontrivial

factorisation would have two factors of degree 1 in y, and the discriminant

$$(\lambda x^2 - \tau x + \bar{\lambda})^2 - 4x^2 = (\lambda x^2 - (\tau+2)x + \bar{\lambda})(\lambda x^2 - (\tau-2)x + \bar{\lambda})$$

would be a square in $\mathbb{C}[x]$. The two quadratic factors have no common root (their difference is $4x$, and neither vanishes at 0), so both would be squares, i.e. $(\tau+2)^2 = 4|\lambda|^2 = (\tau-2)^2$, forcing $\tau = 0$. Hence G is irreducible.

$F = 0$ contains no torsion coset of positive dimension. On $\{(\zeta_1 t^p, \zeta_2 t^q)\}$ with $(p,q) \neq (0,0)$, F becomes a Laurent polynomial in t: if $p, q \neq 0$ its constant term is $-\tau \neq 0$; if $p = 0$ its t^q-coefficient is $-\zeta_2 \neq 0$; if $q = 0$ its t^p-coefficient is $\lambda\zeta_1 \neq 0$.

The Newton polygon of G is the square with vertices $(0,1), (1,0), (2,1), (1,2)$, of area 2. By the theorem of Beukers and Smyth [4], an irreducible plane curve $f = 0$ that is not a torsion coset has at most $22\,V(f)$ torsion points, where $V(f)$ is the area of the Newton polygon of f. Hence the curve $F = 0$ has at most 44 torsion points. Distinct hit vertices give torsion points with distinct first coordinates, and each hit gives two of them unless $\zeta^j = \zeta^{-j}$. Here $\zeta^j = 1$ is impossible (it would put u on Γ), and $\zeta^j = -1$ means $|u - \zeta^a| = 2$, which happens for at most two vertices, since the circle of radius 2 about u is not Γ. So if $d \leq 2$ hits have $c_j = 2$ and h is the number of hits, then $2(h-d) + d \leq 44$, and $h \leq 22 + d/2 \leq 23$. □

A computer search (Appendix A) found that for $M \leq 120$ the maximum number of hits is 13; we have not tried to optimise the constant 23.

Theorem 10.14 (almost cocircular sets)**.** *Let X be a set of n points in convex position and Γ a circle, and put $w := |X \setminus \Gamma|$. Suppose $n \geq 3w$ and $|X \cap \Gamma| \geq 4$. Then*

$$k(X) \;\geq\; \min\Big\{\frac{n-3w+1}{4},\; m - \frac{w}{2} - 22 + \frac{22}{w+1}\Big\}.$$

In particular $k(X) \geq (n-3w+1)/4$ whenever $n + w \geq 91$.

Proof. Put $C := X \cap \Gamma$, $W := X \setminus \Gamma$, $q := |C| = n - w \geq 4$ and $K' := (q-3)/4 \geq 0$.

Case 1: $D(C) > q/2 + K' = (3q-3)/4$. Then $D(X) \geq D(C)$, and Lemma 10.2 gives $k \geq D(X) - m + 1 > \frac{3(n-w)-3}{4} - \frac{n}{2} + 1 = \frac{n-3w+1}{4}$.

Case 2: $D(C) \leq q/2 + K'$. Since $q > 4K' + 2$, Lemma 10.12 puts C inside the vertex set of a regular M-gon on Γ with $M \leq q + 2K' + 1 = (3q-1)/2$.

(i) *The centre O of Γ is not in X.* Otherwise O is a vertex of $\operatorname{conv} X$, so C lies in a closed half-plane bounded by a line through O, i.e. in a closed semicircle, which contains at most $M/2 + 1 \leq (3q+3)/4 < q$ vertices of the M-gon. So every $u \in W$ satisfies $u \neq O$ and $u \notin \Gamma$, and Lemma 10.13 applies to it.

(ii) Within C every distance graph has maximum degree at most 2, hence at most q edges. So a frequent distance of X that is a chord length of the M-gon (a *chord class*) has at least $n + 1 - q = w + 1$ pairs that meet W. Each such pair is either a pair $\{u, c\}$ with $u \in W$, $c \in C$, and c hit by u, or a pair inside W.

(iii) A frequent distance that is not a chord length has all its pairs meeting W (all distances within C are chord lengths). For a given distance, each $u \in W$ has at most 2 points of Γ at that distance, so at least $n + 1 - 2w$ of these pairs lie inside W.

(iv) By Lemma 10.13 there are at most $23w$ pairs of the first kind in (ii), and there are at most $\binom{w}{2}$ pairs inside W; every pair belongs to one distance class. Writing F_{ch} and F_{nc} for the numbers of frequent chord classes and other frequent classes,

$$(w+1)F_{\mathrm{ch}} + (n+1-2w)F_{\mathrm{nc}} \leq 23w + \tbinom{w}{2}.$$

(v) Since $n \geq 3w$ we have $n + 1 - 2w \geq w + 1$, so the number F of frequent distances satisfies $F \leq (23w + \binom{w}{2})/(w+1) = \frac{w}{2} + 22 - \frac{22}{w+1}$.

(vi) By (Alt), $D(X) \geq m$, so $k = D(X) - F \geq m - \frac{w}{2} - 22 + \frac{22}{w+1}$.

For the last claim, $m \geq (n-1)/2$ gives $m - \frac{w}{2} - 22 + \frac{22}{w+1} > \frac{n-1}{2} - \frac{w}{2} - 22 \geq \frac{n-3w+1}{4}$ when $n + w \geq 91$. □

Remark 10.15. (a) For $w = o(n)$ Theorem 10.14 gives $k \geq (\frac{1}{4} - o(1))n$. So the second question of Problem #132 has a positive answer, in a strong form, for sets in convex position that are almost cocircular. What is missing for general convex sets is a stability statement of the form "a convex n-set with at most $n/2 + K$ distances has all but $o(n)$ of its points on one circle"; Kneser's theorem provides this on a circle (Lemma 10.12), and we know no analogue off it.

(b) The loss of $3w/4$ and the factor $\frac{1}{4}$ (rather than $\frac{1}{2}$) come from the threshold $q > 4K' + 2$ in Lemma 10.12.

11. Open problems

Problem 1.6 of [7]. We now know $9/7 \leq L \leq 15/11$, i.e. $1.2857\ldots \leq L \leq 1.3636\ldots$. The lower bound construction [17] is conjectured there to be optimal. In our proof the constant $\frac{15}{11}$ comes only from the R-term: points y without a Δ_2-partner are only known to satisfy $\deg y + m_y \leq 10$ (Theorem 6.1), and for $\tau < \sqrt{3}/2$ this cannot be improved by local arguments (Remark 6.7). In Region II the bound is already $\frac{4}{3}$. So an improvement of $\frac{15}{11}$ by this method needs a non-local argument for the points of R in Regions I and III, for instance one showing that $\Theta(n)$ points with $\deg y + m_y = 10$ cannot coexist with a large $\mu(\Delta_2)$. Reaching $\frac{9}{7}$ would in addition require a bound on $\mu(\Delta_2)$ in Regions II and III that is sharper than $\frac{3}{2}s - \frac{1}{2}p$ (Remark 8.1).

Genericity. One might hope that excluding four concyclic points forces $\mu(\Delta_2) \leq n + O(1)$, which together with Theorem 9.6 would answer the first question of Problem #132 for such sets. This is false. For $n = 4m \geq 12$, start from $n/2$ points on a circle of radius $\frac{1}{2}$ interleaved with $n/2$ points on a circle of radius R, where $R^2 + R\cos(2\pi/n) = \frac{3}{4}$, and perturb generically within the solution set of the distance equations. This gives convex sets with no four concyclic points and $\mu(\Delta_2) = \frac{5}{4}n$. We have verified these examples to 50 digits for $n = 12, 20, 40$ (all equalities to $5 \cdot 10^{-51}$, with concyclicity determinants at least 10^{-8}). A non-convex family with $\mu(\Delta_2) = \frac{5}{4}n - \frac{3}{2}$ was verified numerically for $n \leq 62$. In both families every distance other than Δ and Δ_2 occurs once, so the first question of Problem #132 still has a positive answer for them, but the second rare distance is not Δ_2. An elementary observation, probably folklore, explains the role of concyclicity in the extremal configurations: if $|ab| = |cd| = \Delta$ and $|ac| = |bd| = \Delta_2$, then a, b, c, d are concyclic. (The triangles abc and dcb are congruent. A half-turn would make $abdc$ a parallelogram with $|ad|^2 + |bc|^2 = 2(\Delta^2 + \Delta_2^2) > 2\Delta^2$, which is impossible, so $abdc$ is an isosceles trapezoid.)

Question 11.1. *If no four points of X are concyclic, is $\mu(\Delta_2) \leq \frac{5}{4}n + O(1)$?*

Proposition 9.10 records one further consequence of the no-four-concyclic hypothesis; the examples above show that such local constraints do not bound $\mu(\Delta_2)$ by $n + O(1)$.

Beyond the two largest distances. In a counterexample to the first question of Problem #132, every distance other than Δ occurs at least $n+1$ times, and in particular $\mu(\Delta_2) \geq n+1$. One might hope that then the third-largest distance Δ_3 is rare. This is false: there is an 8-point set, with coordinates in $\mathbb{Q}(\sqrt{3}, \sqrt{5}, \sqrt{10 + 2\sqrt{5}}, \sqrt{10 - 2\sqrt{5}})$, formed by a regular pentagon and three further points, whose distance multiplicities in decreasing order of distance are $2, 9, 9, 5, 2, 1$; so $\mu(\Delta_2) = \mu(\Delta_3) = 9 > 8$ (checked in exact arithmetic, Appendix A). In this example the fourth

distance occurs 5 times. In computer searches (rings of points, symmetric orbits, lattice and cyclotomic pools; about 3,500 sets with $\mu(\Delta_2) > n$), some Δ_j with $j \le 4$ was always rare.

Question 11.2. *Let $n \ge 5$ and let X be an n-point planar set with at least four distinct distances $\Delta > \Delta_2 > \Delta_3 > \Delta_4 > \dots$. Is $\min\{\mu(\Delta_2), \mu(\Delta_3), \mu(\Delta_4)\} \le n$?*

A positive answer would settle the first question of Problem #132 for all sets with at least four distinct distances.

Convex position.

Conjecture 11.3 (Q-ORD)**.** *For every $n \ge 7$, every Q-colouring of the pairs of $\{0, \dots, n-1\}$ has at least three colour classes of size at most n.*

Q-ORD holds for $n \in \{7, 11, 13, 15, 17\}$ (Theorems 10.8 and 10.9), and it would imply that every set of $n \ge 7$ points in convex position has at least three rare distances. (For even n we have not certified it: our solver runs for $n = 8, 10, 12$ also used Lemma T.) Even with the case split, the hardest cases grow from about half a minute at $n = 13$ to about 12 minutes at $n = 17$, so a proof for all n must be human. Deletion-minimal refutations are not local ($(7, 4, 2)$ already needs 25 of the 35 quadrilaterals), so such a proof will have to be global.

For odd $n = 2m + 1 \ge 7$ (no computation needed), Lemma 10.2 and (Alt) show that a convex set with $k \le 2$ must have $D = m + 1$, $k = 2$ and $m - 1$ frequent distances. Moreover, writing the frequent multiplicities as $2m + 2 + e_d$ and the two rare ones as a, b, the identity $a + b + \sum_d e_d = m + 2$ holds. (Indeed $D = m$ gives R_n with $k = m \ge 3$, $D \ge m + 2$ gives $k \ge 3$, and the identity is $(m-1)(2m+2) + \sum e_d + a + b = m(2m+1)$.) So for odd $n \ge 19$ the question in convex position is about this single kind of configuration; by Theorem 10.14 such a configuration cannot have all but w of its points on one circle if $n \ge \max\{3w + 11, 91 - w\}$. Beyond three, the second question of Problem #132 asks whether $k \to \infty$; this is open even in convex position.

Appendix A. Computational verification

All scripts are in the directory `scripts/` of the accompanying repository https://github.com/g8r-b8/erdos132-lean. None of the computations below is used as a proof step, except the SAT/LRAT certificates and the Lean proof for Theorem 10.8 and the DRAT certificates for Theorem 10.9. The rest are independent checks of steps proved in the text. Tags [Ck] in the text refer to sections of `writeup_a2_checks.py`; tags [II], [III], [LP] and [N] refer to `r4_regionII_checks.py`, `r4_regionIII_checks.py`, `r4_lp_checks.py`, and `r4_nb_exact.py` together with `r4_nb_referee_checks.py`, respectively.

Theorem 1.

- `writeup_a2_checks.py`: 58 assertions in exact rationals and rational interval arithmetic, with one-sided rational bounds for $\sqrt{\cdot}$, π, sin, cos, arcsin. Sections: C1 the LP certificates of Lemmas B.1 and B.2 (now used only in Appendix B); C2 the identity of Lemma 2.2(c) and the relation between τ and t; C3 small-angle facts for $\Delta_2 \ge 10^4$; C4 the Region I windows; C5 the Region II constants ((R1), the mixed T-edge count, and, for Appendix B only, (R2), (R2-sym), the rhombus threshold and the pairing ratio); C6 the corner-lemma angles in Region III; C7 the lens constants; C8 case (c2); C9 the bad-edge totals; C10 the degenerate-regime constants; C11 the small-case packing bound. All pass.
- `r4_regionII_checks.py`: 17 assertions in exact rational arithmetic for Section 5.3: the constants (including the type-(i) count), the LP with the p-term, and the identity and threshold of Lemma 5.3 (on 500 exact rational instances and 2000 random exact rational

instances). It also runs an adversarial floating-point search against Lemma 5.4 ($2 \cdot 10^5$ samples), and asserts in exact arithmetic a sufficient condition for the angle estimate in the proof of Lemma 5.4 on all of Region II (in a formulation slightly different from the one in the text, with minimum margin 0.088 over the samples). (The Lean proof of Lemma C in the 15/11 development, which is in progress, follows the formulation of this script rather than the condition $3\theta_0 + \varepsilon < \pi/2$ used in the text; both are valid.)

- `r4_regionIII_checks.py`: 18 assertions in exact rational arithmetic for Section 5.4: the constants of Lemma 5.5 and of the injectivity step, the corner-lemma angles, the mixed T-edge count < 76, the threshold used in Lemma 5.7, the bounds on arcsin and f and the margin $t + 1/(23\Delta_2) < 3/\pi$ in the arithmetic step, and the thin-cap estimate of Lemma 5.6. Floating-point searches (reported, not used): B1, the slot-killing configuration, with maximum $|y - z| = 0.0691$; B2, the forced step $v_{i+1} = v^*$ of Lemma 5.9, with 0 violations in $2 \cdot 10^5$ samples; B3, the minimum of $|j\alpha_2 - k\alpha_1|\,\Delta_2$, which was $1.7 \cdot 10^{-6} > 0$.
- `r4_lp_checks.py`: exact dual certificates for Lemma 3.5 and Remark 3.6, for all values of $2c$ in the remark, with and without the p-term; it also re-checks the certificates of Lemmas B.1 and B.2.
- `r4_nb_exact.py` (exact fractions and decimals, with rigorous rational bounds for sin, cos, tan): (1) the closed forms of c_K, c_D, the identity $c_D(0) = \tau$ and the monotonicity of c_D used in Remark 6.7; (2) the LP values of Remark 3.6; (3) the constants of Lemmas 6.3 and 6.4: the apex arc is at most $231.19 < 232$, the thin-cone margin is positive, $M \le 468^2$, and the count of Lemma 6.4 for $\varepsilon_0 = 1/50$ is at most $6.88 \cdot 10^7$; (4) the 60-digit verification of the configuration in Remark 6.7.
- `r4_nb_referee_checks.py`: an independent set of exact checks, with rigorous Taylor bounds, of the LP steps and the constants of Theorems 6.1 and 6.2.
- `r4_nb_search.py`: an adversarial floating-point search over local boundaries of K with random corners near a point y (reported, not used). No configuration violating the bounds of Theorems 6.1 and 6.2 in their range of validity was found.
- `r4_nb_ring16_check.py`: consistency with ienjoymath's construction (floating point). For $n = 60, 120, 250$, every point of R has $m_y = 0$ and $\deg y + m_y \le 6$.
- `ref2_numeric.py`: an independent set of 40 exact assertions (second referee pass of the $\frac{54}{37}$ argument) covering the inequalities of `writeup_a2_checks.py`, together with an adversarial floating-point search for (R1), (R2), (R2-sym). The search is reported, not asserted. Its maxima were 0.906, 0.897 and 0.914, against the proved bounds 0.908, 0.918 and 0.948. (Only (R1) is used in the proof of Theorem 1; (R2) and (R2-sym) concern Appendix B.)
- `ref2_ownership.py`: an exhaustive exact check of the ownership/pairing bookkeeping of Appendix B (not used for Theorem 1), in the abstract model the argument uses: rows that are disjoint unions of paths and cycles of length ≥ 3, rungs forming a matching, (R2), (R2-sym), and no two consecutive rhombi. It covers all row structures up to isomorphism with at most 6 vertices per row and all partial rung matchings. There were 2,365,426 admissible configurations without rhombi, with maximum ratio exactly 5/4, and 2,741,183 with rhombi allowed, with maximum ratio exactly 4/3. Mutation tests show that (R2-sym) and the rule excluding two consecutive rhombi are both needed.
- `ref_degen.py`, `degen_check.py`, `ref2_degen.py`: adversarial searches in the degenerate regime, including 5500 configurations built so that three or more circles $\mathrm{C}(w, \Delta_2)$ pass through one vertex of ∂K. The maximum of $\mu(\Delta_2) - n$ found was 0, against the proved bound 1.
- `ref_checks.py`, `a2_checks.py`: random tests of the lens lemma (about 7000 lenses, worst ratio 0.66) and of the exact single-centre strip in the near-resonant regime.

The $\frac{54}{37}$ argument of Appendix B, together with the parts of Sections 2 to 5 and 7 that it shares with the present proof, was checked twice more, independently, in adversarial "referee" passes. The new parts of the proof of Theorem 1 were checked in independent referee passes as follows: Region II (Section 5.3): one; Region III with $\tau \neq \sqrt{3}/2$: one; Region III with $\tau = \sqrt{3}/2$: two; Theorems 6.1 and 6.2: one. Vesztergombi's theorem [23] was checked in the original: its main theorem is $\mu(\Delta_2) \leq \frac{3}{2}n$ for all n, with no additive constant.

Section 9.

- `va1b_exact.py` (sympy, mpmath interval arithmetic): exact computation of the six Case 1 candidates of Proposition 9.2 and their pairwise squared distances, with an exhaustive check that no compatible set of candidates covers all three v_i. It also gives a closed form and a rigorous interval check (6000 subintervals, 0 failures) of the Case 2 branch $|r - v_2| = \Delta$.
- `va1c_sympy.py`: symbolic checks of the identity in Lemma 9.3, of $\lambda(\lambda + 2\cos\gamma) = 0$, of the chord formula, and of the bisector direction in Lemma 9.4.
- `va1c_local.py`: exhaustive search for the configuration of Lemma 9.4 with unit vectors from $\mathbb{Z}[\omega]$ (norm ≤ 147, exact), $\mathbb{Z}[i]$ (norm ≤ 325, exact) and N-th roots of unity ($5 \leq N \leq 60$): 0 hits in about $3.2 \cdot 10^6$ configurations. A control run with weakened hypotheses does find hits.
- `va1_clique.py`, `va1_cyclo.py`, `va1c_random.py`: searches over maximal compatible sets in lattice and cyclotomic pools, plus greedy floating-point constructions. Every set was tested against Lemma 9.1, Proposition 9.2, Theorem 9.5, Theorem 9.6 and Corollaries 9.7 and 9.8: more than 800,000 sets in the first searches and about 81,000 in the second, with 0 violations.

Section 10.

- `ver_classification_k.py`, `ver_hexagons.py`: exact chord-index profiles for the classified extremal sets used in Theorems 10.4 and 10.5 and Proposition 10.3, and strict convexity and multiplicity profiles of the three 3-distance hexagons.
- `counting_parity.py`: the closed forms of Lemma 10.2 for $n < 400$.
- `b3_sat.py`: the CNF encoder (direct and order encodings, with or without Lemma T). `b3_enum.py`: the standalone backtracking enumerator for $n = 7$.
- `b3_validate.py`: encoder validation. Every 7-subset of R_N, $7 \leq N \leq 16$ (11,440 sets), and every 11-subset of R_N, $N \leq 15$, satisfies all constraints under exact chord-index ranks, as do 3000 (for $n = 7$) and 500 (for $n = 11$) random convex polygons. The $(7, 3, 0)$ enumeration returns exactly the R_7 colouring. Every satisfying assignment printed by the SAT route was re-checked against the independent constraint code of `b3_enum.py`.
- Four solvers (CaDiCaL, Glucose 4, MiniSat 2.2, Lingeling) agree on the $n = 7$ and $n = 11$ instances.
- `b3_drup_check.py`: an independent forward RUP checker, used for the $n = 7$, $n = 9$ and $n = 11$ DRUP proofs. Its self-test rejects an empty proof of a satisfiable instance, a proof with one negated lemma, and a truncated proof.
- `b3_drat_trim.py`: regenerates each CNF and Glucose proof deterministically and runs `drat-trim` with LRAT output, followed by `lrat-check`. The CNF, DRUP and LRAT files are not stored in the repository because of their size.

For Theorem 10.9:

- `r1_sat_split.py`: the case-split encoder (Q-only order encoding, as in `b3_sat.py`, plus the case constraints and the symmetry-breaking constraints). `r1_run.py`, `r1_batch.py`, `r1_lowD.py`: build each CNF, solve it with Kissat writing a DRAT proof, check the proof with `drat-trim`, and delete both files. A case is counted as done only when `drat-trim`

instance (n, D, F)	encoding	lemmas	`drat-trim`	`lrat-check`
$(7, 4, 2)$	Q	714	verified	verified
$(11, 2, 0)$, $(11, 3, 0)$, $(11, 4, 0)$	Q+T	3, 8, 56	verified	verified
$(11, 5, 3)$	Q+T	524	verified	verified
$(11, 6, 4)$	Q	35,510	verified	verified
$(13, 2, 0)$, $(13, 3, 0)$, $(13, 4, 0)$, $(13, 5, 0)$	Q	2, 7, 82, 448	verified	verified
$(13, 6, 4)$	Q	8,463	verified	verified
$(13, 7, 5)$	Q	446,686	verified	verified

TABLE 2. Certified refutations for Theorem 10.8.

reports `s VERIFIED`. All $5 + 21 + 28$ cases for $n = 15$ and $6 + 28 + 36$ cases for $n = 17$ were verified; solve times were at most 56 s for $n = 15$ with symmetry breaking (282 s for the case $\{7, 8\}$ without it) and at most 739 s for $n = 17$, and the DRAT proofs were at most 124 MB.

- `r1_sb_test.py`: the orbit test of the symmetry-breaking constraints described after Theorem 10.9.

For Section 10.3:

- `r3_cyc_hits.py`: the maximum number of hits (Lemma 10.13) for $M \le 120$. Every point with at least two hits is an intersection of two circles centred at vertices with chord radii; candidates are scanned in floating point with tolerance 10^{-7}, and the best ones are confirmed exactly in $\mathbb{Q}(\zeta_M)$ by arithmetic modulo the cyclotomic polynomial. The maximum found is 13 (at $M = 60, 120$).
- `r3_almostcoc_check.py`: floating-point sanity check of Theorem 10.14 on regular polygons with holes plus up to 3 adversarial points off the circle ($M \le 60$); the slack against the bound was at least 24 in every case.

Section 11.

- `r2_verify.py`: 50-digit verification of the convex examples with $\mu(\Delta_2) = \frac{5}{4}n$ and no four concyclic points, for $n = 12, 20, 40$. `r2_local_lemmas.py`: symbolic checks of identities used in the local analysis around hull vertices with three Δ_2-neighbours.
- `r5_verify_n8.py`: exact (sympy) verification of the 8-point example with $\mu(\Delta_2) = \mu(\Delta_3) = 9$, including its full multiplicity profile.

Lean. The whole development (all `E132*.lean` files: the convex-position results, the layer bound, Vesztergombi's and Hopf–Pannwitz's bounds, and the $\frac{54}{37}$ bound `erdos132_main`) builds with `lake exe cache get` followed by `lake build`; on a laptop this takes about 6 minutes after the Mathlib cache is downloaded, of which about 4 minutes is `bv_decide` for $n = 13$. The toolchain is Lean `v4.35.0-rc2` with Mathlib.

APPENDIX B. THE 54/37 ARGUMENT FORMALISED IN LEAN

This appendix gives the written counterpart of the formal theorem `erdos132_main` (Section 10.2): the bound $\min\{\mu(\Delta_2), \mu(\delta)\} \le \frac{54}{37}n + C_0$ with $C_0 = 3N_0 + 10^8$. It uses Section 2, the reductions Lemma 3.1, Corollary 3.4 and the small cases of Section 3, Section 4, Region I of Section 5.1, (R1) and Lemma 5.1 and Lemma 5.2 of Section 5, and Section 7. It does not use Lemmas 3.3 and 3.5, Lemmas 5.3 and 5.4, the Region III arguments of Section 5.4, or Section 6. In place of the R-term estimate it uses the trivial bound (E), $\mu(\delta) \le e(S) + 6r$.

Linear combinations.

Lemma B.1. *Suppose $e(S) \le ks + C$ with $0 \le k < 3/2$. Then $\min\{\mu(\Delta_2), \mu(\delta)\} \le \frac{9}{7.5-k}\, n + C$.*

Proof. Put $\lambda = (6-k)/(7.5-k) \in (0,1)$. By Lemma 3.1 and Corollary 3.4, $\min\{\mu(\Delta_2), \mu(\delta)\} \le \lambda \cdot \frac{3}{2}s + (1-\lambda)(ks + C + 6r) \le \frac{9}{7.5-k}(s+r) + C$. For $k = 1, \frac{5}{4}, \frac{4}{3}$ the factor $9/(7.5-k)$ is $\frac{18}{13}, \frac{36}{25}, \frac{54}{37}$.[C1] □

Lemma B.2. *Suppose $e(S) \le s + 2p + C$. Then $\min\{\mu(\Delta_2), \mu(\delta)\} \le \frac{7}{5}n + C$.*

Proof. By Lemma 3.2 and Corollary 3.4, $\min \le \frac{4}{5}(\frac{3}{2}s - \frac{1}{2}p) + \frac{1}{5}(s + 2p + C + 6r) = \frac{7}{5}s + \frac{6}{5}r + \frac{C}{5} \le \frac{7}{5}n + C$.[C1] □

Region I gives $e(S) \le s + C_{\text{bad}}$, hence $\frac{18}{13}n + C$ by Lemma B.1, and so does Region II with $\tau > 1$.

Region II with $\sqrt{3}/2 + 3\beta \le \tau \le 1$. Here $t \le 1$, $t \ge \tau$ and $t - \arcsin\beta \ge 0.886$; there are fewer than 71 mixed T-edges (Lemma 5.1), and each vertex has at most one rung (R1). Rungs are written $z = v + o_{\sigma'}(u_v)$ with a sign $\sigma' \in \{+,-\}$, and $|o_\pm(u) - o_\pm(u')| \le 2\sigma + \angle(u,u')$ for any choice of signs. By the T-step form, for T-steps e_1 at v_0 and e_2 at z_0 in the same direction, $\angle(e_1, e_2) \le \angle(u_{v_0}, u_{z_0}) + \arcsin\beta$.

Lemma B.3 ((R2), proximity)**.** *Let (v_0, z_0) be a rung, v_1 the right pure T-neighbour of v_0 and z_1 the right pure T-neighbour of z_0; put $e_1 = v_1 - v_0$, $e_2 = z_1 - z_0$. If v_1 has a rung (v_1, z'), then $z' = z_1$.*

Proof. $z' - z_1 = (e_1 - e_2) + (o_{\sigma_1}(u_{v_1}) - o_{\sigma_0}(u_{v_0}))$. Since $\angle(u_{v_0}, u_{z_0}) \le \beta$ (good rung) and $\angle(u_{v_0}, u_{v_1}) \le \beta$ (good T-edge), $|z' - z_1| \le (\beta + \arcsin\beta) + \beta + 2\sigma \approx 0.918 < 1$.[C5] Distinct points of X are at least 1 apart. □

Lemma B.4 ((R2-sym))**.** *In the setting of Lemma B.3, if z_1 has a rung (v', z_1), then $v' = v_1$.*

Proof. $v' = z_1 - o_{\sigma'}(u_{v'})$ and $v_1 = z_0 - o_{\sigma_0}(u_{v_0}) + e_1$, so $v' - v_1 = (e_2 - e_1) - (o_{\sigma'}(u_{v'}) - o_{\sigma_0}(u_{v_0}))$. The angle between $u_{v'}$ and u_{v_0} is at most 3β, as the sum of three good-edge angles along $v' \to z_1 \to z_0 \to v_0$. Hence $|v' - v_1| \le (\beta + \arcsin\beta) + 3\beta + 2\sigma \le 2\sigma + 6\beta \approx 0.948 < 1$.[C5] □

(R3) *Parallel rungs and rhombi.* Rungs with the same vector c satisfy $c \cdot u_v = -\tau$, so u_v is one of the two unit solutions, which lie at $\pm\theta_0$ from $-c$. Rungs form a matching by (R1), so their K-ends are distinct and by Lemma 2.3 have distinct normals. Hence *no three rungs are parallel*, and two parallel rungs have normals exactly $2\theta_0$ apart. If v_0v_1, v_1z_1, z_1z_0, z_0v_0 are all unit edges (a *rhombus*), then v_1 and z_0 are the two points of $\mathrm{C}(v_0,1) \cap \mathrm{C}(z_1,1)$, so $v_0 + z_1 = v_1 + z_0$ and the rungs (v_0, z_0) and (v_1, z_1) are parallel. Since the T-edge v_0v_1 is good, $2\theta_0 = \angle(u_{v_0}, u_{v_1}) \le \beta$. So *rhombi exist only if* $\tau \ge \cos(\beta/2)$, and $\cos(\beta/2) > \sqrt{3}/2 + 3\beta$.[C5]

Ownership. For a rung $\mathsf{r} = (v, z)$, walk right from v along pure K-row T-edges until reaching the first vertex after v that is a rung vertex, or until the walk stops. Let $k \ge 1$ be the number of vertices visited before stopping, v included. Define d in the same way in the D-row, starting from z. Then r *owns* these $k + d$ vertices. Every other row vertex is *unowned*: it has no rung vertex to its left on its T-path or T-cycle. Ownership is well defined, since each owned vertex belongs to the nearest rung vertex at or to its left, and that rung vertex lies on exactly one rung.

Charging. Charge to r the rung itself and the right T-edge out of each vertex it owns. Charge every other pure T-edge to its unowned left endpoint. Every rung and pure T-edge is charged exactly once, and every vertex has at most one right T-edge. Hence $\text{edges}(\mathsf{r}) \le 1 + k + d$, with equality only if both walks end at a rung vertex ("both reach"), and otherwise $\text{edges}(\mathsf{r}) \le k + d$; unowned vertices carry at most one edge each.

Both-reach gaps are $(1,1)$ *or* $(\geq 2, \geq 2)$. If both walks reach and $k = 1$, then z_0 has a right neighbour z_1 and v_1 is a rung vertex; by Lemma B.3 its rung is (v_1, z_1), so $d = 1$. If both reach and $d = 1$, then v_0 has a right neighbour v_1 and z_1 is a rung vertex; by Lemma B.4 its rung is (v_1, z_1), so $k = 1$. A $(1,1)$ gap is a rhombus $v_0v_1z_1z_0$ (by (R1), $z_1 \neq z_0$ and $v_1 \neq v_0$). A cycle with a single rung vertex gives a both-reach gap with k equal to the cycle length, which is at least 3.

Ratios. A rhombus gap has ratio edges/vertices $= 3/2$; a both-reach gap with $k, d \geq 2$ has ratio at most $5/4$; every other gap has ratio at most 1. If $\tau < \cos(\beta/2)$ there are no rhombi, so $e(S) \leq \frac{5}{4}s + C_{\text{bad}} + 71$ and Lemma B.1 gives $\frac{36}{25}n + C$.

Pairing. If $\tau \geq \cos(\beta/2)$, pair each rhombus rung $\mathsf{r}_0 = (v_0, z_0)$ with its successor $\mathsf{r}_1 = (v_1, z_1)$. The rung r_1 is not itself a rhombus rung: otherwise $\mathsf{r}_0, \mathsf{r}_1, \mathsf{r}_2$ would be three parallel rungs (here $v_2 \neq v_0$ because T-cycles have length at least 3). The pairing is injective, because v_0 is the unique left T-neighbour of v_1. If r_1 owns m vertices, the pair has ratio at most

$$\max\Big\{\frac{4+m}{2+m} : m \geq 4\Big\} \cup \Big\{\frac{3+m}{2+m} : m \geq 2\Big\} = \frac{4}{3}.^{[C5]}$$

Hence $e(S) \leq \frac{4}{3}s + C_{\text{bad}} + 71$, and Lemma B.1 gives $\frac{54}{37}n + C$. This is the only place where $\frac{54}{37}$ is attained.

Region III. By Lemma 5.2 and the angle bound $\min\{2\theta_0, \pi/2 - \theta_0\} > 2\pi/7$,[C6] at most 6 K-vertices with a rung have Δ_2-degree at least 2; every other K-vertex with a rung lies in Q. A K-vertex has at most 2 rungs, and (T1) bounds all T-edges, mixed ones included, by s. Hence

$$e(S) \leq s + 2p + 12 + C_{\text{bad}},$$

and Lemma B.2 gives $\frac{7}{5}n + C$. (A "hexagonal strip" around a single centre has average δ-degree 3 locally, so this regime genuinely needs a weighted bound such as Lemma B.2 for this argument.)

regime ($\delta = 1$, $\beta = 1/100$)	bound on $e(S)$	bound on $\min\{\mu(\Delta_2), \mu(\delta)\}$
degenerate: $\Delta > 1.94\Delta_2$, $n \geq 4$	—	$n + 1$ (Proposition 7.1)
nondegenerate, $\Delta_2 < 10^4$	—	$3n < 3N_0$ (Section 3)
I: $\tau \leq \sqrt{3}/2 - 2\beta$ (incl. $\tau \leq 0$)	$s + C$	$\frac{18}{13}n + C$
III: $\sqrt{3}/2 - 2\beta < \tau < \sqrt{3}/2 + 3\beta$	$s + 2p + C$	$\frac{7}{5}n + C$
II: $\sqrt{3}/2 + 3\beta \leq \tau < \cos(\beta/2)$	$\frac{5}{4}s + C$	$\frac{36}{25}n + C$
II: $\cos(\beta/2) \leq \tau \leq 1$	$\frac{4}{3}s + C$	$\frac{54}{37}n + C$
II: $\tau > 1$ ($\Leftrightarrow t > 1$)	$s + C$	$\frac{18}{13}n + C$

TABLE 3. The regimes of the $\frac{54}{37}$ argument; here $C < 10^8$.

Assembly. Every constant in the last column is at most $\frac{54}{37}$,[C1] and $n + 1 \leq \frac{54}{37}n + 1$. With $C_{\text{bad}} + 71 + 12 < 10^8$ (Proposition 4.3),[C9] the bound holds with $C_0 = 3N_0 + 10^8$.

Remark B.5. The two least standard parts of this argument are the ownership/pairing bookkeeping and the incidence count in Step 4 of Section 7. Both were re-derived independently, and both were also checked by computer in abstract models (Appendix A). The bookkeeping check is exhaustive up to $6+6$ row vertices, and both (R2-sym) and the "no three parallel rungs" fact turn out to be necessary for it. For this argument the bottleneck is the near-resonant regime $\tau \geq \cos(\beta/2)$, where rhombi occur; Lemma 5.4 removes it in the proof of Theorem 1.

References

1. Edward Altman, *On a problem of P. Erdős*, Amer. Math. Monthly **70** (1963), 148–157.
2. ———, *Some theorems on convex polygons*, Canad. Math. Bull. **15** (1972), 329–340.
3. Gilles Audemard and Laurent Simon, *Predicting learnt clauses quality in modern SAT solvers*, Proceedings of IJCAI 2009, 2009, pp. 399–404.
4. Frits Beukers and Chris J. Smyth, *Cyclotomic points on curves*, Number Theory for the Millennium, I (Urbana, IL, 2000), A K Peters, Natick, MA, 2002, pp. 67–85.
5. Armin Biere, Tobias Faller, Katalin Fazekas, Mathias Fleury, Nils Froleyks, and Florian Pollitt, *CaDiCaL, Gimsatul, IsaSAT and Kissat entering the SAT Competition 2024*, Proceedings of SAT Competition 2024: Solver, Benchmark and Proof Checker Descriptions, Department of Computer Science Report Series B, no. B-2024-1, University of Helsinki, 2024, pp. 8–10.
6. Thomas F. Bloom, *Erdős Problem #132*, https://www.erdosproblems.com/132, Accessed 23 September 2026. The page records Erdős's $100 offer for "any nontrivial result" with the reference key [Er97e].
7. Felix Christian Clemen, Adrian Dumitrescu, and Dingyuan Liu, *On multiplicities of interpoint distances*, Acta Math. Hungar. **177** (2025), 231–245, arXiv:2505.04283 (v5, 3 Feb 2026).
8. Luís Cruz-Filipe, Marijn J. H. Heule, Jr. Hunt, Warren A., Matt Kaufmann, and Peter Schneider-Kamp, *Efficient certified RAT verification*, Automated Deduction – CADE 26, Lecture Notes in Comput. Sci., vol. 10395, Springer, 2017, pp. 220–236.
9. Leonardo de Moura and Sebastian Ullrich, *The Lean 4 theorem prover and programming language*, Automated Deduction – CADE 28, Lecture Notes in Comput. Sci., vol. 12699, Springer, 2021, pp. 625–635.
10. Paul Erdős and Peter Fishburn, *Intervertex distances in convex polygons*, Discrete Appl. Math. **60** (1995), 149–158.
11. ———, *Multiplicities of interpoint distances in finite planar sets*, Discrete Appl. Math. **60** (1995), 141–147.
12. ———, *Convex nonagons with five intervertex distances*, Geom. Dedicata **60** (1996), 317–332.
13. *Erdős Problem #132 forum thread and proof claims*, https://www.erdosproblems.com/forum/thread/132, 2026, Comments by Z. Rafik (28 Jan 2026), J. Marchetto (5 Jul 2026), ienjoymath (25 Jul 2026); proof claim by E. Beller (23 Aug 2026). Accessed 23 September 2026.
14. Peter Fishburn, *Convex polygons with few vertices*, Tech. Report 92-17, DIMACS, 1992.
15. ———, *Convex polygons with few intervertex distances*, Comput. Geom. **5** (1995), 65–93.
16. Heinz Hopf and Erika Pannwitz, *Aufgabe Nr. 167*, Jahresber. Deutsch. Math.-Verein. **43** (1934), 114.
17. ienjoymath, *Comment on Erdős Problem #132 (lower bound* $9/7$ *for Problem 1.6 of Clemen–Dumitrescu–Liu)*, Forum post, https://www.erdosproblems.com/forum/thread/132, 25 July 2026; note at https://github.com/mathematiclover1337/erdos-132-note, 2026.
18. Martin Kneser, *Abschätzung der asymptotischen Dichte von Summenmengen*, Math. Z. **58** (1953), 459–484.
19. lyfar, *rare-distance-fields*, https://github.com/lyfar/rare-distance-fields, 2026, Lean project on rare distances for point sets with coordinates in restricted fields; accessed 23 September 2026.
20. Filip Morić and David Pritchard, *Counting large distances in convex polygons: a computational approach*, arXiv:1103.0412, 2011.
21. Terence Tao and Van H. Vu, *Additive combinatorics*, Cambridge Studies in Advanced Mathematics, vol. 105, Cambridge University Press, 2006.
22. The mathlib Community, *The Lean mathematical library*, Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020), 2020, pp. 367–381.
23. Katalin Vesztergombi, *On large distances in planar sets*, Discrete Math. **67** (1987), 191–198.
24. ———, *The two largest distances in finite planar sets*, Discrete Math. **150** (1996), 379–386.
25. Nathan Wetzler, Marijn J. H. Heule, and Jr. Hunt, Warren A., *DRAT-trim: Efficient checking and trimming using expressive clausal proofs*, Theory and Applications of Satisfiability Testing – SAT 2014, Lecture Notes in Comput. Sci., vol. 8561, Springer, 2014, pp. 422–429.