

Dark Matter Particle Scattering Processes and the Role of Sommerfeld Enhancement



UNIVERSITY OF TURIN
Department of Physics
Master's Degree in Theoretical Physics

July 21, 2026

Gabriele Cembalo

Dark Matter

Dark Matter: a hypothetical new component of matter. Having suppressed (or null) electromagnetic interactions.

WIMP Hypothesis (Weakly Interacting Massive Particle): non-baryonic DM particles interacting only via weak interactions and having a sufficiently high mass ($m_x \geq$ a few GeV).

Dark Matter Detection: Production in Colliders, Direct detection, Indirect detection: searching for the products of WIMP particle annihilation processes.

Sommerfeld Enhancement: a typical non-relativistic QM phenomenon involving an increase in the annihilation cross-section, compared to the perturbative calculation, between particles in the presence of an interaction potential.

The Hulthén Potential, the cross section and S

An attractive Hulthén potential can be considered together with an approximation of the centrifugal term:

$$V_H(r) = -\frac{\alpha_x \delta e^{-\delta r}}{1 - e^{-\delta r}} \quad , \quad \tilde{V}_l(r) = \frac{l(l+1)}{m_x} \frac{\delta^2 e^{-\delta r}}{(1 - e^{-\delta r})^2}$$

and solving the Schrödinger equation:

$$\left[\frac{d^2}{dr^2} + \frac{m_x^2 v^2}{4} + \frac{m_x \alpha_x \delta e^{-\delta r}}{1 - e^{-\delta r}} - \frac{l(l+1) \delta^2 e^{-\delta r}}{(1 - e^{-\delta r})^2} \right] \chi_l(r) = 0.$$

This leads to:

$$\chi_l(t) = \frac{1}{2} \left| \frac{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)}{\Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)} \right| t^{l+1} (1-t)^{iac} {}_2F_1(\lambda_+^l, \lambda_-^l, \omega, t)$$

The Hulthén Potential, the cross section and S

This yields the phase shifts:

$$\delta_l = \arg \left(\frac{i \Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)} \right)$$

and the cross-section:

$$\sigma_l = \frac{16\pi}{m_x^2 v^2} \sum_{l=0}^{\infty} (2l+1) \sin^2 \left\{ \arg \left(\frac{i \Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)} \right) \right\}.$$

The Hulthén Potential, the cross section and S





The Sommerfeld factor is:

$$\begin{aligned} S_l &= \left| \frac{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)}{\Gamma(1+l) \Gamma(1+l+2iac)} \right|^2 \\ &= \left| \frac{(1+iac + \sqrt{c-a^2c^2})_l (1+iac - \sqrt{c-a^2c^2})_l}{\Gamma(1+l)(1+2iac)_l} \right|^2 \times \\ &\quad \times \frac{\pi}{a} \frac{\sinh(2\pi ac)}{\cosh(2\pi ac) - \cosh(2\pi \sqrt{a^2c^2 - c})} \end{aligned}$$

The resonant values for S_l can be determined considering the low-velocity limit $v \rightarrow 0$ ($a \rightarrow 0$). Resonance occurs for:

$$m_x = \frac{\pi^2 m_\phi}{6\alpha_x} n^2, \quad n \in \mathbb{N}.$$

The Hulthén Potential, the cross section and S



Conclusions and what I've learnt

What's the point?

- The fundamental nature of Dark Matter remains one of the most significant open questions in contemporary physics.
- WIMP particles are among the leading candidates for DM.
- Sommerfeld enhancement could play a crucial role in the detection of indirect signals from WIMPs.

What have I learnt?

- I've gotten hands-on experience with solving a non-trivial Schrödinger equation, performing asymptotic expansions, and calculating the phase shift and cross section.
- This was my first approach to a model where the theoretical model comes before the experimental evidence.

Thank You

Appendix: table of contents

4 Indications of Dark Matter

- Bullet Cluster
- Rotation Curves

5 WIMP Hypothesis and Scattering Processes of DM Particles

- The Sommerfeld Enhancement
 - $e^+ + e^- \rightarrow 2\gamma$ Annihilation
 - Classical Analogy
- The Hulthén Potential
 - Schrödinger Equation
 - Wavefunction Normalization
 - The cross section
 - Wavefunction Behavior as $x \rightarrow 0$
 - The Sommerfeld factor
 - Plots for $l = 0$
 - Plots for $l = 1$

6 Bibliography

Indications of Dark Matter

Dark Matter: a hypothetical new component of matter. Having suppressed (or null) electromagnetic interactions, its observational detection is highly complex.

Experimental evidence:

- Bullet Cluster
- Galaxy rotation curves
- Gravitational lensing of galaxy clusters
- Cosmic microwave background

Indications of Dark Matter: Bullet Cluster




Figure 1: The Bullet Cluster. The image shows the hot gas (in pink) traced via X-rays, galaxies and stars (the yellow and white areas), and the mass distribution (in blue). Credit: X-ray: NASA/CXC/CfA/M.Markevitch et al.; Optical: NASA/STScI; Magellan/U.Arizona/D.Clowe et al.; Lensing Map: NASA/STScI; ESO WFI; Magellan/U.Arizona/D.Clowe et al.

Indications of Dark Matter: Rotation Curves

Experimentally we observe:

$$M(r) \sim M \quad r \sim \mathcal{O}(10) \text{ kpc.}$$

From Newton's equation:

$$G \frac{mM(r)}{r^2} = m \frac{v_c^2}{r}$$
$$v_c = \sqrt{\frac{GM(r)}{r}}.$$

Therefore, one must assume the existence of a DM component such that $\rho(r) \propto 1/r^2$ for $r \sim \mathcal{O}(10)$ kpc.




Figure 2: Reference: [link](#). Rotation curve of the spiral galaxy **NGC 3198** (located approximately 47 million light-years from Earth).

WIMP Hypothesis and Scattering Processes of DM Particles

WIMP Hypothesis (Weakly Interacting Massive Particle): non-baryonic DM particles interacting only via weak interactions and having a sufficiently high mass ($m_x \geq$ a few GeV).

Dark Matter Detection:

- Production in Colliders
- Direct detection
- Indirect detection: searching for the products of WIMP particle annihilation processes.

Sommerfeld Enhancement: a typical non-relativistic QM phenomenon involving an increase in the annihilation cross-section, compared to the perturbative calculation, between particles in the presence of an interaction potential.

The Sommerfeld Enhancement: $e^+ + e^- \rightarrow 2\gamma$ Annihilation

Annihilation Hamiltonian:

$$H_{ann} = U_0 \delta^3(\mathbf{r}).$$

Annihilation cross-section:

$$\sigma_0 = |\psi_0(0)|^2.$$





Figure 3: Graphical representation of the $e^+ + e^- \rightarrow 2\gamma$ annihilation experiment.

Introducing the interaction potential $V(r)$, we have:

$$\sigma = |\psi(0)|^2 \quad \Rightarrow \quad S = \frac{\sigma}{\sigma_0} = \frac{|\psi(0)|^2}{|\psi_0(0)|^2}.$$

The Sommerfeld Enhancement: Classical Analogy

A planet of mass m interacts gravitationally with the Sun $M_\odot$.



Ignoring gravity:

$$\sigma_0 = \pi R^2.$$

Considering the gravitational interaction $V(r) = -G \frac{mM_\odot}{r}$:

$$\sigma = \pi R^2 \left(1 + \frac{v_{esc}^2}{v^2} \right) \quad \Rightarrow \quad S = \frac{\sigma}{\sigma_0} = 1 + \frac{v_{esc}^2}{v^2}.$$

The Hulthén Potential

An attractive Hulthén potential can be considered together with an approximation of the centrifugal term:

$$V_H(r) = -\frac{\alpha_x \delta e^{-\delta r}}{1 - e^{-\delta r}} \quad , \quad \tilde{V}_l(r) = \frac{l(l+1)}{m_x} \frac{\delta^2 e^{-\delta r}}{(1 - e^{-\delta r})^2}$$

and solving the Schrödinger equation:

$$\left(-\frac{\hbar^2}{2\mu} \Delta + V(\mathbf{r}) \right) \Psi(\mathbf{r}) = E \Psi(\mathbf{r}) \implies \left(\frac{d^2}{dr^2} + \kappa^2 - 2\mu V_{eff.}(r) \right) \chi_l(r) = 0$$

$$\left[\frac{d^2}{dr^2} + \frac{m_x^2 v^2}{4} + \frac{m_x \alpha_x \delta e^{-\delta r}}{1 - e^{-\delta r}} - \frac{l(l+1) \delta^2 e^{-\delta r}}{(1 - e^{-\delta r})^2} \right] \chi_l(r) = 0.$$

where it is convenient to define:

$$a = \frac{v}{2\alpha_x} \quad , \quad c = \frac{\alpha_x m_x}{\delta} \quad \delta = \frac{\pi^2}{6} m_\phi.$$

Hulthén potential: Schrödinger Equation

Indeed, with an attractive Hulthén potential can be considered:

$$V_H(r) = -\frac{\alpha_x \delta e^{-\delta r}}{1 - e^{-\delta r}}$$

and the approximation of the centrifugal term:

$$\tilde{V}_l(r) = \frac{l(l+1)}{m_x} \frac{\delta^2 e^{-\delta r}}{(1 - e^{-\delta r})^2}.$$

To solve the Schrödinger equation, one can define:

$$x = \alpha_x m_x r \quad , \quad a = \frac{v}{2\alpha_x} \quad , \quad c = \frac{\alpha_x m_x}{\delta}$$

so that:

$$\left(-\frac{\hbar^2}{2\mu} \Delta + V(\mathbf{r}) \right) \Psi(\mathbf{r}) = E \Psi(\mathbf{r}) \implies \left(\frac{d^2}{dr^2} + \kappa^2 - 2\mu V_{eff.}(r) \right) \chi_l(r) = 0$$

Hulthén potential: Schrödinger Equation

$$\left[\frac{d^2}{dx^2} + a^2 + \frac{1}{c} \frac{e^{-x/c}}{1 - e^{-x/c}} - \frac{l(l+1)}{c^2} \frac{e^{-x/c}}{(1 - e^{-x/c})^2} \right] \chi_l(r) = 0.$$

By changing the variable to:

$$t = 1 - e^{-x/c}$$

and reparameterizing the wavefunction:

$$\chi_l(x) = t^{l+1} (1-t)^{iac} f(t)$$

we obtain:

$$\left\{ t(1-t) \frac{d^2}{dt^2} + \left[2 + 2l - (2 + 2l + 2iac + 1)t \right] \frac{d}{dt} - \left[l(l+1) + (l+1)(2iac + 1) - c \right] \right\} f(t) = 0$$

Hulthén potential: Schrödinger Equation

where we can define:

$$\begin{cases} \lambda_{\pm}^l = 1 + l + iac \pm \sqrt{c - a^2 c^2} \\ \omega = 2 + 2l \end{cases}$$

such that:

$$\left\{ t(1-t) \frac{d^2}{dt^2} + [\omega - (\lambda_+^l + \lambda_-^l + 1)t] \frac{d}{dt} - \lambda_+^l \lambda_-^l \right\} f(t) = 0.$$

Thus, the solution is:

$$\chi_l(t) = A t^{l+1} (1-t)^{iac} {}_2F_1(\lambda_+^l, \lambda_-^l, \omega, t)$$

The constant A can be fixed by requiring that the part of $\chi_l(x)$ behaving as an outgoing plane wave $e^{i\mathbf{k}\cdot\mathbf{r}}$ for $r \rightarrow \infty$ has a modulus of 1.

Hulthén potential: Wavefunction Normalization

The wavefunction is:

$$\chi_l(t) = A t^{l+1} (1-t)^{iac} {}_2F_1(\lambda_+^l, \lambda_-^l, \omega, t)$$

The constant A can be fixed by requiring that the part of $\chi_l(x)$ behaving as an outgoing plane wave $e^{i\mathbf{k}\cdot\mathbf{r}}$ for $r \rightarrow \infty$ has a modulus of 1.

Using one of the 5 Bolza formulas for the Hypergeometric function, along with ${}_2F_1(a, b, c, z) \rightarrow 1$ as $z \rightarrow 0$, we obtain the behavior for $t \rightarrow 1$

Hulthén potential: Wavefunction Normalization

$(r \rightarrow \infty)$:

$$\begin{aligned} \chi_l(t) = A t^{l+1} (1-t)^{iac} & \left\{ \frac{\Gamma(\omega)\Gamma(\omega - \lambda_+^l - \lambda_-^l)}{\Gamma(\omega - \lambda_+^l)\Gamma(\omega - \lambda_-^l)} \times \right. \\ & \times {}_2F_1(\lambda_+^l, \lambda_-^l, \lambda_+^l + \lambda_-^l - \omega + 1, 1-t) + \\ & + \frac{\Gamma(\omega)\Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l)\Gamma(\lambda_-^l)} (1-t)^{\omega - \lambda_+^l - \lambda_-^l} \times \\ & \left. \times {}_2F_1(\omega - \lambda_+^l, \omega - \lambda_-^l, 1 + \omega - \lambda_+^l - \lambda_-^l, 1-t) \right\}. \end{aligned}$$

Hulthén potential: Wavefunction Normalization

Note that:

$$\Gamma(\omega - \lambda_+^l - \lambda_-^l) = \left(\Gamma(\lambda_+^l + \lambda_-^l - \omega) \right)^*$$

$$\Gamma(\omega - \lambda_+^l) = \left(\Gamma(\lambda_-^l) \right)^*$$

$$\Gamma(\omega - \lambda_-^l) = \left(\Gamma(\lambda_+^l) \right)^*$$

to rewrite the wavefunction as:

$$\begin{aligned} \chi_l(t) = & A \left(\frac{\Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)} \right)^* t^{l+1} (1-t)^{iac} \times \\ & \times {}_2F_1(\lambda_+^l, \lambda_-^l, \lambda_+^l + \lambda_-^l - \omega + 1, 1-t) + A \left(\frac{\Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)} \right) \times \\ & \times t^{l+1} (1-t)^{-iac} {}_2F_1(\omega - \lambda_+^l, \omega - \lambda_-^l, 1 + \omega - \lambda_+^l - \lambda_-^l, 1-t). \end{aligned}$$

Hulthén potential: Wavefunction Normalization

Taking the limit $t \rightarrow 1$ gives:

$$t^{l+1} \rightarrow 1 \quad , \quad (1-t)^{i\alpha} \rightarrow (e^{-x/c})^{i\alpha}$$

and the wavefunction becomes:

$$\begin{aligned} \chi_l(x) \sim A \left(\frac{\Gamma(\omega)\Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l)\Gamma(\lambda_-^l)} \right)^* e^{-iax} + \\ + A \left(\frac{\Gamma(\omega)\Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l)\Gamma(\lambda_-^l)} \right) e^{iax} \end{aligned}$$

Hulthén potential: Wavefunction Normalization

$$\begin{aligned}\chi_l(x) \sim A & \left| \frac{\Gamma(\omega)\Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l)\Gamma(\lambda_-^l)} \right| \times \\ & \times \left[e^{iax} \exp \left\{ i \arg \left(\frac{\Gamma(\omega)\Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l)\Gamma(\lambda_-^l)} \right) \right\} + \right. \\ & \left. + e^{-iax} \exp \left\{ -i \arg \left(\frac{\Gamma(\omega)\Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l)\Gamma(\lambda_-^l)} \right) \right\} \right].\end{aligned}$$

The constant A is determined as:

$$A = \frac{1}{2} \left| \frac{\Gamma(\lambda_+^l)\Gamma(\lambda_-^l)}{\Gamma(\omega)\Gamma(\lambda_+^l + \lambda_-^l - \omega)} \right|.$$

The asymptotic behavior of $\chi_l(x)$ can be rewritten as:

$$\chi_l(x) \sim \cos \left\{ ax + \arg \left(\frac{\Gamma(\omega)\Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l)\Gamma(\lambda_-^l)} \right) \right\}$$

Hulthén potential: Wavefunction Normalization

using $\cos \alpha = \sin \left(\alpha + \frac{\pi}{2} \right)$ and $\arg(zw) = \arg z + \arg w$, it becomes:

$$\chi_l(x) \underset{x \rightarrow \infty}{\sim} \sin \left\{ ax + \arg \left(\frac{i\Gamma(\omega)\Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l)\Gamma(\lambda_-^l)} \right) \right\}.$$

Where we can notice $ax = kr$.

Hulthén potential: The cross section

This leads to:

$$\chi_l(t) = \frac{1}{2} \left| \frac{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)}{\Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)} \right| t^{l+1} (1-t)^{iac} {}_2F_1(\lambda_+^l, \lambda_-^l, \omega, t)$$
$$\chi_l(x) \underset{x \rightarrow \infty}{\sim} \sin \left\{ kr + \arg \left(\frac{i \Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)} \right) \right\}.$$

Knowing the expected behavior from the theory:

$$\chi_l(r) \underset{r \rightarrow \infty}{\sim} \sin \left(kr - \frac{\pi}{2} l + \delta_l \right)$$

one can determine δ_l by computing:

$$\phi_l = \arg \left(\frac{i \Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)} \right) - \arg \left(\frac{i \Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)} \right) \Big|_{\alpha_x \rightarrow 0}.$$



Hulthén potential: The cross section

This yields the phase shifts:

$$\delta_l = \arg \left(\frac{i\Gamma(\omega)\Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l)\Gamma(\lambda_-^l)} \right)$$

and the cross-section:

$$\sigma_l = \frac{16\pi}{m_x^2 v^2} \sum_{l=0}^{\infty} (2l+1) \sin^2 \left\{ \arg \left(\frac{i\Gamma(\omega)\Gamma(\lambda_+^l + \lambda_-^l - \omega)}{\Gamma(\lambda_+^l)\Gamma(\lambda_-^l)} \right) \right\}.$$



Hulthén potential: Wavefunction Behavior as $x \rightarrow 0$

Starting from:

$$\chi_l(t) = \frac{1}{2} \left| \frac{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)}{\Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)} \right| t^{l+1} (1-t)^{iac} {}_2F_1(\lambda_+^l, \lambda_-^l, \omega, t)$$

we can expand it as $x \rightarrow 0$, i.e., as $t = 1 - e^{-x/c} \rightarrow 0$, to find:

$$\chi_l(r) \rightarrow B_l \frac{(\kappa r)^{l+1}}{(2l+1)!!}$$

and specifically the coefficient B_l , which represents the distortion introduced by the potential $V_H(r)$.

Hulthén potential: Wavefunction Behavior as $x \rightarrow 0$

We have:

$$\begin{aligned}\chi_l(x) &\underset{x \rightarrow 0}{\sim} \frac{1}{2} \left| \frac{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)}{\Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)} \right| (1 - e^{-x/c})^{l+1} \cdot 1 \\ &= \frac{1}{2} \left| \frac{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)}{\Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)} \right| \left(1 - \left(1 - \frac{x}{c} + \dots \right) \right)^{l+1} \\ &= \frac{1}{2} \left| \frac{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)}{\Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)} \right| \left(\frac{x}{c} \right)^{l+1}\end{aligned}$$

with the defined parameters, we have $\frac{x}{c} = km_\phi r$, where $k = \pi^2/6$, so:

$$\chi_l(x) \underset{x \rightarrow 0}{\sim} (km_\phi r)^{l+1} \frac{1}{2} \left| \frac{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)}{\Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)} \right|.$$

Hulthén potential: Wavefunction Behavior as $x \rightarrow 0$

Using the parameter definitions again, it can be seen that:

$$\kappa r = \frac{m_x}{2} v r \quad , \quad k m_\phi r = k m_\phi \frac{m_x v}{2} \frac{2}{m_x v} r = \frac{2 k m_\phi}{m_x v} \kappa r = \frac{\kappa r}{a c}$$

allowing us to rewrite:

$$\begin{aligned} \chi_l(x) &\sim \frac{(k m_\phi)^{l+1}}{(\kappa)^{l+1}} (\kappa r)^{l+1} \frac{1}{2} \left| \frac{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)}{\Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)} \right| \\ &\sim \frac{(2l+1)!!}{(ac)^{l+1}} \frac{1}{2} \left| \frac{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)}{\Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)} \right| \frac{(\kappa r)^{l+1}}{(2l+1)!!} . \end{aligned}$$

Hulthén potential: Wavefunction Behavior as $x \rightarrow 0$

Comparing:

$$\chi_l(x) \sim \frac{(2l+1)!!}{(ac)^{l+1}} \frac{1}{2} \left| \frac{\Gamma(\lambda_+^l)\Gamma(\lambda_-^l)}{\Gamma(\omega)\Gamma(\lambda_+^l + \lambda_-^l - \omega)} \right| \frac{(\kappa r)^{l+1}}{(2l+1)!!}$$
$$\chi_l(r) \rightarrow B_l \frac{(\kappa r)^{l+1}}{(2l+1)!!}$$

it follows that:

$$B_l = \frac{(2l+1)!!}{(ac)^{l+1}} \frac{1}{2} \left| \frac{\Gamma(\lambda_+^l)\Gamma(\lambda_-^l)}{\Gamma(\omega)\Gamma(\lambda_+^l + \lambda_-^l - \omega)} \right|.$$

To determine the Sommerfeld factor:

$$S = \frac{|B_l|^2}{|B_l(c \rightarrow 0, ac = u)|^2}$$

Hulthén potential: Wavefunction Behavior as $x \rightarrow 0$

we need to compute $B_l(c \rightarrow 0, ac = u) = B_l(V \rightarrow 0)$, where we have set $ac = \frac{m_x v}{2km_\phi} = u$. We have:

$$\begin{aligned} B_l &= \frac{(2l+1)!!}{(ac)^{l+1}} \frac{1}{2} \left| \frac{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)}{\Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)} \right| \\ &= \frac{(2l+1)!!}{2u^{l+1}} \left| \frac{\Gamma(1+l+iu+ac\sqrt{\frac{1}{a^2c}-1}) \Gamma(1+l+iu-ac\sqrt{\frac{1}{a^2c}-1})}{\Gamma(2l+2) \Gamma(2iu)} \right| \end{aligned}$$

taking the limit $c \rightarrow 0$ and $a \rightarrow \infty$:

$$B_l = \frac{(2l+1)!!}{2u^{l+1}} \left| \frac{\Gamma(1+l+iu+iu) \Gamma(1+l+iu-iu)}{\Gamma(2l+2) \Gamma(2iu)} \right|$$

which ultimately yields:

$$B_l(c \rightarrow 0, ac = u) = \frac{(2l+1)!!}{2(ac)^{l+1}} \left| \frac{\Gamma(1+l+2iac) \Gamma(1+l)}{\Gamma(\omega) \Gamma(2iac)} \right|.$$

Hulthén potential: the Sommerfeld factor

Having determined the two distortion coefficients:

$$B_l = \frac{(2l+1)!!}{(ac)^{l+1}} \frac{1}{2} \left| \frac{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)}{\Gamma(\omega) \Gamma(\lambda_+^l + \lambda_-^l - \omega)} \right|$$
$$B_l(c \rightarrow 0, ac = u) = \frac{(2l+1)!!}{2(ac)^{l+1}} \left| \frac{\Gamma(1+l+2iac) \Gamma(1+l)}{\Gamma(\omega) \Gamma(2iac)} \right|$$

the Sommerfeld factor results in:

$$S_l = \frac{|B_l|^2}{|B_l(c \rightarrow 0, ac = u)|^2}$$
$$= \left(\frac{(2l+1)!!}{2(ac)^{l+1}} \left| \frac{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)}{\Gamma(\omega) \Gamma(2iac)} \right| \right)^2 \left(\frac{2(ac)^{l+1}}{(2l+1)!!} \left| \frac{\Gamma(\omega) \Gamma(2iac)}{\Gamma(1+l+2iac) \Gamma(1+l)} \right| \right)^2$$
$$= \left| \frac{\Gamma(\lambda_+^l) \Gamma(\lambda_-^l)}{\Gamma(1+l) \Gamma(1+l+2iac)} \right|^2.$$

Hulthén potential: the Sommerfeld factor

which, by utilizing:

$$|\Gamma(i\alpha)|^2 = \frac{\pi}{\alpha \sinh(\pi\alpha)} \quad , \quad |\Gamma(1+i\alpha)|^2 = \frac{\pi\alpha}{\sinh(\pi\alpha)} \quad , \quad \alpha \in \mathbb{R}$$

can be rewritten as:

$$S_l = \left| \frac{(1+iac + \sqrt{c-a^2c^2})_l (1+iac - \sqrt{c-a^2c^2})_l}{\Gamma(1+l)(1+2iac)_l} \right|^2 \times \\ \times \left| \frac{\Gamma(1+iac + i\sqrt{a^2c^2-c})\Gamma(1+iac - i\sqrt{a^2c^2-c})}{\Gamma(1+2iac)} \right|^2 .$$

Hulthén potential: the Sommerfeld factor

$$S_l = \left| \frac{(1 + iac + \sqrt{c - a^2 c^2})_l (1 + iac - \sqrt{c - a^2 c^2})_l}{\Gamma(1 + l)(1 + 2iac)_l} \right|^2 \times \\ \times \frac{\pi(ac + \sqrt{a^2 c^2 - c})\pi(ac - \sqrt{a^2 c^2 - c})}{\sinh(\pi(ac + \sqrt{a^2 c^2 - c}))\sinh(\pi(ac - \sqrt{a^2 c^2 - c}))} \cdot \frac{\sinh(2\pi ac)}{\pi(2ac)}.$$

We can make use of:

$$\sinh x \sinh y = \frac{1}{2} (\cosh(x + y) - \cosh(x - y))$$



to obtain the final result:

$$S_l = \left| \frac{(1 + iac + \sqrt{c - a^2 c^2})_l (1 + iac - \sqrt{c - a^2 c^2})_l}{\Gamma(1 + l)(1 + 2iac)_l} \right|^2 \times \\ \times \frac{\pi}{a} \frac{\sinh(2\pi ac)}{\cosh(2\pi ac) - \cosh(2\pi\sqrt{a^2 c^2 - c})}.$$

Hulthén potential: the Sommerfeld factor

The cases for $l = 0$ and $l = 1$ are:

$$S_0 = \frac{\pi}{a} \frac{\sinh(2\pi ac)}{\cosh(2\pi ac) - \cosh(2\pi\sqrt{a^2c^2 - c})} \quad , \quad S_1 = \frac{(1-c)^2 + 4a^2c^2}{1 + 4a^2c^2} S_0$$



Hulthén potential: the Sommerfeld factor

The resonant values for S_l can be determined:

$$S_l = \left| \frac{(1 + iac + \sqrt{c - a^2 c^2})_l (1 + iac - \sqrt{c - a^2 c^2})_l}{\Gamma(1 + l)(1 + 2iac)_l} \right|^2 \times \\ \times \frac{\pi \sinh(2\pi ac)}{a \cosh(2\pi ac) - \cosh(2\pi \sqrt{a^2 c^2 - c})}$$





considering the low-velocity limit $v \rightarrow 0$ ($a \rightarrow 0$). Resonance occurs for:

$$m_x = \frac{\pi^2 m_\phi}{6\alpha_x} n^2, \quad n \in \mathbb{N}.$$





around which the behavior of S is:

$$S = \frac{4\alpha_x k m_\phi}{m_x v^2}.$$

Hulthén potential: plots of the S Factor and σ for $l=0$



Hulthén potential: plots of the S Factor and σ for $l=1$



- [1] Nima Arkani-Hamed et al. “A Theory of Dark Matter”. en. In: *Physical Review D* 79.1 (Jan. 2009). arXiv:0810.0713 [hep-ph], p. 015014. ISSN: 1550-7998, 1550-2368. DOI: [10.1103/PhysRevD.79.015014](https://doi.org/10.1103/PhysRevD.79.015014).
- [2] S. Cassel. “Sommerfeld factor for arbitrary partial wave processes”. en. In: *Journal of Physics G: Nuclear and Particle Physics* 37.10 (Oct. 2010). arXiv:0903.5307 [hep-ph], p. 105009. ISSN: 0954-3899, 1361-6471. DOI: [10.1088/0954-3899/37/10/105009](https://doi.org/10.1088/0954-3899/37/10/105009). URL: <http://arxiv.org/abs/0903.5307>.
- [3] Marco Cirelli, Alessandro Strumia, and Jure Zupan. *Dark Matter*. en. arXiv:2406.01705 [hep-ph]. July 2024. URL: <http://arxiv.org/abs/2406.01705>.

- [4] Francesco Foggetti. “Correlazioni tra diversi tipi di osservazioni nella ricerca di materia oscura”. it. Master thesis. Università del Salento, 2016.
- [5] Ayuki Kamada, Takumi Kuwahara, and Ami Patel. “Quantum Theory of Dark Matter Scattering”. en. In: *Journal of High Energy Physics* 2023.11 (Nov. 2023). arXiv:2303.17961 [hep-ph], p. 105. ISSN: 1029-8479. DOI: [10.1007/JHEP11\(2023\)105](https://doi.org/10.1007/JHEP11(2023)105). URL: <http://arxiv.org/abs/2303.17961>.
- [6] L.D. Landau and E.M. Lifšic. *Meccanica quantistica: teoria non relativistica*. Fisica teorica, Lev D. Landau, Evgenij M. Lifšits. Editori Riuniti, 1976.

- [7] Massimiliano Lattanzi and Joseph Silk. “Can the WIMP annihilation boost factor be boosted by the Sommerfeld enhancement?” en. In: *Physical Review D* 79.8 (Apr. 2009). arXiv:0812.0360 [astro-ph], p. 083523. ISSN: 1550-7998, 1550-2368. DOI: [10.1103/PhysRevD.79.083523](https://doi.org/10.1103/PhysRevD.79.083523). URL: <http://arxiv.org/abs/0812.0360>.
- [8] S.C.A. Nierop. “The Sommerfeld Enhancement”. en. Bachelor thesis. University of Groningen, 2009.
- [9] C. Rossetti. *Metodi matematici della fisica*. Levrotto & Bella, 2000. ISBN: 9788882180607.
- [10] C. Rossetti. *Rudimenti di meccanica quantistica*. Levrotto & Bella, 2011. ISBN: 9788882181512.

- [11] Sean Tulin, Hai-Bo Yu, and Kathryn M. Zurek. “Beyond Collisionless Dark Matter: Particle Physics Dynamics for Dark Matter Halo Structure”. en. In: *Physical Review D* 87.11 (June 2013). arXiv:1302.3898 [hep-ph], p. 115007. ISSN: 1550-7998, 1550-2368. DOI: [10.1103/PhysRevD.87.115007](https://doi.org/10.1103/PhysRevD.87.115007). URL: <http://arxiv.org/abs/1302.3898>.
- [12] B. Zwiebach. *Mastering Quantum Mechanics: Essentials, Theory, and Applications*. MIT Press, 2022. ISBN: 9780262366892.