

Grothendieck's Galois Theory

Gabriele Rastello

October 8, 2020

1 Group actions and adjoints

Definition 1.1. In a category \mathcal{C} an arrow $f: X \rightarrow Y$ is a **strict epimorphism** if it is the joint coequalizer of all the arrows it coequalizes. This means that any arrow $g: X \rightarrow Z$ such that $g \circ x = g \circ y$ for any $x, y: C \rightarrow X$ such that $f \circ x = f \circ y$ there exists a unique arrow $h: Y \rightarrow Z$ such that $h \circ f = g$. Refer to Figure 1.1.




Figure 1.1

Remark 1.2. Strict epimorphisms are coequalizers, thus epimorphisms (as the name implies).

Remark 1.3. If an arrow is both a strict epimorphism and a monomorphism then it is an isomorphism.

Definition 1.4. Let H be a group, A an object of \mathcal{C} and $G = \text{Aut}(A)$ the group of automorphisms of A in \mathcal{C} i.e. the group whose underlying set is the set of isomorphisms of type $A \rightarrow A$ of \mathcal{C} and whose operation is composition in \mathcal{C} . An **action** of H on A is a group homomorphism $H \rightarrow G$.

Notation 1.5. Given an action of a group H on an object A of \mathcal{C} we denote, with a slight abuse of notation, the automorphism of A associated to $h \in H$ by the same symbol h .

Definition 1.6. If H acts on A as defined in 1.4 we define the quotient of A by H in \mathcal{C} to be an element A/H of \mathcal{C} equipped with an arrow $q: A \rightarrow A/H$ such that:

- (1) for all $h \in H$ $q \circ h = q$ holds,
- (2) for any $x: A \rightarrow X$ such that $x \circ h = x$ for all $h \in H$ there exists a unique arrow $\varphi: A/H \rightarrow X$ such that $x = \varphi \circ q$.

See also Figure 1.2.

Remark 1.7. Quotients are defined by a universal property, thus are unique up to unique isomorphism and we can speak of “the” quotient of A by H instead of “a” quotient of A by H .

Notation 1.8. Sometimes we use the sentence “the quotient of A by H ” to refer to the object A/H , some others to the arrow $q: A \rightarrow A/H$; the context should be enough to differentiate between the two cases.




Figure 1.2

Remark 1.9. Consider a quotient $q: A \rightarrow A/H$; by condition (1) above $q \circ h = q = q \circ 1_A$ so q coequalizes all the pairs $(h, 1_A)$, for $h \in H$. If another arrow $x: A \rightarrow X$ coequalizes all the pairs that q does then this arrow is such that $x \circ h = x \circ 1_A = x$ for all $h \in H$ and thus, by condition (2), we have a unique factorization $x = \varphi \circ q$. This proves that all quotients are strict epimorphisms.

Remark 1.10. Let G be a group, \mathbf{GSet} the category of G -sets and G -invariant maps and A an object of \mathbf{GSet} . In this category Definition 1.6 yields the familiar notion of the set of all orbits of an action: A/G is the set of orbits of A .

Remark 1.11. Consider again \mathbf{GSet} . The underlying set of G (that we also denote as G) is a G -set with the action given by left multiplication in G ; we call this the **canonical action** of G on itself. Let $\varphi: G \rightarrow E$ be a G -invariant map; it is easy to see that such a φ , by virtue of being G -invariant, is determined uniquely by the value $\varphi(e)$, where e is the neutral element of G .

Let now E be a transitive G -set i.e. a set upon which the action of G is transitive i.e. such that $E/G = \{*\}$. Fix an $x \in E$ and let φ_x be the G -invariant map defined by $\varphi_x(e) = x$; we argue that $\varphi_x: G \rightarrow E$ makes E into a quotient of G by the subgroup

$$H = \text{Fix}(x) = \{g \in G: gx = x\}.$$

Indeed by using the definition of H and the fact that φ is G -invariant we have

$$(\varphi \circ h)(e) = \varphi(h(e)) = h(\varphi(e)) = h(x) = x.$$

for all $h \in H$. Moreover let $g: G \rightarrow F$ satisfy (1) of Definition 1.6; as we discussed above g is entirely determined by the image of e so we obtain (2) defining an arrow $f: E \rightarrow F$ by $f(x) = g(e)$. The situation is depicted in Figure 1.3.

Trivially G is a transitive G -set and for any $g \in G$ $G/\text{Fix}(g)$ is transitive as well so we have that an object $E \in \mathbf{GSet}$ is transitive if and only if it is isomorphic to G/H where $H = \text{Fix}(x)$ for (any) $x \in E$.




Figure 1.3

For the rest of the section fix a category \mathcal{C} , an object $A \in \mathcal{C}$ and let $G = \text{Aut}(A)$.

Proposition 1.12. Consider a subgroup $H \leq G$ and an object $X \in \mathcal{C}$. H acts on the hom-set $[A, X]$ as follows¹:

$$\begin{aligned}
H \times [A, X] &\longrightarrow [A, X] \\
(h, x) &\longmapsto h \cdot x = x \circ h.
\end{aligned}$$

Remark 1.13. Assume that the action $G \times [A, X] \rightarrow [A, X]$ is transitive and let \mathbf{GSet}^t be the category of transitive G -sets (a subcategory of $G\text{set}$). Then we have a functor

$$\begin{array}{ccc}
[A, -]_G: \mathcal{C} & \longrightarrow & \mathbf{GSet}^t \\
X & & [A, X]_G \\
\downarrow f & \longmapsto & \downarrow f_* \\
Y & & [A, Y]_G
\end{array}$$

where we indicate with $[A, X]_G$ the hom-set $[A, X]$ upon which G acts as described in Proposition 1.12 and f_* is post-composition with f . It is easy to check that f_* is indeed G -invariant.

Remark 1.14. Consider an object $E \in \mathbf{GSet}^t$, pick an element $x_0 \in E$ and let $H = \text{Fix}(x_0) \leq G$ (the choice of x_0 is irrelevant as E is transitive). Moreover assume that \mathcal{C} has quotients of A by any subgroup of G .

By what we observed in Remark 1.11 we have a bijection between elements of E and arrows of type $G \rightarrow E$. Consider then $f \in [A, X]$ and its corresponding arrow $\varphi: G \rightarrow [A, X]_G$; we claim that f factors through A/H if and only if φ factors through $E \cong G/H$ (see Figure 1.4). Indeed f factors if and only if $f \circ h = f$ for all $h \in H$, by using the fact that φ is G -invariant we obtain

$$\varphi(h(e)) = h(\varphi(e)) = h(f) = f \circ h = f$$

and, since φ is uniquely determined by $\varphi(e)$, $\varphi \circ h = \varphi$ for all $h \in H$; this happens if and only if φ factors through $E \cong G/H$.

¹Since an action as of Definition 1.4 is a map that sends elements of a group to arrows it is, in this case, equivalent to give the definition of an action by uncurrying.

This gives us, for each $X \in \mathcal{C}$ and $E \in \mathbf{GSet}^t$, a bijection

$$[A/H, X] \cong \mathbf{GSet}^t(E, [A, X]_G)$$

natural in X in E (see Section 2). Thus, if \mathcal{C} has quotients of A by subgroups of G we have an adjunction $A \times_G - \dashv [A, -]_G$ where $A \times_G E = A/H$ for $H = \text{Fix}(x_0)$ and $x_0 \in E$ as above².




Figure 1.4

Our main problem will be that of finding conditions on \mathcal{C} that make this adjunction into an equivalence of categories.

2 Appendixes

2.1 Naturality of $[A/H, X] \cong \mathbf{GSet}^t(E, [A, X]_G)$

Fix a category \mathcal{C} , an element $A \in \mathcal{C}$ and let $G = \text{Aut}(A)$. Let's indicate with ψ_{EX} the bijection between $[A/H, X]$ and $\mathbf{GSet}^t(E, [A, X]_G)$ described in Remark 1.14; we shall prove that it is natural in both $X \in \mathcal{C}$ and $E \in \mathbf{GSet}^t$.

Naturality in X . Given an arrow $f: X \rightarrow Y$ of \mathcal{C} we want to prove that ψ_{EX} are the components of a natural transformation $[A/H, -] \Rightarrow \mathbf{GSet}^t(E, [A, -]_G)$ i.e that the following diagram commutes.

$$\begin{array}{ccc} [A/H, X] & \xrightarrow{\psi_{EX}} & \mathbf{GSet}^t(E, [A, X]_G) \\ \downarrow f_* & & \downarrow (f_*)_* \\ [A/H, Y] & \xrightarrow{\psi_{EY}} & \mathbf{GSet}^t(E, [A, Y]_G) \end{array}$$

Recall that here $H = \text{Fix}(x_0)$ for $x_0 \in E$. Pick any $x \in [A/H, X]$ and let $q: A \rightarrow A/H$ be the quotient arrow (the quotient exists because \mathcal{C} is assumed to have all quotients by subgroups of G); then chasing x through the diagram down the two possible ways yields two arrows in \mathbf{GSet}^t of type $E \rightarrow [A, Y]_G$. Since E is transitive we have $E \cong G/H$ and this means that arrows of type $E \rightarrow [A, Y]_G$ are determined uniquely by the image of $e \in E \cong G/H$ (refer to the right part of Diagram 1.4). Keeping this in mind the following computations shows that the square commutes.

$$((f_*)_* \circ \psi_{EX})(x)(e) = (f_* \circ \psi_{EX}(x))(e) = f_*(\psi_{EX}(x)(e)) = f_*(x \circ q) = f \circ x \circ q,$$

²We use the notation $A \times_G -$ for the left adjoint in "honor" of the tensor-hom adjunction.

$$(\psi_{EY} \circ f_*)(x)(e) = \psi_{EX}(f \circ x)(e) = f \circ x \circ q.$$

□

Naturality in E. We want to prove that ψ_{EX} are the components of a natural transformation $[A/-, X] \Rightarrow \mathbf{GSet}^t(-, [A, X]_G)$, but before drawing the naturality square as above we shall note that this case is slightly complicated by the fact that it is not immediately clear how the functor $[A/-, X]$ acts on arrows. Indeed let $f: E \cong G/H \rightarrow F \cong G/H'$ be an arrow of \mathbf{GSet}^t with $H = \text{Fix}(x_0)$, $H' = \text{Fix}(f(x_0))$ and $x_0 \in E$; these choices are justified by the fact that both E and F are transitive G -sets. Now notice that, for $h \in H$, $f(h \cdot x_0) = f(x_0)$ by definition of H and $f(h \cdot x_0) = h \cdot f(x_0)$ by G -invariance of f ; thus $h \in H'$ and thus $H \subset H'$. This means that if we let $q: A \rightarrow A/H, q': A \rightarrow A/H'$ be the quotients in \mathcal{C} then there is a unique arrow $\tilde{f}: A/H \rightarrow A/H'$ such that $\tilde{f} \circ q = q'$. The functor $[A/-, X]$ then sends the arrow $f: E \rightarrow F$ to $\tilde{f}^*: [A/H', X] \rightarrow [A/H, X]$.

Now we want to prove the naturality of the following square.

$$\begin{array}{ccc} [A/H, X] & \xrightarrow{\psi_{EX}} & \mathbf{GSet}^t(E, [A, X]_G) \\ \tilde{f}^* \uparrow & & \uparrow f^* \\ [A/H', X] & \xrightarrow{\psi_{FX}} & \mathbf{GSet}(F, [A, X]_G) \end{array}$$

As before we chase a $x \in [A/H', X]$ down the two possible paths to get two arrows of type $E \rightarrow [A, X]_G$ and prove that they are the same by evaluating them on $e \in E$:

$$(\psi_{EX} \circ \tilde{f}^*)(x)(e) = \psi_{EX}(x \circ \tilde{f})(e) = x \circ \tilde{f} \circ q = x \circ q',$$

$$(f^* \circ \psi_{FX})(x)(e) = (\psi_{FX}(x) \circ f)(e) = \psi_{FX}(x)(f(e)) = \psi_{FX}(x)(e) = x \circ q'.$$

Note that we used that $f(e) = e$, this holds because the action of G on $G/H \cong E$ and $G/H' \cong F$ is just left multiplication and f is G -invariant. □