

NICHOLS ALGEBRAS OVER FINITE GROUPS:

NEW & OLD RESULTS

Examples: $S(V)$, $\wedge V$, $U_q(n)$

Ingredient:

Braided vector space


(V, c)

V vector space / k

$c \in GL(V^{\otimes 2})$

braiding, i.e.

$$(c \otimes id)(id \otimes c)(c \otimes id) = (id \otimes c)(c \otimes id)(id \otimes c)$$



Braid group

$$B_n = \langle \sigma_i, i=1, \dots, n-1 \mid \begin{array}{l} \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \\ \sigma_i \sigma_j = \sigma_j \sigma_i \quad |i-j| > 1 \end{array} \rangle$$

c gives rise to a representation $\forall n \geq 2$

$$\rho_n: B_n \longrightarrow GL(V^{\otimes n})$$

$$\sigma_i \longmapsto id^{\otimes i-1} \otimes c \otimes id^{n-i-3}$$

Matsumoto: $\exists M: \mathcal{S}_n \rightarrow B_n$.

The vector space

$$\bigoplus_{n \geq 0} V^{\otimes n}$$

$$V^{\otimes 0} = K$$

Supports 2 graded algebra structures:

$$T_! V \cong K \langle x_1, \dots, x_{\dim V} \rangle \text{ tensor algebra}$$

$T_* V$ cotensor algebra:

$$(V_1 \otimes \dots \otimes V_m) \cdot_* (V_{m+1} \otimes \dots \otimes V_{m+n})$$

$$:= \sum_{\sigma \in \bigsqcup_{m,n} \mathcal{L}_{m,n}} f_{\text{sign}}(\sigma) (v_1 \otimes \dots \otimes v_{m+n})$$

shuffles


$$\mathcal{L}_{m,n} = \left\{ \sigma \in \mathcal{S}_{m+n} \text{ st } \begin{array}{l} \sigma(i) < \sigma(j) \text{ if } 1 \leq i < j \leq m \\ \text{or} \\ m+1 \leq i < j \leq m+n \end{array} \right\}$$

$$T_! V \xrightarrow{\exists! \Omega \text{ algmap extending id}_V} T_* V$$

$$T_! V \searrow \quad \swarrow T_* V$$

$T_{!*} V = \text{NICHOLS ALGEBRA}$

eg $m=2$ $n=1$



id



(123)



(23)

$$(V_1 \otimes V_2) \cdot_* V_3 = V_1 \otimes V_2 \otimes V_3 + H(123) \cdot (V_1 \otimes V_2 \otimes V_3)$$

$$(id \otimes c)(V_1 \otimes V_2 \otimes V_3) =$$

$$= V_1 \otimes V_2 \otimes V_3 + (c \otimes id)(id \otimes c)(V_1 \otimes V_2 \otimes V_3) \\ + (id \otimes c)(V_1 \otimes V_2 \otimes V_3)$$

Why do we care?

CLASSIFICATION OF HOPF ALGEBRAS

Andruskiewitsch
Schneider

MAUE'S CONJECTURE

2017

Ellenberg-Tran-
Westerland

QUESTIONS:

- For which (V, c)
is $T_{1*}(V)$ finite dim?
- For which (V, c)
is $T_{1*}(V)$ finitely presented?

SPECIAL FAMILY OF BYS:

BVS of Group type

Given:

G group

$$V \in {}_{\mathbb{K}G}\text{-mod}$$

$$V = \bigoplus_{g \in G} V_g$$

$$h \cdot V_g \subseteq V_{hgh^{-1}}$$

} \mathcal{YD}
module

Then \exists

$$c: V \otimes V \rightarrow V \otimes V$$

$$v \otimes w \mapsto g \cdot w \otimes v, \quad v \in V_g$$

ANSWERS:

G abelian:
full answer

Andruskiewitsch,
Schneider,
Heckenberger,
Angiono, ...

G non-abelian: open

Examples of fd^e Nichols algebras are rare

CONJECTURE

G simple
nonabelian

\Rightarrow ~~\exists~~ finite dim^e Nichols
algebras over G

The conjecture is confirmed for:

- ALTERNATING GROUPS

- SPORADIC GROUPS

(up to a few classes in F_{22} , B, M)

Andruskiewitsch

Fantino,

Graña, Vendramin, 2011

Beltrán Cubillos, 2020

- SIMPLE GROUPS OF LIE TYPE

$$q = p^m$$

• $PSL_n(q)$: $n > 3$ or $n = 3, q > 2$

• $PSp_{2n}(q)$: $n > 2$ or $n = 2, q > 3$

• $P\Omega_{4n}^+(2^m)$

• $E_7(2^m), E_8(2^m), F_4(2^m), G_2(2^m)$

• ${}^{(3)}D_4(2^m), P\Omega_{4n}^-(2^m)$

Andruskiewitsch - C. García
2015-2023

- SUZUKI AND REE GROUPS C. - Costantini, 2021


For the remaining groups: if $T_{!*}(V)$

is finite dimensional \Rightarrow support of the

grading of V is a conjugacy class of

elements of order coprime to p (A-C-G)

It can be useful to consider more general BVS



How? (Graña)
RACKS + COCYCLES

\cong isomorphic

\sim in Hilbert series

RACK

X set, with binary operation $: X \times X \xrightarrow{\triangleright} X$

such that: ① $x \triangleright -$ bijective $\forall x$

$$\textcircled{2} \quad x \triangleright (y \triangleright z) = (x \triangleright y) \triangleright (x \triangleright z) \quad \forall x, y, z$$

EXAMPLE $X \subseteq G$ conj classe

$$\triangleright = \text{conjugation}$$

$$x \triangleright y = xyx^{-1} \text{ for } x, y \in X$$

RACK COCYCLE

$$q: X \times X \rightarrow \mathbb{C}^n$$

$$q(x, y \triangleright z) q(y, z) = q(x \triangleright y, x \triangleright z) q(x, z)$$

A pair (X, q) gives a BVS:

$$V = kX \otimes \mathbb{C}^n, \quad c_q \in GL(V^{\otimes 2})$$

$$c_q((x \otimes v), (y \otimes w)) = ((x \triangleright y) q(x, y) w) \otimes (x \otimes v)$$

EXAMPLE : $G = S_n$ $n \geq 3$
 $X = \mathcal{O} = \{(i j) \mid i \neq j\}$

$$q^+(\tau, (i j)) = \begin{cases} 1 & \text{if } (\tau(i) - \tau(j))(i - j) > 0 \\ -1 & \text{otherwise} \end{cases}$$

- q^+ is of group type (Milinski-Schneider)

$$\mathcal{O} = \mathcal{O}_{(12)} \quad C_{S_n}(12) \cong S_2 \times S_{n-2}$$

$$\rho = \text{sgn} \boxtimes \text{id} \quad \text{on } \mathbb{C}$$

$$V = \text{Ind}_{C_{S_n}(12)}^{S_n} (\rho) = \mathbb{C} S_n \otimes_{\mathbb{C} C_{S_n}(12)} \mathbb{C}_\rho$$

$$\text{grading: } V_{\sigma(12)\sigma^{-1}} = \mathbb{C} \sigma \otimes_{\mathbb{C} C_{S_n}(12)} \mathbb{C}_\rho$$

- $q^- \equiv -1$

q^- is NOT of group type

FOMIN - KIRILLOV ALGEBRAS:

(MOTIVATION: SCHUBERT CALCULUS)

For an integer $n \geq 3$ denote by \mathcal{E}_n the algebra (of type A_{n-1}) with generators $x_{(ij)}$, where $1 \leq i < j \leq n$, and relations

$$x_{(ij)}^2 = 0,$$

$$x_{(ij)}x_{(jk)} = x_{(jk)}x_{(ik)} + x_{(ik)}x_{(ij)},$$

$$x_{(jk)}x_{(ij)} = x_{(ik)}x_{(jk)} + x_{(ij)}x_{(ik)},$$

$$x_{(ij)}x_{(kl)} = x_{(kl)}x_{(ij)},$$

$$\text{for } 1 \leq i < j \leq n,$$

$$\text{for } 1 \leq i < j < k \leq n,$$

$$\text{for } 1 \leq i < j < k \leq n,$$

$$\text{for any distinct } i, j, k, l.$$

These are the degree 2 relations of $T_{!*}(\mathbb{Q}, q^+)$!!

• For $n=3,4,5$

$$\mathcal{E}_n \cong T_{!*}(\mathbb{Q}, q^+) \otimes \text{fd}^p$$

Milinski-Schneider, F-K, Garcia-Garcia Iglesias, Graña, Roos

$n \geq 6$??

$$\text{Vendramin: } T_{!*}(\mathbb{Q}, q^+) \sim T_{!*}(\mathbb{Q}, -1)$$

Bazlov generalized the construction of FK algebras to arbitrary finite Coxeter groups

G. Harel - C. : Vendramin's result holds for all Bazlov's algebras

RK: Hopefully the pair $(\mathbb{Q}, -1)$ can be attacked by geometric methods using

Kapranov-Schectman, C-Esposito-Rubio y Degrossi