

# REPRESENTATIONS OF $G$ -POSETS AND CANONICAL BRAUER INDUCTION

Robert Boltje (joint work with Nariel Monteiro)

University of California, Santa Cruz

Groups and their actions: algebraic, geometric and combinatorial aspects. Levico Terme, June 3–7, 2024

## Overview

- 1  $G$ -posets and their representations
- 2 The canonical Brauer induction formula
- 3 Categorification of the canonical Brauer induction formula

## 1. $G$ -posets and their representations

- $G$ : finite group.
- $k$ : commutative ring.

## 1. $G$ -posets and their representations

- $G$ : finite group.
- $k$ : commutative ring.

**Definition** A  $G$ -poset  $X$  is a poset  $(X, \leq)$  on which  $G$  acts via poset automorphisms, i.e., if  $x \leq y$  then  $gx \leq gy$ . For  $x \in X$ ,  $G_x$  denotes the stabilizer.

## 1. $G$ -posets and their representations

- $G$ : finite group.
- $k$ : commutative ring.

**Definition** A  $G$ -poset  $X$  is a poset  $(X, \leq)$  on which  $G$  acts via poset automorphisms, i.e., if  $x \leq y$  then  $gx \leq gy$ . For  $x \in X$ ,  $G_x$  denotes the stabilizer.

**Example** The set of subgroups of  $G$  together with the conjugation action of  $G$ .

If  $X$  is a  $G$ -poset, one can form a category  $\mathcal{C}(X)$  as follows:

- Objects: the elements of  $X$ .
- $\text{Hom}_{\mathcal{C}(X)}(x, y) := \{g \in G \mid x \leq gy\}$ .
- Composition:  $x \xrightarrow{g} y \xrightarrow{h} z = x \xrightarrow{gh} z$   
( $x \leq gy, y \leq hz \Rightarrow x \leq gy \leq g(hz) = (gh)z$ ).
- $\text{id}_x = x \xrightarrow{1} x$ .

If  $X$  is a  $G$ -poset, one can form a category  $\mathcal{C}(X)$  as follows:

- Objects: the elements of  $X$ .
- $\text{Hom}_{\mathcal{C}(X)}(x, y) := \{g \in G \mid x \leq gy\}$ .
- Composition:  $x \xrightarrow{g} y \xrightarrow{h} z = x \xrightarrow{gh} z$   
( $x \leq gy, y \leq hz \Rightarrow x \leq gy \leq g(hz) = (gh)z$ ).
- $\text{id}_x = x \xrightarrow{1} x$ .

Note that  $\text{End}_{\mathcal{C}(X)}(x) = G_x^{\text{op}}$ , the stabilizer of  $x$  in  $G$ , with the opposite multiplication  $(g, h) \mapsto hg$ . In particular any endomorphism is an isomorphism.

**Definition** A **representation** of a  $G$ -poset  $X$  over  $k$  is a functor  $F: \mathcal{C}(X)^{\text{op}} \rightarrow {}_k\text{mod}$ . Representations of  $X$  over  $k$  form an abelian category  $\mathcal{P}_k(X)$ . Note that for any  $g \in G$  and  $x \leq y$  in  $X$  one has commutative diagrams

$$\begin{array}{ccc}
 gy & \xrightarrow{g} & y \\
 \uparrow 1 & & \uparrow 1 \\
 gx & \xrightarrow{g} & x
 \end{array}
 \quad \xrightarrow{F} \quad
 \begin{array}{ccc}
 F(gy) & \xleftarrow{c_{g,y}} & F(y) \\
 \downarrow r_{gx}^{gy} & & \downarrow r_x^y \\
 F(gx) & \xleftarrow{c_{g,x}} & F(x)
 \end{array}$$

Moreover,  $F(x)$  is a  $kG_x$ -module.



**Example** Let  $X$  be the set of subgroups of  $G$  endowed with  $G$ -conjugation and let  $V \in {}_kG\text{Mod}$ . One can form the representation  $H \mapsto V^H := \{v \in V \mid hv = v \text{ for all } h \in H\}$ . This defines a functor

$$\mathcal{I}: {}_kG\text{mod} \rightarrow \mathcal{P}_k(X).$$

**Example** Let  $X$  be the set of subgroups of  $G$  endowed with  $G$ -conjugation and let  $V \in {}_kG\text{Mod}$ . One can form the representation  $H \mapsto V^H := \{v \in V \mid hv = v \text{ for all } h \in H\}$ . This defines a functor

$$\mathcal{I}: {}_kG\text{mod} \rightarrow \mathcal{P}_k(X).$$

The restriction maps are inclusions and the conjugation map  $c_{g,H}$  is the application of  $g$  on  $V^H$ .

**Definition** Let  $X$  be a  $G$ -poset. The **incidence algebra**  $A_k(X) = A(X)$  over  $k$  is defined as the free  $k$ -module with basis elements

$$(x, g, y) \quad (\text{where } x \leq gy)$$

**Definition** Let  $X$  be a  $G$ -poset. The **incidence algebra**  $A_k(X) = A(X)$  over  $k$  is defined as the free  $k$ -module with basis elements

$$(x, g, y) \quad (\text{where } x \leq gy)$$

and multiplication defined by

$$(x, g, y) \cdot (y', h, z) := \begin{cases} (x, gh, z) & \text{if } y = y', \\ 0 & \text{if } y \neq y'. \end{cases}$$

**Definition** Let  $X$  be a  $G$ -poset. The **incidence algebra**  $A_k(X) = A(X)$  over  $k$  is defined as the free  $k$ -module with basis elements

$$(x, g, y) \quad (\text{where } x \leq gy)$$

and multiplication defined by

$$(x, g, y) \cdot (y', h, z) := \begin{cases} (x, gh, z) & \text{if } y = y', \\ 0 & \text{if } y \neq y'. \end{cases}$$

This is also the category algebra  $k\mathcal{C}(X)^{\text{op}}$ . If  $X$  is finite,  $A(X)$  has the identity element

$$1_{A(X)} = \sum_{x \in X} e_x,$$

where  $e_x = (x, 1, x) = \text{id}_x$ .

**Proposition** *If  $X$  is a finite  $G$ -poset then one has a category equivalence*

$$\mathcal{P}_k(X) \cong_{A_k(X) \text{ mod }}.$$

*The simple  $A_k(X)$ -modules are parametrized by  $G$ -orbits of pairs  $(x, [V])$ , where  $x \in G$  and  $V$  is a simple  $kG_x$ -module.*

**Proposition** *If  $X$  is a finite  $G$ -poset then one has a category equivalence*

$$\mathcal{P}_k(X) \cong A_k(X) \bmod .$$

*The simple  $A_k(X)$ -modules are parametrized by  $G$ -orbits of pairs  $(x, [V])$ , where  $x \in G$  and  $V$  is a simple  $kG_x$ -module.*

**Theorem** (Linckelmann 2004) *If  $k$  is a field and  $X$  is a finite  $G$ -poset then the incidence algebra  $A_k(X)$  is quasi-hereditary.*

**Proposition** *If  $X$  is a finite  $G$ -poset then one has a category equivalence*

$$\mathcal{P}_k(X) \cong {}_{A_k(X)}\text{mod}.$$

*The simple  $A_k(X)$ -modules are parametrized by  $G$ -orbits of pairs  $(x, [V])$ , where  $x \in G$  and  $V$  is a simple  $kG_x$ -module.*

**Theorem** (Linckelmann 2004) *If  $k$  is a field and  $X$  is a finite  $G$ -poset then the incidence algebra  $A_k(X)$  is quasi-hereditary.*

Consequence: The category  ${}_{A_k(X)}\text{mod}$  has many special properties. For instance, every finitely generated  $A_k(X)$ -module has a finite projective resolution.



**Proposition** *If  $X$  is a finite  $G$ -poset then one has a category equivalence*

$$\mathcal{P}_k(X) \cong A_k(X)\text{mod}.$$

*The simple  $A_k(X)$ -modules are parametrized by  $G$ -orbits of pairs  $(x, [V])$ , where  $x \in G$  and  $V$  is a simple  $kG_x$ -module.*

**Theorem** (Linckelmann 2004) *If  $k$  is a field and  $X$  is a finite  $G$ -poset then the incidence algebra  $A_k(X)$  is quasi-hereditary.*

Consequence: The category  $A_k(X)\text{mod}$  has many special properties. For instance, every finitely generated  $A_k(X)$ -module has a finite projective resolution.

**Proposition** (B.-Monteiro 2004) *One can explicitly determine the central idempotents of  $A_k(X)$  in terms of the central idempotents of the various group algebras  $kG_x$ .*

## 2. The canonical Brauer induction formula

In this section,  $k = \mathbb{C}$ .

$R(G) :=$  ring of virtual characters of  $G =$  Grothendieck ring of  $\mathbb{C}_G\text{mod}$ .

$\hat{G} := \text{Hom}(G, \mathbb{C}^\times) \subseteq R(G)$ .

## 2. The canonical Brauer induction formula

In this section,  $k = \mathbb{C}$ .

$R(G) :=$  ring of virtual characters of  $G =$  Grothendieck ring of  $\mathbb{C}_G\text{mod}$ .

$\hat{G} := \text{Hom}(G, \mathbb{C}^\times) \subseteq R(G)$ .

**Theorem (Brauer 1947)** *For every  $\chi \in R(G)$  there exist  $H_i \leq G$ ,  $\varphi_i \in \hat{H}_i$ ,  $n_i \in \mathbb{Z}$ ,  $i = 1, \dots, r$ , such that*

$$\chi = \sum_{i=1}^r n_i \cdot \text{ind}_{H_i}^G(\varphi_i).$$

## 2. The canonical Brauer induction formula

In this section,  $k = \mathbb{C}$ .

$R(G) :=$  ring of virtual characters of  $G =$  Grothendieck ring of  $\mathbb{C}G\text{mod}$ .

$\hat{G} := \text{Hom}(G, \mathbb{C}^\times) \subseteq R(G)$ .

**Theorem (Brauer 1947)** *For every  $\chi \in R(G)$  there exist  $H_i \leq G$ ,  $\varphi_i \in \hat{H}_i$ ,  $n_i \in \mathbb{Z}$ ,  $i = 1, \dots, r$ , such that*

$$\chi = \sum_{i=1}^r n_i \cdot \text{ind}_{H_i}^G(\varphi_i).$$

Consider the set

$$\mathcal{M}_G := \{(H, \varphi) \mid H \leq G, \varphi \in \hat{H}\}.$$

It is a  $G$ -poset via  $(K, \psi) \leq (H, \varphi) : \iff K \leq H$  and  $\psi = \varphi|_K$ , together with the  $G$ -conjugation action  $(g, (H, \varphi)) \mapsto {}^g(H, \varphi) = ({}^gH, {}^g\varphi)$ .

Consider the free abelian group  $R_+(G)$  with basis  $G \setminus \mathcal{M}_G = \{[H, \varphi]_G\}$ .

Consider the free abelian group  $R_+(G)$  with basis  $G \setminus \mathcal{M}_G = \{[H, \varphi]_G\}$ .  
Dress 1971:  $R_+(G)$  is a commutative ring and there exist natural maps

$$\text{res}_H^G: R_+(G) \rightarrow R_+(H) \quad \text{for all } H \leq G.$$

Consider the free abelian group  $R_+(G)$  with basis  $G \setminus \mathcal{M}_G = \{[H, \varphi]_G\}$ .  
Dress 1971:  $R_+(G)$  is a commutative ring and there exist natural maps

$$\text{res}_H^G: R_+(G) \rightarrow R_+(H) \quad \text{for all } H \leq G.$$

Brauer's induction theorem says that the map

$$b_G: R_+(G) \rightarrow R(G), \quad [H, \varphi]_G \mapsto \text{ind}_H^G(\varphi),$$

is surjective.

Consider the free abelian group  $R_+(G)$  with basis  $G \setminus \mathcal{M}_G = \{[H, \varphi]_G\}$ .  
 Dress 1971:  $R_+(G)$  is a commutative ring and there exist natural maps

$$\text{res}_H^G: R_+(G) \rightarrow R_+(H) \quad \text{for all } H \leq G.$$

Brauer's induction theorem says that the map

$$b_G: R_+(G) \rightarrow R(G), \quad [H, \varphi]_G \mapsto \text{ind}_H^G(\varphi),$$

is surjective.

Also, the diagram

$$\begin{array}{ccc} R_+(G) & \xrightarrow{b_G} & R(G) \\ \text{res}_H^G \downarrow & & \downarrow \text{res}_H^G \\ R_+(H) & \xrightarrow{b_H} & R(H) \end{array}$$

commutes.



**Definition** (B. 1990) A **canonical Brauer induction formula** is a family of maps  $a_G: R(G) \rightarrow R_+(G)$ , one for each finite group  $G$ , such that  $b_G \circ a_G = \text{id}_{R(G)}$  and  $a_G$  commutes with restrictions to subgroups.

**Definition** (B. 1990) A **canonical Brauer induction formula** is a family of maps  $a_G: R(G) \rightarrow R_+(G)$ , one for each finite group  $G$ , such that  $b_G \circ a_G = \text{id}_{R(G)}$  and  $a_G$  commutes with restrictions to subgroups.

- Snaithe 1988: constructed such a map  $a_G$  (only on  $R_{\geq 0}(G)$ , not additive).

**Definition** (B. 1990) A **canonical Brauer induction formula** is a family of maps  $a_G: R(G) \rightarrow R_+(G)$ , one for each finite group  $G$ , such that  $b_G \circ a_G = \text{id}_{R(G)}$  and  $a_G$  commutes with restrictions to subgroups.

- Snaithe 1988: constructed such a map  $a_G$  (only on  $R_{\geq 0}(G)$ , not additive).
- B. 1990: Canonical induction formulas are uniquely determined up to a normalization. The most obvious normalization leads to **the canonical Brauer induction formula**, explicitly given by

$$a_G(\chi) = \sum_{\substack{(H_0, \varphi_0) < \dots < (H_n, \varphi_n) \\ \text{mod } G}} (-1)^n (\chi|_{H_n}, \varphi_n) [H_0, \varphi_0]_G.$$

**Definition** (B. 1990) A **canonical Brauer induction formula** is a family of maps  $a_G: R(G) \rightarrow R_+(G)$ , one for each finite group  $G$ , such that  $b_G \circ a_G = \text{id}_{R(G)}$  and  $a_G$  commutes with restrictions to subgroups.

- Snaith 1988: constructed such a map  $a_G$  (only on  $R_{\geq 0}(G)$ , not additive).
- B. 1990: Canonical induction formulas are uniquely determined up to a normalization. The most obvious normalization leads to **the canonical Brauer induction formula**, explicitly given by

$$a_G(\chi) = \sum_{\substack{(H_0, \varphi_0) < \dots < (H_n, \varphi_n) \\ \text{mod } G}} (-1)^n (\chi|_{H_n, \varphi_n}) [H_0, \varphi_0]_G.$$

Thus, if  $\chi$  is afforded by  $V \in \mathbb{C}G \text{ mod}$  then

$$\chi = \sum_{\substack{(H_0, \varphi_0) < \dots < (H_n, \varphi_n) \\ \text{mod } G}} (-1)^n \text{ind}_{H_0}^G [V^{(H_n, \varphi_n)}],$$

where  $V^{(H, \varphi)} := \{v \in V \mid hv = \varphi(h)v \text{ for all } h \in H\}$ .

**Definition** (B. 1990) A **canonical Brauer induction formula** is a family of maps  $a_G: R(G) \rightarrow R_+(G)$ , one for each finite group  $G$ , such that  $b_G \circ a_G = \text{id}_{R(G)}$  and  $a_G$  commutes with restrictions to subgroups.

- Snaith 1988: constructed such a map  $a_G$  (only on  $R_{\geq 0}(G)$ , not additive).
- B. 1990: Canonical induction formulas are uniquely determined up to a normalization. The most obvious normalization leads to **the canonical Brauer induction formula**, explicitly given by

$$a_G(\chi) = \sum_{\substack{(H_0, \varphi_0) < \dots < (H_n, \varphi_n) \\ \text{mod } G}} (-1)^n (\chi|_{H_n, \varphi_n}) [H_0, \varphi_0]_G.$$

Thus, if  $\chi$  is afforded by  $V \in \mathbb{C}G \text{ mod}$  then

$$\chi = \sum_{\substack{(H_0, \varphi_0) < \dots < (H_n, \varphi_n) \\ \text{mod } G}} (-1)^n \text{ind}_{H_0}^G [V^{(H_n, \varphi_n)}],$$

where  $V^{(H, \varphi)} := \{v \in V \mid hv = \varphi(h)v \text{ for all } h \in H\}$ .

- Symonds 1991: geometric interpretation of this formula.

**Definition** (B. 2001) The category  $\mathbb{C}_G\text{mon}$  of finite  $G$ -line bundles over  $\mathbb{C}$  is defined as follows:

**Definition** (B. 2001) The category  $\mathbb{C}_G\text{mon}$  of **finite  $G$ -line bundles** over  $\mathbb{C}$  is defined as follows:

- Objects: Pairs  $(M, \mathcal{L})$  with  $M \in \mathbb{C}_G\text{mod}$  and  $\mathcal{L} = \{L_1, \dots, L_n\}$  a set of 1-dimensional  $\mathbb{C}$ -subspaces of  $M$  that are permuted by  $G$  and satisfy  $M = L_1 \oplus \dots \oplus L_n$ .

**Definition** (B. 2001) The category  $\mathbb{C}_G\text{mon}$  of **finite  $G$ -line bundles** over  $\mathbb{C}$  is defined as follows:

- Objects: Pairs  $(M, \mathcal{L})$  with  $M \in \mathbb{C}_G\text{mod}$  and  $\mathcal{L} = \{L_1, \dots, L_n\}$  a set of 1-dimensional  $\mathbb{C}$ -subspaces of  $M$  that are permuted by  $G$  and satisfy  $M = L_1 \oplus \dots \oplus L_n$ .

Every  $L_i$  has a stabilizing pair  $(H_i, \varphi_i) \in \mathcal{M}_G$ . For  $(H, \varphi) \in \mathcal{M}_G$  set

$$M((H, \varphi)) := \bigoplus_{\substack{L_i \in \mathcal{L} \\ (H_i, \varphi_i) = (H, \varphi)}} L_i \quad \text{and} \quad M^{((H, \varphi))} := \bigoplus_{\substack{L_i \in \mathcal{L} \\ (H_i, \varphi_i) \geq (H, \varphi)}} L_i.$$



**Definition** (B. 2001) The category  $\mathbb{C}_G\text{mon}$  of **finite  $G$ -line bundles** over  $\mathbb{C}$  is defined as follows:

- Objects: Pairs  $(M, \mathcal{L})$  with  $M \in \mathbb{C}_G\text{mod}$  and  $\mathcal{L} = \{L_1, \dots, L_n\}$  a set of 1-dimensional  $\mathbb{C}$ -subspaces of  $M$  that are permuted by  $G$  and satisfy  $M = L_1 \oplus \dots \oplus L_n$ .

Every  $L_i$  has a stabilizing pair  $(H_i, \varphi_i) \in \mathcal{M}_G$ . For  $(H, \varphi) \in \mathcal{M}_G$  set

$$M((H, \varphi)) := \bigoplus_{\substack{L_i \in \mathcal{L} \\ (H_i, \varphi_i) = (H, \varphi)}} L_i \quad \text{and} \quad M^{((H, \varphi))} := \bigoplus_{\substack{L_i \in \mathcal{L} \\ (H_i, \varphi_i) \geq (H, \varphi)}} L_i.$$

- $\text{Hom}_{\mathbb{C}_G\text{mon}}(M, N)$  is the set of  $f \in \text{Hom}_{\mathbb{C}G}(M, N)$  satisfying

$$f(M^{((H, \varphi))}) \subseteq N^{((H, \varphi))} \quad \text{for all } (H, \varphi) \in \mathcal{M}_G.$$

**Definition** (B. 2001) The category  $\mathbb{C}_G\text{mon}$  of **finite  $G$ -line bundles** over  $\mathbb{C}$  is defined as follows:

- Objects: Pairs  $(M, \mathcal{L})$  with  $M \in \mathbb{C}_G\text{mod}$  and  $\mathcal{L} = \{L_1, \dots, L_n\}$  a set of 1-dimensional  $\mathbb{C}$ -subspaces of  $M$  that are permuted by  $G$  and satisfy  $M = L_1 \oplus \dots \oplus L_n$ .

Every  $L_i$  has a stabilizing pair  $(H_i, \varphi_i) \in \mathcal{M}_G$ . For  $(H, \varphi) \in \mathcal{M}_G$  set

$$M((H, \varphi)) := \bigoplus_{\substack{L_i \in \mathcal{L} \\ (H_i, \varphi_i) = (H, \varphi)}} L_i \quad \text{and} \quad M^{((H, \varphi))} := \bigoplus_{\substack{L_i \in \mathcal{L} \\ (H_i, \varphi_i) \geq (H, \varphi)}} L_i.$$

- $\text{Hom}_{\mathbb{C}_G\text{mon}}(M, N)$  is the set of  $f \in \text{Hom}_{\mathbb{C}G}(M, N)$  satisfying

$$f(M^{((H, \varphi))}) \subseteq N^{((H, \varphi))} \quad \text{for all } (H, \varphi) \in \mathcal{M}_G.$$

$\mathbb{C}_G\text{mon}$  is a  $\mathbb{C}$ -linear additive category, but not abelian.

**Proposition** (B. 2001) *Every indecomposable object in  $\mathbb{C}_G\text{mon}$  is of the form  $\text{Ind}_H^G(\mathbb{C}_\varphi) = \mathbb{C}G \otimes_{\mathbb{C}H} \mathbb{C}_\varphi$  for some  $(H, \varphi) \in \mathcal{M}_G$ , uniquely determined up to conjugation, and the Grothendieck group of  $\mathbb{C}_G\text{mon}$  is  $R_+(G)$ .*

**Proposition** (B. 2001) *Every indecomposable object in  $\mathbb{C}_G\text{mon}$  is of the form  $\text{Ind}_H^G(\mathbb{C}_\varphi) = \mathbb{C}G \otimes_{\mathbb{C}H} \mathbb{C}_\varphi$  for some  $(H, \varphi) \in \mathcal{M}_G$ , uniquely determined up to conjugation, and the Grothendieck group of  $\mathbb{C}_G\text{mon}$  is  $R_+(G)$ . Moreover, the forgetful functor  $\mathbb{C}_G\text{mon} \rightarrow \mathbb{C}_G\text{mod}$  induces the map  $b_G: R_+(G) \rightarrow R(G)$ ,  $[H, \varphi]_G \mapsto \text{ind}_H^G(\varphi)$ .*

**Proposition** (B. 2001) *Every indecomposable object in  $\mathbb{C}_G\text{mon}$  is of the form  $\text{Ind}_H^G(\mathbb{C}_\varphi) = \mathbb{C}G \otimes_{\mathbb{C}H} \mathbb{C}_\varphi$  for some  $(H, \varphi) \in \mathcal{M}_G$ , uniquely determined up to conjugation, and the Grothendieck group of  $\mathbb{C}_G\text{mon}$  is  $R_+(G)$ . Moreover, the forgetful functor  $\mathbb{C}_G\text{mon} \rightarrow \mathbb{C}_G\text{mod}$  induces the map  $b_G: R_+(G) \rightarrow R(G)$ ,  $[H, \varphi]_G \mapsto \text{ind}_H^G(\varphi)$ .*

**Definition** The functors  $\mathcal{I}: \mathbb{C}_G\text{mod} \rightarrow \mathcal{P}(\mathcal{M}_G)$  and  $\mathcal{J}: \mathbb{C}_G\text{mon} \rightarrow \mathcal{P}(\mathcal{M}_G)$  are defined by

$$\mathcal{I}(V) = \left( V^{(H, \varphi)} \right)_{(H, \varphi) \in \mathcal{M}_G} \quad \text{and} \quad \mathcal{J}(M) := \left( M^{((H, \varphi))} \right)_{(H, \varphi) \in \mathcal{M}_G}.$$

**Proposition** (B. 2001) *Every indecomposable object in  $\mathbb{C}_G\text{mon}$  is of the form  $\text{Ind}_H^G(\mathbb{C}_\varphi) = \mathbb{C}G \otimes_{\mathbb{C}H} \mathbb{C}_\varphi$  for some  $(H, \varphi) \in \mathcal{M}_G$ , uniquely determined up to conjugation, and the Grothendieck group of  $\mathbb{C}_G\text{mon}$  is  $R_+(G)$ . Moreover, the forgetful functor  $\mathbb{C}_G\text{mon} \rightarrow \mathbb{C}_G\text{mod}$  induces the map  $b_G: R_+(G) \rightarrow R(G)$ ,  $[H, \varphi]_G \mapsto \text{ind}_H^G(\varphi)$ .*

**Definition** The functors  $\mathcal{I}: \mathbb{C}_G\text{mod} \rightarrow \mathcal{P}(\mathcal{M}_G)$  and  $\mathcal{J}: \mathbb{C}_G\text{mon} \rightarrow \mathcal{P}(\mathcal{M}_G)$  are defined by

$$\mathcal{I}(V) = \left( V^{(H, \varphi)} \right)_{(H, \varphi) \in \mathcal{M}_G} \quad \text{and} \quad \mathcal{J}(M) := \left( M^{((H, \varphi))} \right)_{(H, \varphi) \in \mathcal{M}_G}.$$

**Proposition** (B. 2001)  *$\mathcal{I}$  and  $\mathcal{J}$  are fully faithful embeddings of  $\mathbb{C}_G\text{mod}$  and  $\mathbb{C}_G\text{mon}$  into the full subcategory  $\mathcal{P}'(\mathcal{M}_G)$  of  $\mathcal{P}(\mathcal{M}_G)$  consisting of those functors  $F$ , such that  $h \in H$  acts on  $F(H, \varphi)$  via multiplication with  $\varphi(h)$ .*

**Proposition** (B. 2001) *Every indecomposable object in  $\mathbb{C}_G\text{mon}$  is of the form  $\text{Ind}_H^G(\mathbb{C}_\varphi) = \mathbb{C}G \otimes_{\mathbb{C}H} \mathbb{C}_\varphi$  for some  $(H, \varphi) \in \mathcal{M}_G$ , uniquely determined up to conjugation, and the Grothendieck group of  $\mathbb{C}_G\text{mon}$  is  $R_+(G)$ . Moreover, the forgetful functor  $\mathbb{C}_G\text{mon} \rightarrow \mathbb{C}_G\text{mod}$  induces the map  $b_G: R_+(G) \rightarrow R(G)$ ,  $[H, \varphi]_G \mapsto \text{ind}_H^G(\varphi)$ .*

**Definition** The functors  $\mathcal{I}: \mathbb{C}_G\text{mod} \rightarrow \mathcal{P}(\mathcal{M}_G)$  and  $\mathcal{J}: \mathbb{C}_G\text{mon} \rightarrow \mathcal{P}(\mathcal{M}_G)$  are defined by

$$\mathcal{I}(V) = \left( V^{(H, \varphi)} \right)_{(H, \varphi) \in \mathcal{M}_G} \quad \text{and} \quad \mathcal{J}(M) := \left( M^{((H, \varphi))} \right)_{(H, \varphi) \in \mathcal{M}_G}.$$

**Proposition** (B. 2001)  *$\mathcal{I}$  and  $\mathcal{J}$  are fully faithful embeddings of  $\mathbb{C}_G\text{mod}$  and  $\mathbb{C}_G\text{mon}$  into the full subcategory  $\mathcal{P}'(\mathcal{M}_G)$  of  $\mathcal{P}(\mathcal{M}_G)$  consisting of those functors  $F$ , such that  $h \in H$  acts on  $F(H, \varphi)$  via multiplication with  $\varphi(h)$ . Moreover, every object in  $\mathcal{J}(\mathbb{C}_G\text{mon})$  is projective in  $\mathcal{P}'(\mathcal{M}_G)$ .*

### 3. Categorification of the canonical Brauer induction formula

Again,  $k = \mathbb{C}$  throughout this section.



### 3. Categorification of the canonical Brauer induction formula

Again,  $k = \mathbb{C}$  throughout this section.

**Theorem** (B. 2001) *Let  $V \in \mathbb{C}_G\text{mod}$ . Then  $\mathcal{I}(V)$  has a finite projective resolution in  $\mathcal{P}'(\mathcal{M}_G)$  in terms of objects from  $\mathcal{J}(\mathbb{C}_G\text{mon})$ .*

### 3. Categorification of the canonical Brauer induction formula

Again,  $k = \mathbb{C}$  throughout this section.

**Theorem** (B. 2001) *Let  $V \in \mathbb{C}_G\text{mod}$ . Then  $\mathcal{I}(V)$  has a finite projective resolution in  $\mathcal{P}'(\mathcal{M}_G)$  in terms of objects from  $\mathcal{J}(\mathbb{C}_G\text{mon})$ .*

*Moreover, if  $\mathcal{J}(M_*)$  is this projective resolution of  $V$  then  $a_G([V]) = \sum_{i \geq 0} (-1)^i [M_i]$  in  $R_+(G)$ . In particular, one has a commutative diagram*

$$\begin{array}{ccc} \mathbb{C}_G\text{mod} & \rightarrow & K^b(\mathbb{C}_G\text{mon}) \\ \downarrow [-] & & \downarrow [-] \\ R(G) & \xrightarrow{a_G} & R_+(G) \end{array}$$

### 3. Categorification of the canonical Brauer induction formula

Again,  $k = \mathbb{C}$  throughout this section.

**Theorem** (B. 2001) *Let  $V \in \mathbb{C}_G\text{mod}$ . Then  $\mathcal{I}(V)$  has a finite projective resolution in  $\mathcal{P}'(\mathcal{M}_G)$  in terms of objects from  $\mathcal{J}(\mathbb{C}_G\text{mon})$ .*

*Moreover, if  $\mathcal{J}(M_*)$  is this projective resolution of  $V$  then  $a_G([V]) = \sum_{i \geq 0} (-1)^i [M_i]$  in  $R_+(G)$ . In particular, one has a commutative diagram*

$$\begin{array}{ccc} \mathbb{C}_G\text{mod} & \rightarrow & K^b(\mathbb{C}_G\text{mon}) \\ \downarrow [-] & & \downarrow [-] \\ R(G) & \xrightarrow{a_G} & R_+(G) \end{array}$$

**Remark** For given  $V \in \mathbb{C}_G\text{mod}$  one can find an  $M_*$  of length  $\leq$  longest strictly ascending chain in the set of subspaces  $V^{(H,\varphi)} \neq 0$ .

**Question:** Are the objects of  $\mathcal{J}(\mathbb{C}_G\text{mon})$  also projective in  $\mathcal{P}(\mathcal{M}_G)$ ?

**Question:** Are the objects of  $\mathcal{J}(\mathbb{C}_G \text{mon})$  also projective in  $\mathcal{P}(\mathcal{M}_G)$ ?

Set

$$\varepsilon := \sum_{(H, \varphi) \in \mathcal{M}} \frac{1}{|H|} \sum_{h \in H} \varphi(h^{-1}) \cdot ((H, \varphi), h, (H, \varphi)) \in A(\mathcal{M}_G).$$

Then  $\varepsilon$  is an idempotent and  $\mathcal{P}'(\mathcal{M}_G)$  corresponds under the equivalence  $\mathcal{P}(\mathcal{M}_G) \cong {}_{A(\mathcal{M}_G)}\text{mod}$  to the full subcategory of  ${}_{A(\mathcal{M}_G)}\text{mod}$  consisting of those modules on which  $\varepsilon$  acts as identity.

**Question:** Are the objects of  $\mathcal{J}(\mathbb{C}_G \text{mon})$  also projective in  $\mathcal{P}(\mathcal{M}_G)$ ?

Set

$$\varepsilon := \sum_{(H, \varphi) \in \mathcal{M}} \frac{1}{|H|} \sum_{h \in H} \varphi(h^{-1}) \cdot ((H, \varphi), h, (H, \varphi)) \in A(\mathcal{M}_G).$$

Then  $\varepsilon$  is an idempotent and  $\mathcal{P}'(\mathcal{M}_G)$  corresponds under the equivalence  $\mathcal{P}(\mathcal{M}_G) \cong {}_{A(\mathcal{M}_G)}\text{mod}$  to the full subcategory of  ${}_{A(\mathcal{M}_G)}\text{mod}$  consisting of those modules on which  $\varepsilon$  acts as identity.

**Proposition** (B.-Monteiro 2024) *One has  $A(\mathcal{M}_G)\varepsilon \subseteq \varepsilon A(\mathcal{M}_G)$ . In particular,  $A(\mathcal{M}_G)\varepsilon = \varepsilon A(\mathcal{M}_G)\varepsilon$  and  $a\varepsilon = \varepsilon a\varepsilon$  for all  $a \in A(\mathcal{M}_G)$ .*

**Question:** Are the objects of  $\mathcal{J}(\mathbb{C}_G \text{mon})$  also projective in  $\mathcal{P}(\mathcal{M}_G)$ ?

Set

$$\varepsilon := \sum_{(H, \varphi) \in \mathcal{M}} \frac{1}{|H|} \sum_{h \in H} \varphi(h^{-1}) \cdot ((H, \varphi), h, (H, \varphi)) \in A(\mathcal{M}_G).$$

Then  $\varepsilon$  is an idempotent and  $\mathcal{P}'(\mathcal{M}_G)$  corresponds under the equivalence  $\mathcal{P}(\mathcal{M}_G) \cong {}_{A(\mathcal{M}_G)}\text{mod}$  to the full subcategory of  ${}_{A(\mathcal{M}_G)}\text{mod}$  consisting of those modules on which  $\varepsilon$  acts as identity.

**Proposition** (B.-Monteiro 2024) *One has  $A(\mathcal{M}_G)\varepsilon \subseteq \varepsilon A(\mathcal{M}_G)$ . In particular,  $A(\mathcal{M}_G)\varepsilon = \varepsilon A(\mathcal{M}_G)\varepsilon$  and  $a\varepsilon = \varepsilon a\varepsilon$  for all  $a \in A(\mathcal{M}_G)$ . As a consequence, left  $A(\mathcal{M}_G)$ -modules on which  $\varepsilon$  acts as identity are literally the same thing as left  $\varepsilon A(\mathcal{M}_G)\varepsilon$ -modules. Moreover, projective objects of  $\mathcal{P}'(\mathcal{M}_G)$  are also projective in  $\mathcal{P}(\mathcal{M}_G)$ .*

**Proposition** (B.-Monteiro 2024) *The idempotent  $\varepsilon$  satisfies a property that ensures that the algebra  $\varepsilon A(\mathcal{M}_G)\varepsilon$  is again quasi-hereditary.*



**Proposition** (B.-Monteiro 2024) *The idempotent  $\varepsilon$  satisfies a property that ensures that the algebra  $\varepsilon A(\mathcal{M}_G)\varepsilon$  is again quasi-hereditary.*

**Future work** (in progress with Nariel Monteiro) Let  $V \in \mathbb{C}_G\text{mod}$  and let  $P_*$  be a projective resolution of  $\mathcal{I}(V)$  in  $\mathcal{P}(\mathcal{M}_G)$ .

**Proposition** (B.-Monteiro 2024) *The idempotent  $\varepsilon$  satisfies a property that ensures that the algebra  $\varepsilon A(\mathcal{M}_G)\varepsilon$  is again quasi-hereditary.*

**Future work** (in progress with Nariel Monteiro) Let  $V \in \mathbb{C}_G\text{mod}$  and let  $P_*$  be a projective resolution of  $\mathcal{I}(V)$  in  $\mathcal{P}(\mathcal{M}_G)$ . Applying any functor  $\mathcal{P}(\mathcal{M}_G) \rightarrow \mathcal{A}$  (where  $\mathcal{A}$  is abelian) to  $P_*$  and then taking homology, will result in an invariant of  $V$ .

**Proposition** (B.-Monteiro 2024) *The idempotent  $\varepsilon$  satisfies a property that ensures that the algebra  $\varepsilon A(\mathcal{M}_G)\varepsilon$  is again quasi-hereditary.*

**Future work** (in progress with Nariel Monteiro) Let  $V \in \mathbb{C}_G\text{mod}$  and let  $P_*$  be a projective resolution of  $\mathcal{I}(V)$  in  $\mathcal{P}(\mathcal{M}_G)$ . Applying any functor  $\mathcal{P}(\mathcal{M}_G) \rightarrow \mathcal{A}$  (where  $\mathcal{A}$  is abelian) to  $P_*$  and then taking homology, will result in an invariant of  $V$ .

**Question** Does one obtain this way new interesting invariants of  $V$ ?

**Proposition** (B.-Monteiro 2024) *The idempotent  $\varepsilon$  satisfies a property that ensures that the algebra  $\varepsilon A(\mathcal{M}_G)\varepsilon$  is again quasi-hereditary.*

**Future work** (in progress with Nariel Monteiro) Let  $V \in \mathbb{C}_G\text{mod}$  and let  $P_*$  be a projective resolution of  $\mathcal{I}(V)$  in  $\mathcal{P}(\mathcal{M}_G)$ . Applying any functor  $\mathcal{P}(\mathcal{M}_G) \rightarrow \mathcal{A}$  (where  $\mathcal{A}$  is abelian) to  $P_*$  and then taking homology, will result in an invariant of  $V$ .

**Question** Does one obtain this way new interesting invariants of  $V$ ?

**Example** For  $F \in \mathcal{P}(\mathcal{M}_G)$  and fixed  $(H, \varphi) \in \mathcal{M}_G$  consider the functor

$$F \mapsto F(H, \varphi) / \sum_{(H', \varphi') \leq (H, \varphi)} r_{(H, \varphi)}^{(H', \varphi')}(F(H', \varphi')) \in \mathbb{C}_{G(H, \varphi)}\text{mod}.$$

Thank You