

A brief introduction to Algebraic K-theory

Gabriel Angelini-Knoll

December 11, 2020

Preface

These are notes (in progress) for a 2 hour per week course in [algebraic K-theory](#) taught at the Freie Universität Berlin in Winter 2020/21. They will continue to be updated regularly. Please feel free to reach out at my email address gak@math.fu-berlin.de if you notice any typos or errors. These notes are hosted on my personal [website](#).

These notes draw from the tome of Charles Weibel, known as the K-book [24], Algebraic K-theory and its applications by Jonathon Rosenberg [16], Algebraic K-theory by Srinivas [19], as well as the original papers in the subject by Quillen [14, 15], Segal [17], and Waldhausen [21, 22]. In other words, I do not claim originality of any results or proofs. The goal is to give a brief survey of constructions of algebraic K-theory, fundamental theorems in algebraic K-theory, and applications to geometric topology, algebraic geometry, and number theory. The notes focus on the case of algebraic K-theory of rings, which are the common thread in the applications to each of these three subjects.

These notes briefly survey the field of algebraic K-theory from the time period 1950-1985. The advantage of focusing on this time period is that no previous knowledge of $(\infty, 1)$ -categories is required and for those interested in modern constructions of algebraic K-theory, most of the essential ideas already existed in the work of Quillen, Segal, and Waldhausen.

It is assumed that students in this course have a firm background in the basics of algebra, linear algebra, category theory, and topology. Additional knowledge of geometry topology, algebraic geometry, and number theory is useful for understanding certain examples, but certainly not required for the bulk of the material. Previous knowledge of the theory of simplicial sets will certainly be helpful, but a short section on such material is included as an appendix for those unfamiliar.

Conventions

Throughout, by a ring we will mean an associative ring with unit. We will always specify that a ring is non-unital when we want to consider it without unit.

Let Mod_R be the category of left modules over a ring R . When $R = \mathbb{Z}$, we simply write Ab for this category and refer to it as the category of abelian groups and when R is a field k , we write Vec_k for this category and refer to it as the category of vector spaces over a field k . We let $M(R)$ denote the skeleton of the category of finitely generated left R modules and we let $P(R)$ denote the skeleton of the category of finitely generated projective left modules over a ring R . When k is a commutative ring, let $\text{Rep}_k(G)$ be the skeleton of the category of finitely generated $k[G]$ -modules. Usually, we only consider this in the case k is a field.

Throughout, by a space we mean a compactly generated weak Hausdorff space and we write Top for the category of compactly generated weak Hausdorff spaces. Write CW for the category of CW complexes and CW^f for the category of finite CW complexes. When X is a space, let $VB_{\mathbb{R}}(X)$ denote the skeleton of the category of real vector bundles over X and let $VB_{\mathbb{C}}(X)$ denote the skeleton of the category of complex vector bundles over X .

Let Set be the category of sets and let Fin be the skeleton of the category of finite sets. Let G be a finite group and let Fin_G the skeleton of the category of finite G -sets. When R is a ring, let $\text{Rep}_R(G)$ be the skeleton of the category of finite dimensional $k[G]$ -modules.

Chapter 1

Introduction

The 0-th algebraic K-theory group K_0 was first defined by Grothendieck in the late 1950's in order to generalize the Riemann-Roch Theorem to varieties [4]. The name K-theory comes from German word *Klassen* meaning classes and the reason for this name will be more clear after reading Section 2.1.¹ Even earlier, in the early 1950's, Whitehead studied the simply homotopy of of a finite CW complex and constructed an obstruction to two spaces which are homotopy equivalent being simple homotopy equivalent. It was later understood that this class lived in the first algebraic K-theory group K_1 of an integral group ring. It was then shown that these two algebraic K-theory groups could be related by a localization sequence and that there should in fact be a related group K_i for all integers i extending this localization sequence to the left and right.

Milnor constructed the group K_2 of a ring as the center of the Steinberg group of a ring, inspired in part by a theorem of Matsumoto [10], and used this to motivate his definition of the higher algebraic K-theory groups, now known as Milnor K-theory group, in 1970 [12]. However, as we will see, this theory is not a rich invariant in the sense that for finite fields the Milnor K-theory groups K_n^M vanish for $n \geq 2$.

In 1972, Quillen defined higher algebraic K-theory groups using the $+$ -construction [14]. One of his key insights was that the algebraic K-theory groups should be defined as the homotopy groups of a space. In the same year [15], Quillen defined higher algebraic K-theory groups for a category equipped with notion of exact sequences called an *exact category*. This allowed for much broader input, in particular recovering examples of interest in algebraic geometry.

In 1974 [17], Segal defined the algebraic K-theory of a symmetric monoidal

¹Though Grothendieck famously spent most of his life in France, he was in fact born in Berlin, Germany in 1939. Perhaps this is why Grothendieck chose the letter K from the German word *Klassen* rather than the French, but this is not well documented.

category. This notion is sensitive to the symmetric monoidal structure, so it is not a special case of Quillen's Q -construction unless the symmetric monoidal structure is the direct sum in an additive category. Quillen's Q -construction is also not a special case of Segal's construction. One of Segal's motivations was to give new constructions of infinity loop spaces, which were known to represent cohomology theories by Brown representability [5].

In 1978, Waldhausen extended Quillen's Q -construction further so that the input could be a category with cofibrations and weak equivalences [21]. This allowed one to define the algebraic K -theory of spaces. This new definition extended the applications of algebraic K -theory to manifold theory [23].

Since 1985, there have been several new constructions of algebraic K -theory using the theory of $(\infty, 1)$ -categories. These constructions have proven quite useful for demonstrating universal properties of algebraic K -theory. For example, in 2016, Barwick defined a version of Waldhausen's algebraic K -theory construction for small Waldhausen quasicategories in [1] and proved that algebraic K -theory may be considered as a homology theory, in an abstract sense, on the quasi-category of small Waldhausen quasi-categories. Blumberg–Gepner–Tabuada [3] prove that the connective algebraic K -theory of a small stable quasi-category is the universal additive invariant and non-connective algebraic K -theory of a small stable quasi-category is the universal localizing invariant. Additionally, Gepner–Groth–Nikolaus [7] prove universal properties of the algebraic K -theory of symmetric monoidal quasi-categories. However, we will not discuss these more constructions further in the present notes.

Chapter 2

Classical Algebraic K-theory

We begin by studying the groups K_0 and K_1 . These two groups arose independently in the 1950's from entirely different contexts. Later, it was proven they are related by a localization sequence.

The group K_2 of a ring was then defined by Milnor, inspired by work of Matsumoto [10], and Milnor used this to motivate his definition of higher algebraic K-theory groups K_*^M now known as Milnor K-theory. However, these groups are not as rich an invariant as the higher algebraic K-theory groups that we will discuss in the next chapters, due to Quillen [14, 15]. We will explicitly prove this in the case of finite fields.

At the start, I want to emphasize that there are really two flavors of algebraic K-theory: algebraic K-theory of symmetric monoidal categories and algebraic K-theory categories with a notion of exact sequences, such as exact categories. The two flavors of algebraic K-theory agree when we consider symmetric monoidal categories with respect to the coproduct and algebraic K-theory of categories with exact sequences in which these exact sequences split. For example, this is the case for the category of finitely generated projective R modules. Since the category of finitely generated projective R modules will be our central example throughout and the distinction may not always be clear.

I also want to emphasize that none of the results in this chapter are new and our treatment in this chapter is almost entirely contained in chapters I-III of Weibel's K-book [24]. In [24], Weibel goes into significantly more depth on this subject than we attempt to do here. We also point the reader towards books of Bass [2] and Milnor [13] from the time period before 1972, which give an even more thorough treatment of what was known at the time about algebraic K-theory groups. The first four chapters of Rosenberg's [16] are also a great reference for this material and he much more depth than I do in this first chapter.

2.1 The Grothendieck group

In order to give a very general definition of K_0 , we will first briefly set up the theory of monoidal categories and monoids in a monoidal category.

Monoidal categories are an abstraction of the properties enjoyed by the category of abelian groups Ab with respect to the tensor product $\otimes_{\mathbb{Z}}$ and the integers \mathbb{Z} . In particular, the category Ab is equipped with a functor

$$\otimes_{\mathbb{Z}}: \text{Ab} \times \text{Ab} \rightarrow \text{Ab}$$

and a unit \mathbb{Z} object in the sense that there are isomorphisms

$$M \otimes_{\mathbb{Z}} \mathbb{Z} \cong M \cong \mathbb{Z} \otimes_{\mathbb{Z}} M$$

for any abelian group M , which are natural in M . The tensor product is also associative

$$(M \otimes N) \otimes L \cong M \otimes (N \otimes L),$$

where this associativity is natural in M , N and L . There is also a factor swap map

$$B_{M,N}: M \otimes N \rightarrow N \otimes M$$

which is also natural in M and N . In addition, each of these pieces of data satisfy certain commutative diagrams. This data is abstracted to the definition of a symmetric monoidal category, which also applies in many other contexts.

Definition 2.1.1. A *symmetric monoidal category* \mathcal{C} consists of a category \mathcal{C} a functor

$$\otimes: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$$

and a unit $1_{\mathcal{C}}$ together with four natural isomorphisms:

1. an associator

$$a_{-,=,\equiv}: (- \otimes =) \otimes \equiv \xrightarrow{\cong} - \otimes (= \otimes \equiv)$$

2. a left unitor

$$\lambda_-: 1 \otimes (-) \xrightarrow{\cong} (-),$$

3. a right unitor

$$\rho_-: (-) \otimes 1 \xrightarrow{\cong} (-),$$

and

4. a braiding

$$B_{-,=}: (-) \otimes (=) \xrightarrow{\cong} (=) \otimes (-).$$

These natural transformations must satisfy the triangle identity

$$\mathrm{id}_x \otimes \lambda_y \circ a_{x,1_{\mathcal{C}},y} = \rho_x \otimes \mathrm{id}_y \quad (2.1.2)$$

and pentagon identity

$$a_{w,x,y \otimes z} \circ a_{w \otimes x,y,z} = \mathrm{id}_w \otimes a_{x,y,z} \circ a_{w,x \otimes y,z} \circ a_{w,x,y} \otimes \mathrm{id}_z \quad (2.1.3)$$

the hexagon identities

$$a_{y,z,x} \circ B_{x,y \otimes z} \circ a_{x,y,z} = \mathrm{id}_y \otimes B_{x,z} \circ a_{y,x,z} \circ B_{x,y} \otimes \mathrm{id}_z \quad (2.1.4)$$

$$a_{y,z,x}^{-1} \circ B_{x,y \otimes z} \circ a_{x,y,z} = \mathrm{id}_y \otimes B_{x,z} \circ a_{y,x,z} \circ B_{x,y} \otimes \mathrm{id}_z \quad (2.1.5)$$

and the “squaring to identity” axiom

$$B_{y,x} \circ B_{x,y} = \mathrm{id}_{x \otimes y}. \quad (2.1.6)$$

We succinctly write $(\mathcal{C}, \otimes, 1_{\mathcal{C}})$ for all of this data.

If \mathcal{C} has all of the other structure except that it is not equipped with a braiding natural transformation $B_{-,=}$ satisfying (2.1.4), (2.1.5), and (2.1.6), then we say \mathcal{C} is a *monoidal category*.

Examples 2.1.7. The category $P(R)$ is a symmetric monoidal category with respect to \oplus denoted $(P(R), \oplus, 0)$ and it is a monoidal category with respect to \otimes_R , denoted $(P(R), \otimes_R, R)$. Moreover, when R is a commutative ring then the monoidal category $(P(R), \otimes_R, R)$ is in fact a symmetric monoidal category.

The category $VB_k(X)$ for $k = \mathbb{R}$ or $k = \mathbb{C}$ is a symmetric monoidal category with Whitney sum \oplus and it is a monoidal category with tensor product \otimes denoted $(VB_k(X), \oplus, 0)$ and $(VB_k(X), \otimes, k)$ where k here denotes the trivial one dimensional k vector bundle.

The categories Set (respectively Fin) are symmetric monoidal categories with respect to the coproduct $(\mathrm{Set}, \amalg, \emptyset)$ (respectively $(\mathrm{Fin}, \amalg, \emptyset)$, and with respect to the product $(\mathrm{Set}, \times, *)$ (respectively $(\mathrm{Fin}, \times, *)$). Similarly, Fin_G is a symmetric monoidal category with respect to the coproduct $(\mathrm{Fin}_G, \amalg, \emptyset)$ and the product $(\mathrm{Fin}_G, \times, *)$. Let k be a general field. The category $\mathrm{Rep}_k(G)$ is a symmetric monoidal category with respect to the direct sum $(\mathrm{Rep}_k(G), \oplus, 0)$ and it is a monoidal category with respect to tensor product $(\mathrm{Rep}_k(G), \otimes_k, k)$.

We now discuss monoids in a general symmetric monoidal category.

Definition 2.1.8. A (unital) monoid M in a symmetric monoidal category \mathcal{C} is an object M in \mathcal{C} equipped with an operation

$$\mu: M \otimes M \rightarrow M$$

and a unit map

$$\eta: 1_{\mathcal{C}} \rightarrow M$$

from the unit object $1_{\mathcal{C}}$ in \mathcal{C} to M satisfying:

1. the associativity axiom

$$\begin{array}{ccc}
 M \otimes M \otimes M & \xrightarrow{\mu \times 1} & M \otimes M \\
 \downarrow 1 \times \mu & & \downarrow \mu \\
 M \otimes M & \xrightarrow{\mu} & M
 \end{array} \quad (2.1.9)$$

and

2. the unitality axiom

$$\begin{array}{ccccc}
 M & \xrightarrow{\eta \times M} & M \otimes M & \xleftarrow{1 \times \eta} & M \\
 & \searrow & \downarrow \mu & \swarrow & \\
 & & M & &
 \end{array} \quad (2.1.10)$$

If in addition, the commutativity axiom

$$\begin{array}{ccc}
 M \otimes M & \xrightarrow{\tau} & M \otimes M \\
 & \searrow \mu & \swarrow \mu \\
 & & M
 \end{array} \quad (2.1.11)$$

is satisfied, we say that M is a *commutative monoid* in \mathcal{C} .

When $(\mathcal{C}, \otimes_{\mathcal{C}}, 1_{\mathcal{C}}) = (\text{Set}, \times, *)$ we will simply refer to (unital) monoids and commutative monoids in Set as monoids and commutative monoids.

Each of the examples $(P(R), \oplus, 0)$, $(M(R), \oplus, 0)$, $VB_k(X)$, $(\text{Fin}, \amalg, \emptyset)$, $(\text{Fin}, \times, *)$, $(\text{Fin}_G, \amalg, \emptyset)$, $(\text{Fin}_G, \times, *)$, $(\text{Rep}_k(G), \oplus, 0)$, and $(\text{Rep}_k(G), \otimes_k, k)$ may actually be regarded as commutative monoids by applying the forgetful functor from small categories to sets.

If $(M, +, 0)$ is a commutative monoid with operation $+$ and $(M, \times, 1)$ is a monoid with respect to a second operation \times such that $(M, +, \times, 0, 1)$ forms a ring without additive inverses, then we say that M is a semi-ring. In fact, $(P(R), \oplus, \otimes_R, 0, R)$, $(VB_k(X), \oplus, \otimes, 0, k)$, $(\text{Fin}, \amalg, \times, \emptyset, *)$, $(\text{Fin}_G, \amalg, \times, \emptyset, *)$, and $(\text{Rep}_{\mathbb{C}}(G), \oplus, \otimes_{\mathbb{C}[G]}, 0, \mathbb{C}[G])$ are all examples of semi-rings.

We are now prepared to discuss our definition algebraic K-theory K_0 .

Construction 2.1.12. Given a commutative monoid M we form the Grothendieck group completion of M , denoted M^{gp} as follows. We define an equivalence relation on elements $(m, n) \in M \times M$. We define an equivalence relation

$$(m, n) \sim (m + p, n' + p)$$

for any $p \in M$. We then define $M^{\text{gp}} := M \times M / \sim$.

Exercise 2.1.13. Check that $(m, n) \sim (m + p, n' + p)$ is an equivalence relation.

Note that, by construction the abelian group M^{gp} has the universal property that given a map of commutative monoids $M \rightarrow A$, where A is an abelian group, then the map $M \rightarrow A$ factors as

$$\begin{array}{ccc} M & \longrightarrow & M^{\text{gp}} \\ & \searrow & \downarrow \\ & & A. \end{array} \quad (2.1.14)$$

In other words, there is an adjunction given by the isomorphism

$$\text{Hom}_{\text{CMon}}(M, A) \cong \text{Hom}_{\text{Ab}}(M^{\text{gp}}, A)$$

natural in M and A . In particular, the construction M^{gp} is functorial in M .

Alternatively, we could let $F(M)$ be the free abelian group on symbols $[m]$ where $m \in M$. We can then quotient by the subgroup $R(M)$ of $F(M)$ generated by the relations $[m + n] - [m] - [n]$. This construction also clearly satisfies the universal property 2.1.14. Consequently, we may give a different definition of M^{gp} that agrees with the previous construction up to natural isomorphism

Definition 2.1.15. Given a commutative monoid M , define

$$M^{\text{gp}} := F(M)/R(M)$$

where $F(M)$ and $R(M)$ are as defined above.

For $m \in M$ we will write $[m]$ for a general element in M^{gp} .

Definition 2.1.16. Let R be a ring. Then we define the 0-th algebraic K-theory group of R

$$K_0^{\oplus}(R) := (P(R), \oplus, 0)^+$$

where we regard the set of isomorphism classes of subgroups of $P(R)$ as commutative monoid via \oplus and 0. In fact, since

$$(P(R), \oplus, \otimes_R, 0, R)$$

is a semi-ring, then $K_0(R)$ is a ring. When R is a commutative ring then $K_0(R)$ is also a commutative ring.

In fact, this is a special case of a more general construction.

Definition 2.1.17. Let $(\mathcal{C}, \otimes, 1)$ be a skeletally small symmetric monoidal concrete category with symmetric monoidal skeleton $\text{sk } \mathcal{C}$. Then we may regard $\text{sk } \mathcal{C}$ as a commutative monoid in Set with respect to \otimes and 1 and define

$$K_0^{\otimes}(\mathcal{C}) := (\text{sk}(\mathcal{C}), \otimes 1)^+.$$

Moreover, if

$$(\mathrm{sk}\mathcal{C}, \oplus, \otimes_{\mathcal{C}}, 0_{\mathcal{C}}, 1_{\mathcal{C}})$$

is a semi-ring. Then $K_0^{\oplus}(\mathcal{C})$ is a ring.

This general construction allows us to recover many examples of interest.

Examples 2.1.18. The 0-th complex topological K-theory of X is

$$KU^0(X) \cong K_0^{\oplus}(VB_{\mathbb{C}}(X))$$

and the 0-th real topological K-theory of X is

$$KO^0(X) \cong K_0^{\oplus}(VB_{\mathbb{R}}(X)).$$

In fact these are both rings because $(VB_k(X), \oplus, \otimes, 0, k)$ is a semi-ring when $k = \mathbb{C}$ or $k = \mathbb{R}$.

The Burnside ring of a finite group G is

$$A(G) = K_0^{\Pi}(\mathrm{Fin}_G)$$

where the ring structure comes from the fact that $(\mathrm{Fin}_G, \Pi, \times, \emptyset, *)$ is a semi-ring.

Let k be a field. The representation ring of G is

$$R_k(G) = K_0^{\oplus}(\mathrm{Rep}_k(G))$$

where the ring structure comes from the fact that $(\mathrm{Rep}_k(G), \oplus, \otimes_k, 0, \mathbb{C})$ is a semi-ring.

We finish with some basic computations. First, note that $(\mathbb{N}, +, 0)$ is a commutative monoid and its Grothendieck group completion is clearly

$$\mathbb{N}^+ = \mathbb{Z}.$$

Notice that there is always map of commutative monoids

$$\begin{aligned} \mathbb{N} &\rightarrow P(R). \\ n &\mapsto R^n \end{aligned}$$

and by functoriality of the Grothendieck construction, a group homomorphism

$$\mathbb{Z} \rightarrow K_0(R). \tag{2.1.19}$$

We say that R satisfies the left invariant basis property, denoted *IBP*, if R^n and R^m are not isomorphic whenever $n \neq m$. In this case, the rank of a free left R module does not depend on a choice of basis. All commutative rings satisfy this property and integral group rings $\mathbb{Z}[G]$ all satisfy the invariant basis property.

Example 2.1.20. The ring of k -linear endomorphisms $\text{End}_k(k)$ does not satisfy the IBP property.

Exercise 2.1.21. Prove that there is an isomorphism of $\text{End}_k(k)$ modules

$$\text{End}_k(k) \cong \text{End}_k(k) \oplus \text{End}_k(k),$$

verifying the claim in Example 2.1.20.

Lemma 2.1.22. When R satisfies the IBP, then the map

$$\mathbb{Z} \rightarrow K_0(R)$$

induced by the map $n \mapsto R^n$ is injective.

Definition 2.1.23. We define the reduced K_0 group to be the cokernel of the map $\mathbb{Z} \rightarrow K_0(R)$ and denote it $\tilde{K}_0(R)$.

Proposition 2.1.24. When k is a field, then

$$\tilde{K}_0(k) = 0.$$

Proof. The rank of a vector space gives a map of commutative monoids

$$P(k) \rightarrow \mathbb{N}$$

sending $[k^n]$ to n , which is an isomorphism of commutative monoids. \square

Exercise 2.1.25. Prove that when R is a principle ideal domain, then

$$\tilde{K}_0(R) = 0.$$

Exercise 2.1.26. Prove that when R is a local ring, then

$$\tilde{K}_0(R) = 0.$$

The invariant $\tilde{K}_0(R)$ has interesting applications to geometry and number theory. For example, when G is a group and $R = \mathbb{Z}[G]$ is the associated integral group ring, then we define the 0-th Whitehead group

$$Wh_0(G) := \tilde{K}_0(\mathbb{Z}[G]).$$

We will simply state a result of Wall's that shows that the 0-th Whitehead group is an interesting invariant in topology. We say that a topological space X is dominated by a CW complex K if there is a map $K \rightarrow X$ with right homotopy inverse.

Theorem 2.1.27 (Wall's finiteness obstruction). *Suppose that X is dominated by a finite CW complex K and let $G = \pi_1 X$. Then there is an associated obstruction class $w(X) \in Wh_0(G)$ such that $w(X) = 0$ if and only if X is homotopy equivalent to a finite CW complex.*

Remark 2.1.28. Note that we know by CW approximation that X is homotopy equivalent to a CW complex, so it suffices to consider the case when X is also a CW complex, however it is not at all clear that a homotopy retract of a finite CW complex is again finite CW complex.

Reduced K_0 also has applications to number theory. First, we recall a definition from commutative algebra.

Definition 2.1.29. A Dedekind domain R is an integral domain such that for all nontrivial ideals $J \subset I \subset R$, there exists an ideal K such that $IK = J$.

Remark 2.1.30. In a Dedekind domain R every ideal can be written as a product of prime ideals. However, it may not be able to be written as a product of prime ideals in a unique way. If every ideal can be written as a product of prime ideals in a unique way, then R is a PID. Every PID is also clearly a PID.

Definition 2.1.31. The *ideal class group* of a Dedekind domain R is the quotient

$$Cl(R) = \{I : I \subset R\} / \sim$$

where I ranges over all ideals in R and the equivalence relation states that $I \sim J$ if there exist $x, y \in R$ such that $xI = yJ$ as subsets of R . The group structure is given by the product of ideals.

Exercise 2.1.32. Check that this is in fact an equivalence relation and that $Cl(R)$ is an abelian group.

The ideal class group measures the failure of a Dedekind domain to be a UFD, or in other words, the failure of a Dedekind domain to be a PID.

Again, we will not prove the following result, but we record it as another important application of K_0 .

Theorem 2.1.33. *When R is a Dedekind domain, then there is an isomorphism*

$$\tilde{K}_0(R) \cong Cl(R).$$

The class group measures the failure of unique prime factorization. In other words, when R is also a UFD then $\tilde{K}_0(R) = 0$. To see that unique prime factorization can fail, simply consider the ring $\mathbb{Z}[\sqrt{-5}]$. In this ring,

$$(1 - \sqrt{-5})(1 + \sqrt{-5}) = 6 = 2 \cdot 3.$$

Since $\mathbb{Z}[\sqrt{-5}]$ is a Dedekind domain, we observe the following.

Lemma 2.1.34. *There is an isomorphism*

$$K_0(\mathbb{Z}[\sqrt{-5}]) = \mathbb{Z} \oplus \mathbb{Z}/2.$$

The result Theorem 2.1.33 is a special case of a more general result, which we again state without proof.

Theorem 2.1.35. *Let R be a commutative ring of Krull dimension ≤ 1 , then there is an isomorphism*

$$\text{rank} \oplus \det : K_0(R) \cong [\text{Spec}(R), \mathbb{Z}] \oplus \text{Pic}(R).$$

2.2 The Whitehead group $Wh_1(G)$

In the 1940's and 1950's, Whitehead developed the theory of simple homotopy types. We say that a finite CW complex Y has the same simple homotopy type as a finite CW complex X if they are homotopy equivalent and each homotopy can be described in terms of elementary expansions and collapses.

More precisely, let (K, L) be finite CW pair. Then we write $K \searrow^e L$ and say that K collapses to L via an elementary collapse if the following hold:

1. K is build from L by attaching two cells; i.e. $K = L \cup e_{n-1} \cup e_n$ where $e_1, e_2 \notin K$.
2. there exists a pair (D^n, D^{n-1}) and a map

$$\psi: D^n \rightarrow K$$

such that

$$\begin{array}{ccc} S^{n-1} & \xrightarrow{\quad} & L \\ \downarrow & & \downarrow \\ D^{n-1} & \xrightarrow[\psi|_{D^{n-1}}]{} & L \cup e_{n-1} \end{array}$$

and

$$\begin{array}{ccc} S^n & \xrightarrow{\quad} & L \\ \downarrow & & \downarrow \\ D^{n-1} & \xrightarrow[\psi]{} & L \cup e_{n-1} \end{array}$$

such that the closure $\text{cl}(\delta D^n - D^{n-1}) \subset L^{n-1}$, where L^{n-1} is the n -skeleton of L .

In this situation, we also write $L \nearrow^e K$ and say that L expands to K via an elementary expansions.

Whitehead defined a group which encoded the obstruction to two homotopy equivalent finite CW complexes having the same simple homotopy type. Suppose X and Y are CW complexes and there is a homotopy equivalence

$$X \xrightarrow{\sim} Y.$$

Then clearly this homotopy equivalence induces an isomorphism $\pi_1 X \cong \pi_1 Y$. We would like to know whether X and Y are simple homotopy equivalent. There is an obstruction to this, which lies in a group $Wh(\pi_1 X)$, which is an abelian group that depends only on the group $\pi_1 X$. It was later noted that this group can be defined in terms of algebraic K-theory.

Let R be an associative ring and let $GL_n R$ be the group of invertible $n \times n$ matrices with coefficients in R . There is an inclusion

$$GL_n R \subset GL_{n+1} R$$

given by

$$A \mapsto \begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix}$$

We can then form the union (the colimit) to define

$$GL(R) = \bigcup_{n \geq 1} GL_n(R).$$

In general, if we have a group G we can take the quotient by commutators to define

$$G^{\text{ab}} := G/[G, G].$$

In fact this is a left adjoint to the forgetful functor from abelian groups to groups so it satisfies a universal property, which is encoded in the natural isomorphism

$$\text{Hom}_{\text{Ab}}(G^{\text{ab}}, A) \cong \text{Hom}_{\text{Grp}}(G, A).$$

Definition 2.2.1. Let R be a ring, then we define

$$K_1(R) := GL(R)^{\text{ab}}.$$

In fact, there is a nice description of the commutator $[GL(R), GL(R)]$. Let $E_n(R)$ be the subgroup of $GL_n(R)$ consisting of the $n \times n$ matrices, which are *transvections*. A transvection is the sum of the identity matrix and a matrix with only one nonzero entry, where that nonzero entry does not occur on the diagonal. We write $e_{i,j}(r)$ for this matrix where r is the nonzero entry and it occurs in the i, j -th position where $i \neq j$. We may then define $E(R)$ in the same way that we defined $GL(R)$ as the union

$$E(R) = \bigcup_{n \geq 1} E_n(R).$$

Definition 2.2.2. A group G is *perfect* if

$$G = [G, G].$$

Note that for a perfect group $G^{\text{ab}} = 0$. Such groups are quite interesting from the perspective of topology. For example, a path connected space such that $\pi_1 X$ is a nontrivial perfect group and $\pi_k X = 0$ for all $k > 0$ has the property that its homology is the same as the homology of a point and yet it X is not contractible.

Lemma 2.2.3. When $n \geq 3$, then $E_n(R)$ is a perfect group.

Proof. Whenever i, j, k are distinct, then

$$e_{i,j}(r) = [e_{i,k}(r), e_{k,i}(1)].$$

□

Exercise 2.2.4. Give an example where $E_2(R)$ is not perfect.

Exercise 2.2.5. Verify that, if $g \in GL_n(R)$, the identity

$$\begin{pmatrix} g & 0 \\ 0 & g^{-1} \end{pmatrix} = \begin{pmatrix} 1 & g \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -g^{-1} & 1 \end{pmatrix} \begin{pmatrix} 1 & g \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

holds in $GL_{2n}(R)$.

The following example will be useful in the proof of Whitehead's lemma.

Example 2.2.6. A signed permutation matrix is a matrix that permutes the standard basis on R^n up to a sign. If we write $\{e_1, \dots, e_n\}$ for the standard basis, then a signed permutation acts on the set $\{\pm e_1, \dots, \pm e_n\}$. We observe that, for example, the signed permutation matrix

$$\bar{w}_{1,2} := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = e_{1,2}(1)e_{2,1}(1)e_{1,2}(1)$$

can be written as a product of transvections and therefore it is contained in $E_2(R)$ for any ring R . More generally,

$$\bar{w}_{i,j} \in E_n(R)$$

for $n \geq i, j$. We can then show that cyclic permutations of three basis elements are also contained in $E_n(R)$, since they can be written as $\bar{w}_{jk}\bar{w}_{ij}$. Consequently, every matrix corresponding to an even permutation of basis elements is an element in $E_n(R)$ for some n . Thus, by Exercise 2.2.5 we know that $E_{2n}(R)$ contains the matrix

$$\begin{pmatrix} g & 0 \\ 0 & g^{-1} \end{pmatrix}.$$

The subgroup $E_n(R)$ is not necessarily a normal subgroup in $GL_n(R)$. It is often a normal subgroup in $GL_n(R)$ for sufficiently large n , but even this is too much to ask for in general. When R is a commutative ring, the situation is much easier and $E_n(R)$ is normal in $GL_n(R)$ for $n \geq 3$. Nevertheless, we have the following lemma due to Whitehead which, in particular, implies that $E(R)$ is normal in $GL(R)$.

Lemma 2.2.7 (Whitehead's Lemma). *There is an isomorphism*

$$[GL(R), GL(R)] \cong E(R)$$

Proof. The fact that

$$E(R) \subset [GL(R), GL(R)]$$

follows from Lemma 2.2.3. Conversely, suppose $[A, B] \in [GL_n(R), GL_n(R)]$. Then we can write $[A, B]$ as the product

$$[A, B] = \begin{pmatrix} A & 0 \\ 0 & A^{-1} \end{pmatrix} \begin{pmatrix} B & 0 \\ 0 & B^{-1} \end{pmatrix} \begin{pmatrix} (AB)^{-1} & 0 \\ 0 & AB \end{pmatrix}$$

By Example 2.2.6, we therefore know that $[A, B] \in E(R)$. \square

This gives a new definition of $K_1(R)$.

Definition 2.2.8. Let R be a ring, then we define

$$K_1(R) := GL(R)/E(R).$$

In particular, $K_1(R)$ is a quotient of $GL(R)$ by a perfect normal subgroup. The definition as a quotient by a perfect normal subgroup will be important for the next chapter.

Definition 2.2.9. We define the Whitehead group of a group G as

$$Wh_1(G) := K_1(\mathbb{Z}[G]) / \langle \pm g : g \in G \rangle.$$

where $g \in G$ is regarded as an element in $GL_1(\mathbb{Z}[G]) \subset GL(\mathbb{Z}[G])$ and

$$\langle \pm g : g \in G \rangle$$

denotes the subgroup of $K_1(\mathbb{Z}[G])$ generated by the elements $\pm g \in K_1(\mathbb{Z}[G])$.

Again, we will simply cite a deep result that demonstrates that this group is useful for studying problems in topology.

Theorem 2.2.10 (Whitehead). *Suppose K and L are finite CW complexes and there is a homotopy equivalence $f: K \rightarrow L$ inducing an isomorphism $\pi_1 K \cong \pi_1 L$. Let $G = \pi_1 K$. Then there is an associated class*

$$\tau(f) \in Wh_1(G),$$

called the Whitehead torsion of f , such that $\tau(f) = 0$ if and only if f is a simple homotopy equivalence.

It happens quite often that a homotopy equivalence between finite CW complexes is in fact a simple homotopy equivalence. For an example of non-vanishing Whitehead torsion, consider the Lens spaces

$$L(p, q) = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1|^2 + |z_2|^2 = 1\} / (z_1, z_2) \simeq (\zeta z_1, \zeta^q z_2)$$

where $p, q \geq 1$ are integers and ζ is a primitive p -th root of unity. Then there exists a homotopy equivalence

$$f: L(7, 1) \xrightarrow{\sim} L(7, 2)$$

and one can prove that any such homotopy equivalence cannot be a *simple* homotopy equivalence [3, p.98], giving an example when

$$\tau(f) \neq 0 \in Wh_1(\pi_1(L(7, 1))) = Wh_1(\mathbb{Z}/7).$$

In fact, there are other applications of Whitehead torsion to manifold theory. For those unfamiliar with these constructions in manifold theory, we do not plan to give full definitions as that would be too much of a diversion and these constructions will not be used later. We therefore just provide enough information to state the main results in order to indicate the depth of the subject of algebraic K-theory.

Let (W, M, M') be a triple of compact piecewise linear (PL) manifolds. We say this triple is an *h-cobordism* if W has boundary $M \amalg M'$ and both inclusions $M \subset W$ and $M' \subset W$ are homotopy equivalences. There is therefore a Whitehead torsion class $\tau \in Wh_1(\pi_1 M)$ associated to the inclusion $M \subset W$. We record the following deep result, proven by Mazur [11], without proof.

Theorem 2.2.11 (The s-cobordism theorem). *Given an h-cobordism (W, M, M') of PL-manifolds, with M fixed and $\dim(M) \geq 5$. Then there is a PL homeomorphism of triples*

$$(W, M, M') \cong (M \times [0, 1], M \times \{0\}, M \times \{1\})$$

if and only if $\tau = 0$. Moreover, every element $\tau \in Wh_1(\pi_1 M)$ arises as the Whitehead torsion of some h-cobordism (W, M, M') .

This result can be used to prove a version of the generalized Poincaré conjecture, which had originally proven by Smale [18] before the s-cobordism theorem was known..

Corollary 2.2.12. *Suppose N is a PL manifold with the same homotopy type as a sphere S^n and $n \geq 5$. Then N is PL-homeomorphic to a S^n .*

Proof. Form a PL manifold W by removing two disjoint n -discs D_1 and D_2 from N . Then we produce a PL cobordism $(W, S_1^{n-1}, S_2^{n-1})$ where S_i^{n-1} is the boundary of D_i in N . Since $\pi_1 S^{n-1} = 0$ when $n \geq 5$, we know that the Whitehead torsion $\tau \in Wh_1(0)$ vanishes. Thus, there is a PL homeomorphism $W \cong S^{n-1} \times [0, 1]$ by Theorem 2.2.11 and $N = W \cup D_1 \cup D_2$ is therefore PL homeomorphic to S^n . \square

2.2.1 Relating K_0 and K_1

Finally, we prove that there is a localization sequence relating K_1 and K_0 in certain cases. Let I be an ideal in R and let $GL(I)$ be the kernel of the map

$GL(R) \rightarrow GL(R/I)$. Let $E_n(R, I)$ be the normal subgroup of $E_n(R)$ generated by matrices $e_{i,j}(r)$ such that $r \in I$ and $1 \leq i \neq j \leq n$ and define $E(R, I)$ as the union

$$E(R, I) = \bigcup_{n \geq 1} E_n(R, I).$$

Lemma 2.2.13 (Relative Whitehead Lemma). *The group $E(R, I)$ is normal in $GL(I)$ and*

$$[GL(I), GL(I)] \subset E(R, I).$$

Exercise 2.2.14. Prove Lemma 2.2.13.

Definition 2.2.15. Define the relative K_1 group as

$$K_1(R, I) := GL(I)/E(R, I).$$

We can also define a relative algebraic K-theory group K_0 . Given a ring R and an ideal $I \in R$, we can form the trivial square-zero extension of R by I , denoted $R \oplus I$. We will use an explicit model for this square zero extension as the pullback

$$\begin{array}{ccc} R \times_{R/I} R & \longrightarrow & R \\ \downarrow & & \downarrow \\ R & \longrightarrow & R/I \end{array}$$

where the pullback is

$$R \times_{R/I} R = \{(r_1, r_2) : r_1 - r_2 \in I\}.$$

This pullback has a splitting $R \rightarrow R \times_{R/I} R$ given by sending r to (r, r) and the kernel is I so we see that it is abstractly isomorphic to $R \oplus I$, but the description in terms of the pullback can be useful for relating the quotient $R \oplus I \rightarrow R$ to the quotient $R \rightarrow R/I$.

Definition 2.2.16. We define the relative K_0 group as

$$K_0(R, I) := \ker(K_0(R \oplus I) \rightarrow K_0(R)).$$

The definitions of relative K_1 and relative K_0 are a bit different. This is because $K_0(R, I)$ in fact does not depend on R . If $R \rightarrow S$ is a map of rings and I is mapped isomorphically onto an ideal of S , which we also call I , then $K_0(R, I) \cong K_0(S, I)$. We therefore sometimes simply write

$$K_0(I) := K_0(R, I).$$

The same is not true for $K_1(R, I)$ and in fact there are maps of rings $R \rightarrow S$ where I maps isomorphically onto an ideal of S , also denoted I , and yet

$$K_1(R, I) \not\cong K_1(S, I).$$

It is known that $K_1(R, I)$ is independent of R if and only if $I^2 = I$, or in other words I is idempotent by [20, 14.2].

Given a map of commutative rings $R \rightarrow S$ such that and ideal I in R maps isomorphically onto an ideal of S , the the square

$$\begin{array}{ccc} R & \longrightarrow & S \\ \downarrow & & \downarrow \\ R/I & \longrightarrow & S/I \end{array} \quad (2.2.17)$$

is a pullback of associative rings. We say that the square above is a *Milnor square*.

Construction 2.2.18. Given a Milnor square, we can construct an R -module $M = (M_1, g, M_2)$ from the data of an S -module M_1 and R/I -module M_2 and an isomorphism

$$g: M_2 \otimes_{R/I} S/I \cong S/I \otimes_S M_1$$

of S/I -modules by defining M to be the pullback

$$\begin{array}{ccc} M & \longrightarrow & M_1 \\ \downarrow & & \downarrow \\ M_2 & \longrightarrow & M_1/IM_1 \end{array}$$

in the category of R -modules.

Lemma 2.2.19. *Every finitely generated projective R -module P can be written as*

$$P = (P_1, P_2, g)$$

for some $P_1 \in P(S)$, $P_2 \in P(R/I)$ and

$$g: P_2 \otimes_{R/I} S/I \cong S/I \otimes_S P_1$$

in $P(S/I)$.

In particular, given a matrix $A \in GL_n(S/I) \subset GL_n(S/I)$, we can define a finitely generated projective R -module

$$P = (P_1, P_2, A)$$

where

$$A: P_2 \otimes_{R/I} S/I \cong S/I \otimes_S P_1$$

is our isomorphism and so P_2 is a free R/I -module and P_1 is a free S -module. This defines a map

$$\delta: GL(S/I) \rightarrow K_0(R).$$

Moreover, given a Milnor square this map fits into the following Mayer-Vietoris sequence.

Theorem 2.2.20 (Mayer-Vietoris). *Given a Milnor square (2.2.17), there is an exact sequence*

$$GL(S/I) \rightarrow K_0(R) \rightarrow K_0(R/I) \oplus K_0(S) \rightarrow K_0(S/I)$$

Exercise 2.2.21. If $f: R \rightarrow S$ is a ring map sending I isomorphically onto an ideal of S , also denoted I , then prove that

$$K_0(R, I) \cong K_0(S, I).$$

Hint: Show that $GL(S)/GL(S \oplus I) = 1$. Then prove that if $I \cap J = 0$, then

$$K_0(I + J) = K_0(I) \oplus K_0(J).$$

Use this to prove that there is an exact sequence

$$1 \rightarrow GL(I) \rightarrow GL(R) \rightarrow GL(R/I) \xrightarrow{\delta} K_0(I) \rightarrow K_0(R) \rightarrow K_0(R/I). \quad (2.2.22)$$

Proposition 2.2.23. *There is an exact sequence*

$$K_1(R, I) \rightarrow K_1(R) \rightarrow K_0(R/I) \rightarrow K_0(I) \rightarrow K_0(R) \rightarrow K_0(R/I). \quad (2.2.24)$$

Proof. By Exercise 2.2.21, we know that there is an exact sequence (2.2.22). Passing to quotients by $E(R)$ and $E(R/I)$ gives exactness of the sequence (2.2.24) at $K_1(R/I)$. By Exercise 2.2.21, it therefore suffices to show that the sequence (2.2.24) is exact at $K_0(R)$. Let g be an element of the kernel of the composite

$$GL(R) \rightarrow K_1(R) \rightarrow K_1(R/I).$$

Then we know that the image of g in $GL(R/I)$, is in $E(R/I)$. Write \bar{g} for this element of $E(R/I)$. Since the map $E(R) \rightarrow E(R/I)$ is surjective, there is an element $e \in E(R)$ mapping to \bar{g} . Consequently, ge^{-1} maps to $1 \in E(R/I) \subset GL(R/I)$. So ge^{-1} is in the kernel of $GL(R) \rightarrow GL(R/I)$, which we denoted $GL(I)$. Write $[ge^{-1}]$ for the equivalence class of ge^{-1} in $GL(I)/E(R, I)$.

To summarize, for any $g \in K_1(R)$ in the kernel $K_1(R) \rightarrow K_1(R/I)$, we have produced a well-defined element $[ge^{-1}]$ in $K_1(R, I)$ that maps to $K_1(R)$. Thus, the sequence (2.2.24) is exact at $K_1(R)$. \square

The existence of this sequence was known already in the 1960's [2], but it was not known how to extend the sequence to the left. It was expected that there were groups K_n for all $n \in \mathbb{Z}$ that produce a long exact sequence, but it remained an open question until the 1970's. In Section 2.4, we discuss the first proposed definition of higher algebraic K-theory groups in the early 1970's due to Milnor. This was defined in order to extend the groups K_0 , K_1 , and K_2 , where K_2 is defined in Section 2.3 and the choice of definition of the higher Milnor K-theory groups is clearly inspired by the definition of K_2 .

2.3 The Steinberg group and its center

We now discuss K_2 of a ring and the higher Milnor K-theory groups of a field.

Definition 2.3.1. Let A be a ring. Let $n \geq 3$, then the Steinberg group $St_n(A)$ is the group with generators $x_{i,j}(a)$ for $a \in A$ and $i, j \in \{1, \dots, n\}$ with $i \neq j$ and relations

$$x_{i,j}(a)x_{i,j}(b) = x_{i,j}(a+b) \quad (2.3.2)$$

$$[x_{i,j}(a), x_{k,\ell}(s)] = \begin{cases} 1 & \text{if } j \neq k \text{ and } i \neq \ell \\ x_{i,\ell}(rs) & \text{if } j = k \text{ and } i \neq \ell \\ x_{k,j}(-sr) & \text{if } j \neq k \text{ and } i = \ell, \end{cases} \quad (2.3.3)$$

which are called the *Steinberg relations*.

Exercise 2.3.4. Show that the transvections $e_{i,j}(a)$ in $E_n(A)$ for $n \geq 3$ satisfy the Steinberg relations.

As a consequence of Exercise 2.3.4, there is a canonical surjective group homomorphism

$$St_n(A) \rightarrow E_n(A)$$

for $n \geq 3$ mapping $x_{i,j}(a)$ to $e_{i,j}(a)$. Since the Steinberg relations for n include the Steinberg relations for all $k < n$, there is a canonical inclusion

$$St_{n-1}(A) \rightarrow St_n(A)$$

and we define

$$St(A) = \bigcup St_n(A).$$

Exercise 2.3.5. Prove that a level map $\{A_i\} \rightarrow \{B_i\}$ of sequences of groups, which is a levelwise surjection induces a surjection

$$\operatorname{colim}_i A_i \rightarrow \operatorname{colim}_i B_i.$$

Definition 2.3.6. Let A be a ring. We define $K_2(A)$ to be the kernel of the canonical surjection

$$St(A) \rightarrow E(A).$$

As a consequence of the definition, there is an exact sequence

$$1 \rightarrow K_2(A) \rightarrow St(A) \rightarrow GL(A) \rightarrow K_1(A) \rightarrow 1.$$

As defined, it is not clear that $K_2(A)$ is an abelian group, because $St(A)$ is not necessarily abelian. However, it turns out that it is an abelian group. Moreover, we have the following result of Steinberg, but we omit the proof.

Theorem 2.3.7 (Steinberg). *The group $K_2(A)$ is abelian and it is exactly the center of $St(A)$.*

We end by remarking that the group K_2 really deserves to be called K_2 in the following sense.

Theorem 2.3.8. *Let A be a Dedekind domain with field of fractions F , then there is a long exact sequence*

$$K_2(F) \rightarrow \bigoplus_{\mathfrak{p} \in \mathbb{P}} K_1(A/\mathfrak{p}) \rightarrow K_1(A) \rightarrow K_1(F) \rightarrow \bigoplus_{\mathfrak{p} \in \mathbb{P}} K_0(A/\mathfrak{p}) \rightarrow K_0(A) \rightarrow K_0(F) \rightarrow 0$$

where \mathbb{P} is the set of prime ideals in A .

We will prove this as a consequence of the localization sequence and dévissage due to Quillen [15] in 1972 later in the course. However, it is important to know that the localization sequence as stated in Theorem 2.3.8 had already been proven by Milnor in 1971 and it appears in [13].

2.4 Milnor K-theory of fields

Milnor extended the definition of K_2 to higher algebraic K-theory groups of fields, now known as Milnor K-theory groups, in the 1970's.

Let k be a field. We define the tensor algebra of the group of units k to be

$$T(k^\times) = \bigoplus_{i \geq 0} (k^\times)^{\otimes i}$$

where $(k^\times)^{\otimes 0} = \mathbb{Z}$. This is also known as the free associative algebra on k^\times . Write $\ell(x)$ for an element in k^\times in degree 1 corresponding to $x \in k^\times$. We can then define Milnor K-theory of a field as a graded ring all at once.

Definition 2.4.1. The Milnor K-theory groups of a field k are

$$K_*^M(k) := T(k^\times) / (\ell(x) \otimes \ell(1-x) : 1 \neq x \in k^\times)$$

Note that the ideal generated by the elements $\ell(x) \otimes \ell(1-x)$ is a homogeneous ideal and therefore it makes sense to form the quotient in graded rings.

It is clear that

$$\begin{aligned} K_0^M(k) &= \mathbb{Z} = K_0(k) \text{ and} \\ K_1^M(k) &= k^\times = K_1(k) \end{aligned}$$

for any field k . By a theorem of Matsumoto, $K_2^M(k) \cong K_2(k)$.

Theorem 2.4.2 (Matsumoto). *There is an isomorphism*

$$K_2^M(k) \cong K_2(k)$$

for any field k .

This motivated Milnor's definition of higher algebraic K-theory groups. Note that

$$\begin{aligned} K_0(\mathbb{F}_q) &= K_0^M(\mathbb{F}_q) = \mathbb{Z} \text{ and} \\ K_1(\mathbb{F}_q) &\cong \mathbb{F}_q^\times \end{aligned}$$

which is a cyclic group of order $q - 1$. In light of this, the following result gives a complete calculation of the Milnor K-theory of finite fields.

Proposition 2.4.3. *The Milnor K-theory groups $K_n^M(\mathbb{F}_q)$ vanish for all $n \geq 2$. Consequently, there is an isomorphism of graded rings*

$$K_*^M(\mathbb{F}_q) \cong \mathbb{Z} \oplus \mathbb{F}_q^\times$$

where $\mathbb{Z} \oplus \mathbb{F}_q^\times$ is the trivial square zero extension of \mathbb{Z} by the cyclic group \mathbb{F}_q^\times .

Proof. We first show that

$$\left(\mathbb{F}_q^\times \otimes \mathbb{F}_q^\times / (x \otimes (1 - x) : x \neq 1, 0) \right) = 1.$$

We write $(x \otimes y) \cdot (z \otimes w)$ for the group operation and 1 for the unit. Note that \mathbb{F}_q^\times is cyclic of order $q - 1$ and consequently $\mathbb{F}_q^\times \otimes \mathbb{F}_q^\times$ is also cyclic of order $q - 1$. This cyclic group is generated by $x \otimes x$ whenever x is a generator of \mathbb{F}_q^\times .

We split into two cases. If q is even, then we know $2x \otimes x = 0$ in $\mathbb{F}_q \otimes \mathbb{F}_q$. So $x \otimes x = x \otimes -x$. We also know that $x \otimes -x = x \otimes 1$ in $K_2^M(\mathbb{F}_q)$ by the relations and since 1 is the identity in \mathbb{F}_q^\times , the element $x \otimes 1$ is trivial in the group $K_2^M(\mathbb{F}_q)$. In other words, we conclude that

$$x \otimes x = 1$$

for all elements $x \otimes x \in K_2^M(\mathbb{F}_q)$ where x is a generator of \mathbb{F}_q^\times . This implies that the group $K_2^M(\mathbb{F}_q)$ is trivial. In fact, essentially the same argument implies that $K_n^M(\mathbb{F}_q) = 0$ for $n > 2$.

When q is odd, we still know that $x \otimes -x = x \otimes 1$ is trivial and consequently, we have skew-symmetry

$$(x \otimes y) \cdot (y \otimes x) = (x \otimes -xy) \cdot (y \otimes -xy) = xy \otimes -xy = 1$$

in $K_2^M(\mathbb{F}_q)$. This immediately implies that $(x \otimes x)^2 = 1$ and more generally one can show that

$$(x \otimes x)^{mn} = x^m \otimes x^n$$

when m, n are odd. The set of odd powers of elements in $K_2^M(\mathbb{F}_q)$ is exactly the same as the non-squares, by construction. If there exists a non-square u such that $1 - u$ is also a non-square in \mathbb{F}_q , then all elements are divisible by the element $u \otimes (1 - u) = 0$, or in other words all elements are trivial. To find such a u , we note that there is an involution $u \mapsto 1 - u$ on the set $\mathbb{F}_q = \{0, 1\}$. and the set $\mathbb{F}_q - \{0, 1\}$ consists of $(q - 1)/2$ non squares, but only $(q - 3)/2$ squares. In other words, there are strictly less squares than non-squares and there must be an orbit of the C_2 -action that is completely contained in the non-squares. Again, essentially the same proof implies that $K_n^M(\mathbb{F}_q) = 0$ for $n > 2$. \square

We include this result in order to indicate that Milnor K-theory, though very interesting in its own right, is this is not the richest invariant. We will see a different construction of higher algebraic K-theory groups where the algebraic K-theory of fields are nontrivial in arbitrarily high degrees.

Chapter 3

Group completion algebraic K-theory

As we hinted at in the beginning of the course, there are really two main flavors of algebraic K-theory: group completion algebraic K-theory and algebraic K-theory of a category with a notion of exact sequences. Here we describe a construction of group completion algebraic K-theory, due to Quillen [14], known as the $+$ -construction. We then give a sketch of the computation of algebraic K-theory of finite fields due to Quillen. In the following section, we define algebraic K-theory of a symmetric monoidal category via the $S^{-1}S$ construction of Quillen and Segal, see [8].

3.1 The $+$ -construction

This section draws heavily from [19, Chp. 2]. We gave a construction of the classifying space of a (topological) group in Definition B.1.31. In particular, we can consider the infinite general linear group $GL(R)$ of a ring R and form its classifying space $BGL(R)$. This has the property that

$$\pi_k(BGL(R)) = \begin{cases} GL(R) & \text{if } k = 1 \\ 0 & \text{otherwise.} \end{cases}$$

and any other CW complex with this property will be homotopy equivalent to the model for $BGL(R)$ we gave in Definition B.1.31.

In particular, we note that if we want to form a space $K(R)$ with $\pi_1(K(R)) = K_1(R)$ then we will have to attach cells to $BGL(R)$ to kill the perfect normal subgroup $E(R)$ in $GL(R)$. We would also like to not change the space $BGL(R)$ too much, so we attach cells so that the resulting CW complex $BGL(R)^+$ has the property that

$$H_*(BGL(R)^+, BGL(R); L) \cong 0$$

for any $\mathbb{Z}[K_1(R)]$ -module L . In fact, the space $BGL(R)^+$ can be characterized by this property. In particular, the induced map

$$H_*(BGL(R); \mathbb{Z}) \rightarrow H_*(BGL(R)^+; \mathbb{Z})$$

is an isomorphism. We will give a different characterization of the $+$ -construction, which uses the following definition.

Definition 3.1.1. An H -space is topological space X equipped with continuous maps

$$\mu: X \times X \rightarrow X$$

and

$$e: * \rightarrow X$$

such that the diagram

$$\begin{array}{ccccc} X & \xrightarrow{e \times \text{id}_X} & X \times X & \xleftarrow{\text{id}_X \times e} & X \\ & \searrow \text{id}_X & \downarrow \mu & \swarrow \text{id}_X & \\ & & X & & \end{array}$$

commutes up to homotopy, in the sense that

$$\mu \circ (e \times \text{id}_X) \simeq \text{id}_X \simeq \mu \circ (\text{id}_X \times e).$$

If in addition X is a commutative group object in the homotopy category of topological spaces $\text{ho}(\text{Top})$, then we say that X is a commutative H -group.

Example 3.1.2. Let X be a based topological space. Recall that $\Omega X = \text{Map}_*(S^1, X)$, the space of pointed maps from $S^1 \rightarrow X$, has an operation

$$\Omega X \times \Omega X \rightarrow \Omega X$$

given by concatenation of loops and a unit

$$e: * \rightarrow \Omega X$$

which sends the basepoint to the constant loop at the basepoint of X . This does not satisfy unitality on the nose, but it does satisfy unitality up to homotopy so ΩX is an H -space.

Definition 3.1.3. The space $BGL(R)^+$ is the initial H -space receiving a map from $BGL(R)$, in other words, whenever there is a map $BGL(R) \rightarrow Y$ where Y is an H -space, then the map factors as

$$\begin{array}{ccc} X & \longrightarrow & BGL(R)^+ \\ & \searrow & \downarrow \\ & & Y. \end{array}$$

Remark 3.1.4. Note that $\pi_1(Y)$ is abelian whenever Y is an H -space by the Eckman-Hilton argument. Therefore, $\pi_1(BGL(R)^+)$ must be the abelianization of $\pi_1(BGL(R)) = GL(R)$ as desired.

We may therefore think of the $BGL(R)^+$ as the H -spacification of $BGL(R)$. This specifies the space $BGL(R)^+$ up to homeomorphism, but we need to know that such a space exists. We therefore give an explicit construction. First, we give our definition of the algebraic K-theory space of a ring R

Definition 3.1.5. Given a ring R , the algebraic K-theory space of R is

$$K(R) := K_0(R) \times BGLR^+$$

where $K_0(R)$ is regarded as a discrete space. Moreover,

$$K_n(R) := \pi_n(K(R)).$$

We now give an explicit construction of X^+ for a general path connected based space X of the homotopy type of a CW complex such that $[\pi_1 X, \pi_1 X]$ is a perfect normal subgroup. In fact, there is a plus construction associated to any perfect normal subgroup N of $\pi_1(X)$, but we leave that generalization to the reader. In fact, this is the context in which the +-construction was originally defined by Kervaire in [9], before work of Quillen [14]. Kervaire studied this construction in the context of n -manifolds that are homology spheres, so manifolds M such that $\pi_1(M)$ is a perfect subgroup and M^+ is homotopy n -sphere for $n \geq 3$.

Construction 3.1.6. Let X be a path connected based space of the homotopy type of a CW complex such that $[\pi_1 X, \pi_1 X]$ is a perfect normal subgroup. We will construct a relative CW complex (X^+, X) by attaching cells to X so that $\pi_1(X^+) = H_1(X; \mathbb{Z})$ and the relative homology $H_*(\tilde{X}^+, \tilde{X}; \mathbb{Z})$ is trivial where \tilde{X} is the universal cover of X^+ and \tilde{X} is the pullback of the universal cover along the map $X \rightarrow X^+$. We can actually accomplish this by only attaching 2-cells and 3-cells to X to form X^+ .

Write $N = [\pi_1 X, \pi_1 X]$. Choose a minimal set of generators e_α of the subgroup N of $\pi_1 X$. Choose a basepoint preserving loop $\gamma_\alpha: S^1 \rightarrow X$ for each homotopy class $e_\alpha \in \pi_1 X$. We can then form X_1 as the the pushout

$$\begin{array}{ccc} \coprod_{\alpha} S^1 & \xrightarrow{\coprod \gamma_\alpha} & X \\ \downarrow & & \downarrow \\ \coprod_{\alpha} S^1 & \longrightarrow & X_1 \end{array}$$

and by construction $\pi_1(X_1) = \pi_1(X)/N$ as desired. However, we note that by the exact sequence in homology

$$0 \rightarrow H_2(X; \mathbb{Z}) \rightarrow H_2(X_1; \mathbb{Z}) \rightarrow H_1(\coprod_{\alpha} S^1; \mathbb{Z}) \xrightarrow{0} H_1(X; \mathbb{Z}) \xrightarrow{\cong} H_1(X_1; \mathbb{Z})$$

the map $H_2(X) \rightarrow H_2(X_1)$ is not an isomorphism as desired.

Let $\tilde{X}_1 \rightarrow X_1$ be the universal cover of X_1 and let \hat{X} be the pullback

$$\begin{array}{ccc} \hat{X} & \longrightarrow & \hat{X}_1 \\ \downarrow & & \downarrow \pi \\ X & \longrightarrow & X_1. \end{array}$$

Note that \tilde{X}_1 is formed from \hat{X} by attaching 2-cells as well and \tilde{X}_1 is connected so \hat{X} is also connected. We therefore observe that \hat{X} is a Galois covering space of X corresponding to the normal subgroup N in $\pi_1(X)$ and associated Galois group $\pi_1(X)/N$. For each 2-cell a_α in the relative CW complex (X_1, X) , we can pullback to a collection of 2-cells $\pi^{-1}(a_\alpha)$ of the relative CW complex (\tilde{X}_1, \hat{X}) and $\pi_1(X)$ acts transitively on these 2-cells with stabilizer group $\pi_1 X/N$. Therefore, $H_*(\tilde{X}_1, \hat{X}; \mathbb{Z})$ is a free $\mathbb{Z}[\pi_1 X/N]$ -module on generators $[\tilde{a}_\alpha]$, where \tilde{a}_α is a lift of a 2-cell a_α of (X_1, X) to (\tilde{X}_1, \hat{X}) . We then apply the Hurewicz homomorphism vertically to the long exact sequence in homotopy associated to the pair (\tilde{X}_1, \hat{X}) to form the commutative diagram

$$\begin{array}{ccccccc} \pi_2(\hat{X}) & \longrightarrow & \pi_2(\tilde{X}_1) & \longrightarrow & \pi_2(\tilde{X}_1, \hat{X}) & \longrightarrow & \pi_1(\hat{X}_1) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ H_2(\hat{X}; \mathbb{Z}) & \longrightarrow & H_2(\tilde{X}; \mathbb{Z}) & \longrightarrow & H_2(\tilde{X}_1, \hat{X}; \mathbb{Z}) & \longrightarrow & H_1(\hat{X}; \mathbb{Z}). \end{array}$$

Since N is perfect, $H_1(\hat{X}; \mathbb{Z}) = N/[N, N] = 0$. Also, since $\pi_1(\tilde{X}_1) = \pi_0(\tilde{X}_1) = 0$, the Hurewicz theorem implies that $\pi_2(\tilde{X}) \cong H_2(\tilde{X}; \mathbb{Z})$. We know therefore know the composite map

$$\pi_2(\tilde{X}_1) \rightarrow H_2(\tilde{X}_1; \mathbb{Z}) \rightarrow H_2(\tilde{X}_1, \hat{X}; \mathbb{Z})$$

is surjective. For each $[\tilde{a}_\alpha]$ we choose a lift $[\tilde{f}_\alpha]$ to $\pi_2(\tilde{X}_1)$ represented by a basepoint preserving map $\tilde{f}_\alpha: S^2 \rightarrow \tilde{X}_1$. Define

$$f_\alpha = \pi \circ \tilde{f}_\alpha$$

and let X^+ be the pushout

$$\begin{array}{ccc} \coprod_{\alpha} S^2 & \xrightarrow{f_\alpha} & X_1 \\ \downarrow & & \downarrow \\ \coprod_{\alpha} D^3 & \longrightarrow & X^+. \end{array}$$

Remark 3.1.7. At various points in the construction, we made choices. We leave it to the reader to check that the choices we made do not matter in the

sense that a different choice would lead to a space that is homeomorphic to the space X^+ .

We now have to check that this construction has the desired properties. First, note that we only attached 3-cells to X_1 to form X^+ , so

$$\pi_1(X^+) \cong \pi_1(X)/N$$

as desired.

Write \tilde{X}^+ for the universal cover of X^+ . Then we have a diagram

$$\begin{array}{ccccc} \hat{X} & \longrightarrow & \tilde{X}_1 & \longrightarrow & \tilde{X}^+ \\ \downarrow & & \downarrow & & \downarrow \\ X & \longrightarrow & X_1 & \longrightarrow & X^+ \end{array}$$

where each square in the diagram is a pullback. Again, since X^+ was formed from X by only attaching only 2-cells and 3-cells, we know that \tilde{X}^+ is formed from \hat{X} by attaching only 2-cells and 3-cells. Consequently, the relative cellular chains $C_*(\tilde{X}^+, \hat{X}; \mathbb{Z})$ are concentrated in two degrees

$$0 \rightarrow C_3(\tilde{X}^+, \hat{X}) \xrightarrow{d} C_2(\tilde{X}^+, \hat{X}) \rightarrow 0. \quad (3.1.8)$$

We then observe that there are isomorphisms

$$C_3(\tilde{X}^+, \hat{X}) \cong C_3(\tilde{X}^+, \tilde{X}_1) \cong H_3(\tilde{X}^+, \tilde{X}_1; \mathbb{Z})$$

$$C_2(\tilde{X}^+, \hat{X}) \cong C_2(\tilde{X}_1, \hat{X}) \cong H_2(\tilde{X}_1, \hat{X}; \mathbb{Z})$$

and the boundary map d in (3.1.8) is exactly the boundary map

$$d: H_3(\tilde{X}^+, \tilde{X}_1; \mathbb{Z}) \rightarrow H_2(\tilde{X}_1, \hat{X}; \mathbb{Z})$$

associated to the long exact sequence of the triple $(\tilde{X}^+, \tilde{X}_1, \hat{X})$. Therefore, the map d from (3.1.8) factors as

$$d: H_3(\tilde{X}^+, \tilde{X}_1; \mathbb{Z}) \xrightarrow{\partial} H_2(\tilde{X}_1; \mathbb{Z}) \xrightarrow{j} H_2(\tilde{X}_1, \hat{X}; \mathbb{Z})$$

where ∂ is the boundary map in the long exact sequence for the pair $(\tilde{X}^+, \tilde{X}_1)$ and j is induced by the canonical map of pairs $(\tilde{X}_1, \emptyset) \rightarrow (\tilde{X}_1, \hat{X})$.

We then examine the diagram

$$\begin{array}{ccccccc} \dots & \longrightarrow & \pi_3(\tilde{X}^+) & \longrightarrow & \pi_3(\tilde{X}^+, \tilde{X}_1) & \longrightarrow & \pi_2(\tilde{X}_1) \\ & & \downarrow & & \downarrow & & \downarrow \\ \dots & \longrightarrow & H_3(\tilde{X}^+; \mathbb{Z}) & \longrightarrow & H_3(\tilde{X}^+, \tilde{X}_1; \mathbb{Z}) & \xrightarrow{\partial} & H_2(\tilde{X}_1; \mathbb{Z}) \end{array}$$

where the vertical maps are the Hurewicz maps and the horizontal maps are the long exact sequence of a pair. Given an element

$$[\tilde{b}_\alpha] \in C_3(\tilde{X}^+, \tilde{X}_1; \mathbb{Z}) \cong H_3(\tilde{X}, \tilde{X}_1; \mathbb{Z})$$

corresponding to a 3-cell \tilde{b}_α of the relative CW complex (\tilde{X}^+, \tilde{X}) . The attaching map $\tilde{f}_\alpha: S^2 \rightarrow \tilde{X}_1$ of the 3-cell of $(\tilde{X}^+, \tilde{X}_1)$ corresponds to a class $[\tilde{f}_\alpha]$ of $\pi_2(\tilde{X}_1)$ and maps to a class $[\tilde{f}_\alpha]$ in $H_2(\tilde{X}_1; \mathbb{Z})$. So by inspection $[\tilde{b}_\alpha]$ maps to $[\tilde{f}_\alpha]$ under the map

$$\delta: H_3(\tilde{X}^+, \tilde{X}_1; \mathbb{Z}) \rightarrow H_2(\tilde{X}_1; \mathbb{Z}).$$

In addition, the map

$$j: H_2(\tilde{X}_1; \mathbb{Z}) \rightarrow H_2(\tilde{X}_1, \hat{X}; \mathbb{Z})$$

sends $[\tilde{f}_\alpha]$ to $[\tilde{a}_\alpha]$.

By the same argument as before, $H_3(\tilde{X}^+, \tilde{X}_1; \mathbb{Z})$ is a free $\mathbb{Z}[\pi_1(X)/N]$ -module on generators $[\tilde{b}_\alpha]$ and similarly $H_2(\tilde{X}^+, \hat{X})$ is a free $\mathbb{Z}[\pi_1(X)/N]$ -module on generators $[\tilde{a}_\alpha]$ where α ranges over the same indexing set I . Thus, the map d is an isomorphism of $\mathbb{Z}[\pi_1(X)/N]$ -modules and consequently $C_*(\tilde{X}^+, \hat{X})$ is acyclic.

Moreover, for any $\mathbb{Z}[\pi_1(X)/N]$ -module L , the chains $C_*(\tilde{X}^+, \hat{X}; L)$ whose homology is $H_*(X^+, X; L)$ are defined to be

$$C_*(\tilde{X}^+, \hat{X}; L) \cong L \otimes_{\mathbb{Z}[\pi_1(X)/N]} C_*(\tilde{X}^+, \hat{X}; \mathbb{Z})$$

and consequently $C_*(\tilde{X}^+, \hat{X}; L)$ is also an acyclic chain complex. In particular,

$$H_*(X; \mathbb{Z}) \rightarrow H_*(X^+; \mathbb{Z})$$

is an isomorphism. Moreover, we have shown the following.

Lemma 3.1.9. *For any $\mathbb{Z}[\pi_1 X]$ -module L , the groups*

$$H^*(X^+, X; L) \cong 0.$$

We still need to show that X^+ satisfies the universal property of Definition (3.1.3). The proof will be by obstruction theory. First, we give a general setup of obstruction theory.

Let (X, A) be a based relative CW complex with base point $x_0 \in A$. Assume that X and A are path connected. Given a continuous pointed map $f: A \rightarrow Y$ where Y is path connected,

1. when does there exist an extension F

$$\begin{array}{ccc} A & \xrightarrow{f} & Y \\ \downarrow i & \nearrow F & \\ X & & \end{array}$$

such that the diagram commutes, where $i: A \rightarrow X$ is the canonical inclusion?

2. Given two such extensions $F, F': X \rightarrow Y$ such that $F|_A = F'|_A = f$, when is there a homotopy $H: X \times I \rightarrow Y$ rel A from F to F' ?

Write X_n for the n -skeleton of X . Then we can always extend

$$\begin{array}{ccc} A & \xrightarrow{f} & Y \\ \downarrow i & \nearrow F_1 & \\ X_1 & & \end{array}$$

because if $g: \coprod S^0 \rightarrow A$ is the attaching map for attaching the 1-cells to A to form X_1 , when composing this map with f

$$\coprod S^0 \rightarrow A \rightarrow Y$$

necessarily produces a null homotopy map, because Y is path connected. therefore, we can let F_1 be any null homotopic basepoint preserving map $F_1: X_1 \rightarrow Y$. Note that $\pi_1(X_2) \cong \pi_1(X)$, so if it is possible to extend further

$$\begin{array}{ccc} A & \xrightarrow{f} & Y \\ \downarrow & \nearrow F_2 & \\ X_1 & & \\ \downarrow & \nearrow & \\ X_2 & & \end{array} \quad (3.1.10)$$

then there would be a homomorphism θ in a commuting diagram

$$\begin{array}{ccc} \pi_1 A & \xrightarrow{f} & \pi_1 Y \\ \downarrow & \nearrow \theta & \\ \pi_1 X & & \end{array} \quad (3.1.11)$$

In fact, this is an if and only if, so if there is a group homomorphism $\theta: \pi_1 X \rightarrow \pi_1 Y$ making the diagram commute, then there exists an extension F_2 making the diagram (3.1.10) commute.

Now suppose A extends to X_2 and fix a group homomorphism $\theta: \pi_1 X \rightarrow \pi_1 Y$ such that the diagram (3.1.11) commutes. Now suppose we want to

extend A to X_3

$$\begin{array}{ccc} A & \xrightarrow{f} & Y \\ \downarrow & \nearrow F_3 & \\ X_3 & & \end{array} \quad (3.1.12)$$

compatible with the extensions F_1 and F_2 , which amounts to asking that on π_1 the diagram (3.1.12) is the diagram (3.1.11) up to isomorphism. Recall that $\pi_1(Y)$ acts on $\pi_i(Y)$ for $i \geq 1$, so $\pi_i(Y)$ is a $\mathbb{Z}[\pi_1 Y]$ -module for all $i \geq 2$ (and all $i \geq 1$ if Y is an H -space, as is the case in our main example of interest). Since we have a group homomorphism $\theta: \pi_1(X) \rightarrow \pi_1(Y)$ by assumption, this induces a map $\mathbb{Z}[\pi_1 X] \rightarrow \mathbb{Z}[\pi_1 Y]$ of group rings and therefore we may regard $\pi_i(Y)$ as a $\mathbb{Z}[\pi_1 X]$ -module $\theta^* \pi_i(Y)$ via restriction.

We therefore proceed by induction up the skeleta of X . Suppose we have an extension $F_n: X_n \rightarrow Y$ of f to X_n for some $n \geq 2$ such that $\pi_1(F_n) = \theta: \pi_1(X_n) = \pi_1 X \rightarrow \pi_1 Y$. Then we can define an obstruction cocycle giving an obstruction class

$$\text{Ob}^{n+1}(\theta, F_n) \in H^{n+1}(X, A, \theta^* \pi_n(Y))$$

in the cohomology of the pair (X, A) with coefficients in the $\mathbb{Z}[\pi_1 X]$ -module $\theta^* \pi_n(Y)$. The vanishing of this obstruction class is a necessary and sufficient condition for F_n to extend further to $F_{n+1}: X_{n+1} \rightarrow Y$ such that $F_{n+1}|_{X_n} = F_n$. When $H^n(X, A, \theta^* \pi_n(Y)) \cong 0$ for all $n \geq 2$, then f extends to X .

The second question may actually be regarded as a special case of the first for the pair $(X \times I, X \times \{0, 1\} \cup A \times I)$. Recall that by Van Kampen

$$\pi_1(X \times \{0, 1\} \cup A \times I) \cong \pi_1(X) *_{\pi_1(A)} \pi_1(X)$$

and the map

$$\pi_1(X \times \{0, 1\} \cup A \times I) \rightarrow \pi_1(X)$$

is the canonical quotient. Defining $\tilde{f}: X \times \{0, 1\} \cup A \times I \rightarrow Y$ by

$$\begin{aligned} \tilde{f}|_{X \times 0} &= F \\ \tilde{f}|_{X \times 1} &= F' \\ \tilde{f}|_{A \times I} &= f \circ \pi_1(x, t) \end{aligned}$$

where $\pi_1: A \times I \rightarrow A$ is projection onto the first factor. Then we can say that F is homotopic to F' rel A if and only if the map \tilde{f} extends to a map $\tilde{F}: X \times I \rightarrow Y$. Again, \tilde{f} will extend to the 2-skeleton of $X \times I$ if $\tilde{\theta} = \pi_1(F) = \pi_1(F'): \pi_1 X \rightarrow \pi_1 Y$ by the calculation of fundamental groups above. For $n \geq 1$, the obstructions lie in

$$H^{n+2}(X \times I, X \times \{0, 1\} \cup A \times I; \tilde{\theta}^* \pi_{n+1}(Y))$$

but by inspection of the suspension homomorphism these groups are isomorphic to

$$H^{n+1}(X, A; \tilde{\theta}^* \pi_{n+1}(Y)).$$

We therefore have an obstruction class

$$\text{Ob}^{n+1}(\theta, F, F') \in H^{n+1}(X, A; \tilde{\theta}^* \pi_{n+1}(Y))$$

which is trivial if and only if there is exists a homotopy $H: F \simeq F' \text{ rel } A$. If $H^n(X, A; \tilde{\theta}^* \pi_n(Y)) \cong 0$ for $n \geq 2$, then there always exists a homotopy $H: F \simeq F' \text{ rel } A$.

We now summarize in our special case.

Theorem 3.1.13. *Let X be a based path connected space. The construction X^+ of (3.1.6) has the universal property that for any based path connected commutative H -group Y with a map $X \rightarrow Y$, then there exists a lift h , that is unique up to homotopy*

$$\begin{array}{ccc} X & \longrightarrow & X^+ \\ & \searrow & \downarrow h \\ & & Y \end{array}$$

Proof. Since Y is path connected, there is no obstruction to extending to the 1-skeleton of X^+ (the one skeleton of X^+ is X anyways, so there is nothing to show). The obstruction to extending X to the 2-skeleton of X^+ is the existence of a group homomorphism $\theta: \pi_1(X^+) \rightarrow \pi_1(Y)$ such that the diagram

$$\begin{array}{ccc} \pi_1(X) & \longrightarrow & \pi_1(X^+) \\ & \searrow & \downarrow \theta \\ & & \pi_1(Y) \end{array}$$

commutes. This group homomorphism exists by the universal property of $\pi_1(X^+) = \pi_1(X)/[\pi_1(X), \pi_1(X)] =: \pi_1(X)^{\text{ab}}$ since $\pi_1(Y)$ is necessarily an abelian group. Finally, the obstructions to existence of an extension $h: X^+ \rightarrow Y$ and uniqueness of such a map live in $H^*(X^+, X; \theta^* \pi_* Y)$, but these groups vanish by Lemma 3.1.9. Therefore, there exists an extension $X^+ \rightarrow Y$ and this extension is unique up to homotopy rel X as desired. \square

Remark 3.1.14. Note that we still need to show that X^+ is in fact an H -space. We will do this shortly. First, we give an example.

Example 3.1.15. Consider the colimit

$$\Sigma := \bigcup_{n \geq 0} \Sigma_n$$

of Σ_n along the inclusions $\Sigma_n \rightarrow \Sigma_{n+1}$ sending an automorphism of the finite set $\{1, \dots, n\}$ to the corresponding automorphism of the finite set $\{1, \dots, n +$

1} fixing $n + 1$. Then the inclusion of the alternating $A_n \rightarrow \Sigma_n$ is compatible and produces an inclusion of $A = \cup_{n \geq 0} A_n$ in Σ with quotient $\mathbb{Z}/2$. The group A is in fact a perfect normal subgroup of Σ and we can form the +-construction with respect to this perfect normal subgroup. This gives a space $B\Sigma^+$ such that $\pi_1(B\Sigma^+) \cong \mathbb{Z}/2$. In fact, we have the following theorem of Barratt-Priddy-Quillen.

Theorem 3.1.16 (Barratt-Priddy-Quillen). *There is an isomorphism*

$$\pi_k^s(S^0) \rightarrow \pi_k(\mathbb{Z} \times B\Sigma^+)$$

for $k \geq 0$, where $\pi_k^s S^0$ is the k -th stable homotopy groups of spheres.

Remark 3.1.17. We should think of $\mathbb{Z} \times B\Sigma^+$ as an explicit model for the algebraic K-theory space $K(\text{Fin})$ of the category of finite sets. We will prove this later.

Theorem 3.1.18. *The space $BGL(R)^+$ is an commutative H-group and the map $BGL(R)^+ \rightarrow Y$ is a map of H-spaces whenever Y is an H-space.*

We first define a multiplication

$$\mu: BGL(R)^+ \times BGL(R)^+ \rightarrow BGL(R)^+.$$

As we noted in the appendix, geometric realization commutes with finite products, so we first define an operation

$$\mu_0: GL(R) \times GL(R) \rightarrow GL(R)$$

and then define a map

$$B\mu_0 \circ h: BGL(R) \times BGL(R) \rightarrow BGL(R)$$

where h is the homotopy equivalence

$$h: BGL(R) \times BGL(R) \xrightarrow{\sim} B(GL(R) \times GL(R)).$$

We define $\mu_0(A, B)$ for matrices $A, B \in GL(R)$ by the formula

$$\mu_0(A, B) = (c_{i,j})$$

for $0 \leq i, j \leq n + m$ where

$$c_{i,j} = \begin{cases} a_{k,\ell} & \text{if } i = 2k - 1, j = 2\ell - 1 \\ b_{k,\ell} & \text{if } i = 2k, j = 2\ell \\ 0 & \text{otherwise} \end{cases}$$

where $A = (a_{k,\ell})$ for $0 \leq k, \ell \leq n$ and $B = (b_{k,\ell})$ for $0 \leq k, \ell \leq m$. We leave the following results as simple exercises.

Exercise 3.1.19. The map μ_0 is a group homomorphism.

Exercise 3.1.20. There is a homotopy equivalence

$$h': BGL(R)^+ \times BGL(R)^+ \rightarrow (BGL(R) \times BGL(R))^+.$$

We therefore have an induced map

$$\mu: BGL(R)^+ \times BGL(R)^+ \rightarrow BGL(R)^+$$

We now construct an inverse. First, let $u: \mathbb{N} \rightarrow \mathbb{N}$ be an injective self-map of the set of positive integers. Define

$$u_\bullet: GL(R) \rightarrow GL(R)$$

where $A = (a_{i,j})$ by

$$u_\bullet(A) = \begin{cases} a_{i,j} & \text{if } (i,j) = (u(i), u(j)) \\ \delta_{i,j} & \text{otherwise} \end{cases}$$

where $\delta_{i,j} = 1$ if $i = j$ and 0 otherwise.

Exercise 3.1.21. The map u_\bullet is a group homomorphism.

Lemma 3.1.22. For each u_\bullet , there is an induced homotopy equivalence

$$u^+: BGL(R)^+ \rightarrow BGL(R)^+.$$

Proof. By Exercise 3.1.21, the map u_\bullet induces a map $u^+: BGL(R)^+ \rightarrow BGL(R)^+$. This induces a $K_1(R)$ -equivariant self-map

$$\bar{u}: BE(R)^+ \rightarrow BE(R)^+$$

of the universal cover $\tilde{BGL}(R)^+ = BE(R)^+$. Since $BE(R)^+$ is simply connected, it suffices to check that

$$H_*(\bar{u}; \mathbb{Z}): H_*(BE(R)^+; \mathbb{Z}) \rightarrow H_*(BE(R)^+; \mathbb{Z})$$

is an isomorphism by the Whitehead theorem. To see this, we note that \bar{u} is induced by the map $u_\bullet|_{E(R)}: E(R) \rightarrow E(R)$, which is equal to conjugation by some matrix C so the induced map $H_*(\bar{u})$ is also given by conjugation by C , but inner conjugation induces the identity map on group homology with integer coefficients. \square

Exercise 3.1.23. Let M be the monoid of injective self maps of \mathbb{N} under composition, then $K_0(M) \cong 0$.

Consequently, any u^+ is homotopic to the identity via basepoint preserving maps.

We say group homomorphisms $f, g: G \rightarrow GL(R)$ are pseudo-conjugate if there exists $u \in M$, from Exercise 3.1.23, such that either $u_\bullet \circ f$ and g or f and $u_\bullet \circ g$ are conjugate by an element in $GL(R)$.

Proposition 3.1.24. *If $f, g: G \rightarrow GL(R)$ are pseudo-conjugate, then the induced maps $f^+, g^+: BG \rightarrow BGL(R)^+$ are homotopy as maps preserving the basepoints*

Proof. For any map $f: G \rightarrow GL(R)$ write f' for

$$f'(-) = \mu(f(-), 1).$$

Then f' is homotopic to f by a base point preserving homotopy because $f' = v_\bullet \circ f$, where $v \in M$ is defined by $v(s) = 2s - 1$.

Suppose $f, g: G \rightarrow GL(R)$ satisfy $g(-) = \alpha(u_\bullet \circ f(-))\alpha^{-1} := (u_\bullet \circ f(-))^\alpha$ for $\alpha \in GL(R)$. Letting $\beta = \mu(\alpha, \alpha^{-1}) \in E(R)$, then $\mu(g, 1) = \mu(u_\bullet \circ f, 1)^\beta$. So the induced maps

$$BG \rightarrow BGL(R)$$

are homotopic such that under a homotopy, the image of the base point of BG is a loop homotopic to $[\beta] \in \pi_1 BGL(R) = GL(R)$. The induced maps $BG \rightarrow BGL(R)^+$ are homotopic preserving the base points since $[\beta] = 0 \in \pi_1(BGL(R)^+)$. \square

We can now prove that $BGL(R)^+$ is a commutative H -space with operation

$$\mu: BGL(R)^+ \times BGL(R)^+ \rightarrow BGL(R)^+$$

and identity given by inclusion of the basepoint $e: * \rightarrow BGL(R) \rightarrow BGL(R)^+$.

Proof of Theorem 3.1.18. Let $v, w \in M$ of Exercise 3.1.23 be given by $v(s) = 2s - 1$ and $w(s) = 2s$ for all $s \in \mathbb{N}$. Then for any $A \in GL(R)$, we have that $v_\bullet \circ A = \mu(A, 1)$ and $w_\bullet \circ A = \mu(1, A)$. By construction, the composite

$$BGL(R)^+ \times * \rightarrow BGL(R)^+ \times BGL(R)^+ \xrightarrow{\mu^+} BGL(R)^+$$

is homotopic to v^+ and the the composite

$$* \times BGL(R)^+ \rightarrow BGL(R)^+ \times BGL(R)^+ \xrightarrow{\mu^+} BGL(R)^+$$

is homotopic to w^+ . So the inclusion $e: * \rightarrow BGL(R)^+$ is a two-sided identity up to pointed homotopy. Consequently, $BGL(R)^+$ is an H -space.

There exists $v', w' \in M$ such that for any $x, y, z \in GL(R)$

$$\mu(x, y) = v' \circ \mu(y, x)$$

$$\mu(\mu(x, y), z) = w' \circ \mu(x, \mu(y, z))$$

so $BGL(R)^+$ is homotopy commutative and homotopy associative. Since it is path-connected and it is the homotopy type of a CW complex it is in fact a commutative H -group by [19, Cor. A.47]. \square

Chapter 4

The Q-construction and fundamental theorems

In this chapter, we first setup and prove Quillen's famous Theorem A and Theorem B. These will be used in a key way for proving theorems about the Q-construction of algebraic K-theory. We then give Quillen's Q-construction of algebraic K-theory and prove some basic properties. We then prove some of the fundamental theorems of algebraic K-theory using this construction. The primary reference for this material is the original paper of Quillen [15].

4.1 Quillen's Theorem A and Theorem B

Recall from Definition B.1.30, that given a small category \mathcal{C} , we can form an associated simplicial set $N_\bullet \mathcal{C}$.

Definition 4.1.1. Let \mathcal{C} be a small category and let $N_\bullet \mathcal{C}$ be the simplicial set with n -simplices

$$N_n \mathcal{C} = \text{Fun}([n], \mathcal{C})$$

then we define the *classifying space* of \mathcal{C} to be

$$B\mathcal{C} := |N_\bullet \mathcal{C}|.$$

In this section, we will study the homotopy theory of small categories, by which we mean the homotopy theory of spaces that arise as the classifying space of a small category. Throughout this section, we assume all categories are *small* categories unless otherwise specified. The following definition will be used in several places.

Definition 4.1.2. The comma category $S \downarrow T$ associated to a pair of functors

$$S: \mathcal{A} \rightarrow \mathcal{C} \leftarrow \mathcal{B}: T$$

has objects $(a \in \mathcal{A}, b \in \mathcal{B}, \alpha: S(A) \rightarrow T(B))$ and morphisms $(a, b, \alpha) \rightarrow (a', b', \alpha')$ given by triples

$$f: a \rightarrow a'$$

$$g: b \rightarrow b'$$

and a commuting diagram

$$\begin{array}{ccc} S(a) & \xrightarrow{S(f)} & S(a') \\ \alpha \downarrow & & \downarrow \alpha' \\ T(b) & \xrightarrow{T(g)} & T(b') \end{array}$$

where composition is defined in the evident way.

Examples 4.1.3. Given a category \mathcal{C} and an object $X \in \mathcal{C}$, we define the slice category \mathcal{C}/X to be the comma category corresponding to the span of categories

$$\mathcal{C} \xrightarrow{\text{id}_{\mathcal{C}}} \mathcal{C} \xleftarrow{X} [0]$$

and we define the coslice category $X \backslash \mathcal{C}$ to be the comma category associated to the span of categories

$$[0] \xrightarrow{X} \mathcal{C} \xleftarrow{\text{id}_{\mathcal{C}}} \mathcal{C}.$$

We also define f/Y , where $f: \mathcal{C} \rightarrow \mathcal{C}'$ is a functor and Y is an object in \mathcal{C}' to be the comma category associated to the span of categories

$$\mathcal{C} \xrightarrow{f} \mathcal{C}' \xleftarrow{Y} [0]$$

and similarly we define $X \backslash f$ to be the comma category associated to the span of categories

$$[0] \xrightarrow{X} \mathcal{C}' \xleftarrow{f} \mathcal{C}$$

The following lemmas will be of fundamental importance.

Lemma 4.1.4. *A natural transformation $\eta: f \Rightarrow g$ between functors*

$$f, g: \mathcal{C} \rightarrow \mathcal{D}$$

induces a homotopy

$$H: \mathcal{B}\mathcal{C} \times I \rightarrow \mathcal{D}$$

between Bf and Bg .

Proof. The triple (f, g, η) defines a functor

$$(f, g, \eta): \mathcal{C} \times [1] \rightarrow \mathcal{D}$$

sending $(c, 0)$ to $f(c)$, $(c, 1)$ to $g(c)$ and $(c, 0 \rightarrow 1)$ to $\eta: f(c) \rightarrow g(c)$. It is defined on morphisms in the evident way. We then apply the nerve and note that

$$N_\bullet(\mathcal{C} \times [1]) = N_\bullet(\mathcal{C}) \times \Delta^1$$

and by a result of Milnor geometric realization commutes with finite products so the functor (f, g, η) induces a continuous map

$$H: BC \times I \rightarrow BD$$

such that $H|_{BC \times \{0\}} = Bf$ and $H|_{BC \times \{1\}} = Bg$ as desired. \square

As a consequence of this, we can also prove the following lemma of fundamental importance.

Lemma 4.1.5. *If a functor*

$$f: \mathcal{C} \rightarrow \mathcal{D}$$

has a right or a left adjoint, then f induces a homotopy equivalence

$$Bf: BC \simeq BD.$$

Proof. Suppose f has a right adjoint g . The proof when f has a left adjoint is essentially the same so we omit it. Then there are natural transformations

$$\eta: \text{id}_{\mathcal{C}} \rightarrow g \circ f \text{ and } \epsilon: f \circ g \rightarrow \text{id}_{\mathcal{D}}$$

which induce homotopies

$$H_1: BC \times I \rightarrow BC \text{ and } H_2: BD \times I \rightarrow BD$$

from id_{BC} to $Bg \circ Bf$ and from $Bf \circ Bg$ to id_{BD} respectively. \square

Definition 4.1.6. We say a category \mathcal{C} has an initial object 0 if for all objects $c \in \mathcal{C}$ there exists a unique morphism $0 \rightarrow c$. Similarly, we say that \mathcal{C} has a terminal object 1 if for all objects $c \in \mathcal{C}$ there exists a unique morphism $c \rightarrow 1$.

Lemma 4.1.7. *If \mathcal{C} has an initial or terminal object then BC is contractible.*

Proof. The existence of an initial object in \mathcal{C} is equivalent to the existence of a left adjoint to the functor $[0] \rightarrow \mathcal{C}$ sending 0 to the initial object 0 of \mathcal{C} . Similarly, the existence of a terminal object is equivalent to the existence of a right adjoint to the functor $[0] \rightarrow \mathcal{C}$ sending 0 to the terminal object 1 of \mathcal{C} . \square

We first give a characterization of $\pi_1(BC)$, but we will not give a complete proof. See [19, Lemma 6.1.] for a careful proof of most of the statements. First, we give a definition.

Definition 4.1.8. We say that a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is *morphism inverting* if for all morphisms f in \mathcal{C} , then $F(f)$ is an isomorphism.

Theorem 4.1.9. *There is an equivalence of categories between the category of coverings of BC , denote $\text{Cov}(BC)$ and the category of morphism inverting functors $\mathcal{C} \rightarrow \text{Set}$, which we denote $\text{Fun}^{\text{inv}}(\mathcal{C}, \text{Set})$. Moreover,*

$$\pi_1(\mathcal{C}, X) \cong \text{Hom}_{\mathcal{C}}(X, X)^{\text{gp}}$$

where $\text{Hom}_{\mathcal{C}}(X, X)^{\text{gp}}$ is group completion of the monoid $\text{Hom}_{\mathcal{C}}(X, X)$.

We will just sketch the proof. Given a covering space $E \rightarrow BC$, we consider the fiber $E(X)$ over an object $X \in \mathcal{C}$ regarded as a 0-cell of the CW complex BC . A map $f: X \rightarrow X'$ induces a bijection $E(f): E(X) \cong E(X')$ of sets satisfying $E(f \circ g) = E(f) \circ E(g)$. Thus, we may regard E as a functor

$$E: \mathcal{C} \rightarrow \text{Set}$$

that sends all maps f to bijections $E(f)$ as desired.

In the other direction, given a functor $F: \mathcal{C} \rightarrow \text{Set}$ we post-compose with the inclusion $\text{Set} \rightarrow \text{Cat}$ sending a set S to the category with objects S and only identity morphisms, and denote this functor F by abuse of notation. Then we define the $[0] \setminus F$ to be the comma category associated to the functors

$$[0] \xrightarrow{[0]} \text{Cat} \xleftarrow{F} \mathcal{C}$$

where the left functor sends 0 to the category $[0]$. In other words, unpacking the definition, $[0] \setminus F$ has objects specified by a pair $(X, * \rightarrow F(X))$ and morphisms $f: X \rightarrow X'$ along with a basepoint preserving map $F(X) \rightarrow F(X')$.

Lemma 4.1.10. *The forgetful functor $[0] \setminus F \rightarrow \mathcal{C}$ induces a covering*

$$B([0] \setminus F) \rightarrow BC$$

when $F: \mathcal{C} \rightarrow \text{Set}$ is morphism inverting.

Proof. This follows from [6, Appendix I 3.2]. □

There is a way to formally add inverses to all morphisms in a small category \mathcal{C} to form a groupoid $G = \mathcal{C}[\text{Arr}(\mathcal{C})^{-1}]$ [6, Chapter I, 1.1]. The group $\pi_1(BC, X)$ is then the automorphisms of X in this groupoid G . Moreover, the category of morphism inverting functors

$$\mathcal{C} \rightarrow \text{Set}$$

is equivalent to the category of functors

$$G \rightarrow \text{Set}$$

which we refer to as the category $G\text{-Set}$ of G -sets. When BC is path connected with a 0-simplex X , then the category of coverings of BC is equivalent to the category $G_X\text{-Set}$ of G_X -sets in the traditional sense where G_X is the automorphism group of X in the groupoid $G = \mathcal{C}[\text{Arr}(\mathcal{C})^{-1}]$.

Definition 4.1.11. We say a category \mathcal{I} is a *filtered category* if it is nonempty and for any two objects x, y there exists an object z and morphisms $x \rightarrow z$ and $y \rightarrow z$ and for any two morphisms $f, g: x \rightarrow y$ there exists a morphism $h: y \rightarrow z$ such that $h \circ f = h \circ g$.

Proposition 4.1.12. *Given a functor*

$$I \rightarrow \text{Cat}$$

from a filtered category I to the category of small categories Cat with limit \mathcal{C} , where we denote \mathcal{C}_i the functor evaluated at i , then

$$\text{colim}_i \pi_n BC_i \simeq \pi_n BC$$

for all $n \geq 0$.

If in addition, $BC_i \rightarrow BC_j$ is a homotopy equivalence for all maps $i \rightarrow j$ in I , then there is a homotopy equivalence

$$BC_i \simeq BC$$

for all i .

Corollary 4.1.13. *Any filtered category I is contractible.*

Theorem 4.1.14 (Theorem A). *If $f: \mathcal{C} \rightarrow \mathcal{C}'$ is a functor and the category $A \setminus f$ is contractible for all objects $A \in \mathcal{C}'$, then*

$$Bf: BC \rightarrow BC'$$

is a homotopy equivalence. Similarly, when f/Y is contractible for all objects $A \in \mathcal{C}'$, then the map Bf is a homotopy equivalence.

We will just prove the case $A \setminus f$ since the other version is formally dual in a precise sense. First, we will need the following definition.

Definition 4.1.15. We also need the following variant of the twisted arrow category, denoted $\text{Tw}(f)$ where $f: \mathcal{C} \rightarrow \mathcal{C}'$ is a functor. The objects in $\text{Tw}(f)$ are the same as the objects of the comma category associated to

$$\mathcal{C}' \xrightarrow{\text{id}} \mathcal{C}' \xrightarrow{F} \mathcal{C}$$

and the morphisms from $(x, y, \alpha: x \rightarrow F(y))$ to $(x', y', \alpha': x' \rightarrow F(y'))$ consist of maps $u: x \rightarrow x'$ and $w: y' \rightarrow y$ such that the diagram

$$\begin{array}{ccc} x & \xrightarrow{\alpha} & F(y) \\ u \downarrow & & \uparrow F(w) \\ x' & \xrightarrow{\alpha'} & F(y') \end{array}$$

commutes. Composition is defined by evident vertical composition of squares. When $f: \mathcal{C}' \rightarrow \mathcal{C}'$ is the identity $\text{id}_{\mathcal{C}'}$, we simply write $\text{Tw}(\mathcal{C}') := \text{Tw}(\text{id}_{\mathcal{C}'})$ and we observe that this is the usual twisted arrow category.

The following construction will also be integral to the proof.

Construction 4.1.16. Given a bisimplicial space, or in other words a functor

$$X_{\bullet, \bullet}: \Delta^{\text{op}} \times \Delta^{\text{op}} \rightarrow \text{Top}$$

we can form the geometry realizations

$$|X_{\bullet, n}|: \Delta^{\text{op}} \rightarrow \text{Top}$$

and

$$|X_{n, \bullet}|: \Delta^{\text{op}} \rightarrow \text{Top}$$

for each $[n]$ in Δ^{op} and these are functorial in the variable $[n]$. One can also form a simplicial object $\Delta^* X_{\bullet, \bullet}$ with n -simplices $X_{n, n}$ by precomposition with the diagonal

$$\Delta^* X_{\bullet, \bullet}: \Delta^{\text{op}} \xrightarrow{\Delta} \Delta^{\text{op}} \times \Delta^{\text{op}} \rightarrow \mathcal{C}.$$

Lemma 4.1.17. *There are homeomorphisms*

$$\begin{aligned} |[n] \mapsto |X_{n, \bullet}|| &\cong |[n] \mapsto |X_{\bullet, n}|| \\ &\cong |\Delta^* X_{\bullet, \bullet}| \end{aligned}$$

Proof of Theorem A.. Let $f: \mathcal{C} \rightarrow \mathcal{C}'$ be a functor. Consider the diagram of categories

$$(\mathcal{C}')^{\text{op}} \xleftarrow{\pi_1} \text{Tw}(f) \xrightarrow{\pi_2} \mathcal{C}$$

where $\pi_1(x, y, \alpha: x \rightarrow f(y)) = x$ and $\pi_2(x, y, \alpha: x \rightarrow f(y)) = y$ and these functors are defined on morphisms and composition in the evident way. We form a bisimplicial set $T_{p, q}$ with p, q simplices given by pairs

$$(y_p \rightarrow \cdots \rightarrow y_0 \rightarrow f(x_0), x_0 \rightarrow \cdots x_q)$$

where $y_p \rightarrow \cdots \rightarrow y_0 \rightarrow f(x_0)$ is an object in $\text{Fun}([p], (\mathcal{C}'/f(x_0))^{\text{op}})$ and $x_0 \rightarrow \cdots x_q$ is an object in $\text{Fun}([q], \mathcal{C})$. where face maps in p direction (resp. q direction) composes $x_{i+1} \rightarrow x_i \rightarrow x_{i-1}$ to form a map $x_{i+1} \rightarrow x_{i-1}$ and degeneracies are given by inserting an identity, in the same way that we defined the nerve of a category. By forgetting the first component, there is therefore an evident map

$$T_{p, q} \rightarrow N_q \mathcal{C}$$

of bisimplicial sets, where we regard $N_q \mathcal{C}$ is constant in p . Upon geometric realizations, using any of the three equivalent constructions, we produce a map

$$B\pi_2: B\text{Tw}(f) \rightarrow B\mathcal{C}.$$

If we first, geometrically realize with respect to p , then we produce a map of simplicial spaces defined on q -simplices by

$$\coprod_{x_0 \rightarrow \dots x_q} B(\mathcal{C}'/f(x_0)^{\text{op}}) \rightarrow \coprod_{x_0 \rightarrow \dots x_q} *$$

Since $\mathcal{C}'/f(x_0)$ has a final object and $B\mathcal{D} \cong B\mathcal{D}^{\text{op}}$, we know that $B(\mathcal{C}'/f(x_0)^{\text{op}})$ is contractible this map is a weak equivalence. Since both sides are Reedy cofibrant (we will not go into detail on this point) this map induces a weak equivalence

$$BTw(f) \simeq BC$$

We similarly have a functor

$$T_{p,q} \rightarrow N_{\bullet}((\mathcal{C}')^{\text{op}}).$$

Taking geometric realization with respect to q , we produce a map

$$\coprod_{y_p \rightarrow \dots \rightarrow y_0} B(y_0 \setminus f) \rightarrow \coprod_{y_p \rightarrow \dots y_0} * = N_p(\mathcal{C}')^{\text{op}}.$$

By assumption, $B(y_0 \setminus f)$ is contractible so the map is a weak equivalence.

$$BTw(f) \rightarrow B(\mathcal{C}')^{\text{op}} \cong B(\mathcal{C}').$$

This shows that BC' and BC are homotopy equivalent, but we still want to know that the functor $f: \mathcal{C} \rightarrow \mathcal{C}'$ induces a homotopy equivalence. Note that the construction $Tw(f)$ is functorial in functors to \mathcal{C}' , so we have a functor

$$Tw(f) \rightarrow Tw(\text{id}_{\mathcal{C}'}) = Tw(\mathcal{C}')$$

and the projection maps are also natural so there is a commutative diagram

$$\begin{array}{ccccc} (\mathcal{C}')^{\text{op}} & \xleftarrow{\pi_1} & Tw(f) & \xrightarrow{\pi_2} & \mathcal{C} \\ \downarrow \text{id}_{\mathcal{C}'} & & \downarrow & & \downarrow f \\ (\mathcal{C}')^{\text{op}} & \xleftarrow{\pi_1} & Tw(\mathcal{C}') & \xrightarrow{\pi_2} & \mathcal{C}' \end{array}$$

The same argument as before proves that the functor $B\pi_1: BTw(\mathcal{C}') \rightarrow BC'$ is a homotopy equivalence by simply observing that

$$B(y_0 \setminus \text{id}_{\mathcal{C}'}) = B(y_0 \setminus \mathcal{C}')$$

which clearly has an initial object. Therefore, we observe that the functor $f: \mathcal{C} \rightarrow \mathcal{C}'$ induces a homotopy equivalence $BF: BC \rightarrow BC'$ as desired. \square

The following special case is also of interest. For this special case, we need a definition.

Definition 4.1.18. We say $f: \mathcal{C} \rightarrow \mathcal{C}'$ is *pre-fibered* if and only if the subcategory $f^{-1}(y)$, which is the full subcategory of \mathcal{C} whose objects are x such that $f(x) = y$, equipped with a canonical functor

$$f^{-1}(y) \rightarrow y \backslash f$$

sending x to $(x, \text{id}: f(x) \rightarrow y)$ has a right adjoint. We say f is *pre-cofibered* if the same canonical functor has a left adjoint. Denote the right adjoint (resp. left adjoint) by (x, v) maps to v^*x (resp. (x, v) maps to v_*X) when f is pre-fibered (resp. cofibered). Then for any map $v: y \rightarrow y'$ in \mathcal{C}' , there is a functor

$$v^*: f^{-1}(y) \rightarrow f^{-1}(y')$$

(respectively there is a functor $v_*: f^{-1}(y) \rightarrow f^{-1}(y')$). We say f is *fibered* if for every pair of composable morphism v, w in \mathcal{C}' there is a natural isomorphism

$$v^*w^* \rightarrow (v \circ w)^*$$

of functors (resp. we say f is *cofibered* if there is a natural isomorphism $(v \circ w)_* \rightarrow v_* \circ w_* \rightarrow$). We call v^* base change and v_* cobase change.

Corollary 4.1.19. Suppose $f: \mathcal{C} \rightarrow \mathcal{C}'$ is either pre-fibered or pre-cofibered and suppose that $f^{-1}(y)$ is contractible for all $y \in \mathcal{C}'$, then Bf is a homotopy equivalence.

We first introduce some terminology.

Definition 4.1.20. Given a map of based topological spaces $f: (E, e) \rightarrow (B, b)$, we can form the pullback

$$\begin{array}{ccc} F(f) & \longrightarrow & B^I \\ \downarrow & & \downarrow \\ E \times * & \xrightarrow{f \times b} & B \times B \end{array}$$

which we call the *homotopy fiber* of the map f . We say $f: E \rightarrow B$ is a *quasi-fibration* if the map $f^{-1}(b) \rightarrow F(f)$ is a weak equivalence, meaning it induces an isomorphism on all homotopy groups for all basepoints.

Construction 4.1.21. Since the homotopy fiber sequence $F(f) \rightarrow E \rightarrow B$ induces a long exact sequence in homotopy groups

$$\cdots \rightarrow \pi_{i+1}(B, b) \rightarrow \pi_i(F(f), \bar{e}) \rightarrow \pi_i(E, e) \rightarrow \pi_i(B, b) \rightarrow \cdots$$

where $\bar{e} = (e, \text{const}_b)$, we have a long exact sequence in homotopy groups

$$\cdots \rightarrow \pi_{i+1}(B, b) \rightarrow \pi_i(f^{-1}(b), e) \rightarrow \pi_i(E, e) \rightarrow \pi_i(B, b) \rightarrow \cdots$$

as well whenever $E \rightarrow B$ is a quasi-fibration.

The following is a generalization of this construction that will be useful for our setup.

Definition 4.1.22. Given a diagram

$$\begin{array}{ccc} X & \longrightarrow & Y \\ \downarrow & & \downarrow \\ Z & \longrightarrow & W \end{array} \quad (4.1.23)$$

in Top , define the homotopy pullback to be the pullback of the diagram

$$\begin{array}{ccc} hPB & \longrightarrow & W^I \\ \downarrow & & \downarrow \\ Z \times Y & \longrightarrow & W \times W \end{array}$$

and we say (4.1.23) is a homotopy pullback diagram if $X \rightarrow hPB$ is a homotopy equivalence.

As a special case note that, when $Z = *$, then (4.1.23) is a homotopy pullback diagram when $Y \rightarrow W$ is a quasi-fibration, in fact the diagram (4.1.23) is a homotopy pullback whenever Z is contractible and $Y \rightarrow W$ is homotopy equivalent to quasi-fibration.

Theorem 4.1.24 (Theorem B). *Let $f: \mathcal{C} \rightarrow \mathcal{C}'$ be a functor such that for every map $v: y \rightarrow y'$ the induced functor $v \backslash f: y \backslash f \rightarrow y' \backslash f$ is a homotopy equivalence. Then for every $y \in \mathcal{C}'$, there is a homotopy pullback*

$$\begin{array}{ccc} By \backslash f & \longrightarrow & BC \\ \downarrow & & \downarrow \\ By \backslash \mathcal{C}' & \longrightarrow & BC' \end{array}$$

where $By \backslash \mathcal{C}'$ is contractible because the category $y \backslash \mathcal{C}'$ has an initial object. Moreover, there is a long exact sequence in homotopy groups

$$\cdots \rightarrow \pi_{i+1}(BC', y) \rightarrow \pi_i(y \backslash f, \bar{x}) \rightarrow \pi_i(BC, x) \rightarrow \pi_i(BC', y) \rightarrow \cdots$$

where $\bar{x} = (x, \text{id}_y: y \rightarrow f(x))$.

Remark 4.1.25. Again there is a dual formulation in terms of the categories f/y , but we leave the formulation and the proof to the reader.

First, we need three lemmas about quasi-fibrations.

Lemma 4.1.26. *Let $p: E \rightarrow B$ be a continuous map, and let $U, V \subset B$ be open subsets such that $U \cup V = B$. Suppose $p|_{p^{-1}(U)}$, $p|_{p^{-1}(V)}$, and $p|_{p^{-1}(U \cap V)}$ are quasi-fibrations, then p is a quasi-fibration.*

Proof. We know $U \cap V$ is the pullback of

$$U \hookrightarrow B \leftarrow V$$

so $\text{Fib}(p^{-1}(U \cap V) \rightarrow U \cap V)$ is the pullback of $\text{Fib}(p^{-1}(U) \rightarrow U)$ and $\text{Fib}(p^{-1}(V) \rightarrow V)$ along $\text{Fib}(p)$. If $x \in B$ then there is a weak equivalence $p^{-1}(V) \cap p^{-1}(x) \simeq_{we} \text{Fib}(p^{-1}(V) \rightarrow V)$ if $x \in V$ and there is a weak equivalence $p^{-1}(U) \cap p^{-1}(x) \simeq_{we} \text{Fib}(p^{-1}(U) \rightarrow U)$ if $x \in U$ by assumption. Also, if $x \in U \cap V$, then there is a weak equivalence $p^{-1}(U \cap V) \cap p^{-1}(x) \simeq_{we} \text{Fib}(p^{-1}(U \cap V) \rightarrow U \cap V)$ by assumption. Therefore,

$$\begin{aligned} p^{-1}(x) &= p^{-1}(B) \cap p^{-1}(x) \\ &= (p^{-1}(U) \cup p^{-1}(V)) \cap p^{-1}(x) \\ &= (p^{-1}(U) \cap p^{-1}(x)) \cup_{p^{-1}(U \cap V) \cap p^{-1}(x)} (p^{-1}(V) \cap p^{-1}(x)) \\ &\simeq_{we} \text{Fib}(p^{-1}(U)) \cup_{\text{Fib}(p^{-1}(U \cap V))} \text{Fib}(p^{-1}(V)) \\ &\simeq_{we} \text{Fib}(p). \end{aligned}$$

as desired. \square

Lemma 4.1.27. *Let $p: E \rightarrow B$ be a continuous map onto B , let $B' \subset B$ be a subspace and let $E' = p^{-1}(B')$. Suppose there is a fiber preserving deformation; i.e. suppose there are homotopies*

$$D: E \times I \rightarrow E \text{ and } d: B \times I \rightarrow B$$

such that $D(-, 0) = \text{id}_E$, $d(-, 0) = \text{id}_B$, $D_t(E') = E'$, $d_t(B') = B'$ and $D_1(E) \subset E'$ and $d_1(B) \subset B'$.

Additionally, assume that $p^{-1}(x) \rightarrow p^{-1}(d_1(x))$ is a weak equivalence for all $x \in B$ then p is a quasi-fibration.

Lemma 4.1.28. *Let $p: E \rightarrow B$ be continuous map. Assume that B is CW complex with n -skeleton B_i and assume that $p|_{p^{-1}(B_i)}$ is a quasi-fibration for each i , then p is a quasi-fibration.*

Proof. Any compact subset of B lies inside of some B_i so any compact subset of E lies inside of some $p^{-1}B_i = E_i$. Consequently, given $x \in B_i$ and $y \in p^{-1}(x)$, then the homotopy groups of the pair $E, p^{01}(x)$ satisfy

$$\begin{aligned} \pi_n(E, p^{-1}(x); y) &\cong \text{colim}_j \pi_n(p^{-1}(B_j), p^{-1}(x); y) \\ &\cong \text{colim}_j \pi_n(B_j, x) \cong \pi_n(B, x). \end{aligned}$$

since each map $E_i \rightarrow B_i$ is a quasi-fibration. \square

Proposition 4.1.29. *Suppose I is a small category and $X: I \rightarrow \mathbf{Top}$ is a functor. We form a simplicial space by*

$$\coprod_{i_0 \rightarrow i_1 \rightarrow \dots \rightarrow i_n} X(i_0)$$

where the face and degeneracy maps are induced by functoriality of X and the nerve construction of I , which we index over. There is an obvious map of simplicial spaces from this simplicial space to the nerve of I given on n -simplices by

$$\coprod_{i_0 \rightarrow i_1 \rightarrow \dots \rightarrow i_n} X(i_0) \rightarrow \coprod_{i_0 \rightarrow i_1 \rightarrow \dots \rightarrow i_n} *.$$

Suppose that $X_i \rightarrow X_j$ is a weak equivalence for all maps $i \rightarrow j$ in I , then the map induced on geometric realizations

$$\pi: X_I \rightarrow BI$$

is a quasi-fibration where

$$X_I := |[n] \mapsto \coprod_{i_0 \rightarrow i_1 \rightarrow \dots \rightarrow i_n} X(i_0)|$$

Proof. We need to check that the canonical map $\pi^{-1}(i) \rightarrow \text{Fib}(\pi)$ is a homotopy equivalence. It suffices to check that when restricting to the p -skeleton of BI ; i.e. we need to check that the map $\pi^{-1}((BI)_p) \rightarrow (BI)_p$ is a quasi-fibration for each $p \geq 0$. We proceed by induction up the skeleta. The inductive hypothesis holds because on zero 0-cells the map is a trivial fibration. We consider the map from the diagram

$$\begin{array}{ccc} \coprod_{i_0 \rightarrow \dots \rightarrow i_p \in ND_p(N_\bullet I)} X_{i_0} \times \delta\Delta^p & \longrightarrow & \pi^{-1}((BI)_{p-1}) \\ \downarrow & & \downarrow \\ \coprod_{i_0 \rightarrow \dots \rightarrow i_p \in ND_p(N_\bullet I)} X_{i_0} \times \Delta^p & \longrightarrow & \pi^{-1}((BI)_p) \end{array}$$

to the diagram

$$\begin{array}{ccc} \coprod_{i_0 \rightarrow \dots \rightarrow i_p \in ND_p(N_\bullet I)} \delta\Delta^p & \longrightarrow & \pi^{-1}((BI)_{p-1}) \\ \downarrow & & \downarrow \\ \coprod_{i_0 \rightarrow \dots \rightarrow i_p \in ND_p(N_\bullet I)} \Delta^p & \longrightarrow & \pi^{-1}((BI)_p), \end{array}$$

where $ND_p(N_\bullet I)$ denotes the non-degenerate p -simplices of $N_\bullet I$, in order to do an induction up the skeleta. We assume that we know that $p|_{(BI)_p}$ is a quasi-fibration.

Let U be the space formed from $(BI)_p$ by removing the barycenter of each p -cell. Let $V = (BI)_p - (BI)_{p-1}$. It suffices, by Lemma 4.1.26 to show that $\pi|_U$, $\pi|_V$ and $\pi|_{U \cap V}$ are all quasi-fibrations. This is clear for $\pi|_V$ and $\pi|_{U \cap V}$ because each map is homeomorphic to a trivial fibration.

By our inductive hypothesis, we can assume that $\pi|_{BI_{p-1}}$ is a quasi-fibration. To apply Lemma 4.1.27, we construct a fiber preserving deformation D of $\pi|_U$ to $\pi|_{BI_{p-1}}$ by considering the radial deformation of Δ^p without its barycenter to $\delta\Delta^p$. Thus, we just need to check that, if our deformation D takes $x \in U$ to $x' \in BI_{p-1}$, then the map $g^{-1}(x) \rightarrow g^{-1}(x')$ induced by D is a weak equivalence. It suffices to consider $x \in BI_p - BI_{p-1}$. Consequently, we can let x come from the interior point z of Δ^p , where the interior of Δ^p is $\Delta^p - \delta\Delta^p$. Suppose this Δ^p corresponds to the non-degenerate p -simplex $s = (i_0 \rightarrow \dots \rightarrow i_p)$. Our radial deformation takes z to some open face of Δ^p , say the open face with vertices $j_0 < j_1 < \dots < j_q$, where $\{j_0, \dots, j_q\} \subset \{0, \dots, p\}$. Then $g^{-1}(x) = X_{i_0}$ and $g^{-1}(x') = X_k$ where $k = i_{j_0}$ and the map $X_{i_0} = g^{-1}(x) \rightarrow g^{-1}(x') = X_k$ is the map induced by $i_0 \rightarrow i_k$ coming from the face of s (note that this is some composite of the face maps of $N_\bullet I$, which are all given by composing adjacent composable morphisms). Since we assumed that these maps are weak equivalences, the proof is complete. \square

Corollary 4.1.30. *Under the hypotheses of Theorem 4.1.24, the induced map*

$$B\pi_2: BTw(f) \rightarrow B(\mathcal{C}')^{\text{op}}$$

is a quasi-fibration for any functor $f: \mathcal{C} \rightarrow \mathcal{C}'$.

Proof. Consider the functor $X: (\mathcal{C}')^{\text{op}} \rightarrow \text{Top}$ sending y to $X(y) = By \setminus f$. Then since for $v: y \rightarrow y'$ the induced map $Bv \setminus f$ is a homotopy equivalence we can apply Proposition 4.1.29. In this case $X_{(\mathcal{C}')^{\text{op}}} = BTw(f)$ as we noted earlier so the proposition implies that

$$BTw(f) \rightarrow B(\mathcal{C}')^{\text{op}}$$

is a quasi-fibration. \square

Proof of Theorem B. We will prove Theorem 4.1.24 assuming Lemma 4.1.30. Consider the following diagram as in the proof of Theorem 4.1.14

$$\begin{array}{ccccc} y \setminus f & \longrightarrow & Tw(f) & \xrightarrow{\simeq} & \mathcal{C} \\ \downarrow & & \downarrow f' & & \downarrow f \\ y \setminus \mathcal{C} & \longrightarrow & Tw(\mathcal{C}') & \xrightarrow{\simeq} & \mathcal{C}' \\ \downarrow \simeq & & \downarrow \simeq & & \\ [0] & \longrightarrow & \mathcal{C}' & & \end{array}$$

where the maps indicated with \simeq are maps inducing homotopy equivalences on classifying spaces, as we have already discussed. The vertical map

$$BTw(f) \rightarrow BTw(\mathcal{C}') \rightarrow BC'$$

is $B\pi_2$, which is homotopy equivalent to the map $Bf: BC \rightarrow BC'$ in the top right square as we have already shown in the proof of Theorem 4.1.14. Since $BTw(f) \rightarrow B(\mathcal{C}')^{\text{op}}$ is a quasi-fibration by Lemma 4.1.30, we know that the two squares on the left are homotopy pullbacks. This implies the theorem. \square

Chapter 5

The S -construction and fundamental theorems

Appendix A

Fundamentals

A.1 Categories

Category theory will be of fundamental importance in studying algebraic K-theory. We recall the basic notions here.

Definition A.1.1. A category \mathcal{C} consists of

1. a class of objects denoted $\text{ob}(\mathcal{C})$
2. for each pair of objects c, c' a set $\mathcal{C}(c, c')$ of morphisms from c to c' , and
3. for any triple c, c', c'' a map of sets

$$- \circ -: \mathcal{C}(c', c'') \times \mathcal{C}(c', c'') \rightarrow \mathcal{C}(c, c'')$$

4. For each object c an element

$$\text{id}_c \in \mathcal{C}(c, c).$$

satisfying

$$(f \circ g) \circ h = f \circ (g \circ h)$$

for each triple of maps

$$(f, g, h) \in \mathcal{C}(c''', c'') \times \mathcal{C}(c'', c') \times \mathcal{C}(c', c)$$

and

$$\text{id}_{c'} \circ f = f = f \circ \text{id}_c$$

for each map $f: c \rightarrow c'$ in $\mathcal{C}(c, c')$, where each of these identities are functorial in the appropriate sense.

Example A.1.2. Given a category \mathcal{C} , let \mathcal{C}^{op} be the category whose objects are the same as the objects in \mathcal{C} and there is a unique morphism $f^{\text{op}}: b \rightarrow a$ for every morphism $f: a \rightarrow b$ in \mathcal{C} .

Remark A.1.3. Note that we use the convention that all categories are locally small by requiring that $\mathcal{C}(c, c')$ is a set rather than a proper class. We say a category is a small category if, in addition, it has a set of objects $\text{ob}(\mathcal{C})$.

Definition A.1.4. The skeleton $\text{sk}\mathcal{C}$ of a category \mathcal{C} is the full subcategory of \mathcal{C} consisting of one object for each isomorphism class of objects in \mathcal{C} . We say that \mathcal{C} is *skeletally small* if $\text{sk}\mathcal{C}$ is a small category.

Example A.1.5. The category of sets and set maps, denoted Set , is an example of a category that is not a small category. The category of finite sets and maps of finite sets is skeletally small and we write Fin for its skeleton.

Example A.1.6. Let \mathcal{C} be a small category. Then there is a category $\text{Arr}(\mathcal{C})$ whose objects are maps $f: a \rightarrow b$ in \mathcal{C} and a morphism from $a \rightarrow b$ to $c \rightarrow d$ is a commuting square

$$\begin{array}{ccc} a & \longrightarrow & b \\ \downarrow & & \downarrow \\ c & \longrightarrow & d \end{array}$$

in \mathcal{C} . Composition is defined by vertical composition of squares. We call this category the *arrow category*.

Definition A.1.7. A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ associates to each object $c \in \text{ob}\mathcal{C}$ an object $F(c) \in \text{ob}\mathcal{D}$, to each morphism $f: c \rightarrow c'$ a morphism $F(f): F(c) \rightarrow F(c')$ in \mathcal{D} such that $F(f \circ g) = F(f) \circ F(g)$.

Example A.1.8. Let Top denote the category of topological spaces and continuous maps. Then any topological space may be regarded as a set by forgetting the topology and any continuous map is in particular a map of sets, so this defines a functor

$$U: \text{Top} \rightarrow \text{Set}$$

called the forgetful functor.

Definition A.1.9. A subcategory \mathcal{C} in \mathcal{D} consists of a category \mathcal{C} and a functor $\iota: \mathcal{C} \rightarrow \mathcal{D}$ such that

$$\mathcal{C}(c, c') \rightarrow \mathcal{D}(\iota(c), \iota(c')) \quad (\text{A.1.10})$$

is injective. In particular, this implies that the set of objects in \mathcal{C} inject in the set of objects of \mathcal{D} .

We say \mathcal{C} is a *full* subcategory if the map (A.1.10) is also surjective. More generally, we say a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is faithful if the associated map

$$\mathcal{C}(c, c') \rightarrow \mathcal{D}(F(c), F(c')) \quad (\text{A.1.11})$$

is injective and we say it is fully faithful if this map is also surjective. In this case, we say that there is a fully faithful embedding of \mathcal{C} in \mathcal{D} since the essential image of \mathcal{C} in \mathcal{D} is necessarily a full subcategory of \mathcal{D} .

Definition A.1.12. Given a pair functors $F, G: \mathcal{C} \rightarrow \mathcal{D}$, then a natural transformation $\gamma: F \Rightarrow G$ associates to an object x in \mathcal{C} a map $\gamma_x: F(x) \rightarrow G(x)$ and to a map $f: x \rightarrow y$ a commutative diagram

$$\begin{array}{ccc} F(x) & \xrightarrow{F(f)} & F(y) \\ \gamma_x \downarrow & & \downarrow \gamma_y \\ G(x) & \xrightarrow{G(f)} & G(y) \end{array} \quad (\text{A.1.13})$$

and to a composable pair of morphisms $g \circ f$ in \mathcal{C} where $f: x \rightarrow y$ and $g: y \rightarrow z$, then there is a commutative diagram

$$\begin{array}{ccccc} F(x) & \xrightarrow{F(f)} & F(y) & \xrightarrow{F(g)} & F(z) \\ \gamma_x \downarrow & & \downarrow \gamma_y & & \downarrow \gamma_z \\ G(x) & \xrightarrow{G(f)} & G(y) & \xrightarrow{G(g)} & G(z). \end{array}$$

Definition A.1.14. The functor category $\text{Fun}(\mathcal{C}, \mathcal{D})$ is the category whose objects are functors $F, G: \mathcal{C} \rightarrow \mathcal{D}$ and whose morphisms are natural transformations $\gamma: F \Rightarrow G$, where composition is given by composition of natural transformations where $\alpha \circ \beta$ is defined on objects by

$$(\alpha \circ \beta)_x := \alpha_x \circ \beta_x$$

and on morphisms $f: x \rightarrow y$ by vertical composition of squares of the form (A.1.13) and similarly for compositions of morphisms.

Definition A.1.15. A factorization system on a small category \mathcal{C} consists of a pair of subcategories (E, M) such that

1. E and M each contain all of the isomorphisms in \mathcal{C} and consequently all objects in \mathcal{C} , and
2. every morphism $f: A \rightarrow C$ in \mathcal{C} can be factored as $f = m \circ e$ where $e: A \rightarrow B$ is a map in E and $m: B \rightarrow C$ is a map in M .
3. this factorization defines a functor

$$\text{Arr}(\mathcal{C}) \rightarrow \text{Arr}(E) \times_{\mathcal{C}} \text{Arr}(M)$$

sending f to (m, e) and a commutative square

$$\begin{array}{ccc} A & \xrightarrow{f} & C \\ g \downarrow & & \downarrow g'' \\ A' & \xrightarrow{f'} & C' \end{array}$$

to the composition of squares

$$\begin{array}{ccccc} A & \xrightarrow{e} & B & \xrightarrow{m} & C \\ g \downarrow & & \downarrow g' & & \downarrow g'' \\ A' & \xrightarrow{e'} & B' & \xrightarrow{m'} & C'. \end{array}$$

Here the category $\text{Arr}(E) \times_{\mathcal{C}} \text{Arr}(M)$ is the pullback in the category of small categories of the diagram

$$\text{Arr}(M) \xrightarrow{t} \mathcal{C} \xleftarrow{s} \text{Arr}(E)$$

where t sends an object $e: A \rightarrow B$ in $\text{Arr}(E)$ to B and s sends an object $m: B \rightarrow C$ in $\text{Arr}(M)$ to B . On morphisms these functors are defined in the evident way. In other words, the category $\text{Arr}(E) \times_{\mathcal{C}} \text{Arr}(M)$ has objects pairs of morphisms (e, m) such that, as maps in \mathcal{C} , they are composable.

A.2 Sets

Definition A.2.1. A partial order on a set P is a binary relation \leq satisfying

1. $x \leq x$ (Reflexivity),
2. if $x \leq y$ and $y \leq x$, then $x = y$ (Anti-symmetry),
3. if $x \leq y$ and $y \leq z$, then $x \leq z$ (Transitivity).

We say x is related to y if either $x \leq y$ or $y \leq x$. A set with a partial order will be called a partially ordered set or a POSet for short.

Any POSet P can be considered as a small category with objects the elements of P and morphism sets

$$P(x, y) = \begin{cases} * & x \leq y \\ \emptyset & \text{otherwise} \end{cases}$$

and composition

$$P(y, z) \times P(x, y) \rightarrow P(x, z)$$

is defined in the obvious way using transitivity.

Definition A.2.2. A total order on a set X is a partial order satisfying the totality axiom: for each $x, y \in X$ either $x \leq y$ or $y \leq x$.

In other words, a totally ordered set is simply a partially ordered set in which any two elements are related.

Example A.2.3. The natural numbers

$$\mathbb{N}_0 = \{0, 1, 2, \dots\}$$

are a totally ordered set with the usual total order. The set $n = \{0, 1, \dots, n\}$ is a finite totally ordered set equipped with the restriction of the total order on \mathbb{N}_0 . When we view the totally ordered set n as a category it can be depicted as

$$0 \leftarrow 1 \leftarrow 2 \leftarrow \dots \leftarrow n.$$

Appendix B

Simplicial methods

B.1 The simplex category

Definition B.1.1. Let Ord denote the category of finite totally ordered sets and maps of finite sets that preserve the total order. Let Δ be the the skeleton of this category with exactly one object for each isomorphism class of objects in Ord . The objects in Δ are

$$[n] = \{0, 1, \dots, n\}$$

for $n \geq 0$ and the maps are nondecreasing maps of finite sets.

Example B.1.2. There is a canonical factorization system (E, M) on the category Δ . The category E consists of all isomorphisms as well as the closure under composition of the maps of the form

$$\sigma_k: [n] \rightarrow [n-1]$$

for $0 \leq k \leq n$ such that

$$\sigma_k(i) = \begin{cases} i & \text{if } i < k \\ k & \text{if } i = k, k+1 \\ i-1 & \text{if } k+1 \leq i \leq n. \end{cases}$$

The category M consists of all isomorphisms as well as the closure under composition of the morphisms

$$\delta_k(i) = \begin{cases} i & \text{if } i < k \\ i+1 & \text{if } k \leq i \leq n. \end{cases}$$

We can therefore depict the category Δ as follows

$$[0] \begin{matrix} \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{matrix} [1] \begin{matrix} \xrightarrow{\quad} \\ \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{matrix} [2] \begin{matrix} \xrightarrow{\quad} \\ \xleftarrow{\quad} \\ \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{matrix} \dots \quad (\text{B.1.3})$$

Definition B.1.4. Given a category \mathcal{C} , a *simplicial object in \mathcal{C}* is a functor

$$\Delta^{\text{op}} \rightarrow \mathcal{C}.$$

This can be described as a collection of objects $\{X[i]\}_{i \geq 0}$ sitting in a diagram

$$X[0] \begin{array}{c} \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{array} X[1] \begin{array}{c} \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{array} X[2] \begin{array}{c} \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{array} \dots \quad (\text{B.1.5})$$

where the maps in the diagram are called the face maps

$$\{\partial_i: X[n+1] \rightarrow X[n]\}_{0 \leq i \leq n+1}$$

and degeneracy maps

$$\{s_i: X[n] \rightarrow X[n+1]\}_{0 \leq i \leq n}$$

and these must satisfy the simplicial identities:

1. $\partial_i \circ \partial_j = \partial_{j-1} \circ \partial_i$ if $i < j$,
2. $s_i \circ s_j = s_j \circ s_{i-1}$ if $i > j$,
3. $\partial_i \circ s_j = \begin{cases} s_{j-1} \circ \partial_i & \text{if } i < j \\ \text{id}_{X_j} & \text{if } i = j \text{ or } i = j + 1 \\ s_j \circ \partial_{i-1} & \text{if } i > j + 1 \end{cases}$

Morphisms of simplicial objects in \mathcal{C} are simply natural transformations of functors $\Delta^{\text{op}} \rightarrow \mathcal{C}$.

Example B.1.6. When $\mathcal{C} = \text{Set}$, we call a simplicial object in Set simply a simplicial set and we write sSet for this category. We could also consider pointed sets Set_* and we will write sSet_* for the category of simplicial objects in pointed sets (or equivalently, pointed simplicial sets). When $\mathcal{C} = \text{Top}$ we refer to the category of simplicial objects in Top simply as simplicial spaces and we denote the category of simplicial spaces by sTop .

Example B.1.7. Given an object $[n] \in \Delta$, we can form

$$\text{Hom}_{\Delta}(-, [n]): \Delta^{\text{op}} \rightarrow \text{Set}.$$

This is clearly a simplicial set. We denote this simplicial set

$$\Delta^n := \text{Hom}_{\Delta}(-, [n]).$$

Exercise B.1.8. Use the Yoneda lemma to show that

$$\text{Hom}_{\text{sSet}}(\Delta^n, X) = X[n].$$

Hint: Write $\iota_n := \text{id} \in \text{Hom}_{\Delta}([n], [n])$. Given a simplicial map $\varphi: \Delta^n \rightarrow X$, associate to this map an n -simplex $\varphi(\iota_n) \in X[n]$. Prove that this gives a bijection.

Definition B.1.9. We define a topological space

$$|\Delta^n| = \{(t_0, \dots, t_n) \in [0, 1]^{n+1} : \sum t_i = 1\}$$

equipped with the subspace topology. In fact, the spaces $|\Delta^n|$ form cosimplicial space

$$|\Delta^\bullet|(-) : \Delta \rightarrow \mathbf{Top}$$

such that $|\Delta^\bullet|([n]) = |\Delta^n|$. The map

$$\theta_* : |\Delta^n| \rightarrow |\Delta^m|$$

induced by $\theta : [n] \rightarrow [m]$ is defined by

$$\theta(t_0, t_1, \dots, t_n) = (t'_0, \dots, t'_m)$$

where

$$t'_i = \begin{cases} 0 & \text{if } \theta^{-1}(i) \neq \emptyset, \\ \sum_{j \in \theta^{-1}(i)} t_j & \text{if } \theta^{-1}(i) = \emptyset. \end{cases}$$

Remark B.1.10. There is also an analogous cosimplicial simplicial set

$$\Delta^n : \Delta \rightarrow \mathbf{sSet}$$

given on n -simplices by

$$\Delta^n = \mathrm{Hom}_\Delta(-, [n])$$

where the cosimplicial structure is given by regarding this as a functor in the second variable.

Definition B.1.11. Given a pair of simplicial sets X, Y , we define the product of simplicial sets $X \times Y$ by letting the n -simplices be simply

$$(X \times Y)[n] = X[n] \times Y[n]$$

and the face and degeneracy maps simply as

$$\partial_i^{X \times Y} = (\partial_i^X, \partial_i^Y) \text{ and } s_i^{X \times Y} = (s_i^X, s_i^Y).$$

Note that the non-degenerate n -simplices of $X \times Y$ are not simply the product of the non-degenerate n -simplices of X with the non-degenerate n -simplices of Y .

Definition B.1.12. We also define an internal hom $\underline{\mathrm{Hom}}(X, Y)$ to be the simplicial set with n -simplices

$$\underline{\mathrm{Hom}}(X, Y)[n] := \mathrm{Hom}_{\mathbf{sSet}}(X \times \Delta^n, Y)$$

and face and degeneracy maps induced by considering Δ^n as a cosimplicial object in \mathbf{sSet} and applying the functor $\mathrm{Hom}_{\mathbf{sSet}}(X \times -, Y)$.

Proposition B.1.13. *The category $(\mathbf{sSet}, \times, \Delta^0)$ is a symmetric monoidal category and there is a natural isomorphism*

$$\mathrm{Hom}_{\mathbf{sSet}}(X \times Y, Z) \cong \mathrm{Hom}_{\mathbf{sSet}}(X, \underline{\mathrm{Hom}}(Y, Z)).$$

Definition B.1.14. We say $f: X \rightarrow Y$ is homotopy equivalent to $g: X \rightarrow Y$ if there is a map of simplicial sets

$$H: X \times \Delta^1 \rightarrow Y$$

such that $H|_{(\iota_0)_*(\Delta^0)} = f$ and $H|_{(\iota_1)_*(\Delta^0)} = g$ where $(\iota_j)_*: \Delta^0 \rightarrow \Delta^1$ is the map of simplicial sets induced by the unique map $\iota_j: [0] \rightarrow [1]$ map in Δ sending 0 to j ; i.e.

$$(\iota_j)_* := \mathrm{Hom}(-, \iota_j): \mathrm{Hom}(-, [0]) \rightarrow \mathrm{Hom}(-, [1]).$$

Let $\partial\Delta^n$ denote the smallest sub-simplicial set of Δ^n generated by the faces $d_j(\iota_n)$ for $0 \leq j \leq n$. where ι_n denotes the element

$$\iota_n = \mathrm{id} \in \mathrm{Hom}_{\Delta}([n], [n]) = \mathrm{Hom}_{\mathbf{sSet}}(\Delta^n, \Delta^n) = (\Delta^n)_n$$

Let Λ_k^n denote the smallest sub-simplicial set of Δ^n generated by the faces $d_i(\iota_n)$ for the face $d_k(\iota_n)$. We say a simplicial set is a Kan complex if there exists a unique lift in any diagram of the form

$$\begin{array}{ccc} \Lambda_k^n & \longrightarrow & X \\ \downarrow & \nearrow \exists! h & \\ \Delta^n & & \end{array}$$

for $0 \leq k \leq n$. Where Λ_k^n is the sub-simplicial set of Δ^n generated by $d_j(\iota_n)$

We can consider the full subcategory of the category of simplicial sets whose objects are the category of Kan complexes.

Let X, Y be Kan complexes and define

$$[X, Y] = \mathrm{Hom}(X, Y) / \simeq$$

where \simeq is the equivalence relation given by simplicial homotopy equivalence.

Definition B.1.15. Define the homotopy category of simplicial sets, denoted $\mathrm{ho}(\mathbf{sSet})$ to be the category whose objects are Kan complexes and morphisms from $X \rightarrow Y$ are $[X, Y]$. We define the homotopy category of Top, denoted $\mathrm{ho}(\mathbf{Top})$ to be the category whose objects are CW complexes and maps are homotopy classes of maps from X to Y , which we also denote $[X, Y]$.

Construction B.1.16. Let X be an object in Top. We define

$$\mathrm{sing}(X) = \mathrm{Hom}_{\mathbf{Top}}(|\Delta^\bullet|, X).$$

This is clearly a simplicial set. One can check that it is also a Kan complex when X is a CW complex.

We will now define geometric realization, more generally, of a simplicial space.

Construction B.1.17. Let X_\bullet be an object in \mathbf{sTop} . Then we define

$$|X_\bullet| := \text{coeq} \left(\coprod_{f: i \rightarrow j \in \Delta^{\text{op}}} X_i \times |\Delta^j| \rightrightarrows \coprod_{[n] \in \Delta^{\text{op}}} X_n \times |\Delta^n| \right)$$

$\Pi \text{id}_{X_i} \times |\Delta^f| \quad \quad \quad \Pi X_f \times \text{id}_{|\Delta^j|}$

where the coproducts are equipped with the coproduct topology and coequalizer is equipped with the quotient topology. Sometimes, we simply write this as

$$|X| = \left(\coprod_{n \geq 0} |\Delta^n| \times X_n \right) / \sim$$

where \sim is the equivalence relation generated by

$$(x, \partial_i y) \sim (\delta_i x, y) \text{ and } (x, s_i y) \simeq (\sigma_i x, y).$$

One can check that these two spaces are the same.

Example B.1.18. When X is a simplicial set, we define $|X|$ in the same way by equipping $X[n]$ with the discrete topology for all $n \geq 0$.

Exercise B.1.19. The geometric realization of a simplicial set is always a CW complex.

We can therefore consider the geometric realization as taking values in compactly generated weak Hausdorff spaces. For the next result of Milnor, it is important to note that all constructions take place in this category.

Theorem B.1.20 (Milnor). *Geometric realization commutes with finite limits. In particular, there is a homeomorphism*

$$|X_\bullet \times Y_\bullet| \cong |X_\bullet| \times |Y_\bullet|.$$

The following is one of the fundamental theorems of simplicial sets.

Theorem B.1.21. *There is a natural isomorphism $\gamma_{-, -}$ of functors $\mathbf{sSet}^{\text{op}} \times \mathbf{Top} \rightarrow \mathbf{Set}$ defined on objects by*

$$\gamma_{X, Y}: \text{Hom}_{\mathbf{Top}}(|X|, Y) \cong \text{Hom}_{\mathbf{sSet}}(X, \text{sing}(Y)).$$

Construction B.1.22. There is an alternate way to construct the geometric realization of a simplicial set X that is less intuitive, but makes the proof of Theorem B.1.21 quite easy. Let $\Delta \downarrow X$ denote the category whose objects are

maps of simplicial sets $x: \Delta^n \rightarrow X$ where $n \geq 0$ and whose maps $x \rightarrow y$ are commuting triangles

$$\begin{array}{ccc} \Delta^n & \xrightarrow{x} & X \\ \theta_* \downarrow & \nearrow y & \\ \Delta^m & & \end{array}$$

where $\theta: [n] \rightarrow [m]$ is a map in Δ . Composition is defined in the evident way. Then we can define $|X|$ to be the colimit

$$|X| = \operatorname{colim}_{\Delta \downarrow X} |\Delta^n|$$

in the category of topological spaces.

Exercise B.1.23. Show that the definition of $|X|$ in Construction B.1.22 is the same as in Construction B.1.17 for simplicial sets up to homeomorphism.

Proof of Theorem B.1.21. Recall that $\operatorname{Hom}_{\operatorname{Top}}(-, Y)$ sends limits in $\operatorname{Top}^{\operatorname{op}}$, or in other words colimits in Top , to limits in Set . Therefore,

$$\operatorname{Hom}_{\operatorname{sSet}}(|X|, Y) = \operatorname{Hom}_{\operatorname{sSet}}(\operatorname{colim}_{\Delta \downarrow X} |\Delta^n|, Y) \quad (\text{B.1.24})$$

$$\cong \lim_{\Delta \downarrow X} \operatorname{Hom}_{\operatorname{sSet}}(|\Delta^n|, Y) \quad (\text{B.1.25})$$

$$= \lim_{\Delta \downarrow X} \operatorname{sing}(Y)_n \quad (\text{B.1.26})$$

$$= \lim_{\Delta \downarrow X} \operatorname{Hom}_{\operatorname{sSet}}(\Delta^n, \operatorname{sing}(Y)) \quad (\text{B.1.27})$$

$$\cong \operatorname{Hom}_{\operatorname{sSet}}(\operatorname{colim}_{\Delta \downarrow X} \Delta^n, \operatorname{sing}(Y)) \quad (\text{B.1.28})$$

$$\cong \operatorname{Hom}_{\operatorname{sSet}}(X, \operatorname{sing}(Y)) \quad (\text{B.1.29})$$

where the equality (B.1.24) is the definition of geometric realization, the isomorphism (B.1.25) follows because $\operatorname{Hom}_{\operatorname{Top}}(-, Y)$ send limits in $\operatorname{Top}^{\operatorname{op}}$ to limits in Set , as remarked above, the equality (B.1.26) holds by definition of $\operatorname{sing}(Y)$, the equality (B.1.27) holds by Exercise (B.1.8), the isomorphism (B.1.28) hold because $\operatorname{Hom}_{\operatorname{Top}}(-, Y)$ sends limits in $\operatorname{Top}^{\operatorname{op}}$ to limits in Set . The last isomorphism holds because X is a cocone for the functor from $\Delta \downarrow X \rightarrow \operatorname{sSet}$ sending $\Delta^n \rightarrow X$ to Δ^n . \square

Recall that Cat denotes the category of small categories. Let $[n]$ be the small category

$$0 \rightarrow 1 \rightarrow 2 \rightarrow \cdots \rightarrow n$$

described in Example A.2.3. These form a cosimplicial object in Cat by letting the codegeneracy map σ_i be given by composing the morphisms with source i and $i + 1$. and the coface map δ_i given by inserting an identity map in the i -th position.

Definition B.1.30. Given a small category \mathcal{C} , we define a simplicial set by letting the n -simplices be

$$NC[n] := \text{Cat}([n], \mathcal{C}),$$

which should be viewed as a sequence of n composable morphisms, if $n > 0$ and, and simply the objects in \mathcal{C} if $n = 0$. The face maps are given by composing two adjacent morphisms and the degeneracy maps are given by inclusion of the identity morphism. In other words, we simply use the fact that

$$[\bullet]: \Delta \rightarrow \text{Cat}$$

is a cosimplicial object in small categories and therefore, by functoriality,

$$\text{Cat}([n], \mathcal{C})$$

forms as simplicial set.

For example, any discrete group G can be regarded as a small category with one object $*$ and morphism set $G(*, *) = G$. By unpacking the definition, we see that

$$NG[n] = G^n$$

where $G^0 = *$. We can be even more explicit in this case. The face maps

$$\partial_i: G^{n+1} \rightarrow G^n$$

are defined by

$$\partial_i(g_1, \dots, g_{n+1}) = \begin{cases} (g_2, g_3, \dots, g_{n+1}) & \text{if } i = 0 \\ (g_1, \dots, g_i g_{i+1}, \dots, g_{n+1}) & \text{if } 0 < i < n + 1 \\ (g_1, g_3, \dots, g_n) & \text{if } i = n \end{cases}$$

and the degeneracy maps are defined by

$$s_i(g_1, \dots, g_n) = (g_0, g_1, \dots, g_{i-1}, 1, g_i, \dots, g_{n-1})$$

Note, that there was notion special about doing this construction for a discrete group. More generally, given a topological group G we define an object in sTop in the same way and we denote it

$$B(*, G, *): \Delta^{\text{op}} \rightarrow \text{Top}.$$

Definition B.1.31. Given a category \mathcal{C} , we define

$$BC := |NC|.$$

Given a topological group G , we also define

$$BG = |B(*, G, *)|.$$

Remark B.1.32. Of course, when G is a discrete group then

$$NG = B(*, G, *)$$

so there isn't a conflict in notation.

Bibliography

- [1] Clark Barwick, *On the algebraic K-theory of higher categories*, J. Topol. **9** (2016), no. 1, 245–347. MR3465850
- [2] Hyman Bass, *Algebraic K-theory*, W. A. Benjamin, Inc., New York-Amsterdam, 1968. MR0249491
- [3] Andrew J. Blumberg, David Gepner, and Gonalo Tabuada, *A universal characterization of higher algebraic K-theory*, Geom. Topol. **17** (2013), no. 2, 733–838. MR3070515
- [4] Armand Borel and Jean-Pierre Serre, *Le th or me de Riemann-Roch*, Bull. Soc. Math. France **86** (1958), 97–136. MR116022
- [5] Edgar H. Brown Jr., *Cohomology theories*, Ann. of Math. (2) **75** (1962), 467–484. MR138104
- [6] P. Gabriel and M. Zisman, *Calculus of fractions and homotopy theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 35, Springer-Verlag New York, Inc., New York, 1967. MR0210125
- [7] David Gepner, Moritz Groth, and Thomas Nikolaus, *Universality of multiplicative infinite loop space machines*, Algebr. Geom. Topol. **15** (2015), no. 6, 3107–3153. MR3450758
- [8] Daniel Grayson, *Higher algebraic K-theory. II (after Daniel Quillen)*, Algebraic K-theory (Proc. Conf., Northwestern Univ., Evanston, Ill., 1976), 1976, pp. 217–240. Lecture Notes in Math., Vol. 551. MR0574096
- [9] Michel A. Kervaire, *Smooth homology spheres and their fundamental groups*, Trans. Amer. Math. Soc. **144** (1969), 67–72. MR253347
- [10] Hideya Matsumoto, *Sur les sous-groupes arithm tiques des groupes semi-simples d ploy s*, Ann. Sci.  cole Norm. Sup. (4) **2** (1969), 1–62. MR240214
- [11] Barry Mazur, *Relative neighborhoods and the theorems of Smale*, Ann. of Math. (2) **77** (1963), 232–249. MR150786
- [12] John Milnor, *Algebraic K-theory and quadratic forms*, Invent. Math. **9** (1969/70), 318–344. MR260844
- [13] ———, *Introduction to algebraic K-theory*, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1971. Annals of Mathematics Studies, No. 72. MR0349811
- [14] Daniel Quillen, *On the cohomology and K-theory of the general linear groups over a finite field*, Ann. of Math. (2) **96** (1972), 552–586. MR315016
- [15] ———, *Higher algebraic K-theory. I*, Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972), 1973, pp. 85–147. Lecture Notes in Math., Vol. 341. MR0338129
- [16] Jonathan Rosenberg, *Algebraic K-theory and its applications*, Graduate Texts in Mathematics, vol. 147, Springer-Verlag, New York, 1994. MR1282290
- [17] Graeme Segal, *Categories and cohomology theories*, Topology **13** (1974), 293–312. MR353298

- [18] Stephen Smale, *Generalized Poincaré's conjecture in dimensions greater than four*, Ann. of Math. (2) **74** (1961), 391–406. MR137124
- [19] V. Srinivas, *Algebraic K-theory*, second, Modern Birkhäuser Classics, Birkhäuser Boston, Inc., Boston, MA, 2008. MR2371852
- [20] L. N. Vaseršteĭn, *Foundations of algebraic K-theory*, Uspehi Mat. Nauk **31** (1976), no. 4 (190), 87–149. MR0427426
- [21] Friedhelm Waldhausen, *Algebraic K-theory of topological spaces. I*, Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part 1, 1978, pp. 35–60. MR520492
- [22] ———, *Algebraic K-theory of topological spaces. II*, Algebraic topology, Aarhus 1978 (Proc. Sympos., Univ. Aarhus, Aarhus, 1978), 1979, pp. 356–394. MR561230
- [23] ———, *Algebraic K-theory of spaces, a manifold approach*, Current trends in algebraic topology, Part 1 (London, Ont., 1981), 1982, pp. 141–184. MR686115
- [24] Charles A. Weibel, *The K-book*, Graduate Studies in Mathematics, vol. 145, American Mathematical Society, Providence, RI, 2013. An introduction to algebraic K-theory. MR3076731