


C. J. EVONIUK, G. DOS PASSOS GOMES, S. P. HILL, S. FUJITA, K. HANSON,
I. V. ALABUGIN* (FLORIDA STATE UNIVERSITY, TALLAHASSEE, USA AND KYUSHU
UNIVERSITY, FUKUOKA, JAPAN)

Coupling N–H Deprotonation, C–H Activation, and Oxidation: Metal-Free C(sp³)–H Aminations with Unprotected Anilines

J. Am. Chem. Soc. **2017**, *139*, 16210–16221.

Expanded N-Heterocycles through C(sp³)–H Amination



Significance: Expanded N-heterocycles are common frameworks in organic materials with conductive and/or emissive properties. The authors disclose an efficient metal-free C(sp³)–H amination with unprotected anilines.

Comment: This method is applicable to a range of molecular building blocks; both electron-rich and electron-poor substrates are well tolerated. The combination of *t*-BuOK, molecular oxygen and *N,N*-dimethylformamide enables an efficient sequence of deprotonation followed by proton and electron transfer. Alabugin and co-workers describe extensive investigations into both the mechanism, and the properties of the largest N-heterocycles.

SYNFACTS Contributors: Timothy M. Swager, Constantin-Christian A. Voll

Synfacts 2018, 14(02), 0144 Published online: 18.01.2018

DOI: 10.1055/s-0037-1609179; Reg-No.: S13618SF

2018 © THIEME STUTTGART • NEW YORK