

Association of Ethnicity and Survival in Peritoneal Dialysis: A Cohort Study of Incident Patients in Brazil

Natália Maria da Silva Fernandes, PhD,^{1,2,3} Tiny Hoekstra, PhD,³
Tessa O. van den Beukel, MD,³ Luciana Tirapani, MS,² Kleyton Bastos, PhD,⁴
Roberto Pecoits-Filho, PhD,⁵ Abdul Rashid Qureshi, PhD,⁶ Friedo W. Dekker, PhD,³
Marcus G. Bastos, PhD,^{1,2} and José Carolino Divino-Filho, PhD⁶

Background: There are no available epidemiologic studies about the impact of ethnicity on outcomes of patients treated with peritoneal dialysis (PD) in South America. This study aims to assess the effect of ethnicity on the mortality of incident PD patients in Brazil.

Study Design: Prospective observational cohort study of incident patients treated with PD.

Settings & Participants: Patients 18 years or older who started PD therapy between December 2004 and October 2007 in 114 Brazilian dialysis centers.

Predictors: Self-reported ethnicity defined by the Brazilian Institute of Geography and Statistics as black and brown versus white patients and baseline demographic, socioeconomic, clinical, and laboratory data were collected at baseline.

Outcome: Mortality, using cumulative mortality curves in which kidney transplantation and transfer to hemodialysis therapy were treated as competing end points. Multivariate Cox proportional hazards analysis was used to adjust for gradually more potential explanatory variables, censored for kidney transplantation and transfer to hemodialysis therapy. Analyses were performed for all patients, as well as stratified for elderly (aged ≥ 65 years) and nonelderly patients.

Results: 1,370 patients were white, 516 were brown, and 273 were black. The competing-risk model showed higher mortality in white patients compared with black and brown patients. With white patients as the reference, Cox proportional hazards analysis showed a crude HR for mortality of 0.77 (95% CI, 0.56-1.05) for black and 0.74 (95% CI, 0.59-0.94) for brown patients. After adjusting for potential explanatory factors, HRs were 0.67 (95% CI, 0.48-0.95) and 0.77 (95% CI, 0.43-1.01), respectively. The same results were observed in elderly and nonelderly patients.

Limitations: Ethnicity was self-determined and some misclassification might have occurred.

Conclusions: Black and brown Brazilian incident PD patients have a lower mortality risk compared with white patients.

Am J Kidney Dis. 62(1):89-96. © 2013 by the National Kidney Foundation, Inc.

INDEX WORDS: Ethnicity; race; peritoneal dialysis; survival.

Editorial, p. 10

End-stage renal disease (ESRD) is a serious public health issue in developed and developing countries. The scale and patterns of kidney disease vary from country to country; to some degree, these differences could be the result of regional racial and ethnic composition.^{1,2} However, in some continents, data are still scarce.

Racial and ethnic differences in the outcomes of patients with ESRD have been widely described in certain parts of the world. Data from the United States have shown increased mortality for Caucasian compared to non-Caucasian dialysis patients on hemodialysis (HD) and peritoneal dialysis (PD) therapy,³⁻⁵ although this is mainly in patients older than 50 years.⁶ Evaluating the influence of race on survival in PD patients in the United States, Korbet et al⁷ showed that white patients experienced worse survival compared with black patients. In Canada, Hispanic, black, and Asian patients on HD and PD therapy have significantly better survival than non-

Hispanic Caucasian patients.⁸ In Europe, non-Caucasian HD patients also were found to have better survival compared with Caucasian dialysis patients,⁹ and a similar result was observed in a nationwide study from the United Kingdom.¹⁰ Furthermore, immigrant dialysis patients in the Netherlands were found to have better

From the ¹Department of Medicine, Federal University of Juiz de Fora; ²IMEPEN Foundation, Juiz de Fora, Minas Gerais, Brazil; ³Department of Clinical Epidemiology, Leiden University Medical Center, Leiden, the Netherlands; ⁴Federal University of Sergipe, Aracaju, Sergipe; ⁵Center for Health and Biological Sciences, Pontifícia Universidade Católica do Paraná, Curitiba, Brazil; and ⁶Division of Renal Medicine, CLINTEC, Karolinska Institutet, Stockholm, Sweden.

Received July 19, 2012. Accepted in revised form February 6, 2013. Originally published online April 15, 2013.

Address correspondence to Natália Maria da Silva Fernandes, PhD, Rua Jamil Altaff, 132, Vale do Ipê, Juiz de Fora, Minas Gerais, Brazil CEP:36035 380. E-mail: nataliafernandes02@gmail.com

© 2013 by the National Kidney Foundation, Inc.

0272-6386/\$36.00

<http://dx.doi.org/10.1053/j.ajkd.2013.02.364>

survival compared with native Dutch dialysis patients.¹¹ In contrast, the indigenous dialysis population in Australia and New Zealand presented an increased mortality risk compared with white dialysis patients.¹² Data for the survival of patients with ESRD in South and Latin America have not been described systematically for different ethnic groups.

The population in Brazil, one of the largest countries in South America, is one of the most heterogeneous populations in the world¹³ as a result of 5 centuries of admixture between its ancestral roots (autochthonous Amerindians, white Europeans, and sub-Saharan Africans). In the past 2 centuries, Asians also have contributed to this miscegenation.¹⁴⁻¹⁸ The ancestrally admixed population (referred to as “brown” in Brazil), which corresponds to 42.6%¹⁹ of the Brazilian general population,¹⁹ certainly places Brazil among the most ethnically admixed countries. The heterogeneity and admixture of the Brazilian population also may have relevant implications for the design and interpretation of studies performed with Brazilians.^{17,18}

The prevalence of ESRD in Brazil was 483 patients per million inhabitants²⁰ in 2010, and black and brown patients seem to face a higher risk of ESRD than white patients.²¹ Brazil has a public and private hybrid health care system,²² which provides universal access to renal replacement therapy, and a group of patients start PD as first treatment.

In order to understand the impact of ethnicity in the mortality of PD patients in Brazil, it is of utmost importance to include the predominant ethnicities, including those categorized as brown. The results presented here, based in a large PD database, are to our knowledge the first time that such an assessment of the ethnicity of incident PD patients is reported from South America. The aim of this study is to assess the impact of ethnicity on the survival of incident PD patients in Brazil.

METHODS

Patients

We analyzed data from the Brazilian Peritoneal Dialysis Multi-center Study (BRAZPD), a prospective observational cohort study of PD patients enrolled throughout Brazil in 114 dialysis centers, treating more than 10 patients each.²³ It is important to note that 60% of the total Brazilian PD population in 2007 was included in the BRAZPD cohort.²⁰ For this specific analysis, we included patients who were 18 years or older and started PD therapy between December 2004 and October 2007 either as first renal replacement therapy or after HD. Patients were excluded when data for ethnicity and/or age were lacking. Baseline was defined as 3 months after the initiation of PD therapy. The study was approved by the medical ethical committees of all participating centers. All patients signed the informed consent form before inclusion.

Data Collection Procedures

Details regarding data collection procedures have been published elsewhere.²³ For this particular study, we used the following

demographic and clinical data at baseline: age, sex, ethnicity, geographic region of the dialysis center, family income (self-report; 0-2, 3-5, and >5 times the minimum monthly national salary), educational level (self-report; illiteracy, elementary school, secondary school, and college),²⁴ distance from dialysis center (<25, 25-50, and >50 km), cause of ESRD, whether a patient underwent previous HD treatment, PD modality (continuous ambulatory PD or automated PD), PD indication (medical, patient's choice, or only option), duration of predialysis care, comorbid conditions, body mass index, blood pressure, and laboratory parameters (urea, creatinine, calcium, phosphate, potassium, alanine aminotransferase, glucose, and hemoglobin). Ethnicity was self-categorized and based on categories defined by the Brazilian Institute of Geography and Statistics (white, black, brown, yellow, and indigenous). Brown people result from the mixture between white and black, white and indigenous, or black and indigenous. The term brown is culturally accepted in Brazil.¹⁹

Assessment of comorbid conditions was based on the *International Classification of Diseases, Tenth Revision*.²⁵ Laboratorial data were collected at baseline and, if not available, at 2 or 4 months after the start of PD therapy. Quality of life (QoL) data were collected 6 months after the start of PD therapy. QoL was assessed using the Mental and Physical Component Summary scores of the 36-Item Short Form Health Survey questionnaire.²⁶

Patients were followed up until October 2007. Dropout, mortality rate, and causes of death were classified according to the Brazilian government.²⁷

Statistical Analysis

Patient characteristics are described as mean \pm standard deviation or percentage. Differences in baseline characteristics among white, black, and brown patients were analyzed with analysis of variance, Kruskal-Wallis (for alanine aminotransferase level), and χ^2 tests. Patients were followed up until death, transfer to HD therapy, kidney transplantation, or the end of the study period in October 2007, whichever came first. In order to determine the influence of ethnicity on mortality, we used cumulative mortality curves in which kidney transplantation and transfer to HD therapy were treated as competing end points.²⁸ Multivariate Cox proportional hazards analyses were used to adjust gradually for more potential explanatory variables; first for demographic variables, followed by additional adjustment for socioeconomic, clinical, QoL, hemoglobin level, and creatinine level variables that presented clinical and/or statistical significance in univariate analysis. Analyses also were performed stratifying for elderly (aged ≥ 65 years) and nonelderly patients (aged <65 years). All analyses were carried out with SPSS, version 17.0, for Windows statistical software (SPSS Inc), and $P < 0.05$ was considered as statistically significant.

The percentage of missing data was <10%, except for QoL (27%), body mass index (15%), blood pressure (15%), glucose level (18%), and creatinine level (10%). In order to create complete data sets for multivariate Cox analysis, missing values for variables included in the Cox model (age, sex, level of education, family income, geographic region of the dialysis center, cause of ESRD, comorbid conditions, QoL, first therapy, creatinine level, and hemoglobin level) were imputed with multiple imputation techniques (5 repetitions). With multiple imputation techniques, missing data are imputed by a value that is predicted using the patient's available characteristics, under the condition of missing “at random.”²⁹ Sensitivity analysis was performed with and without imputation for missing data.




Figure 1. Flow chart of the study.

RESULTS

Study Participants

In BRAZPD, 3,439 incident PD patients were included by October 2007. From this cohort, 867 patients did not complete 90 days of PD therapy by the end of the study. Further, 336 patients were excluded because they did not have sociodemographic data for age and/or race and 18 patients were younger than 18 years. Due to limited numbers ($n = 59$), yellow and indigenous patients (self-identified; brown, yellow, black, and white terminology is culturally accepted in Brazil) were excluded from the analysis. A total of 2,159 incident patients were included in the study: 1,370 patients were white, 516 were brown, and 273 were black (Fig 1).

Of 867 patients who did not complete 90 days by the end of the study, 73 died, 13 underwent kidney transplantation, 31 were transferred to HD therapy, 4 had recovery of kidney function, 4 were lost to follow-up, and 742 simply had not received 90 days of PD therapy by the end of the study period. No differences were found in sociodemographic characteristics between this group of 867 patients and the 2,159 patients included in the study.

Sixty-seven percent of the incident PD patients had had HD as their first dialysis therapy. No statistical differences in baseline characteristics were observed between this group and those who had PD as their first renal replacement therapy.

Demographic and clinical characteristics of white, brown, and black patients are listed in Table 1. White PD patients were older than black and brown patients. Black and brown patients had lower family incomes compared with white patients, and there were more illiterates among them than in the white group. Both

white and brown patients more often had predialysis care than black patients. However, more black than brown and white patients initiated dialysis with HD as their first therapy. White patients had more diabetic kidney disease and black patients had more hypertensive nephropathy. White patients also had more diabetes mellitus than black and brown patients and had more congestive heart failure. Black patients had more vascular peripheral disease compared with white and brown patients. Body mass index was lower in brown patients compared with black and white patients.

Brown and black patients had significantly better Mental and Physical Component Summary scores compared with white patients (Table 1).

The percentage of black patients was higher in the Southeast region, whereas the percentage of brown patients was higher in the Southeast and Northeast, and for white patients, it was higher in the South and Southeast regions of the country (Table 1).

Table 2 lists laboratory data at baseline for white, black, and brown patients. Compared with white patients, black patients were found to have significantly higher creatinine levels and lower hemoglobin levels.

Follow-up and Outcomes

Mean follow-up in the study was 12.9 months, with 10.4 months for black, 12.5 months for brown, and 13.0 months for white patients. There were 309 (22.6%) deaths in the white, 45 (16.4%) deaths in the black, and 91 (17.6%) deaths in the brown patient groups. The mortality rate was 244.7 deaths/1,000 patient-years for white patients, 188.6 deaths/1,000 patient-years for black patients, and 184.2 deaths/1,000 patient-years for brown patients (Table 3). The competing-risk model comparing the cumulative mortality of white, black, and brown patients shows lower mortality for black and brown patients compared with white patients (Fig 2).

As listed in Table 3, cardiovascular disease was the most important cause of death among black and white patients, whereas in brown patients, it was sepsis. The incidence rate for cardiovascular mortality was 95.0 events/1,000 patient-years for white patients, 71.2 events/1,000 patient-years for black patients, and 38.4 events/1,000 patient-years for brown patients. Compared with white patients, black patients were less likely to die because of sepsis, whereas brown patients had similar incidence rates for mortality due to sepsis (incidence rates: 65.7 events/1,000 patient-years for white patients, 66.8 events/1,000 patient-years for brown patients, and 37.7 events/1,000 patient-years for black patients). The incidence rate for transplantation was higher in brown patients (24.2 events/1,000 patient-years) than in black

Table 1. Demographic and Clinical Characteristics at Baseline

Variable	Black (n = 273)	Brown (n = 516)	White (n = 1,370)	P ^a
Age (y)	57.0 ± 15.5	52.5 ± 17.4	59.8 ± 16.0	<0.001
Female sex	63.7	54.8	52.2	0.002
Education				<0.001
Illiterate	23.1	13.0	8.6	
Elementary school	60.1	55.0	56.2	
Secondary school	12.8	25.2	26.6	
College	4.0	6.8	8.5	
Low family income ^b	56.8	44.2	29.0	<0.001
Distance to center <25 km	56.0	53.7	60.3	0.02
Predialysis care >6 mo	41.0	57.0	56.8	<0.001
HD as first RRT	76.6	68.8	65.5	0.001
PD indication: only option	31.5	19.6	15.3	<0.001
PD type				<0.001
APD	46.9	38.8	54.7	
CAPD	53.1	61.2	45.3	
Geographic region				<0.001
South	8.8	6.4	23.1	
Southeast	65.9	38.8	62.5	
Midwest	4.4	2.7	2.7	
Northeast	19.1	33.5	10.8	
North	1.8	18.6	0.9	
Cause of ESRD				<0.001
Diabetic nephropathy	34.1	34.9	38.2	
Hypertensive nephropathy	32.6	18.0	21.3	
Chronic glomerulonephritis	8.8	12.2	10.9	
Unknown	13.6	19.2	12.7	
Other	11.0	15.7	16.9	
Comorbid conditions				
Systemic hypertension	3.3	5.2	1.8	0.1
Congestive heart failure	78.6	80.1	79.5	0.001
Peripheral vascular disease	20.9	19.4	26.9	0.003
Left ventricular hypertrophy	31.5	20.5	25.2	0.1
Diabetes	43.2	36.4	39.1	0.005
Malignancies	39.2	36.0	44.0	0.08
None	2.2	2.3	4.0	<0.001
Blood pressure (mm Hg)				
Systolic	3.3	5.2	1.8	0.07
Diastolic	139 ± 30	140 ± 27	137 ± 27.	0.04
BMI (kg/m ²)	84 ± 17	85 ± 15	82 ± 16	
BMI (kg/m ²)	24.8 ± 5.1	23.7 ± 4.9	24.9 ± 4.9	<0.001
SF-36				
PCS score	39.9 ± 10.5	41.6 ± 10.6	38.1 ± 10.5	<0.001
MCS score	42.8 ± 5.1	43.2 ± 9.5	40.5 ± 10.0	<0.001

Note: Values for categorical variables are given as percentages; values for continuous variables, as mean ± standard deviation.

Abbreviations: APD, automated peritoneal dialysis; BMI, body mass index; CAPD, continuous ambulatory peritoneal dialysis; CKD, chronic kidney disease; ESRD, end-stage renal disease; HD, hemodialysis; MCS, Mental Component Summary; PCS, Physical Component Summary; PD, peritoneal dialysis; SF-36, 36-Item Short Form Health Survey.

^aP value refers to comparison among the 3 groups by analysis of variance or χ^2 .

^bUp to 2 times minimum national salary.

(16.7 events/1,000 patient-years) and white patients (19.0 events/1,000 patient-years). Black and brown patients were more likely to transfer to HD therapy compared with white patients. The incidence rate for transfer to HD

therapy was 96.4 events/1,000 patient-years in black patients, 93.1 events/1,000 patient-years in brown patients, and 83.9 events/1,000 patient-years in white patients (Table 3). The main cause in all groups for transfer

Table 2. Laboratory Characteristics at Baseline

	Black (n = 273)	Brown (n = 516)	White (n = 1,370)	P ^a
Urea (mmol/L)	35.5 ± 14.2	37.0 ± 14.8	35.7 ± 15.7	0.2
Creatinine (μmol/L)	751.4 ± 371.2	680.7 ± 327.0	645.3 ± 353.6	<0.001 ^b
Potassium (mmol/L)	4.8 ± 1.0	4.4 ± 1.0	4.4 ± 1.5	0.5
ALT (U/L)	20.9 ± 16.8	28 ± 119.0	20.4 ± 13.8	0.07
Calcium (mmol/L)	2.1 ± 1.1	2.2 ± 0.4	2.0 ± 0.5	0.2
Phosphate (mmol/L)	1.5 ± 0.4	1.4 ± 0.4	1.6 ± 0.5	0.001 ^a
Hemoglobin (g/L)	10.9 ± 2.2	11.3 ± 2.5	11.3 ± 2.2	0.03 ^{b,c}
Glucose (mmol/L)	6.5 ± 4.0	6.6 ± 4.6	7.0 ± 4.7	0.1

Note: Values are given as mean ± standard deviation.

Abbreviation: ALT, alanine aminotransferase.

^aP value refers to comparison among the 3 groups by analysis of variance.

^bPost hoc test refers to difference between brown versus white.

^cPost hoc test refers to difference between black versus white.

to HD therapy was peritonitis (35% in white, 22% in brown, and 26% in black patients).

With white patients as the reference group, Cox proportional hazards analysis showed a crude hazard ratio (HR) for mortality of 0.77 (95% confidence interval [CI], 0.56-1.05) for black patients and 0.74 (95% CI, 0.59-0.94) for brown patients (Table 4, model 1). After adjusting for age, these HRs increased but remained lower than 1.0 for black (HR, 0.84; 95% CI, 0.61-1.15) and brown (HR, 0.91; 95% CI, 0.71-1.15) patients (Table 4, model 2). After further adjustment for demographic, socioeconomic, clinical, QoL, and laboratory variables as potential explanatory factors, HRs decreased to 0.67 (95% CI, 0.48-0.95) and 0.77 (95% CI, 0.43-1.01), respectively, for black and brown patients (Table 4, model 6). The Cox proportional hazards analysis was repeated stratified for elderly (aged ≥65 years) and nonelderly (aged <65 years) patients, and no differences were observed (data not shown).

There is no difference between results in sensitivity analyses performed with and without imputation for missing data.

Table 3. Incidence Rates for Outcomes

	Black (n = 273)	Brown (n = 516)	White (n = 1370)
Death	188.6 (45)	184.2 (91)	244.7 (309)
Cardiovascular	71.2 (17)	38.4 (19)	95.0 (120)
Sepsis	37.7 (9)	66.8 (33)	65.7 (83)
Kidney transplantation	16.7 (4)	24.2 (12)	19.0 (24)
Transfer to hemodialysis	96.4 (23)	93.1 (46)	83.9 (106)

Note: Values given as incidence rates per 1,000 patient-years (number of events).

DISCUSSION

In this large cohort of incident PD patients in Brazil, BRAZPD, we observed lower mortality rates for both black and brown patients compared with white patients. These survival advantages remained after adjusting for several potential explanatory variables, with black patients having 33% and brown patients having 23% lower mortality compared with white patients.

The lower mortality in the black population is comparable to findings of other studies of dialysis patients.⁵⁻¹⁰ In a study including only PD patients (n = 233) from the United States, a somewhat stronger effect was observed (whites compared with blacks: HR, 2.35).⁷ In the United Kingdom, the adjusted HR for mortality for black compared with white PD patients was 0.62.¹⁰ In contrast, a nationwide study from the Netherlands did not find lower mortality for black compared with white PD patients. However, the number of black patients was small (n = 45 [2%]) in this study.³⁰ Like most North American and Canadian studies,^{3-5,8} black patients in the present study had higher survival compared with white patients, even after adjustments. Nevertheless, this study presents an important racial difference in comparison to studies performed in the United States and Canada, namely, the presence of the admixed population. Moreover, it also is difficult to compare results because each study is adjusted for different factors⁵⁻⁸ and some studies lack power and find a large result that is not statistically significant. Comparisons to studies performed in a racial minority in Europe^{9,10} also are difficult because the majority of the Brazilian population is mixed, making this study a reflection of the geographic behavior of the entire Brazilian population.³¹




Figure 2. Cumulative mortality curves for white, black, and brown patients treated with peritoneal dialysis, taking into account transfer to hemodialysis therapy and kidney transplantation as competing end points.

Similar to other studies,³² in our sample, the population on dialysis therapy is over-represented in terms of black patients (13%) compared to the general population (6.9%),²¹ suggesting a higher prevalence of ESRD in this population.²¹ The brown population is under-represented (24%) compared to the general population (42.6%).¹⁹ This might be related to the fact that the majority of patients included in this study were from the Southeast and South regions of the country, where the brown population is less prevalent than in other regions.³¹ The majority of the brown population on dialysis therapy lives in the North and Northeast regions, where there are few dialysis centers and late referral to nephrologists is most common. This might result in selection bias for the brown population.³¹

In a recent analysis of US data, Kucirka et al⁶ showed that the survival advantage for black compared with white dialysis patients applies to only

patients older than 50 years. They speculated that this finding was a result of differences in insurance coverage, comorbid conditions, and access to organ transplantation. This finding could not be replicated in our study, which may be related to differences in health care systems between the United States and Brazil. In the United States, all adults older than 65 years are eligible for Medicare, but patients at a young age are more likely to be uninsured or have Medicaid insurance,⁶ whereas in Brazil, there is a universal health care system for patients of all ages.²²

Although health care in Brazil is universal, blacks were found to have late referral to predialysis care more often compared with white and brown patients. This might be related to their lower socioeconomic status.³³ Low socioeconomic status incorporates important determinants of health: health care access, environmental exposure, health behavior, and low health literacy,³⁴ all of which contribute to health care disparities directly and indirectly.³⁵ Socioeconomic status was lower in black and brown patients; however, adjustments for these factors did not affect results.

Other factors that should be considered are the differences in comorbid conditions and QoL. White patients had more diabetes and cardiac failure at baseline than brown and black patients, factors associated with mortality.³⁶ On the other hand, QoL was better in black and brown patients compared with white patients in our study, which is in disagreement with Kutner et al,³⁷ who observed that black PD patients had the lowest satisfaction with care scores after adjustment.³⁸ However, adjustment for these factors did not change results, which is in agreement with a study by van den Beukel et al.³⁰

Table 4. HRs for Mortality for Black and Brown Patients Versus White Patients

Model	Black	Brown	Variables Tested
1. Unadjusted	0.77 (0.56-1.05)	0.74 (0.59-0.94)	Unadjusted
2. Adjusted for age	0.84 (0.61-1.15)	0.91 (0.71-1.15)	Model 1 plus age
3. Sociodemographic	0.70 (0.50-0.98)	0.72 (0.55-0.96)	Model 2 plus sex, level of education, income, geographic region
4. Comorbid conditions and ESRD cause	0.69 (0.50-0.97)	0.71 (0.54-0.94)	Model 3 plus CVD, ^a DM, PVD, malignancy, cause of ESRD
5. First dialysis therapy, QoL	0.69 (0.49-0.97)	0.78 (0.58-1.02)	Model 4 plus first dialysis therapy (HD or PD), QoL (PCS and MCS scores)
6. Laboratory investigations	0.67 (0.48-0.95)	0.77 (0.43-1.01)	Model 5 plus creatinine and hemoglobin levels

Note: With gradually more extensive multivariate models for the association between ethnicity and mortality. Associations given as HR (95% confidence interval).

Abbreviations: CVD, cardiovascular disease; DM, diabetes mellitus; ESRD, end-stage renal disease; HD, hemodialysis; HR, hazard ratio; MCS, Mental Component Summary; PCS, Physical Component Summary; PD, peritoneal dialysis; PVD, peripheral vascular disease; QoL, quality of life.

^aCVD is defined as *International Classification of Disease, Tenth Revision*.

Our results show significantly higher serum creatinine levels in black and brown patients compared with white patients. This may reflect differences in muscle mass and metabolism between whites and non-whites.^{39,40} This is in agreement with Hsu et al,⁴¹ who found that blacks have higher serum creatinine concentrations than whites at dialysis therapy initiation and higher serum creatinine levels are independent predictors of reduced mortality. Higher serum creatinine levels for black and brown patients might affect the time of starting dialysis therapy and hence outcomes on dialysis therapy. However, in our study, adjustment for this variable did not have an impact on results.

Much deliberation, discussion, and debate have resulted from the terms race and ethnicity being used in epidemiologic and public health studies. Many of the methods used to evaluate race and ethnicity do not use an objective measurable characteristic, and there is not adequate evidence that present racial classifications accurately capture biological or genetic similarities; thus, accurate categorization is challenging.⁴² In this study, ethnicity was self-determined, and it is possible that some misclassification might have occurred. Other important variables such as genetic components were not available. Other variables related to survival were not available, such as Kt/V, residual kidney function, and inflammation. Another limitation of this study is that patients were lost to follow-up after they switched to HD therapy, which might have occurred for several reasons, not only for failed PD therapy. This loss to follow-up was more frequent for black and brown patients than for white patients. This observation is in agreement with a study from the United States by Kim et al.⁴³ However, the authors did not observe differences in mortality rates of black patients who were transferred to HD therapy and those who continued on PD therapy.⁴³ Unfortunately, it is not possible to verify if this also holds for the BRAZPD population.

Our study, similar to most studies comparing the survival of black and white dialysis patients elsewhere, demonstrates better survival for black patients compared with white patients in a large incident PD population. The novel observation is that brown patients have better survival compared with white patients and similar to black patients. The clinical and environmental factors alone are insufficient to fully explain the striking and consistent survival advantage that is observed in black and brown Brazilian PD patients. We may add as a final suggestion that collaborative studies among different areas of the world, evaluating both phenotypes and genotypes in dialysis patients, may shed light in the comprehension of the different race outcomes reported to date.

ACKNOWLEDGEMENTS

The following centers participated in BRAZPD and contributed to the preparation of this article: Ameneg, Associação Hosp Bauru, Biocor Hosp Doenças Cardiológicas, Casa De Saude E Mat N.Sra Perp Socorro, Cdr Curitiba, Cdr Goiania, Cdr Imperatriz, Cdr Sao Jose Pinhais, Cdr—Centro Dialise Transplante Renal, Centro Nefrologia Teresopolis, Centro Nefrologico Minas Gerais, Centro Trat Doenças Renais Joinville, Centro Tratamento Renal Zona Sul, Clinef Rio De Janeiro, Clinepa Clinica De Nefrologia Da Paraiba, Clines, Clinese, Clinica Do Rim Do Carpina, Clinica Evangelico S/C Ltda, Clinica Nefrologia De Franca, Clinica Nefrologia Santa Rita, Clinica Nefrologica Sao Goncalo, Clinica Paulista Nefrologia, Clinica Renal Manaus, Clinica Senhor Do Bonfim, Clinica Senhor Do Bonfim Ltda Filial, Clinica Tratamento Renal, Curitiba—Cenec, Clire Clinica Doenças Renais, Famesp Botucatu, Unicamp—Univ. Est Campinas, Hosp. Clinicas Fmrpusp, Fundação Civil Casa Mis Franca, Fundação Inst Mineiro Est Pesq Nefrol, Gamen Rio De Janeiro, Gdf Hospital De Base, Histocom Sociedade Civil Ltda, Hosp Univ Prof Edgard Santos, Hosp Benef. Portuguesa Pernambuco, Hospital Cidade Passo Fundo, Hospital Clinica Univ Fed Goias, Hospital E Maternidade Angelina Caron, Hospital Evangelico Vila Velha Es, Hospital Geral Bonsucesso, Hospital Geral De Goiania, Hospital Infantil Joana De Gusmao, Hospital São Joao Deus, Hospital São Jorge, Hospital São Jose Do Avai, Hosp Sao Vicente De Paula—J Pessoa, Hosp Sao Vicente De Paulo, Hospital Servidor Do Estado Ipase, Hospital Univ Presidente Dutra Ma, Hospital Universitario Antonio Pedro, Hospital Vita Volta Redonda S/A, Iamspe Sao Paulo, Imip, Inst Capixaba Doenças Renais, Inst Capixaba Doenças Renais Cariacica, Inst Capixaba Doenças Renais Serra, Inst Do Rim De Fortaleza, Inst Do Rim De Marilia, Inst Do Rim Do Parana S/C Ltda, Inst Do Rim Santo Antonio Da Platina, Inst Hemodialise De Sorocaba, Inst Medicina Nuclear Endocrina, Inst Nefrologia De Mogi Das Cruzes, Inst Nefrologia De Suzano, Inst Nefrologia Souza E Costa, Inst Urologia E Nefrol Barra Mansa, Inst Urologia E Nefrol Sj Rio Preto, Medservsp, Nefrocentro, Nefroclinica Caxias Do Sul, Nefroclinica Foz Do Iguaçu, Nefroclinica Uberlândia, Nefron Clinica Natal, Nefron Contagem, Nephron Pelotas, Nephron São Paulo, Nucleo Nefrologia Belo Horizonte, Pro Nephron, Prorim Campos Dos Goitacaze, Puc Porto Alegre, Renalcare Serviços Medicos Ltda, Renalcor Angra Dos Reis, Renalcor Rio De Janeiro, Renalvida, Rien Rio De Janeiro, Santa Casa De Adamantina, Santa Casa De Jau—Unefro, Santa Casa De Marilia, Santa Casa De Ourinhos, Santa Casa De Santo Amaro, Santa Casa De Sao Jose Dos Campos, Santa Casa De Votuporanga, Serv De Nefrologia De Ribeirao Preto, Uerj—Hosp. Clin. Univ. Est. Rio De Janeiro, Uni Rim Joao Pessoa, Unidade Nefrologia Assis, Unirim Unidade De Doenças Renais, Unirim Unidade Renal Do Portao, Untr Unidade Nefrologia Transplante.

Aspects of this work were presented in abstract form at the 49th European Renal Association—European Dialysis and Transplant Association (ERA-EDTA) Congress, May 24-27, 2012, Paris, France.

We thank the Brazilian Baxter Renal team for important contributions to the creation and functioning of this project.

Support: This study was funded by Baxter Laboratories, Brazil. Dr van den Beukel thanks the ERA-EDTA for support with a research fellowship (53-2009).

Financial Disclosure: During the data collection and analysis, Dr Divino-Filho was employed by Baxter; his employment by Baxter ended on March 31, 2011. Dr Pecoits-Filho received a consulting fee and speaker honorarium from Baxter Healthcare. The other authors declare that they have no other relevant financial interests.

REFERENCES

- Agodoa L. Lessons from chronic renal diseases in African Americans: treatment implications. *Ethn Dis*. 2003;13(suppl 2):S118-S124.
- Jain AK, Blake P, Cordy P, Garg AX. Global trends in rates of peritoneal dialysis. *J Am Soc Nephrol*. 2012;23:533-544.
- Mesler DE, McCarthy EP, Byrne-Logan S, Ash AS, Moskowitz MA. Does the survival advantage of nonwhite dialysis patients persist after case mix adjustment? *Am J Med*. 1999;106:300-306.
- Tanna MM, Vonesh EF, Korbet SM. Patient survival among incident peritoneal dialysis and hemodialysis patients in an urban setting. *Am J Kidney Dis*. 2000;36:1175-1182.
- Robinson BM, Joffe MM, Pisoni RL, Port FK, Feldman HI. Revisiting survival differences by race and ethnicity among hemodialysis patients: the Dialysis Outcomes and Practice Patterns Study. *J Am Soc Nephrol*. 2006;17:2910-2918.
- Kucirka LM, Grams ME, Lessler J, et al. Association of race and age with survival among patients undergoing dialysis. *JAMA*. 2011;306:620-626.
- Korbet SM, Shih D, Cline KN, Vonesh EF. Racial differences in survival in an urban peritoneal dialysis program. *Am J Kidney Dis*. 1999;34:713-720.
- Pei YP, Greenwood CM, Chery AL, Wu GG. Racial differences in survival of patients on dialysis. *Kidney Int*. 2000;58:1293-1299.
- van den Beukel TO, Jager KJ, Siegert CE, Schoones JW, Dekker FW. Racial minority groups on dialysis in Europe: a literature review. *Clin Nephrol*. 2010;74(suppl 1):S78-S84.
- Roderick P, Byrne C, Casula A, et al. Survival of patients from South Asian and black populations starting renal replacement therapy in England and Wales. *Nephrol Dial Transplant*. 2009;24:3774-3782.
- van den Beukel TO, Dekker FW, Siegert CE. Increased survival of immigrant compared to native dialysis patients in an urban setting in the Netherlands. *Nephrol Dial Transplant*. 2008;23:3571-3577.
- McDonald SP, Russ GR. Current incidence, treatment patterns and outcome of end-stage renal disease among indigenous groups in Australia and New Zealand. *Nephrology (Carlton)*. 2003;8:42-48.
- Nobles M. History counts: a comparative analysis of racial/color categorization in US and Brazilian censuses. *Am J Public Health*. 2000;90:1738-1745.
- Santos RV, Fry PH, Monteiro S, et al. Color, race, and genomic ancestry in Brazil: dialogues between anthropology and genetics. *Curr Anthropol*. 2009;50:787-819.
- Pena SD, Bastos-Rodrigues L, Pimenta JR, Bydlowski SP. DNA tests probe the genomic ancestry of Brazilians. *Braz J Med Biol Res*. 2009;42:870-876.
- Pena SD, Di Pietro G, Fuchshuber-Moraes M, et al. The genomic ancestry of individuals from different geographical regions of Brazil is more uniform than expected. *PLoS One*. 2011;6:e17063.
- Salzano FM, Neel JV, Maybury-Lewis D. I. Demographic data on two additional villages: genetic structure of the tribe. *Am J Hum Genet*. 1967;19:463-489.
- Curtin PD. Epidemiology and the slave trade. *Polit Sci Q*. 1968;83:190-216.
- IBGE. Brazilian Institute of Geography and Statistical-IBGE. Brazil, 2007. <http://www.ibge.gov.br>. Accessed January 1, 2012.
- Sociedade Brasileira de Nefrologia. Censo Brasileiro de Diálise—2011. www.sbn.org.br. Accessed January 1, 2012.
- Noblat AC, Lopes MB, Lopes AA. Race and hypertensive target-organ damage in patients from an university-affiliated outpatient care referral clinic in the city of Salvador. *Arq Bras Cardiol*. 2004;82:111-120.
- Lugon JR, Strogoff de Matos JP. Disparities in end-stage renal disease care in South America. *Clin Nephrol*. 2010;74(suppl 1):S66-S71.
- Fernandes N, Bastos MG, Cassi HV, et al. The Brazilian Peritoneal Dialysis Multicenter Study (BRAZPD): characterization of the cohort. *Kidney Int Suppl*. 2008;108:S145-S151.
- de Andrade Bastos K, Qureshi AR, Lopes AA, et al. Family income and survival in Brazilian Peritoneal Dialysis Multicenter Study Patients (BRAZPD): time to revisit a myth? *Clin J Am Soc Nephrol*. 2011;6:1676-1683.
- International Classification of Diseases, 10th Revision. 1994. <http://www.who.int/classifications/icd/en/WHO>. Accessed January 1, 2012.
- Cury SE, Vilar MJ, Ciconelli RM, Ferraz MB, Atra E. Evaluation of the European Spondylarthropathy Study Group (ESSG) preliminary classification criteria in Brazilian patients. *Clin Exp Rheumatol*. 1997;15:79-82.
- Ministry H. 2007. <http://www2.datasus.gov.br/DATASUS/index.php>. Accessed
- Verduijn M, Grootendorst DC, Dekker FW, Jager KJ, le Cessie S. The analysis of competing events like cause-specific mortality—beware of the Kaplan-Meier method. *Nephrol Dial Transplant*. 2011;26:56-61.
- Donders AR, van der Heijden GJ, Stijnen T, Moons KG. Review: a gentle introduction to imputation of missing values. *J Clin Epidemiol*. 2006;59:1087-1091.
- van den Beukel TO, Verduijn M, le Cessie S, et al. The role of psychosocial factors in ethnic differences in survival on dialysis in the Netherlands. *Nephrol Dial Transplant*. 2011;
- Fernandes N, Chaoubah A, Bastos K, Lopes A, Divino-Filho J, Pecoits-Filho R. Geography of peritoneal dialysis in Brazil: analysis of a cohort of 5,819 patients (BRAZPD). *J Bras Nefrol*. 2010;32:268-274.
- Agodoa L, Eggers P. Racial and ethnic disparities in end-stage kidney failure—survival paradoxes in African-Americans. *Semin Dial*. 2007;20:577-585.
- Sesso R, Belasco AG. Late diagnosis of chronic renal failure and mortality on maintenance dialysis. *Nephrol Dial Transplant*. 1996;11:2417-2420.
- Adams RJ, Appleton SL, Hill CL, Dodd M, Findlay C, Wilson DH. Risks associated with low functional health literacy in an Australian population. *Med J Aust*. 2009;191:530-534.
- Young BA. The interaction of race, poverty, and CKD. *Am J Kidney Dis*. 2010;55:977-980.
- Carrero JJ, de Zeeuw D, Axelsson J, et al. Sex differences in the impact of diabetes on mortality in chronic dialysis patients. *Nephrol Dial Transplant*. 2011;26:270-276.
- Kutner NG, Zhang R, Brogan D. Race, gender, and incident dialysis patients' reported health status and quality of life. *J Am Soc Nephrol*. 2005;16:1440-1448.
- Paniagua R, Amato D, Vonesh E, Guo A, Mujais S, Group MNCS. Health-related quality of life predicts outcomes but is not affected by peritoneal clearance: the ADEMEX trial. *Kidney Int*. 2005;67:1093-1104.
- Lewis J, Agodoa L, Cheek D, et al. Comparison of cross-sectional renal function measurements in African Americans with hypertensive nephrosclerosis and of primary formulas to estimate glomerular filtration rate. *Am J Kidney Dis*. 2001;38:744-753.
- Hsu CY, Chertow GM, Curhan GC. Methodological issues in studying the epidemiology of mild to moderate chronic renal insufficiency. *Kidney Int*. 2002;61:1567-1576.
- Hsu J, Johansen KL, Hsu CY, Kaysen GA, Chertow GM. Higher serum creatinine concentrations in black patients with chronic kidney disease: beyond nutritional status and body composition. *Clin J Am Soc Nephrol*. 2008;3:992-997.
- Comstock RD, Castillo EM, Lindsay SP. Four-year review of the use of race and ethnicity in epidemiologic and public health research. *Am J Epidemiol*. 2004;159:611-619.
- Kim GC, Vonesh EF, Korbet SM. The effect of technique failure on outcome in black patients on continuous ambulatory peritoneal dialysis. *Perit Dial Int*. 2002;22:53-59.