One-anchor structural construction of Standard Model observables, a dark-matter tower, and Newton's constant from $E_8 \times E_8$ Coxeter geometry on a Schoen Calabi–Yau threefold

Eliot Gable*
Independent Researcher
(Dated: May 1, 2026)

We present a one-anchor structural construction for the Standard Model observables treated here — all twenty-six SM parameters, Newton's gravitational constant G, the Planck mass, a discrete seven-mode dark-matter mass tower, and a misalignment-style dark-matter abundance consistency estimate $\Omega_{\rm DM}h^2$ — from a single measured input (the electron rest mass) and the geometry of $E_8 \times E_8$ Coxeter structure on a Schoen Calabi–Yau threefold $X/(\mathbb{Z}_3 \times \mathbb{Z}_3)$, combined with the McKay correspondence between E_8 and the binary icosahedral group. The framework is anchored in a substrate-physical interpretation in which the speed of light is the universal substrate coupling-rate ratio $c = \ell_c/\tau_c$, Planck's constant is the universal substrate threshold magnitude, and general relativity, quantum mechanics, Newton's laws of motion, and Maxwell's equations are treated as macroscopic limits of a single elastic-medium structure $c = \sqrt{\Theta_{\rm restoring}/\mathcal{I}_{\rm inertial}}$. All structural integers are fixed within the framework. Two-anchor self-consistency between the electron and muon anchors is verified at 0.0011 ppm. Across the primary SM observables compared here, the framework remains within current experimental uncertainties, while algebraically derived bookkeeping quantities are reported separately as consistency checks; the framework selects normal neutrino mass ordering, predicts both CP phases as rational multiples of π ($\delta_{\rm CP} = \pi + \pi/10$, $\delta_{\rm CKM} = 11\pi/30$), the weak mixing angle as $\sin^2\theta_W = 2/9 + \Delta_\varphi/h$, the strong coupling as $\alpha_s(M_Z) = 849/7192$, and identifies a conditional geometric route to a vanishing strong CP angle via the $\mathbb{Z}_3 \times \mathbb{Z}_3$ quotient projection, without introducing fitted continuous parameters or invoking a Peccei–Quinn axion. Newton's G is predicted at 2.5 ppm of CODATA 2022, below the present inter-experiment scatter in direct laboratory determinations. Neutrinos and dark matter emerge as a unified Coxeter tower; with the structurally-derived decay constant f_5, misalignment angle $\theta_5 = \arctan\varphi$, and rank-locked sub-harmonic correction $(1 - \operatorname{rank}\alpha_\tau)$ on the dominant order-5 dark-matter mode, the resulting misalignment-style abundance consistency estimate is $\Omega_{\rm DM}h^2 = 0.119982$, consistent with Planck 2018 ($\Omega_{\rm DM}h^2 = 0.1200 \pm 0.001$) at 0.015 %, without introducing any additional fitted cosmological parameter. The framework reduces the Standard Model parameter count from 26 to 1 in the mass-spectrum construction presented here and places the classical, quantum, gravitational, and electromagnetic sectors within the same elastic-substrate foundation. We emphasise that the relationship of the first-principles substrate-physical inputs to the specific structural configuration developed here is one of *strong selection* rather than *forced derivation*: the substrate, heterotic, McKay, and Lie-algebraic inputs narrow the space of admissible configurations sharply enough that this one is the natural landing, but a single missing deductive link, identifying the substrate-physical projection onto the heterotic moduli of $X/(\mathbb{Z}_3 \times \mathbb{Z}_3)$, would be required to upgrade the present construction from selection to forced derivation.

CONTENTS

I. Introduction 2

II. The substrate framework 3
- A. Heterotic $E_8 \times E_8$ on the Schoen quotient 3
- B. The E_8 Coxeter spectrum 4
- C. The McKay correspondence 4
- D. Substrate-tension interpretation 5

III. Substrate foundations: derivation of c, $\hbar$, and the elastic medium 5
- A. The speed of light as a substrate coupling-rate ratio 5
- B. Planck's constant as the universal threshold magnitude 6
- C. Substrate elastic medium 6

IV. Macroscopic limits: GR, QM, Newton's laws, Maxwell's equations 7
- A. Special relativity as substrate Lorentz invariance 7
- B. General relativity as bulk substrate-tension response 7
- C. Quantum mechanics from accumulator + threshold structure 8
- D. Newton's laws from elastic restoring response 9
- E. Maxwell's equations as $U(1)$ cross-term coupling 9
- F. Connection to the $E_8 \times E_8$ Coxeter framework 10

* 10015996+egable@users.noreply.github.com

V. Gateway formula 10

VI. Charged-fermion masses 11
A. Mass formula schema 11
B. Lepton sector 11
C. Quark sector 11
D. Term-by-term decomposition: charged fermions 12

VII. Higgs sector and dark-sector coupling family 13
A. Boson mass-formula schema and structural corrections 13
B. Origin of the dark-sector coupling family 13
C. The Higgs leak: structural closure rule for δ_H 13
D. The W leak: structural closure rule for δ_W 14
E. Generalization to other cycle-2 modes 14
F. Higgs self-coupling 14
G. Term-by-term decomposition: electroweak bosons 14

VIII. Electroweak gauge sector 15
A. Weak mixing angle 15
B. Gauge couplings and Z boson 15
C. Higgs self-coupling 16
D. Strong coupling 16
E. Fine-structure constant 16

IX. Strong CP angle 16

X. Neutrino sector 16
A. Mass eigenvalues from sub-gateway φ-chain modes 16
B. Mass-squared splittings and bounds 17
C. PMNS mixing angles and CP phase 17

XI. Quark mixing (CKM) 17

XII. Dark-matter tower 18
A. The unified neutrino-DM Coxeter tower 18
B. Cosmological cutoffs and viable mass window 18
C. Order-5 dominant mode: couplings and detection prospects 18
D. Geometric-series abundance convergence 18
E. Decay constant and misalignment angle 18
F. Rank-locked sub-harmonic correction 18
G. Term-by-term decomposition: dark-matter tower modes 19
H. Cosmological abundance result 20

XIII. Gravitational sector 20

XIV. Two-anchor verification 20

XV. Results 20

XVI. Predictions and falsification 23

XVII. Discussion 23
A. Reduction of the parameter count 23
B. Common geometric origin 23
C. Comparison to prior work 24
D. Open directions 24
E. Conclusion 24

A. Coxeter step assignments 24

B. Complete structural correction list 24

C. Numerical precision 24

References 25

I. INTRODUCTION

The Standard Model (SM) of particle physics, despite its empirical success, contains twenty-six parameters whose values are not predicted by any underlying principle but determined empirically from observation [1]. These include the three gauge couplings, the Higgs sector parameters (vacuum expectation value and self-coupling), nine charged-fermion Yukawa couplings, three neutrino mass scales, four quark-sector mixing parameters (CKM), four lepton-sector mixing parameters (PMNS), and the QCD vacuum angle θ_{QCD}. Beyond the SM, the existence and mass scale of dark matter constitutes one of the most prominent unresolved puzzles in contemporary physics [2].

This paper presents a framework that structurally constructs all twenty-six SM parameters, plus Newton's gravitational constant, the Planck mass, and a discrete seven-mode dark-matter mass spectrum together with a misalignment-style cosmological abundance estimate, from a single dimensional anchor (the measured electron rest mass) and the geometry of $E_8 \times E_8$ heterotic compactification on a Schoen Calabi–Yau threefold quotient $X/(\mathbb{Z}_3 \times \mathbb{Z}_3)$. The mass-spectrum construction introduces no fitted continuous parameters: every structural integer or discrete coefficient is either fixed directly by the algebra of E_8, the geometry of the Schoen quotient, the McKay correspondence between E_8 and the binary icosahedral group, or adopted as a framework assignment built from those structures. Independent verification through a second mass anchor (the muon) confirms internal consistency to 0.0011 ppm across the entire prediction set.

Two recent results form the empirical foundation:

1. Coldea *et al.* [3] confirmed that the Zamolodchikov E_8 mass spectrum [4] appears in the quantum critical point of $CoNb_2O_6$, with mass ratios in the golden ratio φ and related E_8-derived values.

2. The McKay correspondence [5] establishes a bijection between finite subgroups of SU(2) and the simply-laced ADE Dynkin diagrams. The binary icosahedral group is the McKay dual of E_8, making icosahedral order-5 structure the harmonic seed selected by the framework for the φ-chain.

Combined with a substrate-tension interpretation of the Higgs mechanism [7, 8] on the Schoen fiber-product quotient $X/(\mathbb{Z}_3 \times \mathbb{Z}_3)$ [10, 12, 13], these structural elements combine to produce the one-anchor SM construction analyzed here. Section III summarizes the substrate-physical content of c, $\hbar$, and the elastic-medium identification that anchors the framework. Section IV summarizes how general relativity, quantum mechanics, Newton's laws of motion, and Maxwell's equations arise as macroscopic limits of the same substrate structure that produces the Standard Model parameter spectrum, following the broader framework development in [27].

a. Epistemic status of the construction. We state the scope of the present work plainly. The first-principles substrate-physical derivations summarized in Sec. III (universal coupling-rate ratio $c = \ell_c/\tau_c$, threshold magnitude $\hbar$, the discrete isotropic update rule), together with the established $E_8 \times E_8$ heterotic compactification on the Schoen quotient [12, 13], the McKay correspondence [5], the experimental E_8-spectrum confirmation in $CoNb_2O_6$ [3, 4], and the standard Lie-algebraic data of E_8 (rank, Coxeter number, exponents, root-system order $|\Phi| = h{\cdot}\text{rank}$ [6]), point strongly toward the specific structural configuration developed in this paper: a φ-chain for the icosahedral lepton sector, a $\sqrt{2}$-chain for the octahedral quark/electroweak-boson sector, integer Coxeter-step positions for primary harmonics, half-integer cycle-2 positions for non-rank-locked modes, E_8 exponents as palindromic-pair labels, and the $\mathbb{Z}_3 \times \mathbb{Z}_3$ quotient as the source of the visible/dark-sector split. The volume of consistent phenomenological agreement (lepton masses to ppb, electroweak observables to ppm, all four PMNS and four CKM parameters to percent-level closed forms, etc.) is sufficient to rule out generic alternative configurations within the same substrate-physical and heterotic-compactification setting.

We have not, however, completed the bridge from these first-principles inputs to a forced derivation of the present configuration. A closed deductive chain that selects, e.g., the φ-chain assignment for the lepton sector against the $\sqrt{2}$-chain alternative; the half-integer cycle-2 positions used by the down quark, W boson, charm quark, and top quark; the rank-locked-subspace identification within the Calabi–Yau cohomology; the specific bundle data on X/Γ realising the three-generation $c_3(V)/|\Gamma| = -3$ outcome; the quotient-trace identity required for $\theta_{\text{QCD}} = 0$; and the framework structural assignments flagged at points throughout the construction (the leading 2/9 in $\sin^2\theta_W$, the 4 in the gateway sub-harmonic loss 988/987, the $\alpha_s = 849/7192$ structural form, etc.) is not given here. The relationship of the first-principles inputs to the structural configuration is therefore one of *strong selection* rather than *forced derivation*: the inputs narrow the space of admissible configurations sharply enough that the present one is the natural — arguably the only — consistent landing, but a single missing deductive link, presumably at the level of an explicit characterisation of the substrate-physical projection onto the heterotic moduli of X/Γ, would be required to upgrade the construction from selection to derivation. Identifying that link is the central open problem the framework leaves for future work.

II. THE SUBSTRATE FRAMEWORK

A. Heterotic $E_8 \times E_8$ on the Schoen quotient

The framework treats heterotic $E_8 \times E_8$ string theory [11] compactified on the Schoen Calabi–Yau threefold quotient $X/(\mathbb{Z}_3 \times \mathbb{Z}_3)$ as the structural substrate. The Schoen threefold [10] is the smooth, simply-connected Calabi–Yau fiber product $X = B \times_{\mathbb{P}^1} B'$ of two rational elliptic surfaces over $\mathbb{P}^1$, with Hodge numbers $h^{1,1}(X) = h^{2,1}(X) = 19$ and Euler characteristic $\chi(X) = 0$. In the heterotic standard-model construction, suitably chosen B, B' admit a freely acting $\Gamma = \mathbb{Z}_3 \times \mathbb{Z}_3$ symmetry, and the quotient X/Γ acquires $\pi_1(X/\Gamma) = \Gamma = \mathbb{Z}_3 \times \mathbb{Z}_3$ (while X itself remains simply-connected), enabling Wilson-line breaking of one E_8 factor to Standard-Model gauge structure [12, 13].

a. Hodge numbers and generation count. The Schoen quotient X/Γ has Hodge numbers $h^{1,1}(X/\Gamma) = h^{2,1}(X/\Gamma) = 3$ (the Γ-invariant subspaces of $H^{1,1}(X)$ and $H^{2,1}(X)$, each 19/9-dimensional after the Γ-projection rounds to the freely-acting integer dimensions; see [12] for the explicit representation-theoretic decomposition). The Euler characteristic of the cover, $\chi(X) = 2[h^{1,1}(X) - h^{2,1}(X)] = 0$, divides under the freely-acting Γ to give $\chi(X/\Gamma) = 0$ in absolute Euler-characteristic terms; the *net chirality index* relevant for the standard heterotic generation count is set instead by the rank of the $SU(5)$ (or $SU(4) \times U(1)$) bundle V chosen to break E_8 via Wilson lines, with $|n_g - \bar{n}_g| = |c_3(V)|/|\Gamma|$. The standard Braun–He–Ovrut–Pantev model selects V such that $c_3(V)/|\Gamma| = -3$, yielding three chiral generations as required by the visible Standard Model. The present framework assumes (rather than derives) this generation-count outcome of the bundle choice.

b. Heterotic anomaly cancellation. Heterotic $E_8 \times E_8$ on a Calabi–Yau threefold requires the modified Bianchi identity $dH = (\alpha'/4)\left(\operatorname{tr} R \wedge R - \operatorname{tr} F \wedge F\right)$ to admit a globally defined H, equivalent to the cohomological condition

$$\mathrm{ch}_2(V) \;=\; \mathrm{ch}_2(TX) \;+\; [W], \tag{1}$$

where ch_2 is the second Chern character, V is the visible-sector gauge bundle, TX is the holomorphic tangent bundle of the Schoen threefold, and $[W]$ is an effective 5-brane class in $H^4(X, \mathbb{Z})$ (vanishing in the standard embedding $V = TX$). The present framework assumes that the bundle choice underlying the three-generation count is consistent with Eq. (1) for some admissible 5-brane class $[W]$, as is the case in the Braun–He–Ovrut–Pantev construction [12]. A first-principles identification of the specific $[W]$ used by the framework, together

with the *slope-stability* (μ-stability) of V required by the holomorphic-Yang–Mills equations on the quotient X/Γ [13], is left for future work.

c. Substrate dimensionality and the 10D embedding. The discrete substrate update rule Eq. (4) is specified directly in $3+1$ macroscopic dimensions, while the heterotic $E_8 \times E_8$ string lives in $9+1$. The framework's operational stance is that the six internal dimensions are *algebraically encoded* in the substrate by the $E_8 \times E_8$ structure of the chain construction (in particular, the Coxeter-chain positions k_X and the sector-modifier $\sqrt{2}$- vs. φ-chain assignments substitute for direct propagation in the compactified directions): the substrate is treated as $3+1$D throughout, with the Calabi–Yau / heterotic structure entering only through the algebra controlling the mass spectrum and gauge structure. This is a *framework choice*, not a derivation: a substrate-physical realisation in which the $3+1$D macroscopic substrate is a boundary or low-energy slice of an underlying $9+1$D substrate (analogous in spirit to brane-world or warped-compactification constructions) is consistent with the present construction but not committed to here. In particular, no claim is made about the size, geometry, or stabilisation of the six internal directions beyond what is needed to identify the algebraic content $E_8 \times E_8$ on X/Γ.

One E_8 factor contains the visible Standard Model gauge structure after compactification. The second E_8 factor is the dark sector, communicating with the visible sector only gravitationally except through identified portal couplings.

d. Visible $E_8 \to$ SM symmetry-breaking chain. The visible E_8 is broken to the Standard-Model gauge group $G_{\rm SM} = SU(3)_{\rm color} \times SU(2)_L \times U(1)_Y$ through the standard heterotic chain

$$E_8 \supset E_6{\times}SU(3) \supset SO(10){\times}U(1) \supset SU(5){\times}U(1) \supset G_{\rm SM}, \tag{2}$$

with the final descent realised by Wilson lines along generators of $\pi_1(X/\Gamma) = \mathbb{Z}_3 \times \mathbb{Z}_3$. The intermediate $SU(5)$ rank-4 GUT structure is the natural host for the rank-4 bundle V implied by the three-generation condition $c_3(V)/|\Gamma| = -3$. Alternative chains (e.g. $E_8 \supset SO(16)$) admit consistent compactifications but are not used here: the $SU(5)$-route construction is canonical for the Schoen quotient with this bundle choice.

B. The E_8 Coxeter spectrum

The E_8 Lie algebra has dimension $\dim(E_8) = 248$, rank 8, 240 roots, Coxeter number $h(E_8) = 30$, and Weyl group order $|W(E_8)| = 696{,}729{,}600$. The exponents of E_8 are the eight integers

$$\{m_1, \ldots, m_8\} = \{1, 7, 11, 13, 17, 19, 23, 29\}, \tag{3}$$

palindromically symmetric around $h/2 = 15$ in pairs $(1,29)$, $(7,23)$, $(11,19)$, $(13,17)$, satisfying $m_j + m_{9-j} = h$. The longest Weyl element acts as $w_0 = -1$ on the Cartan subalgebra of E_8 (a property shared by $A_1, B_n, C_n, D_{2n}, E_7, E_8, F_4, G_2$). The harmonic threshold $h/2 = 15$ is the framework's chirality-flip locus for the Coxeter chain and is not itself an exponent.

a. Framework extension beyond the standard Coxeter spectrum. The mass-formula construction in this paper parameterizes *continuous* chain positions $k \in \mathbb{R}_{\geq 0}$, not just the eight integer exponents above. Half-integer positions ($k_d = 6.5$, $k_W = 34.5$, $k_t = 36.5$, etc.) and "cycle-2" positions ($k > h$) are framework-specific extensions that interpolate between integer exponents and re-use the periodicity of the Coxeter element across multiple cycles. These are structural assignments within the framework rather than standard Coxeter-theoretic invariants. The structural "boundary capacity" $h + m_{\max} = 59$ used in the dark-sector coupling family (Sec. VII) is likewise a framework combination of E_8 invariants rather than an established Lie-algebra invariant in its own right.

C. The McKay correspondence

The McKay correspondence [5] establishes a bijection between finite subgroups of $SU(2)$ and the affine simply-laced ADE Dynkin diagrams: cyclic $\mathbb{Z}_n \leftrightarrow \widehat{A}_{n-1}$, binary dihedral $\leftrightarrow \widehat{D}_n$, binary tetrahedral $\leftrightarrow \widehat{E}_6$, binary octahedral $\leftrightarrow \widehat{E}_7$, and **binary icosahedral** $\leftrightarrow \widehat{E}_8$. The icosahedron has 5-fold rotational symmetry as one of its characteristic rotational orders in the $(2,3,5)$-triangle group, with $\varphi = 2\cos(\pi/5) = (1+\sqrt{5})/2$ the characteristic icosahedral angular ratio. This identifies the integer 5 as the framework's preferred icosahedral order associated with the E_8 McKay dual.

a. Two harmonic chains used by the framework. The mass-formula construction below distinguishes a φ-chain (with φ^k primary harmonics, motivated by 5-fold icosahedral recursion at the binary-icosahedral McKay-dual of E_8) from a $\sqrt{2}$-chain (with $(\sqrt{2})^k$ primary harmonics, motivated by 4-fold octahedral recursion at the binary-octahedral McKay-dual of E_7, where $\sqrt{2} = 2\cos(\pi/4)$). Charged leptons and neutrinos populate the φ-chain; quarks and electroweak bosons populate the $\sqrt{2}$-chain. This sector split is a *framework structural assignment* rather than a derivation from E_8 alone — a single E_8 algebra does not naturally decompose into icosahedral and octahedral sub-recursions, and the allocation of leptons to φ and quarks to $\sqrt{2}$ is adopted on phenomenological grounds (matching observed masses with small-integer Coxeter-step assignments). A first-principles derivation of the chain assignment from a specific Calabi–Yau compactification is left for future work.

b. Chain factors are not Coxeter eigenvalues. We emphasise that the harmonic factors φ^k and $(\sqrt{2})^k$ that appear throughout the mass formulas below are *exponential growth/recursion factors* on a discrete chain — successive applications of a fixed scalar step — and are *not* eigenvalues of the Coxeter element of E_8. The latter are the unit-modulus roots of unity $|e^{2\pi i e_j/h}| = 1$ for

$e_j \in \{1, 7, 11, 13, 17, 19, 23, 29\}$, which describe rotational structure on the Coxeter plane and have nothing directly to do with mass-ratio recursion. The exponents e_j enter the mass formulas separately, as integer positions used to label palindromic pairings (e.g. $11 + 19 = h$); the multiplicative factors $\varphi^k, (\sqrt{2})^k$ encode the geometric step magnitude of the chain itself. These are two mathematically distinct uses of E_8 data, and the framework's construction is the joint use of both.

D. Substrate-tension interpretation

The framework adopts a substrate-tension interpretation of the Higgs mechanism, motivated by Anderson's superconducting precursor to gauge-boson mass generation [7] and by Volovik's quantum-liquid / helium-droplet emergent-vacuum program [8, 9]. Within the substrate framework, the Higgs vacuum expectation value v is interpreted as the baseline relaxed tension of the $E_8 \times E_8$ substrate, and massive SM particles are localized deformations of this tension at structurally-determined Coxeter positions. Photons are massless because they correspond to solitonic excitations of the substrate that do not deform its baseline tension.

III. SUBSTRATE FOUNDATIONS: DERIVATION OF c, $\hbar$, AND THE ELASTIC MEDIUM

Before the structural derivation of Standard Model parameters can be presented, we summarize the substrate-physical content of the fundamental constants and equations of motion that govern propagation on the $E_8 \times E_8$ substrate. The full derivation is developed in [27]; here we collect the results that anchor the $E_8 \times E_8$ Coxeter framework to standard physics.

A. The speed of light as a substrate coupling-rate ratio

The substrate is committed by axiom to a positive coupling reach ℓ_c and a positive coupling time τ_c – the elementary spatial extent over which one local-coupling event reaches, and the elementary duration of that event.

a. Derivation of the wave equation. Let $\phi(\vec{x}, t)$ be a substrate-amplitude pattern. The leading-order linear, isotropic, time-reversal-symmetric local-coupling update rule that uses only nearest-neighbor spatial coupling and nearest-neighbor temporal correlation is

$$\begin{aligned}\phi(\vec{x}, t+\tau_c) - 2\phi(\vec{x}, t) + \phi(\vec{x}, t-\tau_c) \\ = B \sum_{i=1}^{3} \big[\, \phi(\vec{x} + \ell_c \hat{e}_i, t) - 2\phi(\vec{x}, t) \\ + \phi(\vec{x} - \ell_c \hat{e}_i, t) \,\big],\end{aligned} \tag{4}$$

for some dimensionless coupling coefficient B. The Taylor expansions

$$\phi(\vec{x}, t \pm \tau_c) = \phi \pm \tau_c \partial_t \phi + \tfrac{1}{2}\tau_c^2 \partial_t^2 \phi + O(\tau_c^3), \tag{5}$$

$$\phi(\vec{x} \pm \ell_c \hat{e}_i, t) = \phi \pm \ell_c \partial_i \phi + \tfrac{1}{2}\ell_c^2 \partial_i^2 \phi + O(\ell_c^3), \tag{6}$$

substituted into Eq. (4) and divided by τ_c^2 in the continuum limit $\tau_c, \ell_c \to 0$, yield

$$\partial_t^2 \phi = B \cdot \frac{\ell_c^2}{\tau_c^2} \nabla^2 \phi. \tag{7}$$

The continuum equation has the form of a relativistic wave equation with squared wave-speed $c^2 = B\, \ell_c^2/\tau_c^2$.

b. Lattice dispersion and emergent Lorentz invariance. The full discrete dispersion relation associated with Eq. (4) is

$$\sin^2\!\Big(\tfrac{\omega\tau_c}{2}\Big) \;=\; B \sum_{i=1}^{3} \sin^2\!\Big(\tfrac{k_i \ell_c}{2}\Big), \tag{8}$$

obtained by substituting $\phi \propto \exp[i(\vec{k} \cdot \vec{x} - \omega t)]$ into Eq. (4). Expanding at small $|\vec{k}|\, \ell_c$ and $\omega\tau_c$ yields $\omega^2 = B\, c^2 |\vec{k}|^2 + O(\ell_c^2 |\vec{k}|^4)$, with the sub-leading term containing a cubic-lattice anisotropy $\sum_i k_i^4 \neq |\vec{k}|^4$. *Exact* Lorentz invariance is therefore recovered only in the continuum limit $\ell_c, \tau_c \to 0$; finite-k modes carry $O(k^2 \ell_c^2)$ corrections to the relativistic dispersion, so the substrate's Lorentz invariance is an emergent long-wavelength symmetry. With $\tau_c \sim t_{\text{Planck}}$ this anisotropy is unobservably small at any laboratory or astrophysical wavelength realised to date, and present cosmic-ray photon-timing constraints (discussed below) are already consistent with the identification.

c. CFL stability and the marginal-luminal choice. With $B = 1$ the discrete update Eq. (4) sits at $c\tau_c/\ell_c = 1$, the *marginal CFL stability point* of an explicit central-difference scheme on a cubic lattice. The standard CFL condition for stability of such a scheme in d spatial dimensions is $c\tau_c/\ell_c \leq 1/\sqrt{d}$ (Courant–Friedrichs–Lewy 1928), so for $d = 3$ the marginal limit is $c\tau_c/\ell_c = 1/\sqrt{3}$. The choice $B = 1$ therefore exceeds the standard subluminal CFL bound by a factor of $\sqrt{3}$ along the lattice diagonal: axial signal propagation is ringing-free at the luminal limit, whereas off-axis (diagonal) modes carry a residual $O(\ell_c^2 k^2)$ instability with growth rate suppressed only by powers of $\ell_c k$. In the substrate-physical interpretation, this is read as the statement that c is the *global* maximum substrate-deformation rate along any axis, while finite-k off-axis modes near the lattice cutoff are unphysical excitations that the substrate does not support: the framework identifies the substrate's continuum limit as the physically realised regime and treats the lattice artefact at $|\vec{k}|\ell_c \sim 1$ as a non-observable ultraviolet boundary, consistent with $\ell_c \sim l_{\text{Planck}}$ being many orders of magnitude below any observed wavelength. A first-principles continuum derivation of $c = \ell_c/\tau_c$ without

invoking a discrete cubic lattice (e.g. a Hamilton–Jacobi formulation on a continuum substrate manifold) is left for future work.

By the conventional choice $B = 1$ we identify the universal scalar

$$\boxed{c \;\equiv\; \frac{\ell_c}{\tau_c},} \tag{9}$$

which absorbs B into the definition of c in terms of the elementary substrate scales (alternative choices of B correspond to a redefinition of either ℓ_c or τ_c by a constant factor). Lorentz invariance fixes c as a frame-independent constant, but does not separately determine B. With this choice, Eq. (7) becomes

$$\tau_c^2\,\partial_t^2\phi = \ell_c^2\,\nabla^2\phi \quad\Longleftrightarrow\quad \frac{1}{c^2}\partial_t^2\phi - \nabla^2\phi \equiv \Box\phi = 0. \tag{10}$$

Reading off, c is the rate at which one elementary substrate-coupling event extends through space per elementary substrate-coupling duration. The framework forbids ℓ_c and τ_c from being separately observable; only their ratio is. As a substrate-physical quantity, c represents not merely the propagation speed of free mode-patterns but the *global maximum rate at which the substrate can deform itself*. This bound is universal: it applies to every kinematic process in the framework, including the apparent "faster-than-light" geometries of proposed warp-drive metrics [29], in which the warp bubble's apparent displacement is a coordinate-frame artifact of substrate-tension redistribution whose underlying local deformation rate remains bounded by c. The cosmological inflationary epoch likewise does not violate this bound: inflation is the rapid expansion of the substrate's spatial extent itself rather than the propagation of any mode-pattern through pre-existing space, and the local rate of substrate deformation per coupling-event remains bounded by c throughout [27]. Ultra-high-energy cosmic-ray photon timing currently provides the tightest *observational* upper limit identified to date, $\tau_c \lesssim 10^{-34}$ s; this is the maximum value of τ_c consistent with present observations rather than a theoretical bound, and future higher-energy or longer-baseline observations may further constrain it. As a separate *framework identification* — not a forced consequence of the substrate axioms in Sec. III but an independently falsifiable scale-fixing postulate — we set

$$\tau_c \;\equiv\; t_{\rm Planck} \;=\; \sqrt{\frac{\hbar G}{c^5}}, \qquad \ell_c \;\equiv\; l_{\rm Planck} \;=\; \sqrt{\frac{\hbar G}{c^3}}, \tag{11}$$

with $c = \ell_c/\tau_c$ identically. Combined with the framework's predicted Newton's constant G (Sec. XIII, agreeing with CODATA 2022 at 2.5 ppm), this gives $\tau_c \approx 5.391 \times 10^{-44}$ s and $\ell_c \approx 1.616 \times 10^{-35}$ m at $\sim$1.25 ppm framework precision (since $t_{\rm Planck} \propto \sqrt{G}$), ten orders of magnitude tighter than the cosmic-ray observational upper limit and falsifiable should τ_c ever be measured at a scale incompatible with $t_{\rm Planck}$.

B. Planck's constant as the universal threshold magnitude

The substrate is continuous in amplitude, configuration, and tension, yet event counts (detector clicks, atomic transitions) are discrete. We introduce a continuously-accumulating coupling amplitude $C(t)$ whose rate of change is set by the local substrate-tension content E of the participating mode-pattern, and which resolves into a discrete information-transfer event each time C crosses a universal threshold magnitude. That magnitude is

$$\boxed{\hbar \;\equiv\; \text{universal substrate threshold magnitude.}} \tag{12}$$

a. Derivation of the Planck–Einstein relation. For a stationary mode-pattern with substrate-tension content E, the accumulator advances at the constant rate

$$\frac{dC}{dt} = E. \tag{13}$$

A complete substrate-cycle event corresponds to C traversing the total per-cycle action $\Delta C = h$ (where h is the Planck-constant-as-action-per-cycle), in elapsed time $T = h/E$. Identifying T with the period of the corresponding mode-pattern gives the angular frequency

$$\omega \;=\; \frac{2\pi}{T} \;=\; \frac{2\pi E}{h} \;=\; \frac{E}{\hbar}, \tag{14}$$

where the reduced Planck constant $\hbar = h/(2\pi)$ is the action-per-radian threshold. This is the Planck–Einstein relation. For a mode-pattern of rest mass m at rest, the substrate-tension content is the rest energy $E = mc^2$, and Eq. (14) yields the Compton angular frequency

$$\omega_C \;=\; \frac{mc^2}{\hbar}, \qquad \lambda_C \;=\; \frac{2\pi c}{\omega_C} \;=\; \frac{h}{mc}. \tag{15}$$

The boxed identification Eq. (12) is therefore most precisely stated as: $\hbar$ is the universal substrate threshold magnitude per radian of cycle phase, equivalently $h = 2\pi\hbar$ per full cycle.

b. Sub-threshold activity. Below the threshold magnitude $\hbar$: virtual-particle activity, vacuum fluctuations, Casimir effect, Lamb shift, and anomalous magnetic moments are real physical signatures of below-threshold accumulator activity that never transfer information. At threshold: information transfers, the event count advances by one. The constant proportionality $\hbar$ in Eq. (14) is therefore the structural content of Planck's constant in the substrate framework.

C. Substrate elastic medium

a. Derivation of the elastic-medium identification. A linear elastic medium with restoring stiffness Θ and inertial response $\mathcal{I}$ obeys the canonical wave equation

$$\mathcal{I}\,\partial_t^2\phi \;=\; \Theta\,\nabla^2\phi, \tag{16}$$

which can be written equivalently as $\partial_t^2\phi = (\Theta/\mathcal{I})\nabla^2\phi$ with wave speed $v_{\text{wave}} = \sqrt{\Theta/\mathcal{I}}$.

Comparing Eq. (16) term-by-term with the d'Alembert equation (10), $\partial_t^2\phi = c^2\nabla^2\phi$, the identification

$$\boxed{c \;=\; \sqrt{\frac{\Theta_{\text{restoring}}}{\mathcal{I}_{\text{inertial}}}}} \tag{17}$$

follows immediately, with $c^2 = \Theta_{\text{restoring}}/\mathcal{I}_{\text{inertial}}$ matching the elastic wave speed. Combined with the substrate coupling-rate identification $c = \ell_c/\tau_c$ from Eq. (9), the dimensional content of Θ and $\mathcal{I}$ becomes

$$\frac{\Theta_{\text{restoring}}}{\mathcal{I}_{\text{inertial}}} = \frac{\ell_c^2}{\tau_c^2}, \tag{18}$$

fixing the ratio of stiffness to inertia in terms of the substrate elementary scales.

This identification is the substrate-physical content of the framework: the $E_8 \times E_8$ heterotic substrate behaves as a Lorentz-invariant elastic medium whose coupling reach and coupling time are set by the geometry of the Schoen Calabi–Yau quotient. The Higgs vacuum expectation value v is interpreted within the framework as the baseline tension of this medium; particle masses are localized deformations of this tension at the Coxeter positions tabulated in Sec. II and quantified by the gateway formula in Sec. V.

IV. MACROSCOPIC LIMITS: GR, QM, NEWTON'S LAWS, MAXWELL'S EQUATIONS

A. Special relativity as substrate Lorentz invariance

a. Derivation chain. Special relativity emerges from substrate first principles as the long-wavelength continuum-limit symmetry of the discrete update rule Eq. (4) (cf. Eq. (8) and the discussion of emergent Lorentz invariance in Sec. III). The argument has three steps.

Step 1: Universal coupling-rate ratio. The d'Alembert equation Eq. (10) requires the universal scalar $c = \ell_c/\tau_c$ to be the same in every inertial frame; if two observers measured different ratios $c_1 \neq c_2$, the $B = 1$ condition forcing isotropy of the wave-equation operator would fail. Hence the constancy of c is forced rather than postulated.

Step 2: Linear transformations preserving c. The most general linear coordinate transformation $(t, \vec{x}) \to (t', \vec{x}')$ that preserves the d'Alembert operator $\Box = (1/c^2)\partial_t^2 - \nabla^2$ is the Lorentz transformation: for boost velocity $\vec{v}$,

$$t' \;=\; \gamma\big(t - \vec{v}\cdot\vec{x}/c^2\big), \tag{19}$$

$$\vec{x}'_{\parallel} \;=\; \gamma\big(\vec{x}_{\parallel} - \vec{v}t\big), \qquad \vec{x}'_{\perp} \;=\; \vec{x}_{\perp}, \tag{20}$$

with $\gamma = 1/\sqrt{1 - v^2/c^2}$. The substrate elementary scales ℓ_c, τ_c are intrinsic Lorentz-invariant constants (the same in every inertial frame); the standard relativistic length contraction and time dilation effects act on extended physical objects (rulers and clocks built from many mode-patterns) as observation-frame artifacts of Eqs. (19)–(20), not on the elementary substrate scales themselves. For an extended object with proper length L_0 and proper time interval T_0 measured in the rest frame, a boosted observer measures

$$L \;=\; L_0/\gamma, \qquad T \;=\; \gamma T_0 \quad \text{(observation-frame intervals)}, \tag{21}$$

recovering the standard Lorentz contraction and time dilation.

Step 3: Mass-energy equivalence. For a substrate mode-pattern with rest substrate-tension content E_0, threshold-crossing rate (Eq. (14)) at rest $\omega_0 = E_0/\hbar$. Under a boost, the threshold-crossing rate transforms as $\omega = \gamma\omega_0$, so the boosted energy is

$$E \;=\; \hbar\omega \;=\; \gamma E_0. \tag{22}$$

For a mode at rest with rest-mass m, $E_0 = mc^2$ (Compton relation, Eq. (15)); for a mode in motion, $E^2 = E_0^2 + (pc)^2 = (mc^2)^2 + (pc)^2$, the standard relativistic dispersion. Length contraction, time dilation, and relativistic mass–energy equivalence all follow as immediate consequences of Eqs. (19)–(22).

The substrate-physical content of special relativity is thus that the universal ratio $c = \ell_c/\tau_c$ is the same in all inertial frames, with elementary length and time scales transforming inversely so the ratio is preserved.

B. General relativity as bulk substrate-tension response

a. Derivation chain. General relativity emerges in the present framework when the substrate's elementary scales ℓ_c, τ_c are allowed to vary with position in response to local stress-energy density. The argument below is a compressed summary of the substrate-side construction developed in [27]; it records the continuum-limit closure used later in the paper rather than a self-contained proof.

Step 1: Tension-modulated proper time and length. Let the local substrate-tension content at spatial position $\vec{x}$ be characterized by a scalar Newtonian-potential analog $\Phi(\vec{x})$. The substrate elementary scales ℓ_c, τ_c remain intrinsic Lorentz invariants, but the *proper time* and *proper length* intervals at position $\vec{x}$ – the physical clock-tick and ruler intervals an observer measures locally – are modulated by the tension content. For a clock at rest at position $\vec{x}$,

$$d\tau_{\text{proper}}(\vec{x}) \;=\; \sqrt{1 + 2\Phi(\vec{x})/c^2}\; dt \tag{23}$$

relative to coordinate-time dt at infinity, while at leading order in Φ/c^2 the proper radial length scales inversely:

$$d\ell_{\text{proper}}(\vec{x}) \;\approx\; \big(1 - \Phi(\vec{x})/c^2\big)\, dr \tag{24}$$

(the standard weak-field-limit Schwarzschild-style scaling, in which g_{00} and g_{ii} scale oppositely at leading order). The modulation is universal across mode species — the substrate-physical content of the equivalence principle — and is consistent with the equivalence-principle precision $\eta < 10^{-15}$ [30]. The local speed-of-light ratio c remains invariant in any local freely-falling inertial frame.

Step 2: Emergence of the Lorentzian metric. Eq. (23) identifies $g_{00} = -(1 + 2\Phi/c^2)$ in the weak-field limit, recovering Newtonian gravity at leading order; Eq. (24) identifies the spatial metric component as $g_{ii} \approx 1-2\Phi/c^2$ (opposite sign at leading order). Combining these with the special-relativistic result of Sec. IV A (local Lorentz invariance at every point in freely-falling inertial frames) extends the position-dependent proper-time/proper-length modulation to the full Lorentzian metric $g_{\mu\nu}(\vec{x}, t)$ on the substrate. The line element becomes

$$ds^2 \;=\; g_{\mu\nu}(x)\, dx^\mu dx^\nu, \tag{25}$$

with $g_{\mu\nu}$ identified at every point as the local tension-modulated metric, and special relativity recovered as the limit $g_{\mu\nu} \to \eta_{\mu\nu}$ when $\Phi \to 0$.

Step 3: Continuum closure. Once the substrate response is encoded in a Lorentzian metric $g_{\mu\nu}$, the continuum theory must relate curvature to the local stress-energy distribution in a generally covariant, local, second-order equation that preserves stress-energy conservation and reduces to the Newtonian limit already identified in Step 1. In four dimensions, the minimal closure with these properties is

$$G_{\mu\nu} \;=\; \frac{8\pi G}{c^4}\, T_{\mu\nu}, \tag{26}$$

the Einstein field equations describing the bulk geometric response to the local stress-energy distribution. Within the substrate framework, Eq. (26) is therefore not taken as an independent starting axiom but as the continuum closure of the tension-modulated metric response; the detailed substrate-side argument is developed in [27].

b. Negative prediction. Gravity in this framework is bulk tension geometry, not a mode-mediated cross-term coupling: there is no graviton mode species, and quantum-gravity-as-gauge-boson-exchange programs are structurally misguided. The Newton's-constant prediction $G^{\text{pred}} = 6.67431691 \times 10^{-11}$ m^{3}kg^{-1}s^{-2} (Sec. XIII) follows from Eq. (26) combined with the Coxeter-spectrum mass scales.

The Newton's constant prediction in Sec. XIII (G to 2.5 ppm of CODATA) is therefore not a coincidence: it is the substrate-physical content of $c^4/(8\pi)$ in Eq. (26) combined with the structural mass scales fixed by the Coxeter spectrum.

At sufficiently high local stress-energy density, the inertial response $\mathcal{I}_{\text{inertial}}$ diverges and the wave equation Eq. (10) degenerates. The unique resolution consistent with substrate continuity and information conservation is *substrate inversion* through the event horizon into a child substrate region [27]. The black-hole event horizon is the single bidirectional substrate boundary between the elastic regime (outside) and the inversion regime (inside). The inversion event is itself a catastrophe-theoretic ADE event [28]: the local degeneration of the substrate's elastic-response operator at $\mathcal{I}_{\text{inertial}} \to \infty$ is governed by the same ADE classification (cf. McKay correspondence, Sec. II) that organizes the $E_8 \times E_8$ Coxeter spectrum used throughout this paper, providing a structural link between the cosmological inversion threshold and the framework's particle-spectrum classification.

Matter and information traversing the inversion threshold are not destroyed: substrate continuity and information conservation [27] require that the stress-energy content falling through the horizon be re-emitted into the connected child substrate region as a relativistic flux of disassembled mode-pattern content — a sub-atomic particle soup at relativistic speeds, with composite mode-patterns (atoms, nuclei, hadrons) reduced to their constituent free mode-patterns by the extreme substrate-tension gradient at the inversion locus. The process is structurally bidirectional: the same substrate boundary that admits infalling content from the parent substrate region admits outgoing content into the child substrate region, with no preferred temporal direction at the boundary itself. This identifies the cosmological "Big Bang" of any one substrate region with the inversion outflow of a parent black hole in another substrate region, and it forces matter–antimatter parity, baryon-number distribution, and the initial cosmological power spectrum of the child region to inherit specific structural properties from the parent's inversion event — properties whose explicit derivation is developed in [27].

C. Quantum mechanics from accumulator + threshold structure

The substrate-amplitude pattern $\phi(\vec{x}, t)$ obeying Eq. (10) is identified within the framework with the wave-function of standard quantum mechanics. The full framework account of measurement, threshold crossing, entanglement, and Bell correlations is developed in [27]; here we record only the continuum-limit equations used later in the paper and the substrate-side interpretation attached to them.

a. Klein–Gordon from substrate restoring response. For a mode-pattern with substrate-tension content $E = mc^2$ at rest, the Compton angular frequency Eq. (15) fixes the natural local restoring scale of the corresponding substrate deformation. The massless d'Alembert equation (10) describes freely propagating tension waves; a localized massive mode requires an additional quadratic restoring term that pins the mode to its rest oscillation scale $\omega_C = mc^2/\hbar$. The minimal Lorentz-invariant con-

tinuum equation with that restoring scale is

$$\frac{1}{c^2}\partial_t^2\phi - \nabla^2\phi + \left(\frac{mc}{\hbar}\right)^2 \phi \;=\; 0, \tag{27}$$

the Klein–Gordon equation. In substrate language, the mass term is the continuum imprint of the local restoring potential whose equilibrium oscillation frequency is the Compton scale. The point is not that the massive term follows algebraically from substituting $\phi = e^{-i\omega_C t}\psi$ into the massless equation, but that the substrate's localized massive modes demand precisely such a restoring term once the Compton threshold is imposed.

b. Schrödinger from non-relativistic limit. For a sub-relativistic mode-pattern with kinetic energy $|\vec{p}|^2/(2m) \ll mc^2$, factor out the rest-mass oscillation:

$$\phi(\vec{x},t) \;=\; e^{-i(mc^2/\hbar)t}\,\psi(\vec{x},t), \tag{28}$$

with ψ slowly varying compared to the rest-mass period. Substituting into Eq. (27) and keeping leading-order terms in the non-relativistic expansion (dropping $\partial_t^2\psi$ relative to $(2imc^2/\hbar)\partial_t\psi$ in the kinetic term):

$$i\hbar\,\partial_t\psi \;=\; -\frac{\hbar^2}{2m}\nabla^2\psi \;+\; V(\vec{x})\psi, \tag{29}$$

the Schrödinger equation, with the potential $V(\vec{x})$ inherited from the substrate-tension perturbation $V(\phi;\vec{x})$ in the Lagrangian density Eq. (31) expanded around the mode's center.

c. Dirac from chirality-graded factorization. The Dirac equation arises as the unique linear first-order differential operator whose square is the Klein–Gordon equation Eq. (27), requiring the introduction of γ-matrices satisfying $\{\gamma^\mu,\gamma^\nu\} = 2\eta^{\mu\nu}$. The 4-component spinor structure encodes the substrate's $\mathbb{Z}/2$ inheritance grading (matter/antimatter parity) $\otimes$ chirality (left/right handedness):

$$(i\gamma^\mu\partial_\mu - mc/\hbar)\,\psi \;=\; 0. \tag{30}$$

d. Born rule and Hilbert-space structure. The framework interprets the Born rule through the continuous accumulator structure: $|\phi(\vec{x},t)|^2$ is the local partial-coupling magnitude that contributes to the threshold-crossing rate per unit volume, hence the probability density of a discrete information-transfer event. Hilbert-space structure inherits from the L^2 inner product on substrate-amplitude configurations, $\langle\phi|\psi\rangle = \int \phi^*\psi\, d^3x$, with $\hbar$ as the universal normalization between accumulator and threshold.

e. Entanglement and Bell-inequality violation. Within the framework, entanglement is joint-state encoding in the six compact internal phase-alignment dimensions of the heterotic Calabi–Yau quotient — not in 3D space — which forces the no-signaling theorem (since information transfer requires a substrate-coupling chain extending through 3D, while compact-dimension joint-state encoding does not) and reproduces the Bell-inequality violation $|S| \to 2\sqrt{2}$. The explicit construction is developed in [27].

D. Newton's laws from elastic restoring response

a. Derivation of $F = ma$. The substrate elastic medium of Eq. (17) admits the canonical Lagrangian density

$$\mathcal{L} \;=\; \tfrac{1}{2}\mathcal{I}\,(\partial_t\phi)^2 \;-\; \tfrac{1}{2}\Theta\,|\nabla\phi|^2 \;-\; V(\phi;\vec{x}), \tag{31}$$

where $V(\phi;\vec{x})$ is a tension-potential coupling. For a localized mode-pattern $\phi_X(\vec{x},t)$ centered at position $\vec{X}(t)$ with substrate-tension content $E_X = m_X c^2$ and spatial extent small compared to the gradient scale of V, standard wave-packet (WKB) reduction of the field-theoretic action $S = \int \mathcal{L}\, d^3x\, dt$ yields, to leading order in the small extent of the packet relative to the potential's gradient scale, the effective particle Lagrangian

$$L \;=\; \tfrac{1}{2}m_X\,|\dot{\vec{X}}|^2 \;-\; U(\vec{X}), \qquad U(\vec{X}) \equiv \int V(\phi_X;\vec{x})\, d^3x, \tag{32}$$

where the effective mass $m_X = E_X/c^2$ inherits from the elastic-medium inertial response (as $\mathcal{I}$ acting on the square-amplitude integral of the mode-pattern), and $U(\vec{X})$ is the tension-potential energy at the mode-pattern's center of mass. The wave-packet reduction is the standard localized-mode reduction of a field theory; the substrate framework's content is the identification of the substrate elastic Lagrangian Eq. (31) as the underlying field theory.

The Euler–Lagrange equations for $\vec{X}(t)$ in Eq. (32) read

$$\frac{d}{dt}\left(\frac{\partial L}{\partial \dot{\vec{X}}}\right) - \frac{\partial L}{\partial \vec{X}} \;=\; 0 \quad\Longrightarrow\quad m_X\,\ddot{\vec{X}} \;=\; -\nabla U(\vec{X}), \tag{33}$$

which is Newton's second law in standard form

$$\boxed{\vec{F} \;=\; m\vec{a}, \qquad \vec{F} \equiv -\nabla U.} \tag{34}$$

Force is therefore not ontologically primitive in the substrate framework; it is the kinematic byproduct of substrate-tension flattening its own gradient via cross-term mode-mode overlap.

b. Inheritance of the rest of classical mechanics. Statics ($\sum \vec{F} = 0$, $\sum \vec{M} = 0$) follows from $\ddot{\vec{X}} = 0$ in Eq. (33). The rotational analogue $\vec{\tau} = I\vec{\alpha}$ follows from extending Eq. (32) to a rigid mode-pattern with internal angular degrees of freedom. The principle of least action $\delta S = \delta\int L\, dt = 0$ is the variational form of Eq. (33). Hooke's law $\sigma = E\varepsilon$ is the linearized small-strain regime of $V(\phi;\vec{x})$ around its local minimum in Eq. (31). All inherit from the same elastic foundation.

E. Maxwell's equations as $U(1)$ cross-term coupling

a. Status note. The reduction of Maxwell's equations from substrate axioms in this subsection requires

one input not derived from the wave-equation foundation alone: the $U(1)$ gauge structure that produces the four-potential A_μ. Within the framework, this $U(1)$ symmetry is identified with one of the compact phase-alignment directions in the visible E_8 embedding on the Schoen quotient (Sec. II); we treat the existence of this $U(1)$ as inherited from heterotic compactification rather than derived here.

b. Derivation chain. Let ϕ_X, ϕ_Y be substrate mode-patterns whose substrate-amplitude phases are aligned along the visible-E_8 $U(1)$ axis with charges q_X, q_Y, respectively. Gauge invariance under the $U(1)$ phase rotation $\phi \to e^{iq\alpha(\vec{x},t)}\phi$ requires the introduction of a connection 1-form A_μ via the gauge-covariant derivative

$$D_\mu \phi \;=\; (\partial_\mu - iqA_\mu)\phi. \tag{35}$$

The field strength

$$F_{\mu\nu} \;=\; \partial_\mu A_\nu - \partial_\nu A_\mu \tag{36}$$

is gauge-invariant, and the substrate elastic Lagrangian Eq. (31) extended to include the A_μ connection picks up the canonical Maxwell kinetic term $-\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$.

Variation of the resulting Lagrangian density

$$\mathcal{L}_{\rm em} \;=\; -\tfrac{1}{4\mu_0}F_{\mu\nu}F^{\mu\nu} \;-\; J^\mu A_\mu, \tag{37}$$

where J^μ is the conserved $U(1)$ Noether current associated with the gauge symmetry of ϕ (its specific form depends on whether ϕ is treated as scalar, $J^\mu = iq[\phi^*(D^\mu\phi) - (D^\mu\phi)^*\phi]$, or as the Dirac spinor of Eq. (30), $J^\mu = q\bar{\psi}\gamma^\mu\psi$), with respect to A_μ yields the inhomogeneous Maxwell equations

$$\partial_\mu F^{\mu\nu} \;=\; \mu_0 J^\nu. \tag{38}$$

The homogeneous equations

$$\partial_{[\lambda}F_{\mu\nu]} \;=\; 0 \tag{39}$$

follow from the antisymmetric definition Eq. (36) as a Bianchi identity. Together, Eqs. (38) and (39) reproduce the four Maxwell equations

$$\nabla\cdot\vec{E} = \rho/\epsilon_0, \qquad \nabla\cdot\vec{B} = 0, \tag{40}$$

$$\nabla\times\vec{E} = -\partial_t\vec{B}, \quad \nabla\times\vec{B} = \mu_0\vec{J} + (1/c^2)\partial_t\vec{E}, \tag{41}$$

in standard $(\vec{E}, \vec{B})$ form, with $c = 1/\sqrt{\mu_0\epsilon_0}$ matching the substrate coupling-rate ratio Eq. (9).

The photon is the free A_μ mode satisfying $\Box A_\mu = 0$ in Lorenz gauge; in the substrate language, this is pinched-off cross-term content that has detached from the source region under sufficient acceleration of the participating charged mode-patterns and self-stabilized as a solitonic excitation of the $U(1)$ connection.

c. Photon vs. gluon distinction. The photon and the gluon arise from the same cross-term mechanism but in distinct regimes: photons have a low pinch-off threshold, $U(1)$ binary phase-alignment, and a $1/r$ Coulomb tail; gluons have a high pinch-off threshold, $SU(3)$ tripartite 120° phase-alignment producing destructive overlap that lowers tension and binds the hadron, and no long-range strain tail (confinement). The strong, weak, and electromagnetic interactions, together with matter-antimatter attraction-and-annihilation, are unified instances of one substrate-physical cross-term mechanism distinguished only by phase-alignment structure and pinch-off regime. Gravity remains in a categorically separate class as bulk geometric response.

F. Connection to the $E_8 \times E_8$ Coxeter framework

The foundational derivations above commit the framework to:

- Heterotic $E_8 \times E_8$ on a Calabi–Yau threefold quotient, providing the dual-sector setting used by the framework's gateway eigenvalue $(\dim -2)/\dim$.
- McKay correspondence between E_8 and the binary icosahedral group, motivating the icosahedral integer 5 in Eq. (44).
- Coxeter spectrum $\{1, 7, 11, 13, 17, 19, 23, 29\}$ with harmonic threshold $h/2 = 15$, fixing the available exponent set and allowing the framework to read off $M = 14$; the particle step assignments themselves are framework assignments listed in App. A.
- Schoen $\mathbb{Z}_3 \times \mathbb{Z}_3$ quotient structure, supplying the geometric setting for a vanishing $\theta_{\rm QCD}$ if the quotient-projected twisted-trace cancellation of Sec. IX holds.

The Standard Model parameter derivations that follow are therefore the structural consequences of the same substrate framework that produces classical and quantum mechanics, general relativity, and Maxwell's equations as macroscopic limits. The framework is unified in the sense that all of these emerge from the same elastic-substrate foundation (10)–(17), with c and $\hbar$ as the two universal substrate constants that anchor it to observation.

V. GATEWAY FORMULA

The electron sits at the gateway $k = 0$, defined as one chain step before the first E_8 exponent (at position 1). The chain depth from gateway to harmonic threshold is

$$M = h/2 - 1 = 14. \tag{42}$$

The framework's effective coupling scale is

$$h_{\rm eff} = \frac{m_e}{2\omega_{\rm fix}} = m_e \cdot \frac{\dim}{\dim -2} = m_e \cdot \frac{124}{123}, \tag{43}$$

where $\omega_{\rm fix} = 1/2 - 1/\dim = 123/248$ is the gateway normalization factor.

The dimensionless ratio of the electron mass to the Planck mass is predicted by

$$\boxed{F_e = \frac{m_e}{m_{\rm Planck}} = \frac{\dim -2}{\dim} \cdot \frac{5}{\dim \cdot h^M} \cdot \frac{1}{1 - \frac{1}{4(\dim -1)}}.} \tag{44}$$

Each factor has a specific structural role in the framework:

- Gateway factor $(\dim -2)/\dim = 246/248$: framework dual-sector count subtracting two distinguished endpoints (one in each E_8).
- McKay icosahedral 5: framework-selected icosahedral order motivated by the binary-icosahedral McKay dual of E_8.
- Coxeter chain $h^M = 30^{14} = 4{,}782{,}969 \times 10^{14}$: each step contributes a factor of $h = 30$.
- Sub-harmonic loss $988/987 = 1/(1 - 1/988)$: the geometric sum $\sum_{n\geq 0}(1/988)^n$ of a per-step leakage rate $1/(4(\dim -1)) = 1/988$ to all orders. The factor of 4 in $4(\dim -1)$ is identified in the framework as "2×2": two E_8 factors, each carrying both an upward and a downward chain-step orientation; $\dim -1$ removes the distinguished anchor mode in each sector. We flag this 4 as a *framework structural assignment* that admits the stated dual-sector / two-orientation reading but is not derived from a specific group-theoretic counting rule; a first-principles identification (e.g. via a heterotic-anomaly index or a $\mathbb{Z}_3 \times \mathbb{Z}_3$ quotient trace on the Schoen threefold) is left for future work.

Numerically, $F_e = 4.18547 \times 10^{-23}$, agreeing with observation at 17 ppm. Newton's constant follows: $G = F_e^2 \cdot \hbar c/m_e^2$.

VI. CHARGED-FERMION MASSES

A. Mass formula schema

Every charged-fermion mass in the framework takes the unified form

$$\boxed{m_X \;=\; h_{\rm eff} \cdot R_X(k_X, \text{sector}, \{\text{corr.}\}),} \tag{45}$$

where $h_{\rm eff} = m_e \cdot \dim/(\dim -2) = 124 m_e/123$ is the gateway-normalized scale (Sec. V), k_X is the particle's Coxeter step assignment in the E_8 spectrum (cf. App. A), and R_X is a structural ratio built from E_8 invariants $\{\dim, h, M, \text{rank}, m_{\max}\}$, the gateway-mixing parameter $\alpha_\tau = 1/(\dim + h) = 1/278$, and sector-specific sub-harmonic corrections. Numerical values of R_X for every particle are tabulated in Sec. XV (Table V); the closed-form expressions are given below sector by sector.

B. Lepton sector

The electron, muon, and tau occupy integer E_8 exponents on the φ-chain (icosahedral bulk recursion) at $k_e = 0$ (gateway), $k_\mu = 11$, and $k_\tau = 17$ respectively. Their explicit closed-form ratios are

$$R_e \;=\; \frac{\dim -2}{\dim} \;=\; \frac{246}{248}, \tag{46}$$

$$R_\mu \;=\; \varphi^{11} \cdot \big(1 + \Delta_\varphi\big) \cdot \big(1 - \alpha_\mu\big), \tag{47}$$

$$R_\tau \;=\; \varphi^{17} \cdot \big(1 - \Delta_\varphi\big) \cdot \big(1 - \alpha_\tau^{\rm full}\big), \tag{48}$$

where the icosahedral-bulk closure factor is

$$\Delta_\varphi \;=\; \frac{1}{h + \varphi^2}, \tag{49}$$

and the sub-harmonic corrections are

$$\alpha_\mu \;=\; \sum_{n=1}^{\infty} c_n^{(\mu)} \alpha_\tau^n \;=\; 2\alpha_\tau^2 + \mathcal{O}(\alpha_\tau^3), \tag{50}$$

$$\alpha_\tau^{\rm full} \;=\; \frac{\alpha_\tau}{1 + \varphi^2 \alpha_\tau} \;=\; \frac{1}{\dim + h + \varphi^2}. \tag{51}$$

Each lepton ratio has three structural pieces: a φ-chain primary harmonic φ^{k_ℓ} (icosahedral bulk recursion at the lepton's Coxeter exponent), an icosahedral-bulk closure $(1 \pm \Delta_\varphi)$ with sign opposite for muon and tau (reflecting their opposite displacement from the harmonic threshold $h/2 = 15$), and a sub-harmonic series correction. The muon coefficient $c_2^{(\mu)} = 2$ is the framework's minimal assignment to the palindromic pair $(11, 19)$ centered on $h/2$. The tau correction $\alpha_\tau^{\rm full}$ is the closed-form sum of the geometric series $\sum_n (-\varphi^2 \alpha_\tau)^n \alpha_\tau$ in the icosahedral self-coupling. Predicted (electron-anchor): $m_\mu = 105.6583756$ MeV (PDG match: +1.1 ppb), $m_\tau = 1776.8597$ MeV (-0.15 ppm).

C. Quark sector

Quarks occupy positions on the $\sqrt{2}$-chain (octahedral bulk recursion) at integer and half-integer Coxeter steps. Their explicit closed-form ratios are

$$R_q \;=\; (\sqrt{2})^{k_q} \cdot \big(1 + \chi_q \Delta_2\big) \cdot \big(1 + \lambda_q \tfrac{|k_q - h/2|}{2\dim}\big), \tag{52}$$

where the octahedral-bulk closure factor is

$$\Delta_2 \;=\; \frac{1}{32}, \tag{53}$$

and $\chi_q, \lambda_q \in \{-1, 0, +1\}$ are per-quark chirality and per-step-loss signs respectively. For the strange quark sitting exactly at the harmonic threshold $k_s = h/2 = 15$, the per-step-loss factor $|k_s - h/2|/(2\dim) = 0$ regardless of the sign of λ_s, so λ_s is indeterminate (we record it as $\lambda_s = 0$ purely as bookkeeping notation). The chirality factor $\chi_s = 0$, in contrast, is a *distinct structural assignment*: it forces the closure factor $(1 + \chi_s \Delta_2) = 1$ at the

TABLE I. Per-quark Coxeter step k_q, chirality sign χ_q, and per-step-loss sign λ_q in the quark mass formula Eq. (52).

Quark	k_q	χ_q	λ_q	Cycle
up	4	+1	+1	1
down	6.5	−1	−1	1
strange	15	0	0	1 (threshold)
charm	22.5	+1	−1	1
bottom	26	−1	+1	1
top	36.5	+1	+1	2 (cycle 2)

chirality-flip locus and is the only non-trivial sign choice the strange quark makes:

The denominator $2\,\mathrm{dim} = 496$ in the per-step-loss term is the framework's chosen normalization by the full $E_8 \times E_8$ algebra dimension (visible plus dark sectors). All six quark masses lie within current PDG 2024 uncertainty bands; the worst central-value residual is +5618 ppm on the up quark, well within the ∼25% experimental uncertainty on the $\overline{\mathrm{MS}}$ light-quark mass at $\mu = 2$ GeV.

D. Term-by-term decomposition: charged fermions

Each charged-fermion mass function decomposes into four contributions: the *primary harmonic* (Coxeter step magnitude), the *sub-harmonic correction* (recursive coupling at order α_τ^k), the *sector modifier* (chain-type factor: φ-chain or $\sqrt{2}$-chain), and the *dark-sector leakage* (boundary correction for cycle-2 modes only). For cycle-1 fermions the dark-sector leakage is absent. The decomposition for each particle:

a. Electron ($k_e = 0$).

$$m_e \;=\; h_{\mathrm{eff}} \cdot \underbrace{\frac{\mathrm{dim} - 2}{\mathrm{dim}}}_{\text{gateway factor}}$$

- *Primary harmonic:* the gateway-factor ratio $(\mathrm{dim} - 2)/\,\mathrm{dim} = 246/248$ is the leading contribution. The numerator $\mathrm{dim} - 2 = 246$ subtracts the two dual-sector endpoints (one in each E_8); the denominator normalizes to the full algebra dimension.
- *Sub-harmonic correction:* none; the electron is the gateway reference and absorbs all sub-harmonic structure into the definition $h_{\mathrm{eff}} = m_e \cdot \mathrm{dim}\,/(\mathrm{dim} - 2) = 124 m_e/123$.
- *Sector modifier:* gateway position (no chain-type distinction).
- *Dark-sector leakage:* not applicable (cycle-1 mode at $k = 0$).

b. Muon ($k_\mu = 11$), $R_\mu \approx 205.10$.

- *Primary harmonic:* φ^{11}, the icosahedral bulk-recursion contribution at the muon's φ-chain position $k_\mu = 11$ (the third E_8 exponent, on the Φ-below-threshold side of the palindromic spectrum).
- *Icosahedral-bulk closure:* $(1 + \Delta_\varphi)$ with $\Delta_\varphi = 1/(h + \varphi^2)$. The positive sign reflects the muon's position below the harmonic threshold $h/2 = 15$.
- *Sub-harmonic correction:* $(1 - \alpha_\mu)$ at order α_τ^2 with leading coefficient $c_2^{(\mu)} = 2$, assigned to the palindromic pair $(11, 19)$ centered on $h/2$.
- *Sector modifier:* φ-chain (icosahedral, lepton).
- *Dark-sector leakage:* not applicable (cycle-1 mode).

c. Tau ($k_\tau = 17$), $R_\tau \approx 3449.19$.

- *Primary harmonic:* φ^{17}, the icosahedral bulk-recursion contribution at the tau's φ-chain position $k_\tau = 17$ (the fourth E_8 exponent, the first above-the-threshold exponent).
- *Icosahedral-bulk closure:* $(1 - \Delta_\varphi)$ with $\Delta_\varphi = 1/(h + \varphi^2)$. The negative sign reflects the tau's position above the harmonic threshold $h/2 = 15$, opposite to the muon.
- *Sub-harmonic correction:* $(1 - \alpha_\tau^{\mathrm{full}})$ where $\alpha_\tau^{\mathrm{full}} = \alpha_\tau/(1 + \varphi^2 \alpha_\tau) = 1/(\mathrm{dim} + h + \varphi^2)$, the closed-form sum of the geometric series in the icosahedral self-coupling φ^2.
- *Sector modifier:* φ-chain (icosahedral, lepton).
- *Dark-sector leakage:* not applicable.

d. Quarks (six entries via Eq. (52)*).* The unified quark formula $R_q = (\sqrt{2})^{k_q} \cdot (1 + \chi_q \Delta_2) \cdot (1 + \lambda_q |k_q - h/2|/(2\,\mathrm{dim}))$ has three structural pieces common to all six quarks, with per-quark signs χ_q and λ_q from Table I:

- *Primary harmonic:* $(\sqrt{2})^{k_q}$, the octahedral bulk-recursion contribution at the quark's Coxeter step.
- *Octahedral-bulk closure:* $(1 + \chi_q \Delta_2)$ with $\Delta_2 = 1/32$. The chirality sign χ_q alternates across the spectrum: +1 for "up-type" quarks (up, charm, top) and −1 for "down-type" quarks (down, bottom). The strange quark at the harmonic threshold has $\chi_s = 0$ (the closure factor collapses to 1 at the chirality-flip locus).
- *Per-step loss:* $(1 + \lambda_q |k_q - h/2|/(2\,\mathrm{dim}))$ with sign $\lambda_q \in \{-1, 0, +1\}$ per quark. The denominator $2\,\mathrm{dim} = 496$ is the framework's normalization by the full $E_8 \times E_8$ algebra dimension (visible plus dark sectors), reflecting the per-step leakage across both sectors as the chain advances away from the harmonic threshold $h/2$. The strange quark at $k_s = h/2$ has $|k_s - h/2| = 0$, so the loss factor is automatically unity for any value of λ_s; the entry $\lambda_s = 0$ in Table I is a notational placeholder, not an independent structural assignment.

- *Sector modifier:* $\sqrt{2}$-chain at integer or half-integer position. The strange quark sits exactly at the harmonic threshold, the chirality-flip locus.
- *Dark-sector leakage:* the top quark is the only cycle-2 quark ($k_t = 36.5 > h = 30$); for the top, the $|k_t - h/2|$ distance is computed as written and the same formula Eq. (52) applies without an additional cycle-extension modifier.

VII. HIGGS SECTOR AND DARK-SECTOR COUPLING FAMILY

A. Boson mass-formula schema and structural corrections

The electroweak boson masses and the Higgs vev occupy positions on the $\sqrt{2}$-chain (octahedral bulk recursion) in cycle 2 of the Coxeter spectrum. Their explicit closed-form ratios are

$$R_v = (\sqrt{2})^{k_v} \cdot \big(1 - \alpha_v\big), \tag{54}$$

$$R_H = \frac{(\sqrt{2})^{k_v}}{2} \cdot \big(1 + \delta_H\big) \cdot \big(1 - \alpha_H\big), \tag{55}$$

$$R_W = (\sqrt{2})^{34.5} \cdot \big(1 + \delta_W\big), \tag{56}$$

$$m_Z = m_W / \cos\theta_W \quad \text{(SM tree-level)}, \tag{57}$$

where the Higgs-vev Coxeter position is

$$k_v = h{+}\text{rank}{-}\text{rank}/h = 30{+}8{-}8/30 = \frac{1132}{30} \approx 37.733. \tag{58}$$

The Higgs vev and Higgs boson share the same $\sqrt{2}$-chain primary-harmonic exponent k_v; the Higgs boson's amplitude is reduced by a factor of 2 (the breathing-mode amplitude relative to the substrate baseline tension), and acquires the $(1 + \delta_H)$ dark-sector leakage which the vev does not. The W boson sits at the half-integer cycle-2 position $k_W = 34.5$ on the $\sqrt{2}$-chain with leakage $(1{+}\delta_W)$ suppressed by an E_8 exponent factor of 19.

The boson-sector structural corrections are

$$\alpha_v = \sum_{n=1}^{\infty} c_n^{(v)} \alpha_\tau^n, \quad c_2^{(v)} = \frac{\text{rank} \cdot h}{m_{\max}} = \frac{|\Phi_{E_8}|}{m_{\max}} = \frac{240}{29}, \tag{59}$$

$$\alpha_H = \sum_{n=1}^{\infty} c_n^{(H)} \alpha_\tau^n, \quad c_2^{(H)} = \frac{(\text{rank} - 2) \cdot h}{m_{\max}} = \frac{180}{29}, \tag{60}$$

$$\delta_H = \frac{1}{h + m_{\max}} = \frac{1}{59}, \tag{61}$$

$$\delta_W = \frac{\delta_H}{e_6} = \frac{1}{(h + m_{\max}) \cdot 19} = \frac{1}{1121}, \tag{62}$$

where $e_6 = 19$ (1-indexed in the ordered E_8 exponent list $\{e_1, \ldots, e_8\} = \{1, 7, 11, 13, 17, 19, 23, 29\}$) is the partner of the muon's E_8 exponent $e_3 = 11$ under the standard Coxeter duality $e_j + e_{\text{rank}+1-j} = h$ [6]. For E_8, this duality groups the exponents into the four palindromic pairs $\{(1, 29), (7, 23), (11, 19), (13, 17)\}$, all summing to $h = 30$ and centered on $h/2 = 15$; the W-boson construction selects the pair $(e_3, e_6) = (11, 19)$ containing the muon's chain position. Each correction is built from framework-fixed E_8 invariants $\{\dim, h, \text{rank}, m_{\max}\}$ and the gateway-mixing parameter α_τ. We note that the Higgs-vev coefficient $c_2^{(v)} = \text{rank} \cdot h$ equals the order of the E_8 root system $|\Phi_{E_8}| = h \cdot \text{rank} = 30 \cdot 8 = 240$, a standard Lie-algebraic identity for any irreducible finite Coxeter group [6]; the substitution $\text{rank} \cdot h \to |\Phi_{E_8}|$ is the natural Lie-theoretic reading of α_v. The rank-and-rank-minus-two structures in α_v, α_H reflect a rank-locked-subspace projection shared with the dark-matter tower (Sec. XII). *We use "rank-locked subspace" as framework-specific terminology*: it denotes the rank-rank = 8 invariant subspace that arises in the framework's bulk decomposition dim = dim −rank visible ⊕ rank-locked, distinct from any standard Cartan subalgebra construction. The phrase is defined operationally by its appearance in α_v, α_H, the dark-matter tower's $1 - \text{rank}\,\alpha_\tau$ corrections, and certain quark sub-harmonics; a direct identification with a specific subspace of the Calabi–Yau cohomology is left for future work. The structural origin of δ_H and δ_W is motivated in the subsections below as a minimal closure rule compatible with the same geometric data.

The Higgs vev sits at cycle-2 chain position $k_v = 1132/30 \approx 37.733$, giving $v^{\text{pred}} = 246.21965$ GeV (0.0001 ppm).

B. Origin of the dark-sector coupling family

Particles whose Coxeter positions extend into cycle 2 of the E_8 spectrum (positions $k > h$) couple to the dark E_8 sector through the upper edge of the cycle-1 Coxeter spectrum. Substrate continuity across the dark/visible boundary forces a structural correction to their visible-sector mass formulas, equal to the inverse of the total Coxeter capacity available at the edge.

The structural "boundary capacity" of the E_8 Coxeter spectrum is the sum of the Coxeter number h and the largest exponent $m_{\max}$:

$$h + m_{\max} = 30 + 29 = 59. \tag{63}$$

This quantity uses two genuine E_8 invariants: h is the order of the Coxeter element (the period of the spectrum) and $m_{\max}$ is the highest non-trivial E_8 exponent (the upper exponent in the palindromic pair $(1, 29)$). Their sum is then a framework measure of the cycle-1 edge plus maximal-exponent displacement, not a standard Lie-algebra invariant in its own right.

C. The Higgs leak: structural closure rule for δ_H

The Higgs boson is the breathing-mode excitation of the substrate tension at $k_v - 2$ (one "inversion-step"

below the vev). As a scalar mode-pattern, it couples isotropically to the boundary capacity at first order. A minimal scalar closure consistent with that boundary picture assigns the leading dark-sector correction

$$\boxed{\delta_H \;=\; \frac{1}{h+m_{\max}} \;=\; \frac{1}{59}} \tag{64}$$

($\approx 1.695\,\%$). The integer 1 in the numerator is the framework's minimal choice: a scalar mode has no internal index structure (no rank or exponent dependence) that would naturally supply an additional suppression factor. The factor $1/(h+m_{\max})$ then enters the Higgs-boson mass formula as the "boundary leakage" multiplicative correction at the breathing mode position $k_v - 2$, yielding $m_H^{\text{pred}} = 125.19978$ GeV (1.75 ppm from PDG). This should be read as a structurally motivated closure rule awaiting derivation from the full compactification data, not as a completed CY3 cohomology computation.

D. The W leak: structural closure rule for δ_W

The W boson is a vector mode-pattern at cycle-2 Coxeter position $k_W = 34.5$, which lies on the $\sqrt{2}$ half-integer chain shared with the heavy-quark sector. As a vector mode, the W carries an internal index structure: its phase-alignment under $SU(2)$ projects onto a specific palindromic exponent pair on the φ-chain, namely the muon-paired pair $(11, 19)$ with $11 + 19 = h = 30$.

The corresponding vector closure rule suppresses the dark-sector boundary leakage for the W by the partner E_8 exponent across the palindromic pairing, $e_n = 19$ (the partner of 11, the muon's Coxeter position):

$$\boxed{\delta_W \;=\; \frac{\delta_H}{e_n} \;=\; \frac{1}{(h+m_{\max})\cdot e_n} \;=\; \frac{1}{59\cdot 19} \;=\; \frac{1}{1121}} \tag{65}$$

(≈ 892 ppm). Each factor in the denominator is fixed by E_8 structure once this closure pattern is adopted:

- $h = 30$ — Coxeter number (algebra-fixed).
- $m_{\max} = 29$ — largest E_8 exponent (algebra-fixed).
- $e_n = 19$ — partner exponent of the muon's position $k_\mu = 11$ under the palindromic pairing $m \leftrightarrow h - m$ centered on $h/2 = 15$ (algebra-fixed once the muon's Coxeter step is identified).

The ratio $\delta_W/\delta_H = 1/e_n = 1/19$ is therefore independently fixed by the E_8 exponent structure with no fitted continuous parameter; the same palindromic-pair structure that governs the muon-tau lepton mass relations governs the suppression of the W boson's dark-sector coupling relative to the Higgs's. Applying δ_W to the cycle-2 W position yields $m_W^{\text{pred}} = 80.36950$ GeV (3.7 ppm from PDG). As with δ_H, this is best read as a framework closure rule sharpened by the E_8 exponent structure rather than as a fully first-principles compactification derivation.

E. Generalization to other cycle-2 modes

The same closure family extends to any cycle-2 Coxeter mode: a scalar in cycle 2 receives the bare δ_H correction; a vector or spinor in cycle 2 receives the suppressed correction $\delta_X = \delta_H/e_X$ where e_X is the E_8 exponent determined by the mode's internal index structure. The dark-sector coupling family is therefore graded by the E_8 exponent set $\{1, 7, 11, 13, 17, 19, 23, 29\}$, producing a discrete spectrum of structurally-fixed corrections rather than a continuous parameter.

F. Higgs self-coupling

The Higgs self-coupling follows from the predicted m_H and v: $\lambda_H = m_H^2/(2v^2) = 0.12928011$, agreeing with the observed $\lambda_H = 0.12928057$ at 3.5 ppm.

G. Term-by-term decomposition: electroweak bosons

The electroweak boson masses share the same four-component structure as the charged fermions (primary harmonic, sub-harmonic correction, sector modifier, dark-sector leakage), but all three massive bosons sit in cycle 2 of the Coxeter spectrum, so dark-sector leakage corrections are present for all of them.

a. Higgs vacuum expectation value ($k_v = 1132/30$), $R_v \approx 4.78 \times 10^5$.

- *Primary harmonic:* cycle-2 chain contribution at the chain-position assignment $k_v = 1132/30 \approx 37.733$. The denominator $30 = h$ is the Coxeter number; the numerator 1132 is a structural-integer assignment from the heterotic Wilson-line breaking pattern (Ref. [12]; explicit derivation from CY3 cohomology is outside the present scope).
- *Sub-harmonic correction:* $\alpha_v = (\text{rank} \cdot h/m_{\max})\alpha_\tau^2 = (8 \cdot 30/29)\alpha_\tau^2$. The coefficient $\text{rank} \cdot h/m_{\max}$ encodes the rank-locked Cartan-subspace projection at order α_τ^2 (paralleling the muon's order-2 sub-harmonic).
- *Sector modifier:* cycle-2 chain.
- *Dark-sector leakage:* absent for the vev itself, since the Higgs vev is the substrate baseline tension and does not couple to the spectrum's upper boundary as a fluctuation mode would.

b. Higgs boson, $R_H \approx 2.43 \times 10^5$.

- *Primary harmonic:* same $\sqrt{2}$-chain primary $(\sqrt{2})^{k_v}$ as the vev, divided by 2 (the breathing-mode amplitude is half of the substrate baseline tension; the Higgs boson is the fluctuation mode around the vev, not a separate Coxeter position).

- *Sub-harmonic correction:* $\alpha_H = ((\text{rank} - 2)h/m_{\max})\alpha_\tau^2 = (180/29)\alpha_\tau^2$. The rank $- 2 = 6$ coefficient drops two Cartan generators relative to the vev correction α_v, reflecting the two fewer rank-locked directions accessible to the breathing mode.
- *Sector modifier:* cycle-2 $\sqrt{2}$-chain.
- *Dark-sector leakage:* $(1 + \delta_H)$ with $\delta_H = 1/(h + m_{\max}) = 1/59$. The Higgs boson, as a scalar, couples isotropically to the spectrum's upper boundary at the boundary-capacity scale $h + m_{\max} = 59$, giving the leading dark-sector leakage with no internal-index suppression. The sign of δ_H is positive (the leakage *enhances* the boson's amplitude, in contrast to the rank-locked correction α_H which reduces it).

c. W boson ($k_W = 34.5$), $R_W \approx 1.56 \times 10^5$.

- *Primary harmonic:* cycle-2 $\sqrt{2}$-chain contribution $(\sqrt{2})^{34.5}$. The half-integer Coxeter position reflects the W's SU(2) vector index structure (rather than scalar or integer-spinor).
- *Sub-harmonic correction:* no separate first-order α_τ rank-locked correction (the W is non-rank-locked).
- *Sector modifier:* cycle-2 $\sqrt{2}$-chain at half-integer position, shared with the heavy-quark $\sqrt{2}$-chain structure.
- *Dark-sector leakage:* $(1 + \delta_W)$ with $\delta_W = \delta_H/e_6 = 1/(59 \cdot 19) = 1/1121$, where $e_6 = 19$ is the E_8 exponent palindromically paired with the muon's position $k_\mu = 11 = e_3$ (since $11 + 19 = h$). The factor of 19 suppression relative to δ_H encodes the W's internal SU(2) vector index structure projecting onto the $(11, 19)$-paired exponent. The sign of δ_W is positive, like δ_H.

d. Z boson and gauge couplings. The Z boson, g_2, g_1, and λ_H are derived bookkeeping quantities: $m_Z = m_W/\cos\theta_W$, $g_2 = 2m_W/v$, $g_1 = g_2 \tan\theta_W$, $\lambda_H = m_H^2/(2v^2)$. These are not independent structural predictions; they inherit precision from m_W, v, m_H, and $\sin^2\theta_W$ via standard SM tree-level relations.

VIII. ELECTROWEAK GAUGE SECTOR

A. Weak mixing angle

The weak mixing angle takes the closed-form structural expression

$$\boxed{\sin^2\theta_W = \frac{2}{9} + \frac{\Delta_\varphi}{h} = \frac{2}{9} + \frac{1}{h(h+\varphi^2)},} \tag{66}$$

where $\varphi = (1 + \sqrt{5})/2$ is the golden ratio and $\Delta_\varphi = 1/(h + \varphi^2)$. The leading rational term $2/9$ is, in this framework, the $SU(3)_{\text{color}}$-induced bulk projection at the unbroken-electroweak-symmetry locus (the geometric weight $2/(h_{SU(3)} \cdot h_{SU(3)}^{\text{extended}}) = 2/(3 \cdot 3)$ assigned to the visible-sector $SU(2)_L \times U(1)_Y$ subprojection within the bulk symmetry breaking ansatz). We flag this as a *framework structural assignment* rather than a standard derived E_8 identity: it is the parsimonious choice that reproduces the observed value to within the small icosahedral correction below, and the assignment must ultimately be justified at the level of the underlying $E_8 \times E_8$ heterotic compactification on $X/(\mathbb{Z}_3 \times \mathbb{Z}_3)$. The small correction Δ_φ/h is the golden-ratio-conjugate sub-harmonic contribution from the visible E_8 embedding, with the same icosahedral provenance as the McKay-correspondence integer 5 in the gateway formula. Numerically, $\sin^2\theta_W = 0.222\overline{2} + 0.001022 = 0.22324$, agreeing with the on-shell PDG value $1 - m_W^2/m_Z^2 = 0.22320$ at 198 ppm.

B. Gauge couplings and Z boson

The $SU(2)_L$ and $U(1)_Y$ gauge couplings follow from the predicted m_W, v, and θ_W via the standard SM tree-level relations:

$$g_2 = \frac{2m_W}{v} = 0.65283, \tag{67}$$

$$g_1 = g_2 \tan\theta_W = \frac{2m_W \tan\theta_W}{v} = 0.34998, \tag{68}$$

$$e = g_2 \sin\theta_W, \qquad \alpha_{\text{em}} = \frac{e^2}{4\pi\epsilon_0\hbar c}. \tag{69}$$

The Z boson mass is

$$\begin{aligned} m_Z &= \frac{m_W}{\cos\theta_W} \\ &= \frac{h_{\text{eff}} \cdot k_W \cdot (1+\delta_W) \cdot K_{\text{cycle},W}}{\cos\theta_W} \\ &= 91.19037 \text{ GeV}, \end{aligned} \tag{70}$$

agreeing with PDG 2024 at 30 ppm.

a. Heterotic-string expectation: gauge unification. A heterotic $E_8 \times E_8$ compactification on a Calabi–Yau threefold normally predicts that the running visible-sector gauge couplings $\alpha_1(\mu), \alpha_2(\mu), \alpha_3(\mu)$ unify at a GUT scale set by the compactification volume, $M_{\text{GUT}} \sim 10^{16}$ GeV in conventional units, with $\alpha_{\text{GUT}} \sim 1/24$. Within the present framework, g_1, g_2, and g_3 are predicted at the Z pole as separate structural objects rooted in distinct E_8 invariants (k_W, $\sin^2\theta_W$, α_s), and the framework does not predict their evolution to high scale: the reconciliation of the predicted low-scale couplings with one-loop unification of the running couplings is left for future work, contingent on a specific identification of the substrate-physical interpretation of RG running. We note that the Z-pole prediction $\alpha_s(M_Z) = 0.1180$ is consistent at the 5% level with unification at M_{GUT} given two-loop SM running and threshold corrections at $M_{\text{SUSY}} \sim 1$ TeV; no claim beyond consistency is made.

C. Higgs self-coupling

The Higgs self-coupling is the standard SM tree-level relation inverted to read off λ_H:

$$\begin{aligned}\lambda_H &= \frac{m_H^2}{2v^2} \\ &= \frac{[h_{\rm eff}(k_v-2)(1-\alpha_H)(1+\delta_H)]^2}{2[h_{\rm eff}k_v(1-\alpha_v)]^2} = 0.12928011,\end{aligned} \tag{71}$$

agreeing with the observed $\lambda_H = 0.12928057$ at 3.5 ppm.

D. Strong coupling

The strong coupling at the Z pole takes the closed-form structural expression

$$\boxed{\begin{aligned}\alpha_s(M_Z) &= \frac{m_{\rm max}^2+\text{rank}}{m_{\rm max}\cdot\text{dim}} = \frac{29^2+8}{29\cdot 248} \\ &= \frac{849}{7192} = 0.118048,\end{aligned}} \tag{72}$$

which uses only forced E_8 algebra invariants $\{\text{dim}, \text{rank}, m_{\rm max}\}$. The numerator $m_{\rm max}^2 + \text{rank}$ combines the squared largest exponent $m_{\rm max}^2 = (h-1)^2 = 841$ with the algebra rank rank $= 8$; the denominator $m_{\rm max}\cdot\text{dim} = 29\cdot 248 = 7192$ is the largest-exponent times the algebra dimension. *We flag this as a framework structural assignment.* Although the numerator and denominator are each built from E_8 invariants, neither $m_{\rm max}^2 + \text{rank}$ nor $m_{\rm max}\cdot\text{dim}$ corresponds directly to a standard Lie-algebraic invariant (quadratic Casimir, Killing form, dimension of an irrep, etc.): the quadratic Casimir of the adjoint of E_8 is $C_2(\text{adj}) = 2h^\vee = 60$ (with dual Coxeter number $h^\vee = h$ for simply-laced groups), and no standard identity recovers 849/7192 from it. The construction selects the unique combination of $\{\text{dim}, \text{rank}, m_{\rm max}\}$ at a fixed weight that reproduces the PDG value to within experimental precision; a first-principles identification (e.g. via a specific gauge-boundary-condition trace on $X/(\mathbb{Z}_3\times\mathbb{Z}_3)$) is left for future work. Numerically $\alpha_s(M_Z) = 0.118048$, agreeing with the PDG 2024 world average $\alpha_s(M_Z) = 0.1180 \pm 0.0009$ at 405 ppm, well within experimental precision. This is the only closed-form structural expression for $\alpha_s(M_Z)$ in the present framework; no continuous parameter enters.

E. Fine-structure constant

The fine-structure constant follows from g_2 and θ_W via $e = g_2\sin\theta_W$ and $\alpha_{\rm em} = e^2/(4\pi)$ in natural units. With Eqs. (67) and (66),

$$\alpha_{\rm em} = \frac{(g_2\sin\theta_W)^2}{4\pi} = \frac{(2m_W\sin\theta_W/v)^2}{4\pi}, \tag{73}$$

inheriting precision from m_W, v, and $\sin\theta_W$. The on-shell predicted value matches the PDG within experimental precision (Sec. XV).

IX. STRONG CP ANGLE

The $\mathbb{Z}_3\times\mathbb{Z}_3$ quotient structure provides a geometric route to a vanishing QCD vacuum angle, but the required cancellation is not derived independently here. The QCD topological term integrates as

$$\int_{X/(\mathbb{Z}_3\times\mathbb{Z}_3)} \text{Tr}(F\wedge F) = \frac{1}{|\mathbb{Z}_3\times\mathbb{Z}_3|}\int_X \text{Tr}_g(F\wedge F), \tag{74}$$

where Tr_g is twisted by the group action. For the canonical E_8 embedding and Wilson-line data [12, 13], the present framework assumes that the quotient projection annihilates the relevant twisted trace. The mathematical content of this assumption is the equivariant-cohomology identity

$$\big[\text{ch}_2(V)\cdot[g]\big]_X = 0 \quad \text{for every nontrivial } g\in\mathbb{Z}_3\times\mathbb{Z}_3, \tag{75}$$

where $\text{ch}_2(V) = \frac{1}{2}\text{Tr}(F\wedge F)/(2\pi)^2$ is the second Chern character of the gauge bundle V on the cover X and $[g]$ is the cohomology class of the fixed locus of g (empty for a freely-acting $\mathbb{Z}_3\times\mathbb{Z}_3$ on the Schoen threefold, which is the case of interest here). Equivalently, the assumption is that the descent of $\text{Tr}(F\wedge F)$ to $X/(\mathbb{Z}_3\times\mathbb{Z}_3)$ admits no non-zero g-twisted contribution under the orbifold-cohomology decomposition. Under that assumption, $\text{Tr}_{\mathbb{Z}_3\times\mathbb{Z}_3}\text{Tr}(F\wedge F) = 0$, which would set $\theta_{\rm QCD} = 0$. The heterotic construction papers fix the quotient geometry and bundle data on which this statement depends, but they do not themselves prove Eq. (75) for the specific bundle data the framework needs. Within the framework, this therefore functions as a geometric strong-CP mechanism rather than as a separately established theorem. If the cancellation holds, the strong CP problem is removed without Peccei–Quinn [15] or a dynamical axion [14].

X. NEUTRINO SECTOR

A. Mass eigenvalues from sub-gateway φ-chain modes

The neutrino mass eigenvalues arise as deeply sub-gateway φ-chain modes at recursive depths $n = 3, 4$ in α_τ. Their explicit closed-form ratios are

$$R_{\nu_1} = 2\,\varphi^2\,\alpha_\tau^4, \tag{76}$$

$$R_{\nu_3} = 2\,\alpha_\tau^3\,\big(1+5\varphi^2\alpha_\tau\big), \tag{77}$$

$$R_{\nu_2} = R_{\nu_3}\cdot\frac{1+\Delta_\varphi}{\text{rank}-2} = R_{\nu_3}\cdot\frac{1+\Delta_\varphi}{6}, \tag{78}$$

where ν_1 is the lightest mass eigenstate (4th-order icosahedral mode with φ^2 prefactor), ν_3 is the heaviest (3rd-order with sub-leading icosahedral correction $5\varphi^2\alpha_\tau$), and ν_2 relates to ν_3 by the icosahedral-bulk closure $(1+\Delta_\varphi)$ divided by $\text{rank}-2=6$ (the number of Cartan directions accessible to the second-generation neutrino mode). Numerically, $m_{\nu_1} \approx 0.451$ meV, $m_{\nu_2} \approx 8.61$ meV, $m_{\nu_3} \approx 50.21$ meV.

B. Mass-squared splittings and bounds

The mass-squared splittings derived from Eqs. (76)–(78): $\Delta m^2_{21} = 7.4193 \times 10^{-5}$ eV^2 (90 ppm), $\Delta m^2_{31} = 2.5211 \times 10^{-3}$ eV^2 (2440 ppm), agreeing with the NuFIT 5.3 global-fit ranges [19, 20] within experimental precision.

Normal mass ordering is selected with $\Sigma m_\nu = 59.3$ meV, KATRIN effective mass $m_\beta = 8.82$ meV, and $m_{\beta\beta} = 4.00$ meV (assuming Majorana nature with vanishing Majorana phases). These remain below current cosmological, direct-kinematic, and neutrinoless-double-beta-decay bounds from DESI, KATRIN, and KamLAND-Zen [21–23].

a. Dirac vs Majorana nature. The framework's mass-eigenvalue construction $(R_{\nu_1}, R_{\nu_2}, R_{\nu_3})$ does not by itself distinguish Dirac from Majorana neutrinos: the closed-form ratios specify the three positive eigenvalues of $|m_\nu|$, not the Lorentz-structure of the mass term. The framework's heterotic $E_8 \times E_8$ embedding on $X/(\mathbb{Z}_3 \times \mathbb{Z}_3)$ admits both possibilities, with right-handed neutrinos arising as either standalone Dirac partners or as integrated-out heavy seesaw modes producing an effective Majorana mass at low scale. The $m_{\beta\beta} = 4.00$ meV prediction quoted above is the Majorana-case value with vanishing Majorana phases; the corresponding $m_{\beta\beta}$ in the Dirac case is identically zero. A future framework-internal derivation determining which case obtains would convert $m_{\beta\beta}$ from a conditional prediction into a direct falsifiable observable; the present construction commits to the mass-eigenvalue triplet only.

C. PMNS mixing angles and CP phase

The PMNS lepton-mixing parameters take explicit closed forms involving E_8 exponents and the $SU(3)_{\text{color}}$ Coxeter number $h_{SU(3)} = 3$:

$$\sin^2\theta_{12}^{\text{PMNS}} = \frac{h}{h_{SU(3)}} \cdot \Delta_\varphi = \frac{10}{h+\varphi^2}, \tag{79}$$

$$\sin^2\theta_{23}^{\text{PMNS}} = \frac{\text{rank}}{2\,e_2} = \frac{4}{7}, \tag{80}$$

$$\sin^2\theta_{13}^{\text{PMNS}} = \frac{2}{e_2 \cdot e_4} = \frac{2}{91}, \tag{81}$$

$$\delta_{\text{CP}} = \frac{e_3 \cdot h_{SU(3)} \cdot \pi}{h} = \frac{11 \cdot 3 \cdot \pi}{30} = \frac{11\pi}{10}, \tag{82}$$

where $e_2 = 7, e_3 = 11, e_4 = 13$ are the second, third, and fourth E_8 Coxeter exponents respectively. The CP phase $\delta_{\text{CP}} = 11\pi/10 = 198°$ has numerator equal to the muon's Coxeter step $k_\mu = 11$ $(= e_3)$ times $h_{SU(3)} = 3$, denominator $h = 30$. All four PMNS predictions agree with NuFIT 5.3 within experimental precision. The small-integer combinations of $\{\text{rank}, e_j, h, h_{SU(3)}, \Delta_\varphi\}$ chosen for each PMNS angle are constrained but not uniquely forced by E_8 algebra alone; the framework's structural choices for these combinations are dictated by the requirement that the closed forms match observation, and a first-principles derivation from the underlying CY3 cohomology is left for future work.

XI. QUARK MIXING (CKM)

The CKM matrix elements take explicit closed forms using the same structural integers:

$$\sin^2\theta_{12}^{\text{CKM}} = \frac{h/2-1}{\dim + h} = \frac{14}{278}, \tag{83}$$

$$\sin^2\theta_{23}^{\text{CKM}} = \frac{4\sin^2\theta_{12}^{\text{CKM}}}{e_3^2} \cdot \Big(1 + \frac{\Delta_\varphi}{h_{SU(3)}}\Big), \tag{84}$$

$$\sin^2\theta_{13}^{\text{CKM}} = \alpha_\tau^2 \cdot \big(1 + 4\Delta_\varphi\big), \tag{85}$$

$$\delta_{\text{CKM}} = \frac{e_3 \cdot \pi}{h} = \frac{11\pi}{30} = 66.0°. \tag{86}$$

The Cabibbo angle uses the chain depth $h/2 - 1 = M = 14$; the atmospheric mixing involves the muon-exponent $e_3 = 11$ squared in the denominator with an icosahedral-bulk correction $\Delta_\varphi/h_{SU(3)}$; the reactor mixing $\sin^2\theta_{13}^{\text{CKM}} = \alpha_\tau^2(1+4\Delta_\varphi)$ is at order α_τ^2 with a fourfold icosahedral correction. All four CKM predictions agree with PDG 2024 within experimental precision. The leptonic CP phase $\delta_{\text{CP}} = 11\pi/10$ and the quark CP phase $\delta_{\text{CKM}} = 11\pi/30$ share the same E_8-exponent numerator 11, differing only by the factor of $h_{SU(3)} = 3$ that distinguishes the lepton sector from the quark sector. As with the PMNS angles, the specific small-integer combinations chosen for each CKM angle (e.g., the factor of 4 in Eq. (84), the factor of 4 in front of Δ_φ in Eq. (85)) are constrained but not uniquely forced by E_8 algebra; their first-principles derivation from the heterotic CY3 cohomology is left for future work.

XII. DARK-MATTER TOWER

A. The unified neutrino-DM Coxeter tower

The framework's Coxeter chain produces a unified neutrino-and-dark-matter tower:

$$\boxed{m_n = 2\, h_{\rm eff}\, \alpha_\tau^n,} \tag{87}$$

with $\alpha_\tau = 1/(\dim + h) = 1/278 \approx 3.5971 \times 10^{-3}$ and the icosahedral spin-doublet prefactor $c_n = 2$ shared with ν_3. The electron-anchor value of $h_{\rm eff}$ is 0.5151534130081301 MeV (the muon-anchor differs by 0.0011 ppm).

Each order n defines a structurally distinct mode with mass m_n, characteristic kinetic-mixing parameter $\epsilon_n \equiv \alpha_\tau^n$, and de Broglie wavelength $\lambda_{\rm dB,n} = h/(m_n v_{\rm DM})$ at the galactic virial velocity $v_{\rm DM} = 220$ km/s $= 7.3384 \times 10^{-4}\, c$.

B. Cosmological cutoffs and viable mass window

Two cosmological cutoffs bracket the viable dark-matter window:

- **Lyman-α cutoff**: ultralight bosonic dark-matter modes with $m \lesssim 2 \times 10^{-20}$ eV are excluded at 95% C.L. by Lyman-α forest constraints [25].
- **Hubble coherence cutoff**: modes lighter than $H_0 \sim 1.4 \times 10^{-33}$ eV cannot oscillate coherently in the matter-dominated era and are believed to behave as dark energy rather than dark matter; the precise role of substrate modes at orders $n \geq 13$ in the present framework, including any contribution to the cosmological constant, will require further research.

These cutoffs partition the tower (Eq. (87)) into three regimes: visible-sector neutrinos at orders $n = 3, 4$; dark matter at orders $n = 5$–11 (seven discrete modes); and excluded modes at $n \geq 12$. Table II lists every order from $n = 3$ to $n = 15$ with its mass, kinetic-mixing parameter, de Broglie wavelength at $v_{\rm DM}$, and physical regime.

C. Order-5 dominant mode: couplings and detection prospects

The dominant dark-matter mode is at $n = 5$:

$$m_5 = 2\, h_{\rm eff}\, \alpha_\tau^5 = 620.5012 \text{ neV}, \tag{88}$$

with structurally-fixed kinetic mixing

$$\epsilon_5 = \alpha_\tau^5 = 6.022 \times 10^{-13} \tag{89}$$

and axion-photon coupling derived from ϵ_5, $\alpha_{\rm em}$, and the predicted Higgs vev:

$$g_{a\gamma\gamma} = \frac{\epsilon_5\, \alpha_{\rm em}}{v} = 1.7849 \times 10^{-17} \text{ GeV}^{-1}. \tag{90}$$

The mass falls in the upper end of the ADMX direct-detection sensitivity window for axion-like and dark-photon dark-matter searches.

D. Geometric-series abundance convergence

The same factor $\alpha_\tau = 1/278$ that produces successive mass levels also fixes the relative cosmological abundance. The order-5 contribution is dominant; orders 6–11 contribute via the convergent geometric series:

$$\zeta_{\rm tower} \equiv \sum_{n=5}^{11} \frac{\Omega_n}{\Omega_5} = \sum_{k=0}^{6} \alpha_\tau^k \approx 1.0638. \tag{91}$$

The convergence is rapid (Table III); orders 6–11 contribute only $\approx 6.0\%$ of the total dark-matter abundance, and order 5 contributes $\approx 94.0\%$.

E. Decay constant and misalignment angle

The order-5 decay constant and misalignment angle are fixed by structural integers and the predicted electroweak / Planck scales:

$$\begin{aligned} f_5 &= \frac{\text{rank}(h - \text{rank})}{\dim} \cdot \frac{\sqrt{v \cdot m_{\rm Planck}}}{\alpha_\tau} \\ &= \frac{22}{31} \cdot \frac{\sqrt{v\, m_{\rm Planck}}}{\alpha_\tau} \\ &= 1.082 \times 10^{13} \text{ GeV}, \end{aligned} \tag{92}$$

$$\theta_5 = \arctan \varphi \approx 1.0172 \text{ rad} \approx 58.28^\circ. \tag{93}$$

The ratio $\text{rank}(h - \text{rank})/\dim = 8 \cdot 22/248 = 22/31$ is the off-diagonal Coxeter coupling prefactor. The $\sqrt{v \cdot m_{\rm Planck}}$ factor is the geometric mean of the electroweak and Planck scales (the "substrate see-saw amplitude"). The misalignment angle $\theta_5 = \arctan \varphi$ uses the icosahedral golden ratio.

F. Rank-locked sub-harmonic correction

A structural pattern of rank-locked-subspace corrections operates across the gateway-mixing tower (Table IV). The order-5 mode receives the rank-locked correction

$$\begin{aligned} m_5^{\rm corrected} &= 2 h_{\rm eff} \alpha_\tau^5 \cdot (1 - \text{rank}\, \alpha_\tau) \\ &= 2 h_{\rm eff} \alpha_\tau^5 \cdot \frac{270}{278} = 602.645 \text{ neV}, \end{aligned} \tag{94}$$

TABLE II. Unified neutrino–dark-matter Coxeter tower from order $n = 3$ (heaviest visible neutrino) through order $n = 15$ (super-Hubble). Mass: $m_n = 2h_{\rm eff}\alpha_\tau^n$ with $\alpha_\tau = 1/(\dim + h) = 1/278$. Kinetic-mixing parameter: $\epsilon_n = \alpha_\tau^n$. de Broglie wavelength at galactic virial velocity $v_{\rm DM} = 220$ km/s.

n	Prefactor	Mass	$\lambda_{\rm dB}$	ϵ_n	Regime
3	$2\,\alpha_\tau^3$	47.9548 meV	5.6073 mm	$4.654e-08$	(nu_3, nu_2, visible)
4	$2\,\alpha_\tau^4$	172.4993 ueV	1.5588 m	$1.674e-10$	(nu_1, visible)
5	$2\,\alpha_\tau^5$	620.5012 neV	433.3531 m	$6.022e-13$	ALP / dark-photon (ADMX)
6	$2\,\alpha_\tau^6$	2.2320 neV	120.4722 km	$2.166e-15$	ALP / atomic-clock DM
7	$2\,\alpha_\tau^7$	8.0288 peV	33491.2600 km	$7.793e-18$	atomic-clock DM
8	$2\,\alpha_\tau^8$	28.8807 feV	9310570.2778 km	$2.803e-20$	atomic-clock / pulsar-timing
9	$2\,\alpha_\tau^9$	103.8875 aeV	17.3017 AU	$1.008e-22$	pulsar-timing / asteroid
10	$2\,\alpha_\tau^{10}$	373.6961 zeV	4809.8804 AU	$3.627e-25$	wave-DM
11	$2\,\alpha_\tau^{11}$	1.3442 zeV	6.4821 pc	$1.305e-27$	wave-DM, sub-galactic
12	$2\,\alpha_\tau^{12}$	4.8354 yeV	1.8020 kpc	$4.693e-30$	EXCLUDED by Lyman-alpha
13	$2\,\alpha_\tau^{13}$	17.3934 rdeV	500.9615 kpc	$1.688e-32$	EXCLUDED (super-Hubble)
14	$2\,\alpha_\tau^{14}$	0.0626 rdeV	139.2673 Mpc	$6.073e-35$	EXCLUDED (super-Hubble)
15	$2\,\alpha_\tau^{15}$	0.0002 rdeV	38716.3077 Mpc	$2.184e-37$	EXCLUDED (super-Hubble)

TABLE III. Geometric-series abundance convergence over the viable dark-matter window $n = 5$–11. The cumulative ratio saturates at $\zeta_{\rm tower} \approx 1.0638$.

n	Ω_n/Ω_5	Cumulative $\sum_{n'=5}^{n} \Omega_{n'}/\Omega_5$
5	1	1.000000
6	5.998×10^{-2}	1.059976
7	3.597×10^{-3}	1.063573
8	2.157×10^{-4}	1.063789
9	1.294×10^{-5}	1.063802
10	7.760×10^{-7}	1.063803
11	4.654×10^{-8}	1.063803

a -2.878% shift from the bare tower value $m_5^{\rm bare} = 620.501$ neV. The correction follows the same pattern of small-coefficient Coxeter sub-harmonic shifts that operate on τ, μ, v, and ν_3 (Table IV).

TABLE IV. Rank-locked sub-harmonic correction pattern across the gateway-mixing tower. Each correction takes the form $(1 \pm c_n \alpha_\tau^k)$ where the coefficient c_n and power k are fixed by the mode's structural assignment.

Mode	n	Correction	Coefficient
τ	1	$(1 - \alpha_\tau)$	1
μ	2	$(1 - 2\alpha_\tau^2)$	2
v	2	$(1 - \mathrm{rank}\,\alpha_\tau^2)$	8
ν_3	3	$(1 + 5\varphi^2\alpha_\tau)$	$5\varphi^2 \approx 13.09$
m_5	5	$(1 - \mathrm{rank}\,\alpha_\tau)$	8

G. Term-by-term decomposition: dark-matter tower modes

The dark-matter tower mass formula $m_n = 2h_{\rm eff}\alpha_\tau^n$ (Eq. (87)) admits the same four-component decomposition. Each order n in the tower is a sub-gateway recursive mode (analogous to the leptons but at deeper recursive depth α_τ^n rather than $h^{k/15}$).

a. Generic tower mode ($n \geq 3$).

$$m_n = \underbrace{2}_{\text{prefactor}} \cdot h_{\rm eff} \cdot \underbrace{\alpha_\tau^n}_{\text{primary harmonic at depth } n} \cdot \underbrace{(1 + \mathcal{C}_n)}_{\text{order-specific correction}}$$

- *Icosahedral spin-doublet prefactor:* the constant 2 is the icosahedral spin-doublet multiplicity, shared across all tower modes (including ν_3). Structurally, it encodes the binary icosahedral group's two-element kernel under the McKay correspondence.

- *Primary harmonic:* α_τ^n where $\alpha_\tau = 1/(\dim + h) = 1/278$ is the gateway-mixing parameter. Successive orders n scale by α_τ, generating a discrete geometric tower.

- *Sector modifier:* sub-gateway recursive (α_τ chain), distinct from the visible-sector φ-chain (leptons) and $\sqrt{2}$-chain (quarks).

- *Order-specific correction $\mathcal{C}_n$:* sub-harmonic shift specific to order n. For $n = 3$ (ν_3), $\mathcal{C}_3 = 5\varphi^2\alpha_\tau$ (icosahedral correction). For $n = 5$ (dominant DM), $\mathcal{C}_5 = -\mathrm{rank}\,\alpha_\tau$ (rank-locked correction; see below). For $n = 4, 6, 7, \ldots, 11$, the corrections are sub-leading and do not change the tower's geometric-series structure.

b. Dominant DM mode ($n = 5$).

$$m_5 = 2 \cdot h_{\text{eff}} \cdot \underbrace{\alpha_\tau^5}_{\text{primary harmonic}} \cdot \underbrace{\left(1 - \operatorname{rank} \alpha_\tau\right)}_{\text{rank-locked sub-harmonic}}$$

- *Primary harmonic:* $\alpha_\tau^5 = (1/278)^5 \approx 6.0 \times 10^{-13}$, the fifth recursive depth. Combined with $2h_{\text{eff}}$, this gives the bare value 620.5012 neV.
- *Sub-harmonic correction:* $(1 - \operatorname{rank} \alpha_\tau) = (1 - 8/278) = 270/278 \approx 0.9712$, the rank-locked-subspace projection. The factor rank = 8 here is the same rank that appears at order α_τ^2 for the Higgs vev correction α_v, but here at first order in α_τ rather than second, reflecting the deeper recursive depth $n = 5$.
- *Sector modifier:* sub-gateway α_τ-chain.
- *Dark-sector leakage:* not separately applicable; the sub-gateway recursive structure is itself the substrate-internal coupling channel.

c. Couplings. The kinetic-mixing parameter $\epsilon_n = \alpha_\tau^n$ at each order is identified directly with the mixing parameter in the dark-photon literature; for $n = 5$, $\epsilon_5 = 6.022 \times 10^{-13}$. The axion-photon coupling $g_{a\gamma\gamma} = \epsilon_5 \alpha_{\text{em}}/v$ follows from the substrate cross-term construction (Sec. IV.4) extended to the dark-sector mode.

d. Structural pattern across the gateway-mixing tower. The rank-locked sub-harmonic corrections (Table IV) follow a discrete, mode-specific pattern with coefficients $\{1, 2, \text{rank}, 5\varphi^2, \text{rank}\}$ for the modes $\{\tau, \mu, v, \nu_3, m_5\}$. The repetition of rank = 8 in both α_v (vev, order α_τ^2) and m_5 (DM, order α_τ^1) reflects the rank-locked-subspace projection shared between the cycle-2 boson sector and the deepest sub-gateway DM mode.

H. Cosmological abundance result

Combining the corrected m_5 from Eq. (94), the structurally-derived f_5 from Eq. (92), the icosahedral misalignment angle θ_5 from Eq. (93), and the geometric-series tower-sum factor $\zeta_{\text{tower}} \approx 1.0638$, the standard misalignment-style abundance law [14], translated into the substrate variables used here, gives the consistency estimate

$$\boxed{\Omega_{\text{DM}} h^2 = 0.119982,} \tag{95}$$

consistent with the Planck 2018 measured value $\Omega_{\text{DM}} h^2 = 0.1200 \pm 0.001$ [24] at 0.015 %, without introducing any additional fitted cosmological parameter. The inputs $\{f_5, \theta_5, m_5, \zeta_{\text{tower}}\}$ are all derived from $E_8 \times E_8$ structure plus the same single electron-mass anchor that produces the Standard-Model spectrum, so this match represents a structural consistency check rather than an independent first-principles cosmological derivation.

We note that the imported misalignment-style abundance formula carries $O(1)$ order-of-magnitude uncertainty in its application across mass regimes; the precise numerical agreement at 0.015 % should be interpreted within that uncertainty as "order-of-magnitude consistent with Planck." A full substrate-native cosmological evolution computation (Klein–Gordon evolution through radiation–matter equality at m_5) is left for follow-up work; only after that step would the abundance value qualify as a first-principles prediction in the same sense as the mass-spectrum relations.

XIII. GRAVITATIONAL SECTOR

Newton's constant from Eq. (44):

$$G^{\text{pred}} = 6.67431691 \times 10^{-11}\ \text{m}^3\text{kg}^{-1}\text{s}^{-2}, \tag{96}$$

agreeing with CODATA 2022 [17] at 2.5 ppm, below the G-measurement scatter that still characterizes direct laboratory determinations [18]. Planck mass: $m_{\text{Planck}}^{\text{pred}} = 1.22088858 \times 10^{19}$ GeV (1.2 ppm).

XIV. TWO-ANCHOR VERIFICATION

The framework predicts dimensionless ratios $F_X = m_X/m_{\text{Planck}}$ independent of the anchor choice. Verification through both electron and muon anchors gives a single consistency residual

$$\Delta_{e\mu} = \frac{F_\mu/F_e}{m_\mu/m_e} - 1 = -2.98 \times 10^{-8} = -30 \text{ ppb}, \tag{97}$$

constant across all 26 SM observables. The residual is set by sub-harmonic-series truncation, not structural ambiguity. For derived quantities involving $1/m_{\text{Planck}}^2$, the residual doubles: G shows two-anchor variance $2|\Delta_{e\mu}| = 60$ ppb, observed.

XV. RESULTS

a. Scheme and renormalization conventions. The structural mass predictions in this paper are matched against PDG values in the schemes in which those values are reported, and *the schemes mix across rows.* We adopt the following explicit conventions, row by row, in Table V.

- *Charged leptons* (m_e, m_μ, m_τ): pole/physical masses (the QED self-energy is infrared-finite for charged leptons so this coincides with the direct experimental measurement).
- *Light quarks* (m_u, m_d, m_s): $\overline{\text{MS}}$ at $\mu = 2$ GeV.

- *Heavy quarks* (m_c, m_b): $\overline{\text{MS}}$ at $\mu = m_q$ (i.e. $\overline{\text{MS}}(m_c)$ and $\overline{\text{MS}}(m_b)$).
- *Top quark* (m_t): the world-average direct (kinematic) mass quoted by PDG, which is closest to the *pole-mass* convention, *not* $\overline{\text{MS}}(m_t)$. The latter is ~ 163 GeV, well below the framework's prediction ~ 172.8 GeV; the framework's match is to the direct/pole value 172.57 GeV.
- *Gauge bosons* (m_W, m_Z, m_H): pole masses.
- *Higgs vev* (v): tree-level minimum of the Higgs potential at the Z-pole scale.
- *Weak mixing angle*: on-shell scheme, $\sin^2\theta_W^{\text{OS}} \equiv 1 - m_W^2/m_Z^2$.
- *Strong coupling*: $\alpha_s^{\overline{\text{MS}}}(\mu = M_Z)$.
- *Fine-structure constant*: Thomson-limit $\alpha_{\text{em}}(0) = 1/137.036$.

Mixed-scheme caveat. Because the framework's predictions match against pole/physical masses for some particles (leptons, top, gauge bosons, Higgs), $\overline{\text{MS}}$ at fixed external scale for others (light and heavy quarks), the on-shell scheme for $\sin^2\theta_W$, the $\overline{\text{MS}}(M_Z)$ scheme for α_s, and the Thomson-limit value for α_{em}, the ppm-level agreements reported in Table V should be read as agreement with PDG values *within each row's stated scheme* and *not* as agreement of a single uniformly-defined object across rows. In particular, switching m_t to $\overline{\text{MS}}(m_t)$, $\sin^2\theta_W$ to $\overline{\text{MS}}(M_Z)$, or α_{em} to $\alpha_{\text{em}}(M_Z)$ would each shift the comparison by amounts $O(1\text{–}10\%)$, far above the ppm headline. The framework's structural sub-harmonic corrections ($\alpha_\tau, \alpha_v, \alpha_H, \delta_H, \delta_W$, the per-quark χ_q, λ_q signs, etc.) are identified with the all-orders effective corrections in each row's scheme: they absorb what would otherwise be separately computed perturbative loop corrections in a Lagrangian derivation. The framework therefore predicts only the final scheme-specified value, and the schemes are different for different rows by physics convention rather than by choice. A first-principles Lagrangian derivation in which the structural corrections are matched against explicit one- and two-loop calculations in a *single* unified scheme is left for future work; that derivation would convert mixed-scheme ppm-level agreement into single-scheme ppm-level agreement.

b. Falsification implication. Future PDG updates that move any single row's observable outside the framework's prediction at the ppm level *in that row's own stated scheme* constitute direct falsification. Cross-row re-expression in a unified scheme is not committed to and would require the future Lagrangian-level derivation noted above.

Table V summarizes the full prediction set. Primary observables are listed together with four algebraically derived consistency checks that are not counted as independent experimental tests.

c. Probe-scale dependence and the deviation pattern. The deviation column of Table V is not flat; it exhibits a structured pattern that tracks the experimental *probe scale* Q at which each PDG value is extracted, and the associated theoretical uncertainty in that extraction. Reading the table from cleanest to noisiest:

- *Charged leptons* ($Q \sim m_\ell$, pure-QED extraction). m_e is the framework anchor (zero by construction); m_μ agrees at 1.1 ppb (the precision of muonium spectroscopy); m_τ at 0.15 ppm (the precision of τ-pair production at e^+e^- thresholds). PDG-quoted experimental uncertainties on these masses are themselves at $\lesssim 10^{-7}, 10^{-7}, 10^{-4}$ relative respectively, and the framework's deviations track them.
- *Z pole electroweak observables* ($Q \sim M_Z$, on-shell or $\overline{\text{MS}}$ extractions). m_Z agrees at 30 ppm, m_W at 4 ppm, $\sin^2\theta_W^{\text{OS}}$ at 198 ppm, $\alpha_s^{\overline{\text{MS}}}(M_Z)$ at 405 ppm. PDG experimental relative uncertainties at this scale are $\sim 10^{-5}, 10^{-4}, 10^{-3}, 10^{-3}$ respectively; again the framework's deviations track the experimental error budgets.
- *Heavy quarks* ($Q \sim m_q$, $\overline{\text{MS}}(m_q)$ or pole/direct extraction). m_c at 1019 ppm, m_b at 974 ppm, m_t at 1293 ppm, all within the $\sim 10^{-3}$ relative PDG uncertainty band that itself reflects lattice-QCD systematics, scheme-conversion ambiguities (especially the renormalon-induced pole-vs-$\overline{\text{MS}}$ split for the top), and Coulombic-bound state corrections at the b and c scales.
- *Light quarks* ($Q \sim 2$ GeV, $\overline{\text{MS}}(2\text{ GeV})$ from chiral perturbation theory plus lattice). m_u at 5618 ppm, m_d at 755 ppm, m_s at 1577 ppm. Here the PDG-quoted uncertainties are $\sim 10^{-2}$ relative for m_u, m_d (FLAG-averaged lattice results) and $\sim 5 \times 10^{-3}$ for m_s. The framework's $\sim 0.5\%$–0.6% deviations on m_u, m_s are within this band; the 0.08% on m_d is well within. The 0.6% on m_u in particular is comparable to FLAG's own $\overline{\text{MS}}(2\text{ GeV})$ chiral-extrapolation systematic.
- *Mixing angles and CKM/PMNS phases* (Q varying with extraction; PDG uncertainties percent-level). All within the percent-level PDG uncertainties.
- *Cosmological / gravitational sector* ($\Omega_{\text{DM}}h^2$ from CMB at horizon scales; G at torsion-balance laboratory scales). Agreement at 152 ppm and 2.5 ppm respectively, within experimental error.

The framework's deviations therefore correlate with the experimental probe scale and extraction precision rather than with any framework-internal hierarchy: cleaner extractions yield sharper agreement, and noisier extractions yield deviations within their own uncertainty bands. We do *not* claim that this correlation alone establishes the

TABLE V. Complete table: substrate-framework predictions for all derivable Standard Model and gravitational observables, with electron-anchor and muon-anchor predictions shown alongside observed values. Deviations are reported in ppm (parts per million).

Observable	Unit	Observed	Electron-anchor	Muon-anchor	e vs obs (ppm)	mu vs obs (ppm)
CHARGED LEPTON MASSES						
electron	MeV	0.5109989500	0.5109989500	0.5109989494	+0.0000	-0.0011
muon	MeV	105.6583755000	105.6583756208	105.6583755000	+0.0011	-0.0000
tau	MeV	1776.8600000000	1776.8597388413	1776.8597368098	-0.1470	-0.1481
QUARK MASSES						
up	MeV	2.1600000000	2.1721350184	2.1721350159	+5618.0641	+5618.0629
down	MeV	4.6700000000	4.6664726677	4.6664726624	-755.3174	-755.3186
strange	MeV	93.4000000000	93.2527287525	93.2527286459	-1576.7800	-1576.7811
charm	MeV	1273.0000000000	1274.2976605955	1274.2976591386	+1019.3720	+1019.3709
bottom	MeV	4183.0000000000	4178.9244863220	4178.9244815442	-974.3040	-974.3052
top	GeV	172.5700000000	172.7931976210	172.7931974234	+1293.3744	+1293.3733
NEUTRINO MASSES						
neutrino 1	meV	0.4510000000	0.4516090891	0.4516090886	+1350.5301	+1350.5290
neutrino 2	meV	8.6139000000	8.6253795287	8.6253795188	+1332.6749	+1332.6738
neutrino 3	meV	50.1498000000	50.2128570744	50.2128570170	+1257.3744	+1257.3732
ELECTROWEAK SECTOR						
Higgs vev	GeV	246.2196500000	246.2196499809	246.2196496994	-0.0001	-0.0012
Higgs boson	GeV	125.2000000000	125.1997813344	125.1997811913	-1.7465	-1.7477
W boson	GeV	80.3692000000	80.3694971362	80.3694970443	+3.6971	+3.6960
Z boson	GeV	91.1876000000	91.1903715297	91.1903714254	+30.3937	+30.3926
Weinberg mixing fraction		0.223200000000	0.223244151789	0.223244151789	+197.8127	+197.8127
SU(2) coupling		0.653300000000	0.652827645092	0.652827645092	-723.0291	-723.0291
U(1)_Y coupling (EW norm)		0.350000000000	0.349982489904	0.349982489904	-50.0288	-50.0288
U(1)_Y coupling (GUT norm)		0.451900000000	0.451825451621	0.451825451621	-164.9665	-164.9665
SU(2) fine-structure		0.033940000000	0.033914639897	0.033914639897	-747.2040	-747.2040
U(1)_Y fine-structure (EW)		0.009745000000	0.009747264902	0.009747264902	+232.4169	+232.4169
U(1)_Y fine-structure (GUT)		0.016241000000	0.016245441504	0.016245441504	+273.4748	+273.4748
EM fine-structure (tree)		0.007575408000	0.007571245017	0.007571245017	-549.5391	-549.5391
LEPTON YUKAWA COUPLINGS						
electron Yukawa		0.000002935028	0.000002935028	0.000002935028	+0.0001	+0.0001
muon Yukawa		0.000606870766	0.000606870767	0.000606870767	+0.0012	+0.0012
tau Yukawa		0.010205763474	0.010205761974	0.010205761974	-0.1469	-0.1469
QUARK YUKAWA COUPLINGS						
up Yukawa		0.000012406407	0.000012476107	0.000012476107	+5618.0642	+5618.0642
down Yukawa		0.000026823112	0.000026802852	0.000026802852	-755.3173	-755.3173
strange Yukawa		0.000536462247	0.000535616364	0.000535616364	-1576.7799	-1576.7799
charm Yukawa		0.007311739193	0.007319192576	0.007319192576	+1019.3721	+1019.3721
bottom Yukawa		0.024025926978	0.024002518423	0.024002518423	-974.3039	-974.3039
top Yukawa		0.991191541612	0.992473523459	0.992473523459	+1293.3745	+1293.3745
HIGGS SELF-COUPLING						
Higgs self-coupling		0.129280565411	0.129280113847	0.129280113847	-3.4929	-3.4929
STRONG COUPLING						
strong coupling		0.118000000000	0.118047830923	0.118047830923	+405.3468	+405.3468
PMNS NEUTRINO MIXING						
PMNS solar mixing fraction		0.303000000000	0.306578869942	0.306578869942	+11811.4520	+11811.4520
PMNS atmospheric mixing fraction		0.572000000000	0.571428571429	0.571428571429	-999.0010	-999.0010
PMNS reactor mixing fraction		0.022030000000	0.021978021978	0.021978021978	-2359.4200	-2359.4200
PMNS CP phase	rad	3.4382986264	3.4557519189	3.4557519189	+5076.1421	+5076.1421
CKM QUARK MIXING						
Cabibbo mixing fraction		0.050640000000	0.050359712230	0.050359712230	-5534.9086	-5534.9086
CKM atmospheric mixing fraction		0.001682000000	0.001681796794	0.001681796794	-120.8121	-120.8121
CKM reactor mixing fraction		0.000014600000	0.000014526054	0.000014526054	-5064.8028	-5064.8028
CKM CP phase	rad	1.1431906601	1.1519173063	1.1519173063	+7633.5878	+7633.5878
QCD THETA ANGLE						
QCD theta angle	rad	0.0000000000	0.0000000000	0.0000000000	exact 0	exact 0
MASSLESS GAUGE BOSONS						
photon	MeV	0.000000	0.000000	0.000000	exact 0	exact 0
gluon 1	MeV	0.000000	0.000000	0.000000	exact 0	exact 0
gluon 2	MeV	0.000000	0.000000	0.000000	exact 0	exact 0
gluon 3	MeV	0.000000	0.000000	0.000000	exact 0	exact 0
gluon 4	MeV	0.000000	0.000000	0.000000	exact 0	exact 0
gluon 5	MeV	0.000000	0.000000	0.000000	exact 0	exact 0
gluon 6	MeV	0.000000	0.000000	0.000000	exact 0	exact 0
gluon 7	MeV	0.000000	0.000000	0.000000	exact 0	exact 0
gluon 8	MeV	0.000000	0.000000	0.000000	exact 0	exact 0
GRAVITATIONAL SECTOR						

TABLE VI. Per-scheme within-scheme prediction table for the pole/direct scheme. Anchored on the electron mass; predictions for muon, tau, top, W, Z, Higgs boson, and Higgs vev computed as $m_X = (R_X/R_{\text{anchor}}) \cdot m_{\text{anchor}}$. $\Delta(\sigma_{\text{comb}})$ folds target and propagated-anchor uncertainty in quadrature. Anchor: electron (observed = 0.5109989500 MeV, relative $\sigma = 0.000294$ ppm).

Particle	Observed	σ_{obs} (ppm)	Predicted	Δ(pred−obs)	Δ (ppm)	Δ (σ_{comb})
electron (anchor)	0.5109989500 MeV	+0.00	0.5109989500 MeV	–	–	–
muon	105.6583755000 MeV	+0.00	105.6583756208 MeV	0.0000001208 MeV	+0.0011	+3.8841
tau	1776.8600000000 MeV	+67.53	1776.8597388413 MeV	-0.0002611587 MeV	-0.1470	-0.0022
top	172.5700000000 GeV	+1738.42	172.7931976210 GeV	0.2231976210 GeV	+1293.3744	+0.7440
W boson	80.3692000000 GeV	+161.75	80.3694971362 GeV	0.0002971362 GeV	+3.6971	+0.0229
Z boson	91.1876000000 GeV	+23.03	91.1903715297 GeV	0.0027715297 GeV	+30.3937	+1.3198
Higgs boson	125.2000000000 GeV	+878.59	125.1997813344 GeV	-0.0002186656 GeV	-1.7465	-0.0020
Higgs vev	246.2196500000 GeV	+24.37	246.2196499809 GeV	-0.0000000191 GeV	-0.0001	-0.0000

TABLE VII. Per-scheme within-scheme prediction table for the $\overline{\text{MS}}$ scheme at $\mu = 2$ GeV (light quarks, single common scale – fully within-scheme). Anchor: down (observed = 4.6700000000 MeV, relative $\sigma = 7.07e + 04$ ppm).

Particle	Observed	σ_{obs} (ppm)	Predicted	Δ(pred−obs)	Δ (ppm)	Δ (σ_{comb})
up	2.1600000000 MeV	+175925.93	2.1737769100 MeV	0.0137769100 MeV	+6378.1991	+0.0336
down (anchor)	4.6700000000 MeV	+70663.81	4.6700000000 MeV	–	–	–
strange	93.4000000000 MeV	+92077.09	93.3232174033 MeV	-0.0767825967 MeV	-822.0835	-0.0071

framework — the residual deviations could in principle be hiding a systematic offset of a single multiplicative scale that happens to be inside all PDG error bars — but it does establish that the framework is not *tuned* to one probe scale at the expense of others. Future PDG updates that shrink the experimental uncertainty in any row will tighten the falsification window in that row.

XVI. PREDICTIONS AND FALSIFICATION

The framework makes pre-registered predictions across multiple sectors, falsifiable by:

1. Inverted neutrino mass ordering in future oscillation data.
2. Cosmological convergence to $\Sigma m_\nu > 70$ meV in future CMB and large-scale-structure analyses.
3. δ_{CP} outside $198^\circ \pm 5^\circ$ in future long-baseline oscillation data.
4. $0\nu\beta\beta$ at effective Majorana mass significantly above 4 meV (assuming Majorana nature) in next-generation neutrinoless-double-beta-decay searches.
5. Neutron EDM above the current upper bound, or a confirmed nonzero value above $\sim 10^{-30}\, e \cdot \text{cm}$ in next-generation EDM searches [16].
6. Newton's G convergence inconsistent with $6.67431691 \times 10^{-11}$ at ppm precision.
7. Dark-matter detection at masses inconsistent with the predicted tower (orders 5–11), or coupling significantly different from $g_{a\gamma\gamma} = 1.78 \times 10^{-17}$ GeV^{-1}.
8. Precision EW measurements converging to $\sin^2\theta_W$ inconsistent with $2/9 + \Delta_\varphi/h$ at ppm precision.

XVII. DISCUSSION

A. Reduction of the parameter count

Within the mass-spectrum construction presented here, the framework reduces the SM parameter count from 26 to 1: the electron rest mass is the sole dimensional input, with the remaining particle-spectrum quantities built from framework-fixed integers of the $E_8 \times E_8$ Coxeter spectrum on the Schoen Calabi–Yau quotient. The dark-matter abundance and strong-CP sectors are more conditional: the former currently closes through a substrate-to-misalignment dictionary, while the latter depends on the twisted-trace cancellation described in Sec. IX.

B. Common geometric origin

Notable unifying features include both CP phases as rational multiples of π ($\delta_{\text{CP}} = \pi + \pi/10$, $\delta_{\text{CKM}} = 11\pi/30$), the dark-sector coupling family graded by E_8 exponents ($\delta_H = 1/59$, $\delta_W = 1/(59 \cdot 19)$), the quotient-based conditional route to a vanishing strong CP angle, the unified neutrino-dark-matter tower, and the strong coupling as the rational 849/7192 from forced E_8 integers.

TABLE VIII. Per-scheme within-scheme prediction table for the $\overline{\text{MS}}$ scheme at $\mu = m_q$ (heavy quarks, per-particle scale – residual cross-scale within the group). Anchor: bottom (observed = 4183.0000000000 MeV, relative $\sigma = 1.67e + 03$ ppm).

Particle	Observed	σ_{obs} (ppm)	Predicted	Δ(pred−obs)	Δ (ppm)	Δ (σ_{comb})
charm	1273.0000000000 MeV	+3613.51	1275.5404247476 MeV	2.5404247476 MeV	+1995.6204	+0.5010
bottom (anchor)	4183.0000000000 MeV	+1673.44	4183.0000000000 MeV	–	–	–

TABLE IX. Supplementary derived neutrino-sector and cosmological observables not covered by the auto-generated complete table. Mass-squared splittings, sum of neutrino masses (Σm_ν), kinematic effective neutrino mass (m_β), neutrinoless double-beta effective mass ($m_{\beta\beta}$), and the dark-matter relic-density estimate $\Omega_{\text{DM}}h^2$. e-anchor values are computed from the predicted neutrino masses; μ-anchor values are obtained by linear (m) or quadratic (m^2) rescaling under the new h_{eff} ratio.

Observable	Observed	e-anchor	μ-anchor	Deviation
Δm^2_{21} (eV^2)	$7.41^{+0.21}_{-0.20} \times 10^{-5}$	$7.4193221245 \times 10^{-5}$	$7.4193221075 \times 10^{-5}$	−90 ppm
Δm^2_{31} (eV^2)	$(2.511 \pm 0.027) \times 10^{-3}$	$2.5211270648 \times 10^{-3}$	$2.5211270590 \times 10^{-3}$	+2440 ppm
Σm_ν (meV)	< 72 [21]	59.2898456922	59.2898456244	within
m_β (meV)	< 450 [22]	8.8276108075	8.8276107974	within
$m_{\beta\beta}$ (meV)	< 28–122 [23]	3.9984876767	3.9984876721	within
$\Omega_{\text{DM}}h^2$	0.1200 ± 0.001 [24]	0.119981783712	0.119981783369	−151.8 ppm

C. Comparison to prior work

The closest empirical precedent is Coldea *et al.* [3]'s confirmation of the E_8 mass spectrum in $CoNb_2O_6$. The closest theoretical precedent is Koide's mass formula [26], which relates the three charged-lepton masses at 10^{-5} precision with no derivation. The present framework matches Koide-level precision on leptons and extends the structural construction to all 26 SM parameters plus dark matter and gravity, while leaving the compactification-level explanation of some closure factors for follow-up work.

The geometric strong-CP mechanism, if the quotient-projected twisted trace vanishes, avoids invoking Peccei–Quinn [15] and so contrasts with the dominant axion-mediated approach.

D. Open directions

The paper still leaves several genuine follow-up tasks. Open directions for follow-up work:

1. Higher-order sub-harmonic precision beyond the current 0.0011 ppm floor.
2. Compactification-level derivation of the cycle-2 closure factors δ_H, δ_W, and the Coxeter-step assignment k_v from CY3 cohomology and Wilson-line data.
3. Substrate-native cosmological evolution for the dark-matter tower, replacing the imported misalignment-style abundance estimate with a full continuum calculation.
4. Independent derivation of the quotient-projected twisted-trace cancellation required for $\theta_{\text{QCD}} = 0$.
5. Cosmological inflationary connection: $n_s = 58/60$ from dual-E_8 Coxeter structure.
6. Beyond-SM extensions: direct DM detection signatures, generation count derivation.

E. Conclusion

We have presented a one-anchor structural construction of the Standard Model, the dark-matter sector, and the gravitational sector — 26 SM parameters, Newton's G, the Planck mass, the discrete dark-matter mass spectrum, and a misalignment-style cosmological dark-matter abundance estimate — from a single measured input (the electron rest mass) and the geometry of $E_8 \times E_8$ heterotic compactification on a Schoen Calabi–Yau quotient $X/(\mathbb{Z}_3 \times \mathbb{Z}_3)$, with McKay-correspondence anchoring. A conditional geometric route to $\theta_{\text{QCD}} = 0$ is identified through the quotient projection. Both CP phases are rational multiples of π. Two-anchor self-consistency between electron and muon anchors is verified at 30 parts per billion across all observables. The framework reduces the SM parameter count from 26 to 1 in the mass-spectrum construction presented here, while leaving the strong-CP cancellation and the dark-matter abundance as sharpened but not yet fully first-principles closures.

Appendix A: Coxeter step assignments

Appendix B: Complete structural correction list

Appendix C: Numerical precision

All numerical evaluations use mpmath at $\geq$ 50-digit decimal precision. Key exact rational values:

TABLE X. Coxeter step assignments. The φ-chain hosts integer E_8 exponent positions; the $\sqrt{2}$-chain hosts integer and half-integer positions. Cycle 2 indicates positions beyond $h = 30$.

Particle	k	Chain type / sector
Electron	0	φ gateway, dark-sector
Muon	11	φ exponent, visible
Tau	17	φ exponent, visible
Up	4	$\sqrt{2}$ int
Down	6.5	$\sqrt{2}$ half
Strange	15	$\sqrt{2}$ midpoint
Charm	22.5	$\sqrt{2}$ half
Bottom	26	$\sqrt{2}$ int
Top	36.5	$\sqrt{2}$ half, cycle 2
W boson	34.5	cycle 2, with δ_W
Z boson	via $m_W/\cos\theta_W$	derived
Higgs vev	1132/30	cycle 2
Higgs boson	$k_v - 2$	breathing mode, with δ_H

TABLE XI. All structural corrections, built from framework-fixed E_8 invariants dim $= 248$, $h = 30$, $M = 14$, rank $= 8$, $m_{\max} = 29$, together with the framework-selected McKay order 5 and golden ratio φ.

Symbol	Definition	Role
Δ_φ	$1/(h+\varphi^2)$	enters $\sin^2\theta_W$
Δ_2	$1/2^5 = 1/32$	2-fold structure
α_τ	$1/(\dim + h) = 1/278$	tau base
$\alpha_\tau^{\text{full}}$	$1/(\dim + h + \varphi^2)$	tau closed-form
α_μ	$2\alpha_\tau^2$	muon
α_v	$(\operatorname{rank} h/m_{\max})\alpha_\tau^2$	Higgs vev
α_H	$((\operatorname{rank}-2)h/m_{\max})\alpha_\tau^2$	Higgs boson
δ_H	$1/(h+m_{\max}) = 1/59$	Higgs dark coupling
δ_W	$1/(59\cdot 19) = 1/1121$	W dark coupling
ε	$1/(2\dim) = 1/496$	per-step quark loss
ω_{fix}	$1/2 - 1/\dim = 123/248$	gateway normalization
k_v	1132/30	Higgs vev position
f_5	$\frac{\operatorname{rank}(h-\operatorname{rank})}{\dim}\frac{\sqrt{vM_{\text{Pl}}}}{\alpha_\tau}$	DM decay constant
θ_5	$\arctan\varphi$	DM misalignment

- $h_{\text{eff}}^{(e)}/m_e = 124/123$;
- $30^{14} = 4{,}782{,}969 \times 10^{14}$;
- $\sin^2\theta_{12}^{\text{CKM}} = 14/278 = 7/139$;
- $\delta_{\text{CKM}} = 11\pi/30$;
- $\delta_{\text{CP}} = 11\pi/10$;
- $\sin^2\theta_W = 2/9 + 1/[h(h+\varphi^2)]$;
- $\alpha_s(M_Z) = 849/7192$;
- $\delta_H = 1/59$, $\delta_W = 1/1121$.

[1] S. Navas *et al.* (Particle Data Group), "Review of Particle Physics," Phys. Rev. D **110**, 030001 (2024). Open-access PDG portal (free, complete review): pdg.lbl.gov/2024. DOI: 10.1103/PhysRevD.110.030001.

[2] G. Bertone and T. M. P. Tait, "A new era in the search for dark matter," Nature **562**, 51–56 (2018). Open-access preprint (free): arXiv:1810.01668. DOI: 10.1038/s41586-018-0542-z.

[3] R. Coldea, D. A. Tennant, E. M. Wheeler, E. Wawrzynska, D. Prabhakaran, M. Telling, K. Habicht, P. Smeibidl, and K. Kiefer, "Quantum Criticality in an Ising Chain: Experimental Evidence for Emergent E_8 Symmetry," Science **327**, 177–180 (2010). Open-access repository copy (free, same content): arXiv:1103.3694. PubMed (free abstract): PMID 20056884. DOI: 10.1126/science.1180085.

[4] A. B. Zamolodchikov, "Integrals of motion and S-matrix of the (scaled) $T = T_c$ Ising model with magnetic field," Int. J. Mod. Phys. A **4**, 4235–4248 (1989). Inspire-HEP record (free abstract, cross-references): inspirehep.net/literature/276727. DOI: 10.1142/S0217751X8900176X.

[5] J. McKay, "Graphs, singularities, and finite groups," Proc. Symp. Pure Math. **37**, 183–186 (Amer. Math. Soc., 1980). Open scan of the original conference paper (free): McKaysOriginalPaper.pdf. DOI of the original: 10.1090/pspum/037/604577.

[6] J. E. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge Studies in Advanced Mathematics **29** (Cambridge University Press, 1990). ISBN 978-0-521-43613-7. Identity $|\Phi| = h \cdot \operatorname{rank}$ for irreducible finite Coxeter groups: §3.18, Eq. (3) (consequence of the exponents satisfying $\sum_j(e_j+1) = |\Phi|/2 + \operatorname{rank}$ with mean exponent $h/2$). Cambridge product page (free abstract): cambridge.org/core/books/reflection-groups-and-coxeter-groups. DOI: 10.1017/CBO9780511623646.

[7] P. W. Anderson, "Plasmons, Gauge Invariance, and Mass," Phys. Rev. **130**, 439–442 (1963). APS abstract page (free abstract): journals.aps.org/pr/abstract/10.1103/PhysRev.130.439. NASA ADS (free abstract + cross-references): ui.adsabs.harvard.edu/abs/1963PhRv..130..439A. DOI: 10.1103/PhysRev.130.439.

[8] G. E. Volovik, *The Universe in a Helium Droplet*, International Series of Monographs on Physics **117** (Oxford University Press, 2003). Oxford Academic landing page (free metadata): academic.oup.com/book/11557. DOI of book: 10.1093/acprof:oso/9780199564842.001.0001.

[9] G. E. Volovik, "Superfluid analogies of cosmological phenomena," Phys. Rep. **351**, 195–348 (2001). Open-access preprint (free): arXiv:gr-qc/0005091. DOI: 10.1016/S0370-1573(00)00139-3.

[10] C. Schoen, "On fiber products of rational elliptic surfaces with section," Math. Z. **197**, 177–199 (1988). EuDML page (free metadata): eudml.org/doc/183725. DOI: 10.1007/BF01215188.

[11] D. J. Gross, J. A. Harvey, E. Martinec, and R. Rohm, "Heterotic String," Phys. Rev. Lett. **54**, 502–505 (1985). Inspire-HEP record (free abstract + cross-references): inspirehep.net/literature/15974. NASA ADS: ui.adsabs.harvard.edu/abs/1985PhRvL..54..502G. DOI: 10.1103/PhysRevLett.54.502.

[12] V. Braun, Y.-H. He, B. A. Ovrut, and T. Pantev, "A standard model from the $E_8 \times E_8$ heterotic superstring," JHEP **06**, 039 (2005). Open-access preprint (free): arXiv:hep-th/0502155. DOI: 10.1088/1126-6708/2005/06/039.

[13] R. Donagi, Y.-H. He, B. A. Ovrut, and R. Reinbacher, "The spectra of heterotic standard model vacua," JHEP **06**, 070 (2005). Open-access preprint (free): arXiv:hep-th/0411156. DOI: 10.1088/1126-6708/2005/06/070.

[14] J. E. Kim and G. Carosi, "Axions and the strong CP problem," Rev. Mod. Phys. **82**, 557–601 (2010). Open-access preprint (free): arXiv:0807.3125. DOI: 10.1103/RevModPhys.82.557; Erratum: Rev. Mod. Phys. **91**, 049902 (2019).

[15] R. D. Peccei and H. R. Quinn, "CP Conservation in the Presence of Pseudoparticles," Phys. Rev. Lett. **38**, 1440–1443 (1977). NASA ADS (free abstract + cross-references): ui.adsabs.harvard.edu/abs/1977PhRvL..38.1440P. Inspire-HEP: inspirehep.net/literature/120612. DOI: 10.1103/PhysRevLett.38.1440.

[16] C. Abel *et al.* (nEDM Collaboration at PSI), "Measurement of the Permanent Electric Dipole Moment of the Neutron," Phys. Rev. Lett. **124**, 081803 (2020). Open-access preprint (free): arXiv:2001.11966. DOI: 10.1103/PhysRevLett.124.081803. Reports $|d_n| < 1.8 \times 10^{-26}\, e \cdot \mathrm{cm}$ at 90% C.L.

[17] P. J. Mohr, D. B. Newell, B. N. Taylor, and E. Tiesinga, "CODATA recommended values of the fundamental physical constants: 2022," Rev. Mod. Phys. **97**, 025002 (2025). Open-access preprint (free): arXiv:2409.03787. NIST free interactive constant database: physics.nist.gov/cuu/Constants. DOI: 10.1103/RevModPhys.97.025002.

[18] C. Rothleitner and S. Schlamminger, "Invited Review Article: Measurements of the Newtonian constant of gravitation, G," Rev. Sci. Instrum. **88**, 111101 (2017). NIST author manuscript (free): nist.gov/publications/invited-review-article-measurements-newtonian-constant-gravitation-g. DOI: 10.1063/1.4994619.

[19] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, T. Schwetz, and A. Zhou, "The fate of hints: updated global analysis of three-flavor neutrino oscillations," JHEP **09**, 178 (2020). JHEP is open access; published version free at: doi:10.1007/JHEP09(2020)178. Preprint: arXiv:2007.14792.

[20] NuFIT Collaboration (I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, T. Schwetz, A. Zhou), "NuFIT 5.3: Three-neutrino fit based on data available in March 2024," `http://www.nu-fit.org/?q=node/278` (free, accessed April 2026).

[21] A. G. Adame *et al.* (DESI Collaboration), "DESI 2024 VI: Cosmological Constraints from the Measurements of Baryon Acoustic Oscillations," JCAP **02**, 021 (2025). JCAP is open access; published version free at: doi:10.1088/1475-7516/2025/02/021. Preprint: arXiv:2404.03002.

[22] M. Aker *et al.* (KATRIN Collaboration), "Direct neutrino-mass measurement based on 259 days of KATRIN data," Science **388**, 180–185 (2025). Open-access preprint (free): arXiv:2406.13516. DOI: 10.1126/science.adq9592. Reports $m_\beta < 0.45$ eV at 90% C.L.

[23] S. Abe *et al.* (KamLAND-Zen Collaboration), "Search for Majorana Neutrinos with the Complete KamLAND-Zen Dataset," Open-access preprint (free): arXiv:2406.11438 (2024). Reports $T^{0\nu}_{1/2} > 3.8 \times 10^{26}$ yr at 90% C.L., $\langle m_{\beta\beta} \rangle < 28$–$122$ meV depending on nuclear matrix element.

[24] N. Aghanim *et al.* (Planck Collaboration), "Planck 2018 results. VI. Cosmological parameters," Astron. Astrophys. **641**, A6 (2020). A&A is open access; published version free at: aanda.org/articles/aa/full_html/2020/09/aa33910-18. Preprint: arXiv:1807.06209. DOI: 10.1051/0004-6361/201833910; Erratum: A&A **652**, C4 (2021).

[25] K. K. Rogers and H. V. Peiris, "Strong Bound on Canonical Ultralight Axion Dark Matter from the Lyman-Alpha Forest," Phys. Rev. Lett. **126**, 071302 (2021). Open-access preprint (free): arXiv:2007.12705. DOI: 10.1103/PhysRevLett.126.071302. Reports $m_a > 2 \times 10^{-20}$ eV at 95% C.L.

[26] Y. Koide, "Fermion-boson two-body model of quarks and leptons and Cabibbo mixing," Lett. Nuovo Cimento **34**, 201–205 (1982). Inspire-HEP record (free abstract): inspirehep.net/literature/176414. DOI: 10.1007/BF02817096; expanded in Y. Koide, Phys. Rev. D **28**, 252 (1983), APS abstract; DOI: 10.1103/PhysRevD.28.252; Phys. Lett. B **120**, 161–165 (1983), publisher abstract. DOI: 10.1016/0370-2693(83)90644-5.

[27] E. Gable, *Exist: The Universal Ontology* (2026). Foundational substrate-physical derivations of c, $\hbar$, general relativity, quantum mechanics, Newton's laws, Maxwell's equations, and the heterotic $E_8 \times E_8$ commitment.

[28] V. I. Arnold, *Catastrophe Theory*, 3rd ed. (Springer-Verlag, Berlin, 1992). Free expository synonym (open access): encyclopediaofmath.org/wiki/Catastrophe_theory (Springer/EMS, free). Springer landing page (free metadata): link.springer.com/book/10.1007/978-3-642-58124-3. DOI of book: 10.1007/978-3-642-58124-3. The simply-laced ADE classification of elementary catastrophes shares its label set with the McKay correspondence used in Sec. II, but the two structures are mathematically distinct.

[29] M. Alcubierre, "The warp drive: hyper-fast travel within general relativity," Classical Quantum Gravity **11**, L73–L77 (1994). Open-access preprint (free): arXiv:gr-qc/0009013. DOI: 10.1088/0264-9381/11/5/001.

[30] P. Touboul *et al.* (MICROSCOPE Collaboration), "MI-

CROSCOPE Mission: Final Results of the Test of the Equivalence Principle," Phys. Rev. Lett. **129**, 121102 (2022). Open-access preprint (free): arXiv:2209.15487. DOI: 10.1103/PhysRevLett.129.121102. Reports Eötvös parameter $\eta(\mathrm{Ti}, \mathrm{Pt}) = (-1.5 \pm 2.3_{\mathrm{stat}} \pm 1.5_{\mathrm{sys}}) \times 10^{-15}$.