Config

```
import json
import os
import pandas as pd
import re

import nltk
import spacy
import sklearn
import torch
import transformers as trf
import umap
import umap.plot
from bertopic import BERTopic
from bertopic.representation import KeyBERTInspired, MaximalMarginalRelevance, OpenAI, PartOfSpeech
from bertopic.vectorizers import ClassTfidfTransformer
from hdbscan import HDBSCAN
from openai import OpenAI
from sentence_transformers import SentenceTransformer
from sklearn.cluster import KMeans
from sklearn.decomposition import PCA
from sklearn.feature_extraction.text import CountVectorizer
from umap import UMAP

# Load up config file
with open('../config.json', 'r') as f:
    config = json.load(f)

# Set up OpenAI client
client = OpenAI(api_key=config['openai_key'])
```

Datasets

Trait Phrases Dataset

```
# Load up data
trait_path = os.path.join('..', config['trait_data'])
trait_df = pd.read_csv(trait_path)

# Add combined_phrase column
trait_df['combined_phrase'] = trait_df['trait'] + ' in ' + trait_df['species']

print("Number of good phrases in the trait dictionary: {:d}".format(trait_df.shape[0]))
trait_df.head()
```

```
Number of good phrases in the trait dictionary: 22729
```

	trait	journal	title	abstract_category
0	FT protein	Science (New York, N.Y.)	Science-censored analysis of quantitative trait loci (QTL) in yeast reveals QTL for severe deleterious mutations.	0
1	FT protein	J Exp Bot	Single-nucleotide polymorphism identification, characterization and association analysis of candidate genes for the seed storage protein content of oilseed rape (Brassica napus) using high-throughput SNP genotyping.	0
2	FT protein	J Exp Bot	Single-nucleotide polymorphism identification, characterization and association analysis of candidate genes for the seed storage protein content of oilseed rape (Brassica napus) using high-throughput SNP genotyping.	0
3	FT protein	Science (New York, N.Y.)	Genetic quantitative trait loci, flowering time, and phenotypic variation in Arabidopsis thaliana.	0
4	FT protein	Science (New York, N.Y.)	Quantitative trait loci, growth, and behavior: how many genes does it take to build a mouse?	0

```
print("Number of good phrases in the trait dictionary: {:d}".format(trait_df.shape[0]))
```

```
Number of good phrases in the trait dictionary: 22729
```

Corpus Dataset

```
# Load up data
corpus_path = os.path.join('..', config['corpus_data'])
corpus_df = pd.read_csv(corpus_path)

print("Number of documents: {:d}".format(corpus_df.shape[0]))
corpus_df.head()
```

```
Number of documents: 111478
```

	PMID	Journal	Title	Abstract	Category
0	7710226	Genetics	A new linkage map of the common fruit fly Drosophila melanogaster.	A genetic map of the common fruit fly Drosophila melanogaster, based on data collected by many laboratories, is presented. The map is based on 1,232 genetic markers (mutations, restriction fragment length polymorphisms, transposable elements, and inversion breakpoints) and has a total length of 1,890 cM. The standard error for the position of a marker on the map averages 1.5 cM. The chromosomal locations of 235 DNA segments (single-copy genomic clones, pseudogenes, and functionless duplications) are also reported.	0
1	7710227	Genetics	Genetic analysis of the Drosophila segmentation gene hairy reveals novel pair-rule expression.	The hairy gene of Drosophila melanogaster encodes a nuclear protein with a basic domain and a helix-loop-helix domain. Mutations in hairy cause segmentation defects and ectopic bristles. We have cloned hairy and used in situ hybridization to RNA to determine the pattern of hairy expression in embryos. hairy RNA is expressed in a complex pattern that changes dynamically during segmentation. Initially, hairy RNA is expressed in a broad anterior domain and in seven stripes. Each stripe is wider than a parasegment and is positioned over the anterior half of the parasegment. Later, hairy RNA is expressed in a complex pattern in the developing central nervous system. Comparison of hairy expression with the expression of other segmentation genes suggests that hairy is a pair-rule gene.	0
2	7710228	Genetics	Mutations affecting the pattern of the larval cuticle in Drosophila melanogaster. I. Zygotic loci on the second chromosome.	Mutations affecting the pattern of the larval cuticle in Drosophila melanogaster. I. Zygotic loci on the second chromosome.	0
3	7710229	Genetics	Mutations affecting the pattern of the larval cuticle in Drosophila melanogaster. II. Zygotic loci on the third chromosome.	Mutations affecting the pattern of the larval cuticle in Drosophila melanogaster. II. Zygotic loci on the third chromosome.	0
4	7710230	Genetics	Genetic fine structure analysis of the rosy locus and adjacent regions in Drosophila melanogaster.	Genetic fine structure analysis of the rosy locus and adjacent regions in Drosophila melanogaster.	0

```
print("Number of good phrases in the trait dictionary: {:d}".format(corpus_df.shape[0]))
```

```
Number of good phrases in the trait dictionary: 111478
```

Pre-processing and other requirements

Down-Pre-Processing Requirements

```
# For this project, you will need to use "entities" and "noun_chunks", and you can ignore the other fields.
doc = nlp("Apple is looking at buying U.K. startup for $1 billion")
print("Apple is looking at buying U.K. startup for $1 billion")

# For loop to print the text and label_ for each entity in the doc
for ent in doc.ents:
    print(ent.text, ent.label_)

# For loop to print the text for each noun chunk in the doc
print("\nNoun chunks:")
for chunk in doc.noun_chunks:
    print(chunk.text)
```

```
Apple is looking at buying U.K. startup for $1 billion
Apple ORG
U.K. GPE
$1 billion MONEY

Noun chunks:
Apple
U.K. startup
$1 billion
```

Tf-Idf Pre-Processing

```
# Load up stopwords and spacy English
nltk.download('stopwords')
stopwords = nltk.corpus.stopwords.words('english')

# Setting up TF-IDF vectorizer. These steps also are the ones highlighted the most on my instructional phase
tf_idf_vectorizer = CountVectorizer(stop_words=stopwords, min_df=10, max_df=0.4, ngram_range=(1,3))
tf_idf = tf_idf_vectorizer.fit_transform(corpus_df['Abstract'])
print("Shape of TF-IDF matrix: {}".format(tf_idf.shape))

# Get vocabulary
tf_idf_vocab = tf_idf_vectorizer.vocabulary_
```

```
[nltk_data] Downloading package stopwords to /root/nltk_data...
[nltk_data]   Package stopwords is already up-to-date!
Shape of TF-IDF matrix: (111478, 1000)
```

	Abstract	Category
0	A genetic map of the common fruit fly Drosophila melanogaster, based on data collected by many laboratories, is presented. The map is based on 1,232 genetic markers (mutations, restriction fragment length polymorphisms, transposable elements, and inversion breakpoints) and has a total length of 1,890 cM. The standard error for the position of a marker on the map averages 1.5 cM. The chromosomal locations of 235 DNA segments (single-copy genomic clones, pseudogenes, and functionless duplications) are also reported.	0
1	The hairy gene of Drosophila melanogaster encodes a nuclear protein with a basic domain and a helix-loop-helix domain. Mutations in hairy cause segmentation defects and ectopic bristles. We have cloned hairy and used in situ hybridization to RNA to determine the pattern of hairy expression in embryos. hairy RNA is expressed in a broad anterior domain and in seven stripes. Each stripe is wider than a parasegment and is positioned over the anterior half of the parasegment. Later, hairy RNA is expressed in a complex pattern in the developing central nervous system. Comparison of hairy expression with the expression of other segmentation genes suggests that hairy is a pair-rule gene.	0
2	Mutations affecting the pattern of the larval cuticle in Drosophila melanogaster. I. Zygotic loci on the second chromosome.	0
3	Mutations affecting the pattern of the larval cuticle in Drosophila melanogaster. II. Zygotic loci on the third chromosome.	0
4	Genetic fine structure analysis of the rosy locus and adjacent regions in Drosophila melanogaster.	0

Spacy Pre-Processing

```
# Load up spacy model
nlp = spacy.load("en_core_web_sm")

# Getting all of the lemmas for each abstract
lemma_data = []
for doc in nlp.pipe(corpus_df['Abstract']):
    lemmas = [token.lemma_ for token in doc if token.is_alpha and not token.is_stop]
    lemma_data.append(' '.join(lemmas))

print("First lemmatized abstract:\n")
print(lemma_data[0])
```

```
First lemmatized abstract:

genetic map common fruit fly drosophila melanogaster base datum collect laboratory present map base 1,232 genetic marker mutation restriction fragment length polymorphism transposable element inversion breakpoint total length 1,890 cm standard error position marker map average 1.5 cm chromosomal location 235 dna segment single copy genomic clone pseudogene functionless duplication report
```

Analysis

Analysis Without Key-Phrases

Task 1

Use sentence-transformer model in this corpus. The figure should be 800*600, with white background color. You will need to generate two word cloud images: 1) save word frequency, and 2) save tf-idf. You may find the following libraries useful for this task: WordCloud, nltk.

1) Frequency Base

```
# Setting word cloud
wordcloud = WordCloud(width=800, height=600, background_color='white').generate(corpus_text)

# Plotting
plt.figure(figsize=(8, 6))
plt.imshow(wordcloud, interpolation='bilinear')
plt.axis('off')
plt.show()
```

2) Tf-idf Base

```
# Get tf-idf scores for each word
tf_idf_scores = dict(zip(tf_idf_vectorizer.get_feature_names_out(), tf_idf.sum(axis=0).tolist()[0]))

# Setting up word cloud
wordcloud = WordCloud(width=800, height=600, background_color='white').generate_from_frequencies(tf_idf_scores)

# Plotting
plt.figure(figsize=(8, 6))
plt.imshow(wordcloud, interpolation='bilinear')
plt.axis('off')
plt.show()
```



Task 2: Topic Modeling

Use sentence-transformer model in this corpus, with the following parameters: n_neighbors=15, n_components=5, min_dist=0.0, metric='cosine'. You will need to generate two word cloud images: 1) save word frequency, and 2) save tf-idf. You may find the following libraries useful for this task: WordCloud, nltk.

```
# Load up embedding model
embedding_model = SentenceTransformer("all-MiniLM-L6-v2")

# Pre-calculating embeddings
embeddings = embedding_model.encode(corpus_df['Abstract'], show_progress_bar=True)
```

```
Batches:   0%|          | 0/3484 [00:00<?, ?it/s]
```

```
# Setting up and training UMAP model
umap_model = UMAP(n_neighbors=15, n_components=5, min_dist=0.0, metric='cosine', random_state=42)
umap_embeddings = umap_model.fit_transform(embeddings)
```

```
# Setting up and training KMeans model
cluster_model = KMeans(n_clusters=50, random_state=42)
y_kmeans = cluster_model.fit_predict(umap_embeddings)
```

```
# Setting up and training BERTopic model
topic_model = BERTopic(hdbscan_model=cluster_model, verbose=True).fit(corpus_df['Abstract'], embeddings)
```

```
2024-04-15 17:57:38,227 - BERTopic - Reduced dimensionality
2024-04-15 17:57:38,239 - BERTopic - Clustered reduced embeddings
```

```
# Getting topic info
topic_model.get_topic_info()
```

	Topic	Count	Name
0	-1	33	-1_and_of_the_in
1	0	3000	0_of_and_the_in
2	1	3000	1_of_and_the_in
3	2	3000	2_of_and_the_in
4	3	3000	3_of_and_the_in
5	4	3000	4_of_and_the_in
6	5	3000	5_of_and_the_in
7	6	3000	6_of_and_the_in
8	7	3000	7_of_and_the_in
9	8	3000	8_of_and_the_in
10	9	3000	9_of_and_the_in
11	10	3000	10_of_and_the_in
12	11	3000	11_of_and_the_in
13	12	3000	12_of_and_the_in
14	13	3000	13_of_and_the_in
15	14	3000	14_of_and_the_in
16	15	3000	15_of_and_the_in
17	16	3000	16_of_and_the_in
18	17	3000	17_of_and_the_in
19	18	3000	18_of_and_the_in
20	19	3000	19_of_and_the_in
21	20	3000	20_of_and_the_in
22	21	3000	21_of_and_the_in
23	22	3000	22_of_and_the_in
24	23	3000	23_of_and_the_in
25	24	3000	24_of_and_the_in
26	25	3000	25_of_and_the_in
27	26	3000	26_of_and_the_in
28	27	3000	27_of_and_the_in
29	28	3000	28_of_and_the_in
30	29	3000	29_of_and_the_in
31	30	3000	30_of_and_the_in
32	31	3000	31_of_and_the_in
33	32	3000	32_of_and_the_in
34	33	3000	33_of_and_the_in
35	34	3000	34_of_and_the_in
36	35	3000	35_of_and_the_in
37	36	3000	36_of_and_the_in
38	37	3000	37_of_and_the_in
39	38	3000	38_of_and_the_in
40	39	3000	39_of_and_the_in
41	40	3000	40_of_and_the_in
42	41	3000	41_of_and_the_in
43	42	3000	42_of_and_the_in
44	43	3000	43_of_and_the_in
45	44	3000	44_of_and_the_in
46	45	3000	45_of_and_the_in
47	46	3000	46_of_and_the_in
48	47	3000	47_of_and_the_in
49	48	3000	48_of_and_the_in
50	49	3000	49_of_and_the_in

```
# Getting topic info
topic_model.get_topic_info()
```

	Topic	Count	Name
0	-1	33	-1_and_of_the_in
1	0	3000	0_of_and_the_in
2	1	3000	1_of_and_the_in
3	2	3000	2_of_and_the_in
4	3	3000	3_of_and_the_in
5	4	3000	4_of_and_the_in
6	5	3000	5_of_and_the_in
7	6	3000	6_of_and_the_in
8	7	3000	7_of_and_the_in
9	8	3000	8_of_and_the_in
10	9	3000	9_of_and_the_in
11	10	3000	10_of_and_the_in
12	11	3000	11_of_and_the_in
13	12	3000	12_of_and_the_in
14	13	3000	13_of_and_the_in
15	14	3000	14_of_and_the_in
16	15	3000	15_of_and_the_in
17	16	3000	16_of_and_the_in
18	17	3000	17_of_and_the_in
19	18	3000	18_of_and_the_in
20	19	3000	19_of_and_the_in
21	20	3000	20_of_and_the_in
22	21	3000	21_of_and_the_in
23	22	3000	22_of_and_the_in
24	23	3000	23_of_and_the_in
25	24	3000	24_of_and_the_in
26	25	3000	25_of_and_the_in
27	26	3000	26_of_and_the_in
28	27	3000	27_of_and_the_in
29	28	3000	28_of_and_the_in
30	29	3000	29_of_and_the_in
31	30	3000	30_of_and_the_in
32	31	3000	31_of_and_the_in
33	32	3000	32_of_and_the_in
34	33	3000	33_of_and_the_in
35	34	3000	34_of_and_the_in
36	35	3000	35_of_and_the_in
37	36	3000	36_of_and_the_in
38	37	3000	37_of_and_the_in
39	38	3000	38_of_and_the_in
40	39	3000	39_of_and_the_in
41	40	3000	40_of_and_the_in
42	41	3000	41_of_and_the_in
43	42	3000	42_of_and_the_in
44	43	3000	43_of_and_the_in
45	44	3000	44_of_and_the_in
46	45	3000	45_of_and_the_in
47	46	3000	46_of_and_the_in
48	47	3000	47_of_and_the_in
49	48	3000	48_of_and_the_in
50	49	3000	49_of_and_the_in

Task 3: Task 2 Classes

Use a BoxClip model on the corpus, with the following parameters: ecos_eps=50, window=5, min_count=5 for each of the top 10 of all words, and the 20 most similar words.

```
# Setting word cloud
wordcloud = WordCloud(width=800, height=600, background_color='white').generate(corpus_text)

# Plotting
plt.figure(figsize=(8, 6))
plt.imshow(wordcloud, interpolation='bilinear')
plt.axis('off')
plt.show()
```


Task 4: Task 3 Word Clusters

Use a BoxClip model on the corpus, with the following parameters: ecos_eps=50, window=5, min_count=5 for each of the top 10 of all words, and the 20 most similar words.

```
# Setting word cloud
wordcloud = WordCloud(width=800, height=600, background_color='white').generate(corpus_text)

# Plotting
plt.figure(figsize=(8, 6))
plt.imshow(wordcloud, interpolation='bilinear')
plt.axis('off')
plt.show()
```


Analysis With Key-Phrases

Task 2

Extract phrases and repeat task 1 and 3. You can take a shortcut at phrase extraction.

```
# Extract noun phrases
abstract_textacy = df_processed['abstract_info_textacy']

# Create a dictionary and corpus for LDA
dictionary = corpora.Dictionary(abstract_textacy)
corpus = [dictionary.doc2bow(text) for text in abstract_textacy]

# Train LDA model
lda_model = gensim.models.LdaModel(corpus, num_topics=10, id2word=dictionary, passes=15)

# Print the topics
topics = lda_model.print_topics(num_words=10)
for topic in topics:
    print(topic)
```

```
(0, '0.000*"genetic_map" + 0.000*"fruit_fly" + 0.000*"drosophila_melanogaster" + 0.000*"base_datum" + 0.000*"collect_laboratory" + 0.000*"present_map" + 0.000*"base_genetic_marker" + 0.000*"mutation_restriction_fragment_length_polymorphism" + 0.000*"transposable_element_inversion_breakpoint" + 0.000*"total_length_1,890_cm"')
(1, '0.000*"genetic_map" + 0.000*"fruit_fly" + 0.000*"drosophila_melanogaster" + 0.000*"base_datum" + 0.000*"collect_laboratory" + 0.000*"present_map" + 0.000*"base_genetic_marker" + 0.000*"mutation_restriction_fragment_length_polymorphism" + 0.000*"transposable_element_inversion_breakpoint" + 0.000*"total_length_1,890_cm"')
(2, '0.000*"genetic_map" + 0.000*"fruit_fly" + 0.000*"drosophila_melanogaster" + 0.000*"base_datum" + 0.000*"collect_laboratory" + 0.000*"present_map" + 0.000*"base_genetic_marker" + 0.000*"mutation_restriction_fragment_length_polymorphism" + 0.000*"transposable_element_inversion_breakpoint" + 0.000*"total_length_1,890_cm"')
(3, '0.000*"genetic_map" + 0.000*"fruit_fly" + 0.000*"drosophila_melanogaster" + 0.000*"base_datum" + 0.000*"collect_laboratory" + 0.000*"present_map" + 0.000*"base_genetic_marker" + 0.000*"mutation_restriction_fragment_length_polymorphism" + 0.000*"transposable_element_inversion_breakpoint" + 0.000*"total_length_1,890_cm"')
(4, '0.000*"genetic_map" + 0.000*"fruit_fly" + 0.000*"drosophila_melanogaster" + 0.000*"base_datum" + 0.000*"collect_laboratory" + 0.000*"present_map" + 0.000*"base_genetic_marker" + 0.000*"mutation_restriction_fragment_length_polymorphism" + 0.000*"transposable_element_inversion_breakpoint" + 0.000*"total_length_1,890_cm"')
(5, '0.000*"genetic_map" + 0.000*"fruit_fly" + 0.000*"drosophila_melanogaster" + 0.000*"base_datum" + 0.000*"collect_laboratory" + 0.000*"present_map" + 0.000*"base_genetic_marker" + 0.000*"mutation_restriction_fragment_length_polymorphism" + 0.000*"transposable_element_inversion_breakpoint" + 0.000*"total_length_1,890_cm"')
(6, '0.000*"genetic_map" + 0.000*"fruit_fly" + 0.000*"drosophila_melanogaster" + 0.000*"base_datum" + 0.000*"collect_laboratory" + 0.000*"present_map" + 0.000*"base_genetic_marker" + 0.000*"mutation_restriction_fragment_length_polymorphism" + 0.000*"transposable_element_inversion_breakpoint" + 0.000*"total_length_1,890_cm"')
(7, '0.000*"genetic_map" + 0.000*"fruit_fly" + 0.000*"drosophila_melanogaster" + 0.000*"base_datum" + 0.000*"collect_laboratory" + 0.000*"present_map" + 0.000*"base_genetic_marker" + 0.000*"mutation_restriction_fragment_length_polymorphism" + 0.000*"transposable_element_inversion_breakpoint" + 0.000*"total_length_1,890_cm"')
(8, '0.000*"genetic_map" + 0.000*"fruit_fly" + 0.000*"drosophila_melanogaster" + 0.000*"base_datum" + 0.000*"collect_laboratory" + 0.000*"present_map" + 0.000*"base_genetic_marker" + 0.000*"mutation_restriction_fragment_length_polymorphism" + 0.000*"transposable_element_inversion_breakpoint" + 0.000*"total_length_1,890_cm"')
(9, '0.000*"genetic_map" + 0.000*"fruit_fly" + 0.000*"drosophila_melanogaster" + 0.000*"base_datum" + 0.000*"collect_laboratory" + 0.000*"present_map" + 0.000*"base_genetic_marker" + 0.000*"mutation_restriction_fragment_length_polymorphism" + 0.000*"transposable_element_inversion_breakpoint" + 0.000*"total_length_1,890_cm"')
```

Task 3: Task 1 Repeat

Repeat task 1 using the following phrases:

```
# Setting word cloud
wordcloud = WordCloud(width=800, height=600, background_color='white').generate(corpus_text)

# Plotting
plt.figure(figsize=(8, 6))
plt.imshow(wordcloud, interpolation='bilinear')
plt.axis('off')
plt.show()
```


Task 4: Task 2 Classes

Repeat task 2 using the following phrases:

```
# Setting word cloud
wordcloud = WordCloud(width=800, height=600, background_color='white').generate(corpus_text)

# Plotting
plt.figure(figsize=(8, 6))
plt.imshow(wordcloud, interpolation='bilinear')
plt.axis('off')
plt.show()
```


Task 5: Task 3 Word Clusters

Repeat task 3 using the following phrases:

```
# Setting word cloud
wordcloud = WordCloud(width=800, height=600, background_color='white').generate(corpus_text)

# Plotting
plt.figure(figsize=(8, 6))
plt.imshow(wordcloud, interpolation='bilinear')
plt.axis('off')
plt.show()
```


Key Phrase Extractor - Rake

```
# Extract phrases
df_processed['abstract_info_rake'] = df_processed['abstract_info'].apply(lambda x: [phrase[0] for phrase in x])

# Print the phrases
df_processed[['abstract_info', 'abstract_info_rake']].head()
```

	abstract_info	abstract_info_rake
0	[(genetic map, 225.0), (common fruit fly drosophila melanogaster, 169.0), (base, 169.0), (collect laboratory, 169.0), (present map, 169.0), (base genetic marker, 169.0), (mutation restriction fragment length polymorphism, 169.0), (transposable element inversion breakpoint, 169.0), (total length 1,890 cm, 169.0), (standard error position marker map average 1.5 cm chromosomal location 235 dna segment single copy genomic clone pseudogene functionless duplication report, 169.0)]	[genetic map, fruit fly drosophila melanogaster, base datum, collect laboratory, present map, base genetic marker, mutation restriction fragment length polymorphism, transposable element inversion breakpoint, total length 1,890 cm, standard error position marker map average 1.5 cm chromosomal location 235 dna segment single copy genomic clone pseudogene functionless duplication report]
1	[(hairy gene drosophila melanogaster encodes nuclear protein basic domain helix loop helix domain, 225.0), (mutation hairy cause segmentation defect ectopic bristle, 169.0), (clone hairy use situ hybridization rna determine pattern hairy expression embryo, 169.0), (hairy rna express broad anterior domain seven stripe, 169.0), (stripe wide parasegment position anterior half parasegment, 169.0), (later hairy rna express complex pattern develop central nervous system, 169.0), (comparison hairy expression expression segmentation gene suggest hairy pair rule gene, 169.0)]	[hairy gene drosophila melanogaster encodes nuclear protein basic domain helix loop helix domain, mutation hairy cause segmentation defect ectopic bristle, clone hairy use situ hybridization rna determine pattern hairy expression embryo, hairy rna express broad anterior domain seven stripe, stripe wide parasegment position anterior half parasegment, later hairy rna express complex pattern develop central nervous system, comparison hairy expression expression segmentation gene suggest hairy pair rule gene]
2	[(mutation affect pattern larval cuticle drosophila melanogaster zygotic locus second chromosome, 225.0)]	[mutation affect pattern larval cuticle drosophila melanogaster zygotic locus second chromosome]
3	[(mutation affect pattern larval cuticle drosophila melanogaster zygotic locus third chromosome, 225.0)]	[mutation affect pattern larval cuticle drosophila melanogaster zygotic locus third chromosome]
4	[(genetic fine structure analysis rosy locus adjacent region drosophila melanogaster, 225.0)]	[genetic fine structure analysis rosy locus adjacent region drosophila melanogaster]

Task 6: Task 1 (Rake)

Repeat task 1 using the following phrases:

```
# Setting word cloud
wordcloud = WordCloud(width=800, height=600, background_color='white').generate(corpus_text)

# Plotting
plt.figure(figsize=(8, 6))
plt.imshow(wordcloud, interpolation='bilinear')
plt.axis('off')
plt.show()
```


Task 7: Task 2 (Rake)

Repeat task 2 using the following phrases:

```
# Setting word cloud
wordcloud = WordCloud(width=800, height=600, background_color='white').generate(corpus_text)

# Plotting
plt.figure(figsize=(8, 6))
plt.imshow(wordcloud, interpolation='bilinear')
plt.axis('off')
plt.show()
```


Task 8: Task 3 (Rake)

Repeat task 3 using the following phrases:

```
# Setting word cloud
wordcloud = WordCloud(width=800, height=600, background_color='white').generate(corpus_text)

# Plotting
plt.figure(figsize=(8, 6))
plt.imshow(wordcloud, interpolation='bilinear')
plt.axis('off')
plt.show()
```

snp
region
qtl
gene
chromosome
study
effect
associated
analysis
pig
genome
genomewide
identified
polymorphism
weight
fat
chicken
sheep
milk
growth
significant
family
production
genetic
carcass
detected
linkage
model
single
haplotype
protein
candidate
mutation
locus
allele
variation
genotyped
cattle
gwas
qtls
level
selection
trait
found
located
total
acid
result
number
variance
animal
cm
genomic
different
quantitative
chromosomal
muscle
breeding
horse
sire
dairy
bovine
reported
disease
expression
control
sample
female
male
frequency
large
revealed
potential
factor
including
selected
human
white
meat
cross
interval
porcine
background
nucleotide
microsatellite
genotype
genotyping
estimated
area
size
trait
affect
resistance
identify
egg
day
feed
respectively
related
mastitis
bone
composition
mapped
resource
explained
average
test
high
detect
score
new
previously
beef
consumption
value
hair
fatty
based
component
objective
multiple
showed
present
genome
response
method
f2
duroc
study
variant
content
qtls
level
gwas
selection
approach
reproductivesignificance
candidate
protein
production
identified
mutation
locus
allele
variation
genotyped
analysis
significant
family
reported
bovine
detect
poor
dairy
test
high
previously
milk
growth
muscle
disease
consumption
chromosome
gene
study
qtl
region
snp
pig
genome
genomewide
identified
polymorphism
weight
fat
chicken
sheep
haplotype
linkage
model
single
detected
quantitative
different
genetic
carcass
result
number
variance
animal
cm
genomic
acid
total
located
pig
cross
including
selected
human
white
meat
porcine
background
nucleotide
microsatellite
genotyped
genotyping
estimated
area
size
trait
affect
resistance
identify
egg
day
feed
respectively
related
mastitis
bone
composition
mapped
resource
explained
average
test
high
detect
score
new
previously
beef
consumption
value
hair
fatty
based
component
objective
multiple
showed
present
genome
response
method
f2
duroc
study
variant
content
qtls
level
gwas
selection
approach
reproductivesignificance
candidate
protein
production
identified
mutation
locus
allele
variation
genotyped
analysis
significant
family
reported
bovine
detect
poor
dairy
test
high
previously
milk
growth
muscle
disease