

Healthy
Environments and
Consumer Safety
Branch

HEALTH EFFECTS OF WINDSOR AIR POLLUTION ON DIABETIC PATIENTS AND PREGNANT WOMEN

Ling Liu, Health Canada October 21, 2005

Canadä



WHY CONDUCT STUDY ON DIABETICS?

- Epidemiological evidence:
 - Air pollution increases the risk of cardiovascular mortality and morbidity, e.g. myocardial infarction
 - Diabetic patients appear to be more susceptible
 - Mortality study conducted in Montreal (Goldberg et al., 2000)
 - CVD hospitalizations in the U.S.
 - Diabetics: Cross-sectional association between PM2.5 and reduced vascular reactivity (O'Neill et al., 2005)
- Uncertainty: Mechanistic evidence is still limited



WHY CONDUCT STUDY ON BIRTH OUTCOMES?

- Epidemiological evidence:
 - Exposure to ambient PM is associated with low birth weight (LBW)
 - Exposure to PM is associated with infant mortality; the association was stronger for LBW infants than for normalbirth-weight infants
- Question: Does exposure to air pollution during pregnancy cause physiological/biochemical changes which subsequently result in adverse birth outcomes?

DIABETICS AND PREGNANT WOMEN ARE VULNERABLE

Diabetic patients:

- Diabetics tend to have higher blood levels of proinflammatory cytokines and oxidative stress - alter cardiovascular system
- Cardiovascular disease, atherosclerosis and nephropathy are major complications of diabetes mellitus

Pregnant women:

- They experience rapid physiological changes during pregnancy
- The developing embryo is sensitive to changes in environment
- Biological changes in maternal blood, e.g. proinflammatory cytokine levels and reactive oxygen species, are associated with adverse birth outcomes
 - pre-term births
 - Low weight & length

HECS

HYPOTHESIS

Exposure to high air pollution Systemic Direct Increased blood Increased proinflammatory brachial oxidative ET-1 cytokines artery stress constriction Endothelial/epithelial inflammation and dysfunction, deposition of lipids on artery wall Atherosclerosis, coronary artery disease, nephropathy etc. in diabetic patients

Adverse birth outcomes

HECS

DIABETES STUDY PROCEDURE

Recruitment:

- 25 patients
- 18-50 year old
- Type I/II diabetes
- No heart & lung conditions
- Non-smokers
- Visited a Windsor clinic once a week for 6-7 weeks, 0.5 hours each time




- Personal PM10
- •EC pollution data
- Spatial pollution data







STUDY PROCEDURE -Pregnant women pilot study



Non-smokers

Recruitment:

- No heart & lung conditions
- Visit Ottawa Hospital once each trimester

