

# Downscaling metawebs: propagation of uncertainties in species distribution and interaction probability

Gabriel Dansereau<sup>1,2,‡</sup> Ceres Barros<sup>3</sup> Timothée Poisot<sup>1,2</sup>

<sup>1</sup> Université de Montréal <sup>2</sup> Québec Centre for Biodiversity Sciences <sup>3</sup> University of British Columbia

‡ Equal contributions

## Correspondance to:

Gabriel Dansereau — [gabriel.dansereau@umontreal.ca](mailto:gabriel.dansereau@umontreal.ca)

# 1 Introduction

2 Here, we present a method to downscale a metaweb in space by developing an explicit spatial probabilistic  
3 metaweb for Canadian mammals. We present how the spatial structure of the downscaled metaweb varies in  
4 space and how the uncertainty of interactions can be made spatially explicit. We further show that the  
5 downscaled metaweb can highlight important biodiversity areas and bring novel ecological insights compared  
6 to community measures.

## 7 Methods

8 Fig. 1 shows a conceptual overview of the methodological steps leading to the downscaled metaweb. The  
9 components were grouped as the inputs (spatial or non-spatial), the localized steps (divided into  
10 single-species-level, two-species-level, and network-level steps), and the final downscaled and spatialized  
11 output. Throughout these steps, we highlight the importance of presenting the uncertainty of both interactions  
12 and their distribution in space. We argue that this requires adopting a probabilistic view and incorporating  
13 variation between scales.

14 [Figure 1 about here.]

## 15 Inputs

16 The inputs were divided into two main categories: the spatial and non-spatial ones (*Inputs* box on Fig. 1).

### 17 Non-spatial inputs

18 The main building block for the interaction data was the metaweb for Canadian mammals from Strydom *et al.*  
19 (2022a), a non-spatial input (represented as nodes and links on Fig. 1). A metaweb contains all the possible  
20 interactions between the species found in a given regional species pool (Dunne 2006). The species list for the  
21 Canadian metaweb was extracted from the International Union for the Conservation of Nature (IUCN) checklist  
22 (Strydom *et al.* 2022a). Briefly, the metaweb was developed using graph embedding and phylogenetic transfer  
23 learning based on the metaweb of European mammals, which is itself based on a comprehensive survey of  
24 interactions reported in the scientific literature (Maiorano *et al.* 2020). The Canadian metaweb is probabilistic,

25 which has the advantage of taking into account that species do not necessarily interact whenever they co-occur  
26 (Blanchet *et al.* 2020). However, the Canadian metaweb is not explicitly spatial: it only gives information on  
27 interactions in Canada as a whole and does not represent networks at specific locations. Local networks, on the  
28 other hand, are realizations from the metaweb resulting from sorting the species and the interactions (Poisot *et*  
29 *al.* 2015). A spatial and localized metaweb is not equivalent to the local networks, as it will have a different  
30 structure and a higher connectance (Strydom *et al.* 2022b). Therefore, producing a spatial metaweb requires  
31 additional steps to account for species composition and interaction variability in space.

## 32 **Spatial inputs**

33 The spatial data used to develop the spatial component of the metaweb were species occurrences and  
34 environmental data. First, we extracted species occurrences from the Global Biodiversity Information Facility  
35 (GBIF; [www.gbif.org](http://www.gbif.org)) for the Canadian mammals after reconciling species names between the Canadian  
36 metaweb and GBIF using the GBIF Backbone Taxonomy (GBIF Secretariat 2021). Doing so, we removed  
37 potential duplicates where species listed in the Canadian metaweb are considered as a single species by GBIF.  
38 We collected occurrences for our species list using the GBIF download API on October 21st 2022 (GBIF.org  
39 2022). We restricted our query to occurrences with coordinates between longitudes 175°W to 45°W and  
40 latitudes 10°N to 90°N. This was meant to collect training data covering a broader range than our prediction  
41 target (Canada only) and include observations in similar environments. Then, since GBIF observations  
42 represent presence-only data and most predictive models require absence data, we generated pseudo-absence  
43 data using the surface range envelope method available in `SimpleSDMLayers.jl` (Dansereau & Poisot 2021).  
44 This method generates pseudo-absences by selecting random non-observed locations within the spatial range  
45 delimited by the presence data (Barbet-Massin *et al.* 2012).

46 We used environmental data and species distribution models (SDMs, Guisan & Thuiller 2005) to predict the  
47 distribution of Canadian mammals across the whole country. The environmental data we used were the 19  
48 bioclimatic variables from CHELSA (Karger *et al.* 2017) and the 12 consensus land cover variables from  
49 EarthEnv (Tuanmu & Jetz 2014). The CHELSA bioclimatic variables (*bio1-bio19*) represent various measures  
50 of temperature and precipitation (e.g., annual averages, monthly maximum or minimum, seasonality) and are  
51 available for land areas across the globe. Therefore, they can be used to capture the climatic tolerance of species  
52 and model habitat suitability in new locations. We used the most recent version, the CHELSA v2.1 dataset  
53 (Karger *et al.* 2021). However, this version also includes bioclimatic data for open water, while we decided here

54 to focus only on land surfaces. We used the previous version, CHELSA v1.2 (Karger *et al.* 2018), which shares  
55 a similar grid but does not cover open water, as a mask to clip the v2.1 data to land surfaces only. The EarthEnv  
56 land cover variables represent classes such as Evergreen broadleaf trees, Cultivated and managed vegetation,  
57 Urban/Built-up, and Open Water. Values range between 0 and 100 and represent the consensus prevalence of  
58 each class in percentage within a pixel. We coarsened both the CHELSA and EarthEnv data from their original  
59 30 arc-second resolution to a 2.5 arc-minute one (around 4.5 km at the Equator) using GDAL/OGR  
60 contributors (2021). This represented a compromise to catch both local variations and broad scale patterns  
61 while limiting computation costs to a manageable level, as memory requirements on localized interactions rise  
62 very quickly.

63 Our selection criteria for choosing an SDM algorithm was to have a method that generated probabilistic results,  
64 including both a probability of occurrence for a species in a specific location and the uncertainty associated  
65 with the prediction. These were crucial to obtaining a probabilistic version of the metaweb as they were used to  
66 create spatial variations in the localized interaction probabilities (see next section). One promising method for  
67 this is Gradient Boosted Trees with a Gaussian maximum likelihood from the `EvoTrees.jl` *Julia* package  
68 (<https://github.com/Evoest/EvoTrees.jl>). This method returns a prediction for every pixel with an average value  
69 and a standard deviation, which we used as a measure of uncertainty to build a Normal distribution for the  
70 probability of occurrence of a given species at all pixels (represented as probability distributions on Fig. 1). We  
71 trained models across the extent chosen for occurrences (longitudes 175°W to 45°W and latitudes 10°N to  
72 90°N), then predicted species distributions only for Canada. We used the 2021 Census Boundary Files from  
73 Statistics Canada (Statistics Canada 2022) to set the boundaries for our predictions.

## 74 **Localized steps**

75 The next part of the method was the localized steps which produce local metawebs in every pixel. This  
76 component was divided into single-species, two-species, and network-level steps (*Localized steps* box on Fig. 1).  
77 The single-species steps represented four possible ways to account for uncertainty in the species distributions  
78 and bring variation to the spatial metaweb. We explored four different options to select a value from the  
79 occurrence distributions obtained in the previous steps (Inputs section): 1) taking the mean from the distribution  
80 as the probability of occurrence (option 1 on Fig. 1); 2) converting the mean value to a binary one using a  
81 specific threshold per species (option 2); 3) sampling a random value within the Normal distribution (option 3);

4) converting the random value into a binary result (option 4). The threshold ( $\tau$  on Fig. 1) used was the value that maximized Youden's  $J$  informedness statistic (Youden 1950), the same metric used by Strydom *et al.* (2022a) at an intermediate step while building the metaweb. The four sampling options were intended to explore how uncertainty and variation in the species distributions can affect the metaweb result and reproduce some of the filterings that create the local network realizations (Poisot *et al.* 2015). We expected thresholding to have a more pronounced effect on network structure as it should reduce the number of links by removing many of the rare interactions (Poisot *et al.* 2016). Meanwhile, we expected random sampling to create spatial heterogeneity compared to the mean probabilities, as including some extreme values should disrupt the potential effects of environmental gradients.

Next, the two-species steps aimed to give the probability of observing a given interaction in a location. For all species pairs, we multiplied the two species' occurrence probability obtained using the sampling options described in the previous paragraph, then multiplied the co-occurrence probability by the interaction probability from the Canadian metaweb. For cases where species in the Canadian metaweb were considered as the same species by the GBIF Backbone Taxonomy (the reconciliation step mentioned earlier), we used the highest interaction probabilities involving the duplicated species.

The network-level steps then created the probabilistic metaweb for the location. We assembled all the local interaction probabilities (from the two-species steps) into a probabilistic network (Poisot *et al.* 2016). We then sampled several random network realizations to represent the potential local realization process (Poisot *et al.* 2015). Finally, this resulted in a distribution of localized networks, which we averaged over the number of simulations to obtain a probabilistic network.

## Outputs

The final output of our method was the spatial probabilistic metaweb, which contains a localized probabilistic metaweb in every cell across the student extent (Outputs box on Fig. 1). This gives us an idea of the possible networks in all locations as the metaweb essentially serves to set an upper bound on the potential interactions (Strydom *et al.* 2022b), but with the added benefit of accounting for co-occurrence probabilities in this case. From there, we can create maps of network properties (e.g. number of links, connectance) measured on the local realizations, display their spatial distribution, and compute some community-level measures such as species richness. We can also calculate the uncertainty associated with the network and community

110 measurements and contrast their spatial distribution (see Supplementary Material).

## 111 **Ecoregions**

112 Since both species composition and network summary values display a high spatial variation and complex  
113 patterns, we simplified the representation of their distribution by grouping sites by ecoregion, as species and  
114 interaction composition have been shown to differ between ecoregions across large spatial scales (Martins *et al.*  
115 2022). To do so, we used the global map of ecoregions from (Dinerstein *et al.* 2017; also used by Martins *et al.*  
116 2022), rasterized it, and clipped it to Canada, which selected 44 different ecoregions. For every measure we  
117 report (e.g. species richness, number of links), we first calculated the measure for every site separately, then we  
118 extracted the median value for each ecoregion. We also measured the within-ecoregion variation by measuring  
119 the 89% interquantile range of the values in each ecoregion (threshold chosen to avoid confusion with  
120 conventional significance tests, inspired by McElreath 2020).

## 121 **Ecological uniqueness**

122 We compared the compositional uniqueness of the networks and the communities to verify if they indicated  
123 different exceptional areas. We measured uniqueness using the local contributions to beta diversity (LCBD,  
124 Legendre & De Cáceres 2013), which identify sites with exceptional composition by quantifying how much one  
125 site contributes to the total variance in the community composition. While many studies used LCBD values to  
126 evaluate uniqueness on local scales or few study sites (for example, da Silva & Hernández 2014; Heino &  
127 Grönroos 2017), recent studies used the measure on predicted species compositions over broad spatial extents  
128 and a large number of sites (Vasconcelos *et al.* 2018; Dansereau *et al.* 2022). LCBD values can also be used to  
129 measure uniqueness for networks by computing the values over the adjacency matrix, which has been shown to  
130 capture more unique sites and uniqueness variability than through species composition (Poisot *et al.* 2017).  
131 Here, we measured and compared the uniqueness of our localized community and network predictions. We  
132 were especially interested in seeing if the sites identified as unique were the same based on the species and the  
133 interactions or if this method allowed identifying areas unique for one element (interactions, for instance) but  
134 not the other. Sites with such mismatches should warrant more investigation to understand the reasons for this  
135 difference.

## 136 Software

137 We used *Julia* v1.9.0 (Bezanson *et al.* 2017) to implement all our analyses. We used packages `GBIF.jl`  
138 (`Dansereau & Poisot 2021`) to reconcile species names using the GBIF Backbone Taxonomy,  
139 `SpeciesDistributionToolkit.jl` to handle raster layers and species occurrences, `EcologicalNetworks.jl`  
140 (`Poisot et al. 2019) to analyse network and metaweb structure, and Makie.jl (Danisch & Krumbiegel 2021) to  
141 produce figures. Our data sources (CHELSA, EarthEnv, Ecoregions) were all unprojected and we did not use a  
142 projection in our analyses, but we displayed the results using a Lambert conformal conic projection more  
143 appropriate for Canada using GeoMakie.jl. All the code used to implement our analyses is available on GitHub  
144 (https://github.com/PoisotLab/SpatialProbabilisticMetaweb) and includes instructions on how to run a smaller  
145 example at a coarse resolution. Note that running our analyses at full scale is resource and memory intensive and  
146 required the use of compute clusters provided by Calcul Québec and the Digital Research Alliance of Canada.`

## 147 Results

148 Fig. 2 shows ecoregion-level measures.

149 [Figure 2 about here.]

150 Fig. 3 shows the LCBD results for the ecoregions.

151 [Figure 3 about here.]

## 152 Discussion

153 Barbet-Massin, M., Jiguet, F., Albert, C.H. & Thuiller, W. (2012). [Selecting pseudo-absences for species](#)  
154 [distribution models: How, where and how many?](#) *Methods in Ecology and Evolution*, 3, 327–338.

155 Bezanson, J., Edelman, A., Karpinski, S. & Shah, V.B. (2017). [Julia: A fresh approach to numerical computing.](#)  
156 *SIAM Review*, 59, 65–98.

157 Blanchet, F.G., Cazelles, K. & Gravel, D. (2020). [Co-occurrence is not evidence of ecological interactions.](#)  
158 *Ecology Letters*, 23, 1050–1063.

159 da Silva, P.G. & Hernández, M.I.M. (2014). [Local and regional effects on community structure of dung beetles](#)  
160 [in a mainland-island scenario](#). *PLOS ONE*, 9, e111883.

161 Danisch, S. & Krumbiegel, J. (2021). [Makie.jl: Flexible high-performance data visualization for Julia](#). *Journal*  
162 *of Open Source Software*, 6, 3349.

163 Dansereau, G., Legendre, P. & Poisot, T. (2022). [Evaluating ecological uniqueness over broad spatial extents](#)  
164 [using species distribution modelling](#). *Oikos*, 2022, e09063.

165 Dansereau, G. & Poisot, T. (2021). [SimpleSDMLayers.jl and GBIF.jl: A framework for species distribution](#)  
166 [modeling in Julia](#). *Journal of Open Source Software*, 6, 2872.

167 Dinerstein, E., Olson, D., Joshi, A., Vynne, C., Burgess, N.D., Wikramanayake, E., *et al.* (2017). [An](#)  
168 [Ecoregion-Based Approach to Protecting Half the Terrestrial Realm](#). *BioScience*, 67, 534–545.

169 Dunne, J. (2006). The network structure of food webs. In: *Ecological Networks: Linking Structure to Dynamics*  
170 *in Food Webs*. pp. 27–86.

171 GBIF Secretariat. (2021). [GBIF Backbone Taxonomy](#).

172 GBIF.org. (2022). [GBIF occurrence download](#).

173 GDAL/OGR contributors. (2021). *GDAL/OGR geospatial data abstraction software library*. Manual. Open  
174 Source Geospatial Foundation.

175 Guisan, A. & Thuiller, W. (2005). [Predicting species distribution: Offering more than simple habitat models](#).  
176 *Ecology Letters*, 8, 993–1009.

177 Heino, J. & Grönroos, M. (2017). [Exploring species and site contributions to beta diversity in stream insect](#)  
178 [assemblages](#). *Oecologia*, 183, 151–160.

179 Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., *et al.* (2017). [Climatologies at](#)  
180 [high resolution for the earth's land surface areas](#). *Scientific Data*, 4, 170122.

181 Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., *et al.* (2018). [Data from:](#)  
182 [Climatologies at high resolution for the earth's land surface areas](#).

183 Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., *et al.* (2021). [Climatologies at](#)  
184 [high resolution for the earth's land surface areas](#).



185 Legendre, P. & De Cáceres, M. (2013). [Beta diversity as the variance of community data: Dissimilarity](#)  
186 [coefficients and partitioning](#). *Ecology Letters*, 16, 951–963.

187 Maiorano, L., Montemaggiore, A., Ficetola, G.F., O'Connor, L. & Thuiller, W. (2020). [TETRA-EU 1.0: A](#)  
188 [species-level trophic metaweb of European tetrapods](#). *Global Ecology and Biogeography*, 29, 1452–1457.

189 Martins, L.P., Stouffer, D.B., Blendinger, P.G., Böhning-Gaese, K., Buitrón-Jurado, G., Correia, M., *et al.*  
190 (2022). [Global and regional ecological boundaries explain abrupt spatial discontinuities in avian frugivory](#)  
191 [interactions](#). *Nature Communications*, 13, 6943.

192 McElreath, R. (2020). *Statistical rethinking: A bayesian course with examples in R and Stan*. Second.  
193 Chapman and Hall/CRC, New York.

194 Poisot, T., Bélisle, Z., Hoebeke, L., Stock, M. & Szefer, P. (2019). [EcologicalNetworks.jl: Analysing ecological](#)  
195 [networks of species interactions](#). *Ecography*, 42, 1850–1861.

196 Poisot, T., Cirtwill, A.R., Cazelles, K., Gravel, D., Fortin, M.-J. & Stouffer, D.B. (2016). [The structure of](#)  
197 [probabilistic networks](#). *Methods in Ecology and Evolution*, 7, 303–312.

198 Poisot, T., Guévenex-Julien, C., Fortin, M.-J., Gravel, D. & Legendre, P. (2017). [Hosts, parasites and their](#)  
199 [interactions respond to different climatic variables](#). *Global Ecology and Biogeography*, 26, 942–951.

200 Poisot, T., Stouffer, D.B. & Gravel, D. (2015). [Beyond species: Why ecological interaction networks vary](#)  
201 [through space and time](#). *Oikos*, 124, 243–251.

202 Statistics Canada. (2022). *Boundary files, reference guide second edition, Census year 2021*. Second edition.  
203 Statistics Canada = Statistique Canada, Ottawa.

204 Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M.J., *et al.* (2022a). [Food web](#)  
205 [reconstruction through phylogenetic transfer of low-rank network representation](#). *Methods in Ecology and*  
206 *Evolution*, n/a.

207 Strydom, T., Bouskila, S., Banville, F., Barros, C., Caron, D., Farrell, M.J., *et al.* (2022b). [Predicting metawebs:](#)  
208 [Transfer of graph embeddings can help alleviate spatial data deficiencies](#).

209 Tuanmu, M.-N. & Jetz, W. (2014). [A global 1-km consensus land-cover product for biodiversity and ecosystem](#)  
210 [modelling](#). *Global Ecology and Biogeography*, 23, 1031–1045.

211 Vasconcelos, T.S., Nascimento, B.T.M. do & Prado, V.H.M. (2018). [Expected impacts of climate change](#)  
212 [threaten the anuran diversity in the Brazilian hotspots](#). *Ecology and Evolution*, 8, 7894–7906.

213 Youden, W.J. (1950). [Index for rating diagnostic tests](#). *Cancer*, 3, 32–35.




Figure 1: Conceptual figure of the workflow to obtain the spatial probabilistic metaweb (Chapter 1). The workflow has three components: the inputs, the localized steps, and the final spatial output. The inputs are composed of the spatial data (data with information in every cell) and the non-spatial data (constant for all of Canada). The localized steps use these data and are performed separately in every cell, first at a single-species level (using distribution data), then for every species pair (adding interaction data from the metaweb), and finally at the network level by combining the results of all species pairs. The final output coming out of the network-level steps contains a spatialized probabilistic metaweb for every cell across the study extent.




Figure 2: (A-B) Example of a community measure (A, expected species richness) and a network one (B, expected number of links). Both measures are assembled from the predicted probabilistic communities and networks, respectively. Values are first measured separately for all sites, then the median value is taken to represent the ecoregion-level value. (C-B) Representation of the 89% interquartile range of values within the ecoregion for expected richness (C) and expected number of links (D).




Figure 3: (A-B) Representation of the ecoregion uniqueness values based on species composition (a) and network composition (b). LCBD values were first computed across all sites and scaled relative to the maximum value observed. The ecoregion LCBD value is the median value for the sites in the ecoregion. (C) Bivariate representation of species and network composition LCBD. Values are grouped into three quantiles separately for each variable. The colour combinations represent the nine possible combinations of quantiles. The species uniqueness (horizontal axis) goes left to right from low uniqueness (light grey, bottom left) to high uniqueness (green, bottom right). The network composition uniqueness goes bottom-up from low uniqueness (light grey, bottom left) to high uniqueness (blue, top left). (D) Probability densities for the ecoregion LCBD values for species and network LCBD (left), highlighting the variability of the LCBD between ecoregions, and the 89% interquartile range of the values within each ecoregion (right), highlighting the variability within the ecoregions.