

Optimizing sampling and monitoring of species interactions within Biodiversity Observation Networks

Gabriel Dansereau^{1,2}, Michael D. Catchen^{1,2}, Ceres Barros^{3,4,5} and Timothée Poisot^{1,2}

¹ Université de Montréal, Département de sciences biologiques, Montréal QC, Canada

² Québec Centre for Biodiversity Science, , Montréal QC, Canada

³ Canadian Forest Service, Natural Resources Canada, Victoria BC, Canada

⁴ Department of Forest Resources Management, University of British Columbia, Vancouver BC, Canada

⁵ Département des sciences du bois et de la forêt, Université Laval, Québec QC, Canada

Correspondence to Gabriel Dansereau — gabriel.dansereau@umontreal.ca

Abstract: Optimal monitoring strategies should be designed to efficiently monitor all essential facets of biodiversity. Yet, species interactions are often overlooked in monitoring designs compared to spatial coverage and species richness, partly due to the inherent difficulty of sampling and monitoring interactions compared to species distributions. Here, we used simulations to test the efficiency of monitoring species interactions within Biodiversity Observations Networks (BONs). We tested several methods for designing and optimizing BONs to monitor species interactions and examined efficiency when increasing the number of monitored sites. We showed that the required sampling effort is considerably higher for interactions, especially when considering that interaction realization and detection realistically depend on species abundance. Thus, expectations to monitor a relevant proportion of interactions within a BON should be lower than expectations to monitor its species. However, from a single-species perspective, optimizing monitoring designs based on a known species range proved an efficient strategy to retrieve its interactions, outperforming designs based on total species or interaction richness. Further efficiency gains required detailed knowledge of where realized interactions occurred, highlighting the need for better knowledge of where interactions are likely to take place and integration of factors influencing interaction realization, such as species abundances. Our results were consistent even when the total species range was over- or underestimated by 10%, which shows a tolerance in the precision of the information required for optimization. Therefore, our results highlight how attainable levels of information can lead to efficiency improvements when designing BONs. With the right target and optimization strategy, available information, such as species ranges, can guide observation network designs to monitor species interactions as central biodiversity components.

Keywords: biodiversity monitoring, species interaction networks, biodiversity observation networks, sampling design

1 Introduction

2 Biodiversity monitoring should encompass all facets of biodiversity. Article 7 of the Convention on
3 Biological Diversity ([United Nations 1992](#)) calls for Parties to “[i]dentify components of biological
4 diversity important for its conservation and sustainable use”, and “[m]onitor, through sampling and
5 other techniques, the components of biological diversity”. Biodiversity Observation Networks
6 (BONs) are promising initiatives to improve monitoring across scales and biodiversity facets. BONs
7 are coordinated monitoring networks designed to harmonize observation systems, assist reporting
8 on international assessments, and reinforce scientific basis in biodiversity monitoring ([Navarro et al.
9 2017](#), [Gonzalez et al. 2023](#)). BONs can be thematic (Marine BON, Freshwater BON), national (China
10 BON, French BON, Colombia BON) or regional (Asia-Pacific, Europe); some represent networks of
11 monitoring sites for systematic monitoring (China BON) while others represent biodiversity data
12 hubs (French BON) ([Xu et al. 2017](#), [Navarro et al. 2017](#)). Additional BONs are currently in the
13 development or planification stages, notably in Europe ([Kissling et al. 2024](#)) and in Canada ([Simard
14 2024](#), [Gonzalez et al. 2025](#)), and could eventually be interlinked into the Global Biodiversity
15 Observation System (GBIOS), a globally connected network providing capacity to monitor
16 biodiversity change ([Gonzalez et al. 2023](#)) and guiding actions needed for targets and goals of the
17 Kunming-Montreal Global Biodiversity framework ([CBD 2022](#)).

18 Recent work highlighted the need for improved sampling designs in BONs and proper adjustment to
19 monitoring goals. First, current sampling designs over large scales are often too sparse, at too coarse
20 a scale, too infrequent, and without temporal replication ([Santana et al. 2025](#)). For example,
21 recommendations for more effective spatial sampling designs in Europe included i) stratified
22 random sampling (with stratification across environmental, anthropogenic and political gradients or
23 classes, ii) incorporation of existing monitoring sites, iii) filling of spatial gaps ([Kissling et al. 2024](#)).
24 Second, ecological monitoring design relies on site selection algorithms more often discussed
25 regarding their capacity to achieve spatial balance and representative samples, and less frequently
26 for achieving an ecological monitoring goal ([Norman and Poisot 2025](#)). Species richness is the most
27 commonly used biodiversity measure in monitoring contexts ([Hillebrand et al. 2018](#)), including in
28 discussions over sampling designs. On one hand, maximizing richness can efficiently capture
29 functional and phylogenetic diversity in conservation networks ([Willig et al. 2023](#)). On the other
30 hand, the use of species richness to measure biodiversity change and guide biological conservation
31 has also been criticized, mainly since trends in species richness do not properly capture local
32 changes, neglecting species identities and functional roles, and do not providing information on
33 persistence of biodiversity and ecosystem services ([Hillebrand et al. 2018](#), [Aguilar et al. 2024](#), [Fletcher
34 Jr. et al. 2025](#)). Thus, there is a need to better adjust BON sampling designs to biodiversity facets
35 beyond species richness.

36 Species interactions are a central component of biodiversity ([McCann 2007](#), [Valiente-Banuet et al.
37 2015](#), [Harvey et al. 2017](#)), critical for ecosystem functioning and health, yet are undervalued in
38 biodiversity monitoring and conservation ([Jordano 2016a](#), [Dansereau et al. 2025](#)). Interactions
39 mediate community and ecosystem responses to change ([Garzke et al. 2019](#)) and can improve our

ability to predict responses to climate change (Abrego et al. 2021). They have their own spatial and macro-ecological dynamics (Baiser et al. 2019, Mestre et al. 2022, Windsor et al. 2023, Dansereau et al. 2024), and may be lost (or change) faster than species (Valiente-Banuet et al. 2015, Doré et al. 2021). Despite a lack of data at global scales and uneven data availability across different interaction types and environments (Hortal et al. 2015, Poisot et al. 2021), we now have sufficient conceptual tools to support the added value of species interactions in designing efficient management actions (Dansereau et al. 2025, Moracho et al. 2025). Therefore, integrating species interactions into existing monitoring programs, such as BONs, is something we need to undertake in the near future.

Prior investigations of interaction sampling did not address how efficiently we should expect to monitor interactions within the BON framework, nor did studies on the effectiveness of BONs pay particular attention to the sampling of species interactions. Instead, much focus has been given on how to assess sampling completeness and diversity metrics (e.g. Chacoff et al. 2012, Chiu et al. 2023) and how sampling effort affects network properties (e.g. McLeod et al. 2021). Recent empirical (Caron et al. 2024) and theoretical (Poisot 2023) results have established that knowing the network structure (i.e. network properties) does not always allow to know the list of interactions (and vice versa), therefore justifying the need to sample interactions as their own biological entities. A central challenge to do so is that the sampling effort required to document interactions is much higher than for species (e.g. 500% vs 64% in Chacoff et al. 2012). Moreover, evaluating sampling completeness for interactions is more complex than for species, as it requires assessing the proportion of forbidden links (i.e. links that cannot occur) due to life history or morphological restrictions, which in turn requires knowledge of species natural history and traits (Jordano 2016b). While some quantitative network properties such as connectance are less affected by sampling completeness, network composition and binary measurements derived from it will vary widely if interactions are missed (e.g. number of links, Poisot et al. 2012, Vizentin-Bugoni et al. 2016).

One way to define expectations for interaction monitoring within BONs is to consider variation in local sampling from an explicitly spatial perspective—that is, quantifying sampling effort from specific sites where interactions are sampled—and use such expectations to improve monitoring efficiency. Previous evaluation of sampling effort relating to interactions were not explicitly spatial in a way that directly ties to biodiversity monitoring: they investigated expected variations using dataset resampling (to measure the effect on network composition and properties, Poisot et al. 2012, Henriksen et al. 2019), focused on the effect of scale (showing that many network structure metrics are robust across spatial scales, Wood et al. 2015), or assessed the ability to capture regional properties (showing that sampling larger to smaller lakes better captures regional metaweb properties, McLeod et al. 2021). We now have frameworks and models to describe and investigate the variation and detection of interaction networks in space and time (Poisot et al. 2015, Cirtwill et al. 2019, Catchen et al. 2023), and therefore the next step is to consider how to optimize monitoring for species interactions. Simulations offer opportunities to explore such optimizations: for example, to anticipate sampling effects on estimated network properties compared to a known reference (Aguiar et al. 2019), determine optimal sampling frequency for monitoring (Daugaard et al. 2024), and examine how monitoring efficiency and trends detection is influenced by the number of monitoring sites under a range of scenarios and monitoring schemes (Ficetola et al. 2018).

In this manuscript, we address three key questions. First, what expectations should we have for monitoring species interactions across space within Biodiversity Observation Networks? Second, what effect does sampling design strategies have on the efficiency with which we can document interactions? Finally, what available information (e.g. from existing monitoring programs) could be used to optimize the design of future BONs for interactions and how precise should such information be to expect efficiency improvements?

We use spatially explicit community simulations to test the efficiency of monitoring species interactions within BONs, and explore various algorithmic and design options for network sampling optimization. We aim to define realistic expectations for monitoring species interactions, in comparison to more commonly evaluated elements such as species richness and species ranges. We examine how to assess and compare monitoring efficiency using simulations. We conclude with recommendations on how this should influence the design of BONs to improve the monitoring of species interactions as an important biodiversity component.

Methods

We used simulations to generate species interaction networks, species ranges, and biodiversity observations networks (BONs) with the goal of evaluating monitoring potential for interaction networks within BONs. We first detail the steps for the general model we used (Figure 1). We then go over the simulations we performed to investigate the monitoring of species interactions within BONs, aiming to 1) establish baseline expectations for the monitoring of interactions; 2) assess opportunities to improve the monitoring efficiency using different site selection algorithms and optimization targets; and 3) evaluate how precisely we need to know a species ranges to use it as a target for monitoring design.

General model

We followed the approach and process-based, spatio-temporal model introduced by [Catchen et al. \(2023\)](#) (building on [Cirtwill et al. 2019](#)) to simulate interaction realization and detection while accounting for spatial variation in species occurrences, implemented in `SpeciesInteractionSamplers.jl`. This model broadly captures the different processes involved into turning a *potential* interaction (i.e. a biologically feasible interaction) into a *realized* one (i.e. an interaction that took place at a specified location), as outlined in [Morales-Castilla et al. \(2015\)](#). To this generative model, we added the design of Biodiversity Observation Networks and the evaluation of sampling efficiency between and across simulations for species interactions (Figure 1).

Step 1: Generating interaction networks with spatial variation

First, we simulated networks with variation in species and interaction composition across a simulated landscape (Figure 1, Step 1), with the intent to replicate the ecological context in which biodiversity monitoring could take place. To do so, we generated species ranges representing the distribution of species across the landscape using neutral landscape models (NLMs, [Gardner et al. 1987](#), [Etherington et al. 2015](#)). NLMs are used to generate landscapes replicating realistic spatially-




Figure 1: Conceptual diagram of the workflow and simulations. We used the spatio-temporal model by [Catchen et al. \(2023\)](#) to generate landscapes with spatial variation in species and interaction composition. We then designed biodiversity observation networks (BONs) across the simulated landscapes using a combination of site selection algorithms (samplers) and optimization targets. We measured the efficiency of the BON designs for sampling species interactions using a comparable measure (the number of sites required to document 80% of the interactions) across all simulations.

autocorrelated distributions and representing useful baselines to explore ecological processes in a spatial context ([Hesselbarth et al. 2024](#)). NLMs have been used to generate realistic landscapes in contexts relating to biodiversity distribution and preservation, for instance to compare avian diversity between conservation approaches in coffee-growing landscapes ([Valente et al. 2022](#)) or to assess the potential of a suggested umbrella species in an ecotone ([Duchardt et al. 2023](#)). NLMs have also been used with species interactions, for example to evaluate the effect of landscape structure in cross-species disease transmission between interacting species ([Forero-Muñoz et al. 2025](#)). Here, we generated species ranges following the implementation in `SpeciesInteractionSamplers.jl` ([Catchen et](#)

al. 2023), which 1) uses the Diamond-Square algorithm (Fournier et al. 1982; also used similarly and described by Forero-Muñoz et al. 2025) to generate a probabilistic spatially-autocorrelated range for a species (default autocorrelation parameter $h=0.85$); 2) draws a threshold based on a Beta distribution (default $\alpha = 10$, $\beta = 10$); 3) converts the probabilistic range into a binary one, considering all sites with values higher than the threshold as presence and those with lower values as absence, and 4) repeating the process independently for all species (here 75 species). In the end, this process generates communities of known composition and species richness at every site in the simulated landscape.

Next, using the spatio-temporal model presented by Catchen et al. (2023), we defined the **feasible interactions** between the species by generating a metaweb using the niche model (Williams and Martinez 2000). The niche model defines interactions by assigning species a niche position and feeding range, similarly to body-size relationships between predator and prey, and generates networks with structural properties similar to empirical food webs (Williams and Martinez 2000, Gravel et al. 2013, Delmas et al. 2019). It is widely used as a generative model, including in the context of investigating interaction sampling (Poisot et al. 2012, Wood et al. 2015).

Combining the species ranges and the metaweb, we verified where species with feasible interactions co-occur, to build a list of **possible interactions** (Catchen et al. 2023) at every location in our landscape. Next, we defined **realized interactions** by conditioning possible interactions to species abundances. Rare and low-abundance species are less likely to interact in a location because they are less likely to encounter one another when compared to abundant species, leading to neutrally forbidden links (Canard et al. 2012, Catchen et al. 2023). The neutral model introduced by Catchen et al. (2023) first generates a distribution of species relative abundances from a log-normal distribution (followed by many empirical communities, Preston 1948; the variance is controlled by a parameter σ). It then defines an interaction rate between species pairs based on the product of their relative abundances, scaled by a fixed realization energy parameter (ϵ) determining the total expected number of interactions. Finally, they draw realized interactions from a Poisson distribution parametrized on the interaction rate (see Catchen et al. 2023 for full details).

Finally, we defined the set of **detected interactions** by also conditioning the detection of realized interactions on species abundances. Even if an interaction is realized in a location, sampling must occur at the moment of interaction for its detection; this is again less likely for low probability events, like interactions between rare occurring species. Catchen et al. (2023) first establish a per-species detection probability conditioned on relative abundance and a fixed scaling parameter (α), compute the interaction-level detection rate as the product of the per-species probability, then draws the detected interactions from a Binomial distribution parametrized on the number of realized interactions and the interaction detection rates.

The result of the general model is a **landscape configuration** with spatial variation in species and interaction composition and detection. Species composition, realized interactions, and detected interactions are based on the process model and known at all locations, allowing to compare the efficacy and efficiency of sampling strategies at retrieving interactions within the landscape configuration.

Step 2: Designing Biodiversity Observation Networks

We designed **Biodiversity Observation Networks (BONs)** across the simulated landscape and measured their efficiency at monitoring the generated species interaction networks. The objective of this step is to select the sites that will form the monitoring network—a BON being a monitoring network for biodiversity, and for which we here focus specifically on species interactions—similar to BON examples with systematic grid designs at national scales (e.g. China BON, [Xu et al. 2017](#), [Yi et al. 2022](#)) and subnational ones (Québec and Alberta in Canada, [Burton et al. 2014](#), [Simard 2024](#)). [Norman and Poisot \(2025\)](#) presented an overview of the design and site selection process accounting for ecological monitoring goals. The design process uses a site selection algorithm to designate candidate sites for the monitoring network. Key features of the site selection algorithms include the ability to balance sites across space for equal representation (see [Benedetti et al. 2017](#), [Kermorvant et al. 2019](#) for reviews), to stratify across regions or populations, to include auxiliary site information (environmental or biological) to weight inclusion probability, and to produce master samples including current monitoring site or from which to select a subset of sites for a specific monitoring project ([Dam-Bates et al. 2018](#), [Robertson et al. 2024](#)).

Based on the landscape configurations generated in Step 1, we can use **site selection algorithms** (generally referred to as *samplers*) to distribute monitoring sites across the landscape, then measure each design's efficiency at monitoring species interactions. Some algorithms can optimize the location of sites given different information such as species ranges, species richness or presence of realized interactions. For example, we can design BONs using the Balanced Acceptance Sampling algorithm ([Robertson et al. 2013](#)) when the aim is to balance sites across our simulated landscapes, given its computational efficiency and its ability to weight site inclusion probabilities using a layer of weights. On the other hand, active learning methods like the Uncertainty Sampling algorithm ([Lewis and Gale 1994](#)) can be used to optimize sampling given information on element uncertainty ([Settles 2009](#)). Instead of focusing on uncertainty, we can apply the algorithm to target the sites that provide the highest information gain for the BON, given a layer of 'information' values to maximize. Examples of active learning relating to biodiversity monitoring and sampling site selection include improving species range estimation from a smaller number of actively selected sites ([Lange et al. 2023](#)) and minimizing field data collection costs based on distances between samples ([Malek et al. 2019](#)).

Step 3: Defining a comparable measure for sampling efficiency

As interaction composition is known at all locations in our simulated landscapes, we can determine the **proportion of interactions** (noted p) of the focal species that can be sampled for every generated BON design (e.g. a BON of 50 sites using a specific site selection algorithm). This proportion can be seen as a 'snapshot' view of a given BON design's sampling efficiency for that simulated landscape. The proportion of sampled interactions should increase with the number of sites in the BON (n), although the exact value of p can fluctuate given randomness in the site selection algorithm. To remove this randomness and ensure a consistent comparison across independent simulations, we state that the **proportion of sampled interactions for a given number of sites in the BON**, noted $p(n)$, follows an accumulation curve described by the equation:

$$p(n) = \frac{n}{a + n} \quad (1)$$

where a is the single parameter describing the shape of the sampling curve. The lower the value for a , the faster we reach a high proportion of monitored interactions. We can identify the value of a yielding the curve with the best fit with a simulation's results, as described in Simulation Study II.

Next, we derived an interpretable efficiency measure based on Equation 1. Although the parameter a can be directly compared across samplers, optimization targets and independent simulations, its value is not intuitive in a monitoring context. Instead, we used the **number of sites required to sample 80% of the interactions**, which can directly be tied to sampling effort and design. Solving Equation 1 for n , we obtain:

$$n_p(a) = \frac{p \cdot a}{1 - p} \quad (2)$$

where $n_p(a)$ represents the number of sites required to sample a proportion of interactions p , following an efficiency curve described by the parameter a . Here, we fixed $p = 0.80$ (80% of the interactions), an arbitrary threshold representing a highly effective monitoring outcome, for which we obtain $n_{0.80} = 4 \cdot a$ using Equation 2. $n_{0.80}$ measures the speed at which interactions are documented, and therefore the sampling efficiency of a BON design within a given landscape. We then used $n_{0.80}$ values to compare site selection algorithms and optimization targets to identify which ones are more efficient—i.e. allow reaching a high proportion of monitored interactions more quickly.

Simulation studies

We performed three simulation studies to investigate the efficiency of monitoring species interactions within BONs. Across our simulations, we used landscapes of 100 x 100 pixels, 75 species, and a connectance of 0.2 to generate metawebs using the niche model. The connectance and number of species are within the range of values of well-characterized empirical food webs (Williams and Martinez 2000; Dunne et al. 2002, 2004; Smith-Ramesh et al. 2017) commonly used for reference (Williams and Martinez 2008, Baiser et al. 2010, Wood et al. 2015). These parameters are also within the range of previous simulation studies on spatial ecological network dynamics, e.g. Thompson and Gonzalez (2017). We used the following parameters for the spatio-temporal model implemented in SpeciesInteractionSamplers.jl (Catchen et al. 2023): landscape autocorrelation of $h = 0.85$, Beta distribution with $\alpha = 10$ and $\beta = 10$ to threshold landscapes, variance of log-normal distribution for relative abundances $\sigma = 1.2$, realization energy $\epsilon = 50,000$, and detection scaling $\alpha = 50$. Exploratory simulations showed that model parameters such as metaweb connectance and landscape autocorrelation did not have an important effect on the proportion of sampled interactions and the accumulation observed when increasing the number of sites in the BON. The activation energy ϵ and detection scaling α parameters determine the total number of expected and detected interactions (Catchen et al. 2023), and therefore influenced the absolute numbers observed. Here, we chose to use a single fixed value to focus on differences between sampling strategies— influenced in the same way by the parameter values—and refer readers to Catchen et al. (2023) for the detailed discussion over the parameters' effect. We implemented our simulations in Julia v1.11.9 (Bezanson et al. 2017). All the code used to implement the simulations and perform the analyses is

archived on Zenodo ([Dansereau 2026](#)). Full computations were run on compute clusters provided by Calcul Québec and the Digital Research Alliance of Canada.

Simulation study I: Establishing baseline expectations for the monitoring of interactions

Our first simulation study aimed to establish baselines for the monitoring of interactions across space, defining expectations for how we could document a metaweb of interactions using a BON. We aimed to represent the case of existing BONs (e.g. Québec, China) designed to cover a wide range of taxa and environmental conditions without specific considerations for species interactions. We tested the case of spatially balanced BONs with evenly spread sites across the landscape configurations, which is generally considered an effective way to cover environmental space ([Kermorvant et al. 2019](#)). We used the Balanced Acceptance Sampling algorithm ([Robertson et al. 2013, 2017](#)) to generate candidate BONs with a number of sites between 1 and 500 at intervals of 10 sites, with 50 replicates for each number of sites, yielding 2,550 BON designs in total. For every design, we used the general model (Step 1) to generate an independent landscape configuration: a specific metaweb, a set of species ranges, an abundance distribution for the species, and realized interactions. In every one, we evaluated the sampling efficiency of the BON design in terms of sampling interactions from the metaweb and in terms of documenting species richness. For species richness, we assumed that all species present at sites in the BON were sampled, then compared the number of species sampled in the BON against the total species richness across the landscape (always 75 species). For interactions, we evaluated each interaction type separately (possible, realized and detected), counting the number of distinct interactions sampled in the BON, then evaluated each number in proportion to the metaweb (specific to the landscape configuration, but generated with the same connectance of 0.2 using the niche model). As a whole, these simulations compared four sampling targets (species richness, possible interactions, realized interactions, and detected interactions), establishing baselines for the monitoring of each.

Simulation study II: Evaluating strategies to optimize the monitoring of interactions for a focal species

Interaction network sampling and monitoring may be targeted at a focal species because of its endangered status (e.g. [Rioux et al. 2022](#), because it is at the centre of conservation or restoration programmes (e.g. BearConnect), or even due to human-nature conflicts (e.g. [Baranowska et al. 2025](#)). Hence, we also investigated sampling efficiency from a focal species perspective, by simulating a virtual observer that would aim to sample all the interactions of a focal species (here known from the metaweb), but might have different information on how to optimize the sampling design. Thus, the idea of our focal sampling exploration is to represent a simplified case with a clear objective (retrieving a species' interactions) and where we do have prior knowledge about the system, either through empirical data or model predictions, which can be used to optimize sampling.

Optimizing focal sampling relies on two elements: the site selection algorithm (sampler) used and the information provided for optimization. A species range represents an attainable level of information an interested observer could have, for instance based on expert knowledge or species distribution models. Therefore, we investigated the case where an observer knows the species range across the landscape (presence or absence) and tries to optimize the BON design to retrieve the interactions. We tested four strategies for comparison: a *Balanced Sampling* spreading sites across

the entire study area (using Balanced Acceptance Sampling, hereafter BAS), a balanced sampling only within the range of the focal species (hereafter called *Balanced Within Range*), a *Weighted Sampling* favouring the species range in site selection among all sites in the study area (adjusting the inclusion probability in BAS), and a *Targeted Sampling* selecting sites with higher ‘information’ values in the layer provided (using Uncertainty Sampling). Next, we tested four information targets for optimization with the Targeted Sampling strategy (all with the Uncertainty Sampling algorithm): the exact *focal species range* (our attainable information knowledge for the observer), a *probabilistic range* for the focal species (representing a less-informed case, similar to the output of a species distribution model), *species richness* across the landscape (to represent a holistic level of information about the system, which an observer might have based on detailed knowledge about biodiversity in a region) and finally the location of *realized interactions* for the focal species across the landscape (to represent a best-case scenario, where an observer would know exactly the right information to target).

We performed 200 independent simulations with independent landscape configurations (as described in Simulation study I). In each simulation, we selected the species with the highest degree (number of interactions) in the metaweb as the focal species for the simulation (as it was previously shown as an efficient strategy for sampling and conservation, [Pires et al. 2017](#), [Aguiar et al. 2019](#)). We then generated BONs of 1 to 500 sites (at interval increments of 10 sites) using the three balanced strategies (Balanced Sampling, Balanced Within Range, Weighted Sampling) and four optimization targets for the Targeted Sampling strategy (focal species range, probabilistic range, species richness, realized interactions), with 50 replicates for every number of sites. For each BON generated, we extracted the number of unique realized interactions sampled across all sites for the focal species. We then performed a grid search across possible values of a (from Equation 1) to identify the value yielding the curve with the best fit to the simulation’s results. To do so, we extracted the median result (proportion of sampled interactions) across replicates for every number of sites in the BONs. We then evaluated 10,000 evenly spaced values for a between e^{-5} and e^{15} and selected the value yielding the curve minimizing the sum of squared errors to the simulation results. We searched for values of a on an exponential scale as it allowed for more gradual control over the curve shape and adjusted the limits to ensure they went beyond the fitted values on both the lower and higher side. After selecting the best-fit value for a , we evaluated the sampling efficiency for the simulation using the $n_{0.80}$ measurement described in Step 3 (number of sites required to document 80% of the interactions).

Finally, we defined a confidence interval for our $n_{0.80}$ efficiency measurement. As the best-fit for a is based on the median result across replicates, we used the confidence interval of the median to define a lower and upper bound for values of a . We calculated the order statistics delimiting an 90% equal-tailed interval using the normal approximation to the binomial distribution ([Gibbons and Chakraborti 2010](#))—e.g., at $n=50$, the 90% confidence interval for the median goes from the 19th and to the 31st value—and extracted the corresponding results from our simulations. We performed separate grid searches for the parameter a using the lower and upper confidence intervals bounds of the median, thus delimiting the lower and upper bounds for a , and subsequently bounds for $n_{0.80}$. This approach returns a confidence interval for the sampling efficiency, which we used to compare

site selection algorithms and optimization targets within a specific landscape configuration, given that the monitoring context is unique in each one (different metawebs and species ranges). Doing so, we identified which site selection algorithms or optimization target were equal and which were more efficient (thus quickly reaching a high proportion of monitored interactions), while also quantifying the magnitude the difference between strategies in a comparable way between independent simulations.

Simulation study III: Assessing the precision required when estimating a species range

Building on the Balanced Within Range strategy from Simulation study II, we assessed how precisely an observer should know the focal species range to achieve an efficient monitoring. To do so, we compared the Balanced Within Range strategy, where the balanced sampling design is only generated across the exact focal species' range, with alternative scenarios where we over- and underestimated the species range for the BON design. We tested estimation offsets from 0 to 50% of the true range size, at 2% increment intervals, generating balanced designs within the over- or underestimated range. For a given landscape configuration, we extracted the true range size, calculated the number of sites to add or remove to reach the offset percentage, then used the non-thresholded probabilistic range (Step 1) to add the absence sites closest to the threshold (for overestimation) or remove the presence sites closest to the threshold (for underestimation).

We followed the same approach as in Simulation study II to design BONs and measure efficiency using the same 200 simulations with independent landscape configurations. We generated BONs of 1 to 500 sites for all the estimation offset ranges (again with 50 replicates at all number of sites) and measured the efficiency at documenting interactions. We then compared the $n_{0.80}$ values of each estimation offset range to the $n_{0.80}$ for the true species range, which we expected would be the most efficient, and used the confidence intervals to determine which estimation offsets differed from the true range.

Results

Simulation study I: Lower sampling expectations for species interactions than species richness

Our results for the baseline spatially-balanced sampling designs highlight a marked difference between the proportions of sampled species and interactions (Figure 2). Sampled species saturated very quickly, with all species sampled with less than 25 sites in a BON across the landscape. Possible interactions, which only depend on co-occurrence of species with a feasible interaction in the metaweb, also saturated quickly reaching over 90% of interactions in the metaweb with 25 sites and increased asymptotically afterwards. In contrast, realized and detected interactions, which account for the impact of species abundances on interaction realization and detectability, reached much lower proportions, both remaining under 25% of the all metaweb interactions at 100 sampling sites and under 50% at 500 sites. While the proportion of sampled realized and detected interactions would keep increasing with a higher number of sites ($n_{0.80}$ around 2100 and 7300, respectively), the speed at which they accumulate is notably lower than for species richness or possible interactions,




Figure 2: Proportion of sampled species and interactions for a given number of sampling sites in a spatially-balanced Biodiversity Observation Network (BON) across the simulated landscape (Simulation Study I). Proportions were calculated with respect to the total species richness and the total number of interactions in the metaweb. Lines indicate the median proportion for a given number of sites, while the shaded bands delimit the 5th and 95th percentiles across replicates (50 independent replicates per number of sites; 2,550 in total).

highlighting the need for a much higher sampling effort for interactions within spatially-balanced BONs.

Simulation study II: Higher efficiency for Targeted Sampling and Balanced Within Range compared to spatially balanced strategies

To better illustrate the results of Simulation Study II, we first showcase the results for a representative landscape configuration (metaweb, species ranges, realized interactions). The conclusions for this landscape configuration are consistent with the most common results across all simulations (200 independent simulations with different landscape configurations), which we summarize right after the showcased one.

Our focal sampling experiment showed the Balanced Within Range and Targeted Sampling strategy to be most efficient at optimizing BON-design based on a known species range and while aiming to maximize the sampling of interactions of a single focal species (Figure 3). Balanced sampling, representing a spatially balanced design across space unrelated to the focal species (similar to the baseline in Simulation Study I), performed the worst among the algorithms tested. Weighted Sampling maintained a better spatial balance across the landscape but performed notably worse

than Targeted Sampling and Balanced Within Range at maximizing the efficiency for the focal species. Targeted Sampling and Balanced Within Range performed similarly.

Optimizing on the known species range, our proxy for an attainable level of information an interested user might have, was more efficient at sampling the focal species' interactions than optimizing based on the probabilistic species range or on species richness, effectively sampling more than half of the species' interactions (Figure 4). In contrast, optimization based on the location of realized interactions (our best-case scenario requiring a higher level information) expectedly performed better, retrieving most of the species' interactions.




Figure 3: Efficiency of site selection strategies at sampling realized interactions for a given focal species within Biodiversity Observation Networks (BONs). This figure compares four sampling strategies: three balanced strategies (across the entire area, across the focal species range only, and using weights to favour the species range) and a targeted strategy using the species ranges as a fixed optimization target (the species range). Solid lines indicate the median proportion for a given number of sites (50 replicates per number of sites), while the shaded bands delimit the 5th and 95th percentile across replicates. The panels on the right illustrate examples of BONs with 50 sites (coloured points) generated by each strategy over the same landscape (species presences in yellow, absence in purple, masked area in white). This showcased example is representative of the most common results across independent simulations.




Figure 4: Efficiency of optimization targets at sampling realized interactions for a given focal species within Biodiversity Observation Networks (BONs). This figure compares four optimization targets with a fixed targeted sampler (Uncertainty Sampling). Solid lines indicate the median proportion for a given number of sites, while the shaded bands delimit the 5th and 95th percentile across replicates. The panels on the right illustrate examples of BONs with 50 sites (coloured points) generated by each strategy over the target landscape (areas in yellow indicate a higher value for targeting, e.g. presence or richness, while areas in purple indicate a lower value).

Comparing optimization options by simulation clearly highlighted a consistently more efficient approach across landscape and metaweb contexts (Figure 5; see also Supp. Mat. for all pair-by-pair comparisons), consistent with the showcased single-simulation example (Figure 3, Figure 4). Targeted Sampling was more efficient than Weighted Sampling in 63% of simulations (lower value for $n_{0.80}$, indicating a faster documentation of interactions), and both were more efficient than Balanced Acceptance in all simulations (Figure 5A). Targeted Sampling and Balanced Within Range were equally efficient in all cases. Among optimization targets used with Targeted Sampling (Figure 5B), optimizing on the focal species range was more efficient than on the probabilistic range or on species richness (78% and 90%, respectively), but always less efficient than optimizing on realized interactions.





Figure 5: Within-simulation comparison of efficiency between samplers and optimization targets for 200 independent simulations. The comparison value is based on the number of sites required to document 80% of the focal species' interactions ($n_{0.80}$), described in Equation 2. Negative values (in orange) indicate that the compared option led to a more efficient sampling than its reference (faster documentation of interactions), while positive values (in green) indicate it required a higher sampling effort. Equal values (in pink) have overlapping confidence intervals. The reference is the same on the two subpanels (Targeted Sampling using the focal species range) and was chosen to simplify the number of comparisons displayed. All one-by-one comparisons are available in Supp. Mat.

Simulation Study III: Tolerance to 10% over- and underestimation of the focal species range

Our results highlight a tolerance in the precision of the species range estimation required for the Balanced Within Range strategy. Overestimating the range size from 0 to 50% led to a linear increase in the value of $n_{0.80}$ compared to the $n_{0.80}$ value obtained using the Balanced Within Range strategy on the exact species range (Figure 6A, right side). The $n_{0.80}$ was equal to the true range until a 10% overestimation offset (based on overlapping confidence intervals), at which point it started to be higher across all simulations, indicating a lesser efficiency (Figure 6B). In contrast, underestimating the species range showed a lesser increase of $n_{0.80}$ from 0 to -50% based on the median result across simulations, with a wider range of possible outcomes (Figure 6A, left side). Importantly, while the $n_{0.80}$ was also equal until a -10% offset and started to decrease beyond, it stayed equal in the majority of simulations (> 70%) even for underestimation of -50% (Figure 6B). In the remaining simulations (< 30%), the $n_{0.80}$ was higher than for the true range (less efficient), except for a few cases where it was lower (more efficient).

a) Comparison of efficiencies between range estimations



b) Summary of comparisons across all simulations




Figure 6: Comparison of efficiencies when over- and underestimating the focal species range for 200 independent simulations. Results are compared to the Balanced Within Range strategy applied on the exact species range, whose $n_{0.80}$ value is used as reference (specific to every simulation). In panel A, we use the over- and underestimation $n_{0.80}$ in proportion to the reference $n_{0.80}$ (y axis) given the range estimation offset (x axis) to highlight the linear trend and association with the offset percentage for overestimation. The dotted line at 1.00 indicates exactly equal $n_{0.80}$. In panel B, we summarize comparisons across all simulations to display the proportion of each comparison outcome (equal, higher, lower). The solid line represents the actual proportion value, while the shaded bands delimit the 90% confidence intervals for the proportion calculated using the Wilson interval method (Brown et al. 2001, implemented in HypothesisTests.jl).

Discussion

What should be realistic expectations for sampling and monitoring of species interactions?

Our simulations highlight that expectations for monitoring species interactions within spatially-balanced, general-purpose BONs should be much lower than for species richness, especially when considering that interaction realization and detection both depend on species abundance (Figure 2). In other words, we can expect the monitoring of species interactions within a BON to require a (much) higher monitoring effort across space than the one required for monitoring species and community composition. Our observations are consistent with previous ones regarding higher sampling effort required to document pollinator networks (Chacoff et al. 2012), yet here framed in the context of BON design and monitoring across space. Overall, these results reinforce the need for optimizing sampling designs for species interactions beyond and in complement to existing monitoring networks.

We can improve interaction coverage with the right optimization method, but efficiently doing so requires a clear and attainable monitoring objective. Adopting a single-species perspective (e.g. focusing on interactions for a keystone or high degree species, Pires et al. 2017, Aguiar et al. 2019) can yield relevant results while relying on attainable levels of information (Figures 3, 4, 5). Through our representative example in Figure 3, we showed how Targeted Sampling and Balanced Within Range outperform spatially-balanced designs to monitor interactions for a focal species, while only relying on an attainable level of information (the species range). Two emerging questions that follow up on this result are: 1) what information is most efficient to target sites in sampling designs; and 2) how precisely do we need to estimate such information to expect efficiency improvements? Our simulation studies offer insights into both, as we detail next before concluding with recommendations for monitoring.

What strategies and available information are most efficient to optimize monitoring for species interactions?

Ecologically-relevant metrics are required to evaluate the performance of monitoring designs, beyond spatial balance (Norman and Poisot 2025). While many sampling algorithms can perform similarly for biodiversity monitoring in certain contexts (Norman and Poisot 2025), our results highlighted better options when we can define a specific target for monitoring (Figure 5). Optimizing on a known species range proved an efficient strategy to retrieve its interactions that was better than optimizing based on more holistic, community-level information such as species richness (Figures 4, 5). This contrasts with earlier accounts where designs based on maximising species richness also proved efficient to conserve multiple biodiversity facets (Willig et al. 2023). Instead, our results highlight that more information about an entire system does not always lead to more efficient monitoring, and that efforts could instead be put towards identifying relevant targets with available information to guide monitoring design.

Optimizing based on the location of the realized interactions was, as expected, even more efficient than targeting the species range (Figures 4, 5). Yet, this represents a best case scenario that is likely beyond reach in an applied context: it assumes that there is already pre-existing knowledge on where species interactions can be observed, which is unlikely to exist for most interactions, species and locations across the globe, regardless of their relevance for monitoring. Nonetheless, it highlights that the most important efficiency gains will come from better knowledge of where interactions are likely to take place. On one hand, better incorporating factors influencing interaction realization and detection, such as species abundance in our simulations, will be crucial and potentially more impactful than improving knowledge or reducing uncertainty over species ranges (Figures 4, 5). Integrating abundances into network evaluations could be based on species regional abundances (Reji Chacko et al. 2025) and would echo similar calls for integration of abundances and processes into conservation (Dornelas et al. 2023, Fletcher Jr. et al. 2025, Tobias et al. 2025). On the other hand, our result also suggest that monitoring will become more efficient over time as we keep gathering knowledge on interactions, especially if local knowledge of the system and species under focus is incorporated. Such opportunities should emerge in recent BON initiatives integrating Indigenous and local community knowledge for an inclusive view of biodiversity change and recognizing multiple ways of knowing (Gonzalez et al. 2025). Overall, we should not wait for perfect knowledge: establishing operational monitoring frameworks to inform biodiversity conservation and management is possible as we keep gathering new information, and better ensures we translate knowledge into action (Buxton et al. 2021, Dansereau et al. 2025).

How precise should estimations be to expect improvements in monitoring efficiency?

A similarly-efficient alternative to targeted sampling is to follow an approach such as our Balanced Within Range strategy, where we generate a spatially balanced design across the area believed to be more informative. Our results from Simulation Study III are encouraging in this sense, as they show we do not need perfect knowledge when using this strategy (Figure 6). The tolerance we observed to 10% over- and underestimation of the focal species range size offers an important margin that changes the information we can use to guide BON design. For instance, our test case of underestimating a species range highlights that we could focus mainly on the most certain parts of a species range prediction (e.g. using conformal predictions along with species distribution models, Poisot 2024) and expect similar monitoring efficiency. On the opposite side, in terms of range size, our overestimation test case is more analogous to a model focusing on the potential distribution rather than the realized distribution.

Recommendations for monitoring

Documenting species interactions extensively requires higher sampling efforts than required to document species ranges. While spatially balanced designs are efficient in general purpose biodiversity monitoring, species interactions may be better served by specifically-optimized monitoring designs. To do so, we show the importance of selecting an appropriate target (e.g. a focal species' interaction) and using attainable information restricting the phenomenon (here the species range) to optimize monitoring designs efficiently. There is a degree of tolerance in the level of exact

information needed (similar to range over- and underestimation), therefore available information should be used as soon as possible to guide monitoring.

Acknowledgements

GD was funded by the NSERC Postgraduate Scholarship – Doctoral (grant ES D – 558643) and the FRQNT doctoral scholarship (grant no. 301750). TP was funded through award no. 223764/Z/21/Z from the Wellcome Trust, by the Discovery Grant program of NSERC, through a donation from the Courtois Foundation. MDC is funded by an IVADO Postdoctoral Fellowship. We thank the members of the *Laboratoire d'Écologie Prédicative et Interprétable pour la Crise de la Biodiversité* in Montréal, QC, and of the Predictive Ecology team at the *Pacific Forestry Centre* in Victoria, BC, for their constructive comments during discussions of this manuscript.

References

- Abrego N, Roslin T, Huotari T, et al (2021) Accounting for Species Interactions Is Necessary for Predicting How Arctic Arthropod Communities Respond to Climate Change. *Ecography* 44:885–896. <https://doi.org/10.1111/ecog.05547>
- Aguiar LMS, Diniz UM, Bueno-Rocha ID, et al (2024) Untangling Biodiversity Interactions: A Meta Network on Pollination in Earth's Most Diverse Tropical Savanna. *Ecology and Evolution* 14:e11094. <https://doi.org/10.1002/ece3.11094>
- Aguiar MAM de, Newman EA, Pires MM, et al (2019) Revealing Biases in the Sampling of Ecological Interaction Networks. *PeerJ* 7:e7566. <https://doi.org/10.7717/peerj.7566>
- Baiser B, Gravel D, Cirtwill AR, et al (2019) Ecogeographical Rules and the Macroecology of Food Webs. *Global Ecology and Biogeography* 28:12925. <https://doi.org/10.1111/geb.12925>
- Baiser B, Russell GJ, Lockwood JL (2010) Connectance Determines Invasion Success via Trophic Interactions in Model Food Webs. *Oikos* 119:1970–1976. <https://doi.org/10.1111/j.1600-0706.2010.18557.x>
- Baranowska W, Bartoszewicz M, Nowak S, et al (2025) Low Contribution of Livestock in the Grey Wolf Diet in the Area with High Availability of Free-Ranging Cattle and Horses. *European Journal of Wildlife Research* 71:48. <https://doi.org/10.1007/s10344-025-01926-3>
- Benedetti R, Piersimoni F, Postiglione P (2017) Spatially Balanced Sampling: A Review and A Reappraisal. *International Statistical Review* 85:439–454. <https://doi.org/10.1111/insr.12216>
- Bezanson J, Edelman A, Karpinski S, Shah VB (2017) Julia: A Fresh Approach to Numerical Computing. *SIAM Review* 59:65–98. <https://doi.org/10.1137/141000671>
- Brown LD, Cai TT, DasGupta A (2001) Interval Estimation for a Binomial Proportion. *Statistical Science* 16:101–133. <https://doi.org/10.1214/ss/1009213286>

569 Burton AC, Huggard D, Bayne E, et al (2014) A Framework for Adaptive Monitoring of the
570 Cumulative Effects of Human Footprint on Biodiversity. *Environmental Monitoring and*
571 *Assessment* 186:3605–3617. <https://doi.org/10.1007/s10661-014-3643-7>

572 Buxton RT, Bennett JR, Reid AJ, et al (2021) Key Information Needs to Move from Knowledge to
573 Action for Biodiversity Conservation in Canada. *Biological Conservation* 256:108983. <https://doi.org/10.1016/j.biocon.2021.108983>

575 Canard E, Mouquet N, Marescot L, et al (2012) Emergence of Structural Patterns in Neutral Trophic
576 Networks. *PLOS ONE* 7:e38295. <https://doi.org/10.1371/journal.pone.0038295>

577 Caron D, Brose U, Lurgi M, et al (2024) Trait-Matching Models Predict Pairwise Interactions across
578 Regions, Not Food Web Properties. *Global Ecology and Biogeography* 33:e13807. <https://doi.org/10.1111/geb.13807>

580 Catchen MD, Poisot T, Pollock LJ, Gonzalez A (2023) The Missing Link: Discerning True from False
581 Negatives When Sampling Species Interaction Networks. <https://ecoevorxiv.org/repository/view/4929/>. Accessed 19 Jan 2023

583 CBD (2022) Decision Adopted by the Conference of the Parties to the Convention on Biological
584 Diversity 15/4. Kunming-montreal Global Biodiversity Framework

585 Chacoff NP, Vázquez DP, Lomáscolo SB, et al (2012) Evaluating Sampling Completeness in a Desert
586 Plant–Pollinator Network. *Journal of Animal Ecology* 81:190–200. <https://doi.org/10.1111/j.1365-2656.2011.01883.x>

588 Chiu C-H, Chao A, Vogel S, et al (2023) Quantifying and Estimating Ecological Network Diversity
589 Based on Incomplete Sampling Data. *Philosophical Transactions of the Royal Society B: Biological Sciences* 378:20220183. <https://doi.org/10.1098/rstb.2022.0183>

591 Cirtwill AR, Eklöf A, Roslin T, et al (2019) A Quantitative Framework for Investigating the
592 Reliability of Empirical Network Construction. *Methods in Ecology and Evolution* 10:902–911. <https://doi.org/10.1111/2041-210X.13180>

594 Dam-Bates P van, Gansell O, Robertson B (2018) Using Balanced Acceptance Sampling as a Master
595 Sample for Environmental Surveys. *Methods in Ecology and Evolution* 9:1718–1726. <https://doi.org/10.1111/2041-210X.13003>

597 Dansereau G (2026) Gabrieldansereau/NetworkMonitoring: V0.1.0. <https://zenodo.org/doi/10.5281/zenodo.18484358>. Accessed 4 Feb 2026

599 Dansereau G, Barros C, Poisot T (2024) Spatially Explicit Predictions of Food Web Structure from
600 Regional-Level Data. *Philosophical Transactions of the Royal Society B: Biological Sciences* 379:20230166. <https://doi.org/10.1098/rstb.2023.0166>

602 Dansereau G, Braga J, Ficetola GF, et al (2025) Overcoming the Disconnect between Species
 603 Interaction Networks and Biodiversity Conservation. *Trends in Ecology & Evolution* 40:840–851.
 604 <https://doi.org/10.1016/j.tree.2025.05.010>

605 Daugaard U, Merkli S, Merz E, et al (2024) The Dependence of Forecasts on Sampling Frequency as a
 606 Guide to Optimizing Monitoring in Community Ecology. *Ecosphere* 15:e4786. [https://doi.org/10.](https://doi.org/10.1002/ecs2.4786)
 607 [1002/ecs2.4786](https://doi.org/10.1002/ecs2.4786)

608 Delmas E, Besson M, Brice M-H, et al (2019) Analysing Ecological Networks of Species Interactions.
 609 *Biological Reviews* 94:16–36. <https://doi.org/10.1111/brv.12433>

610 Dornelas M, Chase JM, Gotelli NJ, et al (2023) Looking Back on Biodiversity Change: Lessons for the
 611 Road Ahead. *Philosophical Transactions of the Royal Society B: Biological Sciences*
 612 378:20220199. <https://doi.org/10.1098/rstb.2022.0199>

613 Doré M, Fontaine C, Thébault E (2021) Relative Effects of Anthropogenic Pressures, Climate, and
 614 Sampling Design on the Structure of Pollination Networks at the Global Scale. *Global Change*
 615 *Biology* 27:1266–1280. <https://doi.org/10.1111/gcb.15474>

616 Duchardt CJ, Monroe AP, Edmunds DR, et al (2023) Using Neutral Landscape Models to Evaluate the
 617 Umbrella Species Concept in an Ecotone. *Landscape Ecology* 38:1447–1462. [https://doi.org/10.](https://doi.org/10.1007/s10980-022-01586-7)
 618 [1007/s10980-022-01586-7](https://doi.org/10.1007/s10980-022-01586-7)

619 Dunne JA, Williams RJ, Martinez ND (2002) Network Structure and Biodiversity Loss in Food Webs:
 620 Robustness Increases with Connectance. *Ecology Letters* 5:558–567. [https://doi.org/10.1046/j.](https://doi.org/10.1046/j.1461-0248.2002.00354.x)
 621 [1461-0248.2002.00354.x](https://doi.org/10.1046/j.1461-0248.2002.00354.x)

622 Dunne JA, Williams RJ, Martinez, Neo D. (2004) Network Structure and Robustness of Marine Food
 623 Webs. *Marine Ecology Progress Series* 273:291–302. <https://doi.org/10.3354/meps273291>

624 Etherington TR, Holland EP, O'Sullivan D (2015) NLMpy: A Python Software Package for the
 625 Creation of Neutral Landscape Models within a General Numerical Framework. *Methods in*
 626 *Ecology and Evolution* 6:164–168. <https://doi.org/10.1111/2041-210X.12308>

627 Ficetola GF, Romano A, Salvidio S, Sindaco R (2018) Optimizing Monitoring Schemes to Detect
 628 Trends in Abundance over Broad Scales. *Animal Conservation* 21:221–231. [https://doi.org/10.](https://doi.org/10.1111/acv.12356)
 629 [1111/acv.12356](https://doi.org/10.1111/acv.12356)

630 Fletcher Jr. RJ, Green RE, Bladon EK, et al (2025) Beyond Species Richness for Biological
 631 Conservation. *Conservation Letters* 18:e13124. <https://doi.org/10.1111/conl.13124>

632 Forero-Muñoz NR, Dansereau G, Viard F, et al (2025) Spatial Landscape Structure Influences Cross-
 633 Species Transmission in a Rabies-like Virus Model. *Microorganisms* 13:416. [https://doi.org/10.](https://doi.org/10.3390/microorganisms13020416)
 634 [3390/microorganisms13020416](https://doi.org/10.3390/microorganisms13020416)

635 Fournier A, Fussell D, Carpenter L (1982) Computer Rendering of Stochastic Models. *Commun ACM*
 636 25:371–384. <https://doi.org/10.1145/358523.358553>

637 Gardner RH, Milne BT, Turnei MG, O'Neill RV (1987) Neutral Models for the Analysis of Broad-Scale
638 Landscape Pattern. *Landscape Ecology* 1:19–28. <https://doi.org/10.1007/BF02275262>

639 Garzke J, Connor SJ, Sommer U, O'Connor MI (2019) Trophic Interactions Modify the Temperature
640 Dependence of Community Biomass and Ecosystem Function. *PLOS Biology* 17:e2006806.
641 <https://doi.org/10.1371/journal.pbio.2006806>

642 Gibbons JD, Chakraborti S (2010) *Nonparametric Statistical Inference*. CRC Press

643 Gonzalez A, O'Connor MI, Bates AE, et al (2025) A Biodiversity Observation Network to Support
644 Conservation Action and Mainstream Knowledge in Canada. *FACETS* 10:1–19. <https://doi.org/10.1139/facets-2024-0206>

646 Gonzalez A, Vihervaara P, Balvanera P, et al (2023) A Global Biodiversity Observing System to Unite
647 Monitoring and Guide Action. *Nature Ecology & Evolution* 1–5. [https://doi.org/10.1038/s41559-](https://doi.org/10.1038/s41559-023-02171-0)
648 [023-02171-0](https://doi.org/10.1038/s41559-023-02171-0)

649 Gravel D, Poisot T, Albouy C, et al (2013) Inferring Food Web Structure from Predator–Prey Body
650 Size Relationships. *Methods in Ecology and Evolution* 4:1083–1090. [https://doi.org/10.1111/2041-](https://doi.org/10.1111/2041-210X.12103)
651 [210X.12103](https://doi.org/10.1111/2041-210X.12103)

652 Harvey E, Gounand I, Ward CL, Altermatt F (2017) Bridging Ecology and Conservation: From
653 Ecological Networks to Ecosystem Function. *Journal of Applied Ecology* 54:371–379. [https://doi.](https://doi.org/10.1111/1365-2664.12769)
654 [org/10.1111/1365-2664.12769](https://doi.org/10.1111/1365-2664.12769)

655 Henriksen MV, Chapple DG, Chown SL, McGeoch MA (2019) The Effect of Network Size and
656 Sampling Completeness in Depauperate Networks. *Journal of Animal Ecology* 88:211–222.
657 <https://doi.org/10.1111/1365-2656.12912>

658 Hesselbarth MHK, Nowosad J, Flamingh A de, et al (2024) Computational Methods in Landscape
659 Ecology. *Current Landscape Ecology Reports* 10:2. <https://doi.org/10.1007/s40823-024-00104-6>

660 Hillebrand H, Blasius B, Borer ET, et al (2018) Biodiversity Change Is Uncoupled from Species
661 Richness Trends: Consequences for Conservation and Monitoring. *Journal of Applied Ecology*
662 55:169–184. <https://doi.org/10.1111/1365-2664.12959>

663 Hortal J, Bello F de, Diniz-Filho JAF, et al (2015) Seven Shortfalls That Beset Large-Scale Knowledge
664 of Biodiversity. *Annual Review of Ecology, Evolution, and Systematics* 46:523–549. [https://doi.](https://doi.org/10.1146/annurev-ecolsys-112414-054400)
665 [org/10.1146/annurev-ecolsys-112414-054400](https://doi.org/10.1146/annurev-ecolsys-112414-054400)

666 Jordano P (2016a) Chasing Ecological Interactions. *PLOS Biology* 14:e1002559. [https://doi.org/10.](https://doi.org/10.1371/journal.pbio.1002559)
667 [1371/journal.pbio.1002559](https://doi.org/10.1371/journal.pbio.1002559)

668 Jordano P (2016b) Sampling Networks of Ecological Interactions. *Functional Ecology* 30:1883–1893.
669 <https://doi.org/10.1111/1365-2435.12763>

670 Kermorvant C, D'Amico F, Bru N, et al (2019) Spatially Balanced Sampling Designs for
671 Environmental Surveys. *Environmental Monitoring and Assessment* 191:524. [https://doi.org/10.](https://doi.org/10.1007/s10661-019-7666-y)
672 1007/s10661-019-7666-y

673 Kissling WD, Breeze TD, Lique C, et al (2024) Towards a Modern and Efficient European
674 Biodiversity Observation Network Fit for Multiple Policies

675 Lange C, Cole E, Van Horn G, Mac Aodha O (2023) Active Learning-Based Species Range
676 Estimation. *Advances in Neural Information Processing Systems* 36:41892–41913

677 Lewis DD, Gale WA (1994) A Sequential Algorithm for Training Text Classifiers. In: *Proceedings of*
678 *the 17th Annual International ACM SIGIR Conference on Research and Development in*
679 *Information Retrieval*. Springer-Verlag, Berlin, Heidelberg, pp 3–12

680 Malek S, Miglietta F, Gobakken T, et al (2019) Optimizing Field Data Collection for Individual Tree
681 Attribute Predictions Using Active Learning Methods. *Remote Sensing* 11:949. [https://doi.org/10.](https://doi.org/10.3390/rs11080949)
682 3390/rs11080949

683 McCann K (2007) Protecting Biostructure. *Nature* 446:29. <https://doi.org/10.1038/446029a>

684 McLeod A, Leroux SJ, Gravel D, et al (2021) Sampling and Asymptotic Network Properties of Spatial
685 Multi-Trophic Networks. *Oikos* 130:2250–2259. <https://doi.org/10.1111/oik.08650>

686 Mestre F, Gravel D, García-Callejas D, et al (2022) Disentangling Food-Web Environment
687 Relationships: A Review with Guidelines. *Basic and Applied Ecology* 61:102–115. [https://doi.org/](https://doi.org/10.1016/j.baae.2022.03.011)
688 10.1016/j.baae.2022.03.011

689 Moracho E, Normand S, Ordonez A (2025) Rebuilding Ecological Networks: A Multi-Scale Approach
690 to Global Restoration. *Biological Conservation* 312:111477. [https://doi.org/10.1016/j.biocon.2025.](https://doi.org/10.1016/j.biocon.2025.111477)
691 111477

692 Morales-Castilla I, Matias MG, Gravel D, Araújo MB (2015) Inferring Biotic Interactions from
693 Proxies. *Trends in Ecology & Evolution* 30:347–356. <https://doi.org/10.1016/j.tree.2015.03.014>

694 Navarro LM, Fernández N, Guerra C, et al (2017) Monitoring Biodiversity Change through Effective
695 Global Coordination. *Current Opinion in Environmental Sustainability* 29:158–169. [https://doi.](https://doi.org/10.1016/j.cosust.2018.02.005)
696 org/10.1016/j.cosust.2018.02.005

697 Norman KE, Poisot T (2025) Site Selection Algorithms for Optimal Ecological Monitoring Design.
698 *Ecological Indicators* 178:114104. <https://doi.org/10.1016/j.ecolind.2025.114104>

699 Pires MM, Marquitti FMD, Guimarães PR (2017) The Friendship Paradox in Species-Rich Ecological
700 Networks: Implications for Conservation and Monitoring. *Biological Conservation* 209:245–252.
701 <https://doi.org/10.1016/j.biocon.2017.02.026>

702 Poisot T (2023) Guidelines for the Prediction of Species Interactions through Binary Classification.
703 *Methods in Ecology and Evolution* 14:1333–1345. <https://doi.org/10.1111/2041-210X.14071>

704 Poisot T (2024) Conformal Prediction Quantifies the Uncertainty of Species Distribution Models

705 Poisot T, Bergeron G, Cazelles K, et al (2021) Global Knowledge Gaps in Species Interaction
706 Networks Data. *Journal of Biogeography* 48:1552–1563. <https://doi.org/10.1111/jbi.14127>

707 Poisot T, Canard E, Mouillot D, et al (2012) The Dissimilarity of Species Interaction Networks.
708 *Ecology Letters* 15:1353–1361. <https://doi.org/10.1111/ele.12002>

709 Poisot T, Stouffer DB, Gravel D (2015) Beyond Species: Why Ecological Interaction Networks Vary
710 through Space and Time. *Oikos* 124:243–251. <https://doi.org/10.1111/oik.01719>

711 Preston FW (1948) The Commonness, And Rarity, of Species. *Ecology* 29:254–283. <https://doi.org/10.2307/1930989>

713 Reji Chacko M, Albouy C, Altermatt F, et al (2025) Species Loss in Key Habitats Accelerates Regional
714 Food Web Disruption. *Communications Biology* 8:988. [https://doi.org/10.1038/s42003-025-08396-](https://doi.org/10.1038/s42003-025-08396-y)
715 *y*

716 Rioux È, Pelletier F, St-Laurent M-H (2022) Trophic Niche Partitioning between Two Prey and Their
717 Incidental Predators Revealed Various Threats for an Endangered Species. *Ecology and*
718 *Evolution* 12:e8742. <https://doi.org/10.1002/ece3.8742>

719 Robertson B, Price C, Reale M (2024) Well-Spread Samples with Dynamic Sample Sizes. *Biometrics*
720 80:ujae26. <https://doi.org/10.1093/biomtc/ujae026>

721 Robertson BL, Brown JA, McDonald T, Jaksons P (2013) BAS: Balanced Acceptance Sampling of
722 Natural Resources. *Biometrics* 69:776–784

723 Robertson BL, McDonald T, Price CJ, Brown JA (2017) A Modification of Balanced Acceptance
724 Sampling. *Statistics & Probability Letters* 129:107–112. <https://doi.org/10.1016/j.spl.2017.05.004>

725 Santana J, Porto M, Morán-Ordóñez A, et al (2025) Large Biodiversity Monitoring Gaps Remain
726 Across Europe. *Conservation Letters* 18:e13134. <https://doi.org/10.1111/conl.13134>

727 Settles B (2009) Active Learning Literature Survey

728 Simard A (2024) Le Réseau de Suivi de La Biodiversité: Un Outil Pour Mesurer l'effet Des
729 Changements Climatiques Sur Notre Patrimoine Naturel. *Natura* 4:13–22

730 Smith-Ramesh LM, Moore AC, Schmitz OJ (2017) Global Synthesis Suggests That Food Web
731 Connectance Correlates to Invasion Resistance. *Global Change Biology* 23:465–473. <https://doi.org/10.1111/gcb.13460>

732

733 Thompson PL, Gonzalez A (2017) Dispersal Governs the Reorganization of Ecological Networks
734 under Environmental Change. *Nature Ecology & Evolution* 1:1–8. [https://doi.org/10.1038/s41559-](https://doi.org/10.1038/s41559-017-0162)
735 *017-0162*

736 Tobias JA, Bullock JM, Dicks LV, et al (2025) Biodiversity Conservation Requires Integration of
737 Species-Centric and Process-Based Strategies. *Proceedings of the National Academy of Sciences*
738 122:e2410936122. <https://doi.org/10.1073/pnas.2410936122>

739 United Nations (1992) Convention on Biological Diversity (Text with Annexes). United Nations,
740 New Yoek

741 Valente JJ, Bennett RE, Gómez C, et al (2022) Land-Sparing and Land-Sharing Provide
742 Complementary Benefits for Conserving Avian Biodiversity in Coffee-Growing Landscapes.
743 *Biological Conservation* 270:109568. <https://doi.org/10.1016/j.biocon.2022.109568>

744 Valiente-Banuet A, Aizen MA, Alcántara JM, et al (2015) Beyond Species Loss: The Extinction of
745 Ecological Interactions in a Changing World. *Functional Ecology* 29:299–307. <https://doi.org/10.1111/1365-2435.12356>

747 Vizentin-Bugoni J, Maruyama PK, Debastiani VJ, et al (2016) Influences of Sampling Effort on
748 Detected Patterns and Structuring Processes of a Neotropical Plant–Hummingbird Network.
749 *Journal of Animal Ecology* 85:262–272. <https://doi.org/10.1111/1365-2656.12459>

750 Williams RJ, Martinez ND (2000) Simple Rules Yield Complex Food Webs. *Nature* 404:180–183.
751 <https://doi.org/10.1038/35004572>

752 Williams RJ, Martinez ND (2008) Success and Its Limits among Structural Models of Complex Food
753 Webs. *Journal of Animal Ecology* 77:512–519. <https://doi.org/10.1111/j.1365-2656.2008.01362.x>

754 Willig MR, Presley SJ, Klingbeil BT, et al (2023) Protecting Biodiversity via Conservation Networks:
755 Taxonomic, Functional, and Phylogenetic Considerations. *Biological Conservation* 278:109876.
756 <https://doi.org/10.1016/j.biocon.2022.109876>

757 Windsor FM, Hoogen J van den, Crowther TW, Evans DM (2023) Using Ecological Networks to
758 Answer Questions in Global Biogeography and Ecology. *Journal of Biogeography* 50:57–69.
759 <https://doi.org/10.1111/jbi.14447>

760 Wood SA, Russell R, Hanson D, et al (2015) Effects of Spatial Scale of Sampling on Food Web
761 Structure. *Ecology and Evolution* 5:3769–3782. <https://doi.org/10.1002/ece3.1640>

762 Xu H, Cao M, Wu Y, et al (2017) Optimized Monitoring Sites for Detection of Biodiversity Trends in
763 China. *Biodiversity and Conservation* 26:1959–1971. <https://doi.org/10.1007/s10531-017-1339-3>

764 Yi J, Ma F, Li J, et al (2022) China Biodiversity Observation Network for Better Global Sustainability
765 Developments. *Eco-Environment & Health* 1:201–203. <https://doi.org/10.1016/j.eehl.2022.11.001>