

Review

The genetics of rheumatoid arthritis

Laura E. Dedmon¹

Abstract

RA is a chronic systemic inflammatory disease that primarily affects the small joints of the hands and feet, and results in a mean reduction in life expectancy of 3–10 years. RA is a multigene disorder with a substantial genetic component and a heritability estimate of 60%. Large-scale Genome-Wide Association Studies (GWAS) and meta-analyses have revealed common disease-associated variants in the population that may contribute cumulatively to RA pathogenesis. This review identifies the most significant genetic variants associated with RA susceptibility to date, with particular focus on the contribution of the HLA class II genes across different ethnic groups. Also discussed are the potential applications of pharmacogenomics to RA management by identifying polymorphisms associated with variation in treatment response or toxicity. The use of genetic variants to guide treatment strategy has the potential to not only reduce National Health Service costs, but also drastically improve patient experience and quality of life.

Key words: rheumatoid arthritis, heritability, genetics, methotrexate, biologics, sequencing, polymorphism, pharmacogenomics

Rheumatology key messages

- Rheumatoid arthritis is associated with many common polymorphisms that contribute cumulatively to disease pathogenesis.
- Response to treatment varies widely between patients and has a genetic component.
- Polymorphisms associated with improved or reduced treatment response could be used to guide clinical practice.

Introduction

RA is a chronic systemic inflammatory disease that primarily affects the small joints of the hands and feet [1]. It has a prevalence of ~1% in European populations and is associated with premature mortality, with a mean reduction in life expectancy of 3–10 years [2]. RA is a multigene disorder with a substantial genetic component and a heritability estimate of 60%, at least 30% of which is likely attributable to genes in the HLA class II family [3]. Large-scale Genome-Wide Association Studies (GWAS) and meta-analyses have revealed common disease-associated variants in the population that may contribute in a cumulative fashion to RA pathogenesis. The identification of variants that are associated with increased RA susceptibility and severity could have diagnostic and clinical applications, not only to enable clinicians to anticipate those at greatest risk of disease, but also to direct drug choice for greatest efficacy and

success in patients. This review identifies the most significant genetic variants associated with RA susceptibility to date, with particular focus on the contribution of the HLA class II genes across different ethnic groups. Also discussed are the potential application of these findings to the treatment of RA, and the use of pharmacogenomics in predicting patient response to RA drugs.

Background

RA is characterized by symmetrical peripheral polyarthritis due to an autoimmune inflammatory response focused in the synovium, and results in deformity of the joints, particularly the MCP, PIP and MTP joints (Fig. 1). Prevalence in women is at least twice that in men, and peak age of onset is in the fifth to sixth decades of life. RA presents initially with peripheral joint inflammation, causing inflammatory pain that is worse in the mornings and eases throughout the day. As the disease progresses it can result in joint damage and deformity, particularly subluxation of the wrist and fingers and ulnar deviation of the MCP joints [5]. RA is also associated with a 2–3-fold increase in cardiovascular disease and presents a significant disease

¹UCL Medical School, London, UK

Submitted 11 January 2020; accepted 3 April 2020

Correspondence to: Laura Dedmon, UCL Medical School, 74 Huntley Street, London WC1E 6DE, UK. E-mail: laura.dedmon.12@ucl.ac.uk

Fig. 1 Hand deformity in RA

The hands of an RA patient showing swelling, subluxation and ulnar drift of the MCP joints, characteristic of late-stage disease. Reprinted with permission from RheumExam Atlas (2015) [4].

burden with higher morbidity and mortality in those affected [5, 6].

The presence of autoantibodies to the Fc region of immunoglobulin G (RF) and to citrullinated cyclic peptide (ACPAs) distinguish seropositive RA from seronegative RA. Patients with RF or ACPAs typically present with more aggressive and severe disease. This often leads to worse disease outcome, particularly in patients who smoke [7–9].

The disease activity in RA is quantified by DAS28, a scoring system based on the number of swollen and tender joints, the amount of inflammation indicated by blood tests, and by the patient's own sense of wellbeing. DAS28 scoring is often used to determine when a referral to a rheumatologist may be necessary [7]. Corticosteroid injections into an acutely swollen joint can rapidly reduce inflammation, and NSAIDs may be used as pain relief, but neither have any effect on disease activity. Early management with DMARDs and biologics can slow disease progression and improve outcomes. DMARDs are the first line of treatment for RA, and include MTX, sulfasalazine and azathioprine. They are frequently used in combination, and can take 6–12 months to produce symptomatic improvement [5]. NICE guidelines mandate that patients may be prescribed biologics only if they continue to present with active RA despite having tried at least two different DMARDs [9]. Biologics include TNF- α inhibitors (infliximab, etanercept, adalimumab), B-cell depleters (rituximab), IL-1/IL-6 inhibitors (tocilizumab), and inhibitors of T-cell co-stimulation (abatacept). They are often used in combination with MTX [5]. Response to DMARDs and biologics is highly variable between patients, and biologics are still prescribed on a largely trial-and-error basis [10].

The genetics of rheumatoid arthritis

It has long been acknowledged that having a family history of RA increases an individual's likelihood of developing the disease, and the heritability of RA has been estimated to be around 60% [1]. Early twin studies and

linkage analyses seeking to identify disease-associated genes consistently highlighted one gene family in particular – the HLA *DRB1* alleles, found within the human MHC on chromosome 6. The HLA region contains class I and class II genes, the latter of which (*HLA-DR*, *DQ* and *DP*) produce the alpha and beta chains of the corresponding MHC class II molecule. This molecule is found on antigen-presenting cells and is responsible for the presentation of extracellular pathogens to T cells, resulting in an immune response [11].

The *HLA-DRB1* alleles constitute the single strongest genetic association for RA, and likely contribute at least 30% of the total genetic component of this disease [12]. The disease-associated alleles have a conserved sequence of five amino acids, termed the 'shared epitope' [13]. The shared epitope hypothesis suggests that certain alleles with this conserved sequence are associated with, and in fact directly contribute to, the pathogenesis of RA by permitting the incorrect presentation of autoantigens to T cells by antigen-presenting cells, resulting in a T-cell-mediated autoimmune response [12, 13]. The alleles of the shared epitope have been strongly associated with RA susceptibility and increased disease severity, as well as with an increased risk of developing extra-articular manifestations [12, 14]. These alleles appear to show a gene dosage effect, with RA susceptibility and severity increasing in proportion with the number of alleles present [15]. Jawaheer *et al.* (1994) showed that twins with the shared epitope, particularly the *HLA-DRB1*04* (*HLA-DR4*) alleles, have increased disease concordance. Twins who are homozygous for the shared epitope have five times higher concordance than those lacking the shared epitope completely. The researchers concluded that the shared epitope is 'the most important factor in determining RA concordance' [16].

Other studies seeking to explain the remaining heritability of RA have identified strongly associated loci outside of the shared epitope, such as those involving *PTPN22*, *CTLA4* and *PADI4* [17–19]. However, the candidate gene approach used by most of these studies has limitations, chiefly that it is wholly dependent on the appropriate selection of genes known (or thought) to contribute to disease pathogenesis, and is therefore reliant on a prior thorough understanding of disease mechanisms. This is difficult for studies of RA, which has a complex pathogenesis and polygenic inheritance pattern, and this strategy will inevitably miss contributions from significant loci. Researchers have therefore turned increasingly to Genome Wide Association Studies (GWAS) to look for common variants that may contribute, in small but cumulative ways, to increased RA susceptibility. GWA studies take advantage of regions of linkage disequilibrium (LD) across the genome. These are blocks of genes with lower rates of recombination than expected, which are inherited as haplotype blocks and can be tagged for single nucleotide polymorphisms (SNPs), common single base pair variations that occur throughout the genome [20]. This allows researchers to observe which polymorphisms are more common in cases than in controls.

In 2007, the Wellcome Trust Case-Control Consortium (WTCCC) carried out an extensive GWA study in the British population to look for common variants associated with susceptibility to seven major diseases, including RA [21]. The study confirmed the established association of the *HLA-DRB1* locus with RA, as well as the 1858 C/T polymorphism of the *PTPN22* gene previously associated by Begovich *et al.* (2004) [17, 21]. Of the non-*HLA-DRB1* loci, the *PTPN22* gene is by far the most significantly associated with RA susceptibility and, together with the *HLA-DRB1* alleles, is thought to be responsible for ~50% of the genetic component of RA [21, 22]. The *PTPN22* protein product is lymphoid protein tyrosine phosphatase, an enzyme found only in immune cells where it plays an important part in reducing the responsiveness of T and B-cell receptors [17]. However, despite the significance of this variant in populations of European descent, it occurs very rarely in Asian populations, and is therefore unlikely to contribute substantially to RA risk amongst those groups [23–25].

The WTCCC GWA study identified and confirmed several other variants with moderate disease-association. An SNP mapping to the alpha and beta chain regions of the IL2 receptor (*IL2RA* and *IL2RB*) is of particular note [21]. The IL2 receptor is responsible for regulating IL2 stimulation of T cells, and is therefore important for inhibiting autoimmune responses. The association with *IL2RA* was later refined and confirmed by Stahl *et al.* (2010) [26]. This SNP, as well as a variant in *STAT4*, has been associated with susceptibility to several other autoimmune-related diseases, including systemic lupus erythematosus and type 1 diabetes, suggesting a common aetiology underlying these autoimmune disorders [10, 21].

Another GWA study in 2007 identified an SNP (rs3761847) significantly linked with the TNF receptor associated factor 1 and complement component 5 gene regions (*TRAF1/C5*) that is associated with increased risk of ACPA-seropositive RA in European populations [27]. Huang *et al.* (2019) identified another polymorphism (rs10818488) associated with RA not only in Europeans, but also Asians and Africans [28]. Subsequent analysis of eQTL data (a method of testing linkage between gene expression phenotype and genetic polymorphisms) strongly suggests that the association stems from the *TRAF1* region, rather than the *C5*, and *TRAF1* is now thought to be the more functionally-relevant gene associated with these variants [29].

The choice of SNPs used to tag regions of the genome is critical, and is determined by the microarray genochip used in the GWA study. Eyre *et al.* (2012) used Immunochip (an autoimmune disease-specific microarray) to perform a meta-analysis, combining data from almost 50 000 individuals. By genotyping for 186 GWAS-confirmed loci, they were able to identify 14 new disease-associated loci, and pinpoint the association signal to a single gene for 19 of the 39 previously associated loci on Immunochip [30].

Through such strategies, researchers may gradually determine the true causal variants of RA susceptibility.

However, certain microarrays are more appropriate for some ethnic groups than others, and failure to account for this variation can lead to unreliable results. Blocks of LD vary in structure between ethnicities, and this is responsible in part for the heterogeneity of associated polymorphisms between populations [31]. Additionally, care must be taken when selecting cohorts for these studies, as cases and controls must be of similar ethnic backgrounds to avoid effects of population stratification [32]. The results of GWAS are only applicable to the population studied, and cannot be reliably extrapolated to others.

To address this issue, researchers have combined data from multiple GWA studies into meta-analyses, which benefit from increased statistical power compared with single GWA studies, and can allow for the detection of subtle associations across ethnic groups. One of the most significant contributions in this field is a meta-analysis by Okada *et al.* (2014), which used 29 880 RA cases and 73 758 controls from European and Asian cohorts to analyse around 10 million SNPs. The researchers discovered 42 novel loci that are associated with increased RA risk at genome-wide significance, 38 of which are significant at a trans-ethnic level [33]. Further GWA studies and meta-analyses have identified more loci, such as *PADI2* and *NFKBIE*, that are consistently associated across ethnicities (see Table 1 for summary) [25, 33, 38].

Meta-analyses have also shed further light on the MHC class II HLA alleles and their association with RA across ethnicities. The prevalence of RA itself is higher in some populations than others, with 1% of Europeans, 0.5% of Asians and Africans, and over 2% of some Native American populations suffering from the disease [40]. As discussed above, the most common SE-coding alleles are those in the *HLA-DR4* group, though RA in the absence of an *HLA-DR4* background appears to be more common in certain ethnic groups, particularly among Americans of African descent and Jewish Israelis [41, 42]. In patients lacking the *HLA-DR4* alleles, variants in other MHC class II-corresponding HLA genes, such as alleles *HLA-DRB1*01* and *HLA-DRB1*10:01*, are associated with increased RA susceptibility (though conferring lower relative risk than *HLA-DR4*) [13]. The *HLA-DPB1*02:01* and *HLA-DRB1*09:01* alleles are independently associated with disease in Japanese populations; while a variant in the *HLA-DOA* gene is associated with disease in Japanese and Europeans [36, 37, 43]. Conversely, alleles *DRB1*13:01* and *HLA-DRB1*13:02* have been found to have a negative association with RA in northern European and Japanese populations, respectively, suggesting they have a protective effect on ACPA-positive RA [44, 45]. Amongst Indigenous North Americans, a population with 2–3 times the prevalence of RA in Europeans and Asians, the *HLA-DRB1*14:02* allele has been shown to be associated with increased risk of ACPA-positive RA, independently of other HLA loci [35, 46, 47]. This allele is very uncommon in Caucasian and Asian ethnicities, but reaches a prevalence of 80% in some Native American populations [47].

TABLE 1 Significant genes associated with RA susceptibility

Gene	Protein Product	Population	Reference
<i>HLA-DRB1</i>	HLA class II histocompatibility antigen, DRB1 beta chain	Trans-ethnic African-American/In digenous North American	[25, 33, 34, 35]
<i>HLA-DPB1</i>	HLA class II histocompatibility antigen, DP beta 1 chain	Japanese	[36]
<i>HLA-DOA</i>	HLA class II histocompatibility antigen, DO alpha chain	Japanese European	[37]
<i>PADI4</i>	Protein-arginine deiminase type-4	Trans-ethnic African-American	[25, 33]
<i>PTPN22</i>	Tyrosine-protein phosphatase non-receptor type 22	Trans-ethnic	[33]
<i>CTLA4</i>	Cytotoxic T-lymphocyte protein 4	Trans-ethnic African-American	[25, 33]
<i>IL2RA</i>	IL-2 receptor subunit alpha	Trans-ethnic African-American	[25, 33]
<i>STAT4</i>	Signal transducer and activator of transcription 4	Trans-ethnic	[33]
<i>TRAF1-C5</i>	TNF receptor-associated factor 1 - Complement C5	Trans-ethnic African	[28, 33]
<i>CD40</i>	C-C chemokine receptor type 6	Trans-ethnic	[33]
<i>CCR6</i>	IFN regulatory factor 5	Trans-ethnic	[33]
<i>IRF4</i>	Transcription regulator protein BACH2	Trans-ethnic	[33]
<i>BACH2</i>	DNA repair protein RAD51 homolog 2	European	[38]
<i>RAD51B</i>	Dipeptidyl peptidase 4	European	[38]
<i>DPP4</i>	RNA binding protein fox-1 homolog 1	Han Chinese	[39]
<i>RBFOX1</i>	Protein-arginine deiminase type-2	African-American	[25]
<i>PADI2</i>	CDK5 regulatory subunit-associated protein 2	African-American	[25]
<i>CDK4RAP2</i>	Protein LBH	Han Chinese European	[39]
<i>LBH</i>	HLA class II histocompatibility antigen, DO alpha chain	Trans-ethnic African-American	[25, 33]
<i>COG6</i>	Non-receptor tyrosine-protein kinase TYK2	Trans-ethnic African-American	[25, 33]
<i>TYK2</i>	Protein-arginine deiminase type-2	Trans-ethnic African-American	[25, 33]
<i>PADI4</i>	Trans-acting T-cell-specific transcription factor GATA-3	Trans-ethnic African-American	[25, 33]
<i>GATA3</i>	Conserved oligomeric Golgi complex subunit 6	European	[30]

A summary of the most significant genes currently associated with RA susceptibility, and the populations within which they are significant. Variants found to be significant across both European and Asian ethnicities are classified as 'trans-ethnic'. Studies demonstrating the broadest association of the variant across ethnicities are referenced in each case.

The substantial contribution of *HLA-DRB1* alleles to the genetic component of RA and the increasing number of non-HLA loci associated with the disease have led researchers to investigate connections between these two groups. Diaz-Gallo *et al.* (2018) found significant enrichment in interactions between *HLA-DRB1* shared epitope alleles and SNPs associated with ACPA-positive RA. They suggested a 'dominion hypothesis', wherein the smaller effects of non-HLA variants on RA susceptibility are amplified through interaction with a central 'hub' of shared epitope alleles [48]. The researchers found that of all the loci, the most highly interactive SNPs were rs2476601 and rs1073958. eQTL analysis revealed that in shared-epitope carriers, the rs2476601 variant in the *PTPN22* gene shows the most significant interaction with the shared epitope alleles, while rs1073958 corresponds to *CDK5RAP2* (previously identified by Jiang L *et al.*) and *PSMD5* (Table 1) [39, 48].

GWA studies have contributed substantially to our understanding of the genetic aspect of RA, allowing researchers to cast a wide net without the restrictions of the candidate gene approach. However, these studies also have limitations integral to their design. While they can detect common variants associated with complex diseases, GWAS fail to identify contributing rare genetic variants [49]. This is often due to a lack of statistical power, as well as the limitations of SNPs in identifying

larger sources of variation (such as copy number variants, deletions or rearrangements) [50]. GWA studies also suffer from a high rate of false positives due to the large number of statistical tests performed on the data and the presence of individuals in the control group who have not yet developed the disease, but will in the future. It is therefore crucial that studies adhere to a very low *P*-value as the threshold for significance – typically taken as less than 5×10^{-8} [32].

Determining the mechanism by which a SNP is associated with disease is difficult, as it may be causal itself, or simply in LD with the true causal variant. A strong association is not sufficient to prove causality, and identifying the truly causative genetic variant presents a challenge in this field. Fine mapping techniques may be of use in analysing disease-associated regions from GWA studies [51]. This involves exploring in detail the structure of each region tagged by a promising SNP, and may allow researchers to begin to identify true causal variants from the staggering number of disease-associated SNPs [52]. Farh *et al.* (2015) succeeded in developing a new statistical algorithm for fine-mapping, allowing them to identify over 5000 candidate causal SNPs for 21 autoimmune diseases [53]. They integrated these data with chromatin and transcription factor binding maps to determine function, and found that the majority of causal variants are non-coding, involved in

enhancer elements, and located near to immune transcription factor binding sites [53]. In fact, 80% of the >100 loci now associated with RA are in non-coding regions of the genome [54]. The researchers proposed that the effects of such variants could produce subtle but significant alterations in immune response that could predispose towards an autoimmune disease [53]. Effects on expression could be mediated through several mechanisms, including regulatory elements affecting promoter and enhancer activity, alternative splicing and chromatin configuration [54]. Such effects are difficult to elucidate and beyond the scope of this review, but are discussed in depth by Ding J *et al.* (2019) in their review of the use of functional genomics in RA [55].

Pharmacogenomics and personalized medicine in the treatment of rheumatoid arthritis

Although the recent progress in identifying variants associated with RA susceptibility has potential to improve understanding of disease pathogenesis, separate large-scale studies investigating variants associated with treatment response will be more useful for advancing the fields of pharmacogenomics and personalized medicine. Personalized medicine seeks to tailor disease management to individuals based on biomarkers associated with drug response or toxicity. Research in pharmacogenomics aims to identify which genetic variants are associated with drug responses or reactions. Success in these areas would be revolutionary for the management of RA, as the majority of drugs currently prescribed for treatment produce highly variable responses in patients.

Several DMARDs are used to treat RA. MTX is considered to be a cornerstone of disease management, but only 46–65% of RA patients respond positively to its use [56, 57]. Its mechanism of action is unknown, but likely involves inhibition of the folate metabolism pathway [58]. Several genetic variants have been identified as associated with variation in clinical response to MTX, the most significant of which are discussed here (Table 2).

One of the most strongly associated variants is a minor allele of *ATIC* (rs4673993). Lee *et al.* (2009) found that the C allele is associated with increased response to MTX in RA patients on MTX monotherapy compared with the T allele, while another study in 2011 showed that the C allele is associated with lower disease activity [59, 60]. This was further supported by a meta-analysis by Chen *et al.* (2017), which showed significant association of this variant across Caucasian and non-Caucasian populations [61]. The *ATIC* C allele is therefore strongly associated with improved response to MTX in RA patients.

Other genetic associations with MTX response or toxicity are less clear and often controversial. Polymorphisms in genes such as *SLC19A1*, *ABCB1* and *MTHFR* have all been shown to be associated with increased, decreased or no change in MTX response or toxicity, making it very difficult to draw conclusions as to the true effects (if any) of these alleles (Table 2) [62–

65, 79–81]. Chen *et al.* (2017) found only one other SNP associated with MTX responsiveness – the 34C>T polymorphism in the *AMPD1* gene [61]. Significant association of the T allele with improved response was found across multiple ethnicities. The variant produces an enzyme (adenosine monophosphate deaminase) with lower activity, though it is unclear how this might affect MTX efficacy [82]. Most recently, Taylor *et al.* (2018) performed a GWA study using >1000 RA cases to evaluate patient response to MTX, and found that the only locus to suggest association was rs168201 in the *NRG3* gene; however, even this failed to achieve genome-wide significance [66]. This study represents the largest GWA study of RA patient response to MTX treatment to date, and its failure to identify any loci of substantial significance suggests that MTX response is highly polymorphic [66]. Larger, more extensive studies will be required to shed light on any true genetic associations.

Other DMARDs used in RA therapy include sulfasalazine and azathioprine. Patient response to sulfasalazine has long been associated with variants in the *NAT2* gene that affect the ability of the N-acetyltransferase 2 enzyme to metabolise sulfasalazine (Table 2) [67, 68, 71]. Homozygotes for variant alleles, including *NAT2*5B*, *NAT2*6A* and *NAT2*7B*, are ‘slow acetylators’, and recent studies have shown that these individuals are more likely to suffer adverse drug reactions from sulfasalazine treatment than those with the wild-type *NAT2*4* allele, and are more likely to show non-compliance [69–71]. Wiese *et al.* (2014) also found a polymorphism (C412A) in the *ABCG2* gene, which is associated with disease remission using sulfasalazine [69]. These findings suggest that *NAT2* and *ABCG2* genotyping could be useful in predicting clinical response and toxicity of sulfasalazine during treatment [70].

Azathioprine is an immunosuppressant that works by inhibiting purine synthesis, particularly in white blood cells. The gene *TPMT* codes for the protein thiopurine methyltransferase (TPMT), which is involved in the metabolism of azathioprine [83]. Multiple studies have shown that patients with variations in the *TPMT* gene (alleles *TPMT*2*, **3A*, **3B*, **3C*, **4*) have deficient TPMT activity, resulting in decreased inactivation of thiopurines, and an increased risk of azathioprine toxicity compared with individuals homozygous for the wild-type *TPMT*1* allele [72, 84, 85] (Table 2). These variants are common, with ~10% of the Caucasian population heterozygous for one allele, and one in 300 homozygous (no TPMT activity) [86]. Patients with deficient TPMT activity are at greater risk of life-threatening myelotoxicity and pancytopenia with standard azathioprine dosing, and current NICE clinical guidelines recommend determining patient genotype or TPMT enzyme activity prior to prescribing azathioprine [72, 87]. The introduction of standard TPMT status assessment as a part of clinical decision-making represents one of the clearest demonstrations of the field of pharmacogenetics at work.

A promising addition to this field is the effect of a variant in the gene *NUDT15* on azathioprine toxicity

TABLE 2 Variants associated with response/toxicity to RA drugs

Drug	Gene	Polymorphism/Allele	Effect	Reference
MTX	<i>ATIC</i>	T675C (rs4673993)	C allele associated with lower disease activity and improved treatment response compared to TT	[59, 60, 61]
	<i>SLC19A1</i>	G80A (rs1051266)	G allele associated with increased risk of MTX-related gastrointestinal toxicity compared to AA	[62]
			No association	[63]
	<i>ABCB1</i>	C3435T (rs1045642)	C allele associated with increased risk of MTX toxicity	[64]
			No association	[65]
	<i>MTHFR</i>	C677T (rs1801133)	T allele associated with increased risk of toxicity after 12 months of MTX treatment	[64]
Sulfasalazine			No association	[63]
	<i>AMPD1</i>	35C>T (rs17602729)	T allele associated with increased response	[61]
			No association	[63]
	<i>NRG3</i>	G>A (rs168201)	Associated with improvement in DAS28 score	[66]
	<i>NAT2</i>	NAT2*5	Associated with increased risk of adverse drug reaction compared to wild-type allele	[67]
		NAT2*6		[68]
Azathioprine		NAT2*7		[69]
				[70]
	<i>ABCG2</i>	C412A (rs2231142)	Associated with reduced efficacy	[71]
			A allele associated with increased rates of remission after 12 months	[69]
	<i>TPMT</i>	TPMT*2 TPMT*3A TPMT*3B TPMT*3C TPMT*4	Associated with increased risk of toxicity	[72]
	<i>NUDT15</i>	T>C (rs116855232)	T allele associated with increased risk of toxicity	[73]
TNF-α inhibitors	<i>TNF</i>	–308 G/A (rs1800629)	A allele associated with poor response to adalimumab, etanercept and infliximab	[74]
				[75]
Rituximab			No association with infliximab	[76]
	<i>FGCR3A</i>	–236 A/C (rs396991)	C allele associated with improved response after 6 months of treatment	[77]
				[78]

A summary of the most significant variants associated with response or toxicity for DMARDs and biologic therapies used in RA.

(Table 2). While *TPMT* variations are relatively common in Caucasian populations, they are rare in Asians. Conversely, variants in *NUDT15* are more common in Asian populations. The T allele in the *NUDT15* gene has been shown to be strongly associated with azathioprine-induced leukopenia in patients with inflammatory bowel disease compared with allele C [73]. The association is strong across both Asian and European ancestry groups, and researchers have estimated that *NUDT15* and *TPMT* variants together account for ‘~50% of severe thiopurine-related hematotoxicity’, suggesting that *NUDT15* genotyping could be a useful tool in the future for determining the risk of adverse drug reactions with azathioprine [88].

Patients who continue to suffer with active RA despite having tried at least two DMARDs are managed using biologics [9]. These drugs are synthesized from

biological sources and vary drastically in efficacy between patients [89]. This variability has been partially associated with specific gene polymorphisms.

TNF- α inhibitors are the first type of biologic given to most RA patients, and act by suppressing the body’s response to TNF, an essential component of the inflammatory response. There are several types, chiefly etanercept, infliximab and adalimumab. They are very effective for some, but as many as 30–40% of patients do not respond to treatment [90]. Several pharmacogenomics studies of TNF- α inhibitors have identified variants that may be associated with treatment response, including the common variant 308 G/A (rs1800629) in the *TNF- α* gene (Table 2) [74, 75]. Two separate meta-analyses identified the A allele as associated with decreased response to adalimumab, etanercept and infliximab in RA patients compared with the G allele [74, 75]. However,

not all studies agree, and a study by Maxwell *et al.* (2008) found that there is no association of this polymorphism with response to infliximab in particular [76]. A large GWA study on this topic in 2018 failed to identify any variants of genome-wide significance capable of predicting change in DAS28 score during the first 3–6 months of treatment [91]. The study did determine that a variant in the *FTO* gene (previously linked to BMI), is associated with improvement in the number of swollen joints in patients treated with infliximab, even when the results are controlled for BMI. Treatment non-compliance is a complicating factor for these studies, but the lack of substantial genetic association with treatment response suggests that response to TNF- α inhibitors is a highly polygenic trait, and greater statistical power is needed to elucidate associated variants [91].

The biologic rituximab is an anti-CD20 monoclonal antibody that depletes B-cell numbers. It is given to patients with severe disease resistant to other drugs. Although rituximab appears to successfully reduce B-cell count in almost all patients, 40–50% of patients still fail to improve [92]. A cohort study of 212 RA patients in 2014 found that RA patients with the V allele (158V/F) of the *FCGR3A* gene have increased response to rituximab, and are less likely to have loss of response to the drug after 4–6 months of treatment (Table 2) [77]. These findings are supported by other smaller studies, suggesting that the *FCGR3A* 158V polymorphism could be useful for determine efficacy of rituximab treatment [77, 78].

The ability to reliably identify genetic polymorphisms that have a substantial effect on treatment response in RA patients would revolutionize the management of this disease. This is particularly true for biologics, which are currently prescribed in a systematic manner based on price, wherein a patient must fail to respond to the first drug before they can try the next [9]. This can be a very slow and frustrating experience for patients, particularly those who fail to respond to multiple drugs and must continue living with uncontrolled RA while potentially experiencing the side effects of each new drug they try. It is also very cost-ineffective – TNF- α inhibitors, for example, are estimated to cost the National Health Service (NHS) over £9000 per patient per year [93]. The ability to better predict which patients are most likely to respond well or poorly would therefore make a profound difference to clinical management of RA, patient quality of life and NHS spending.

RA prognosis is intrinsically linked with early diagnosis and management [94]. While it is not possible to predict with certainty who will develop RA, identifying individuals at higher genetic risk of the disease could help clinicians to better advise patients about lifestyle factors prior to disease onset, and enable closer monitoring to begin treatment immediately at disease onset. Polygenic risk scores are one method of quantifying the cumulative effect of all gene variants associated with a disease. However, because they are specific to the ethnic groups used to identify them originally, clinical use of existing data could result in the neglect of certain populations

(for example, Afro-Caribbean and Middle Eastern) who have not been studied so rigorously for disease-associated variants compared with European and Asian populations. This has the potential to create and/or exacerbate inequalities in healthcare, and more research targeting these under-studied populations is required to ensure equal distribution of resources [95].

Conclusion

Understanding the genetic basis of RA is important for prediction of disease severity, prognosis and response to treatment. Candidate gene and genome-wide studies have succeeded in identifying a large number of polymorphisms associated with increased RA susceptibility in different human populations. Focus on the use of functional studies to determine not only the causal variant in each case, but also the function of the causal variant, the genes affected by the causal variant, and how all of these factors contribute to disease pathogenesis, is perhaps now the more important area for study. This, combined with the use of pharmacogenomics in clinical practice, would represent a tremendous step forward in the management of RA, and would undoubtedly improve patient quality of life.

Acknowledgements

I would like to thank Professor Surjit Kaila Singh Srai of University College London for his support, encouragement and guidance in the publication of this review.

Funding: No specific funding was received from any bodies in the public, commercial or not-for-profit sectors to carry out the work described in this article.

Disclosure statement: The author has declared no conflicts of interest.

References

- MacGregor AJ, Snieder H, Rigby AS *et al.* Characterizing the quantitative genetic contribution to rheumatoid arthritis using data from twins. *Arthritis Rheum* 2000;43: 30–7.
- Toledano E, Candelas G, Rosales Z *et al.* A meta-analysis of mortality in rheumatic diseases. *Reumatol Clin* 2012;8:334–41.
- Deighton CM, Walker DJ, Griffiths ID, Roberts DF. The contribution of HLA to rheumatoid arthritis. *Clin Genet* 2008;36:178–82.
- RheumExam Atlas. Typical Ulnar Deviation of Rheumatoid Arthritis (Pronation). Dr Lori Albert, University of Toronto; 2015. http://www.rheumexamatlas.com/hands_typicalulnardeviation_parti/ (24 December 2019, date last accessed).
- Wilkinson IB, Raine R, Wiles K *et al.* Oxford handbook of clinical medicine. 10th edn Oxford: Oxford University Press, 2017: 546–547.A

- 6 Meune C, Touzé E, Trinquart L, Allanore Y. Trends in cardiovascular mortality in patients with rheumatoid arthritis over 50 years: a systematic review and meta-analysis of cohort studies. *Rheumatology* 2009;48:1309–13.
- 7 Kumar P, Clark M. *Clinical medicine*. 8th edn. Netherlands: Elsevier Ltd; 2012: 516–25.
- 8 Mahdi H, Fisher BA, Kallberg H *et al*. Specific interaction between genotype, smoking and autoimmunity to citrullinated alpha-enolase in the etiology of rheumatoid arthritis. *Nat Genet* 2009;41:1319–24.
- 9 National Institute for Health and Care Excellence. (2018) Rheumatoid arthritis in adults: management. NICE guideline [NG100]. <https://www.nice.org.uk/guidance/ng100/chapter/Recommendations> (7 March 2020, date last accessed).
- 10 Kurko J, Besenyei R, Laki J *et al*. Genetics of rheumatoid arthritis – a comprehensive review. *Clin Rev Allergy Immunol* 2013;45:170–9.
- 11 Transplantation Immunology. HLA genes and Molecules. Faculte de medicine de Geneve. Updated: 2015. http://medweb4.unige.ch/immunologie/home/HSC/transplantation_barriers/structures_recognized/major_histocompatibility_antigens/genetic.php (7 March 2020, date last accessed).
- 12 Van der Helm-van Mil AH, Huizinga TW, Schreuder GM *et al*. An independent role of protective HLA class II alleles in rheumatoid arthritis severity and susceptibility. *Arthritis Rheum* 2005;52:2637–44.
- 13 Gregersen PK, Silver J, Winchester RJ. The shared epitope hypothesis. An approach to understanding the molecular genetics of susceptibility to rheumatoid arthritis. *Arthritis Rheum* 1987;30:1205–13.
- 14 Du Montcel ST, Michou L, Petit-Teixeira E *et al*. New classification of HLA-DRB1 alleles supports the shared epitope hypothesis of rheumatoid arthritis susceptibility. *Arthritis Rheum* 2005;52:1063–8.
- 15 Weyand CM, Goronzy JJ. Disease mechanisms in rheumatoid arthritis: gene dosage effect of HLA-DR haplotypes. *J Lab Clin Med* 1994;124:335–8.
- 16 Jawaheer D, Thomson W, MacGregor AJ *et al*. “Homozygosity” for the HLA-DR shared epitope contributes the highest risk for rheumatoid arthritis concordance in identical twins. *Arthritis Rheum* 1994;37:681–6.
- 17 Begovich AB, Carlton VE, Honigberg LA *et al*. A missense single-nucleotide polymorphism in a gene encoding a protein tyrosine phosphatase (PTPN22) is associated with rheumatoid arthritis. *Am J Hum Genet* 2004;75:330–7.
- 18 Rodríguez MR, Núñez-Roldán A, Aguilar F *et al*. Association of the CTLA4 3' untranslated region polymorphism with the susceptibility to rheumatoid arthritis. *Hum Immunol* 2002;63:76–81.
- 19 Suzuki A, Yamada R, Chang X *et al*. Functional haplotypes of PADI4, encoding citrullinating enzyme peptidylarginine deiminase 4, are associated with rheumatoid arthritis. *Nat Genet* 2003;34:395–402.
- 20 Slatkin M. Linkage disequilibrium—understanding the evolutionary past and mapping the medical future. *Nat Rev Genet* 2008;9:477–85.
- 21 The Wellcome Trust Case Control Consortium. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* 2007;447:661–78.
- 22 Van der Helm-van Mil AH, Wesoly JZ, Huizinga TW. Understanding the genetic contribution to rheumatoid arthritis. *Curr Opin Rheumatol* 2005;17:299–304.
- 23 Kawasaki E, Awata T, Ikegami H, *et al*. Systematic search for single nucleotide polymorphisms in a lymphoid tyrosine phosphatase gene (PTPN22): association between a promoter polymorphism and type 1 diabetes in Asian populations. *Am J Med Genet A*. 2006;140(6):586–593.
- 24 Nabi G, Akhter N, Wahid M *et al*. Meta-analysis reveals PTPN22 1858C/T polymorphism confers susceptibility to rheumatoid arthritis in Caucasian but not in Asian population. *Autoimmunity* 2016;49:197–210.
- 25 Laufer VA, Tiwari HK, Reynolds RJ *et al*. Genetic influences on susceptibility to rheumatoid arthritis in African-Americans. *Hum Mol Genet* 2019;28:858–74.
- 26 Stahl EA, Raychaudhuri S, Remmers EF *et al*. Genome-wide association study meta-analysis identifies seven new rheumatoid arthritis risk loci. *Nat Genet* 2010;42:508–14.
- 27 Carlton VE, Hu X, Chokkalingam AP *et al*. PTPN22 genetic variation: evidence for multiple variants associated with rheumatoid arthritis. *Am J Hum Genet* 2005;77:567–81.
- 28 Huang SC, Hua DJ, Sun QQ *et al*. Associations of *TRAF1/C5* rs10818488 and rs3761847 polymorphisms with genetic susceptibility to rheumatoid arthritis: a case-control study and updated meta-analysis. *Cent Eur J Immunol* 2019;44:159–73.
- 29 Zhu Z, Zhang F, Hu H *et al*. Integration of summary data from GWAS and eQTL studies predicts complex trait gene targets. *Nat Genet* 2016;48:481–7.
- 30 Eyre S, Bowes J, Diogo D *et al*. High-density genetic mapping identifies new susceptibility loci for rheumatoid arthritis. *Nat Genet* 2012;44:1336–40.
- 31 Yamamoto K, Okada Y, Suzuki A, Kochi Y. Genetic studies of rheumatoid arthritis. *Proc Jpn Acad Ser B Phys Biol Sci* 2015;91:410–22.
- 32 Pearson TA, Manolio TA. How to interpret a genome-wide association study. *JAMA* 2008;299:1335–44.
- 33 Okada Y, Wu D, Trynka G *et al*. Genetics of rheumatoid arthritis contributes to biology and drug discovery. *Nature* 2014;506:376–81.
- 34 Raychaudhuri S, Sandor C, Stahl EA *et al*. Five amino acids in three HLA proteins explain most of the association between MHC and seropositive rheumatoid arthritis. *Nat Genet* 2012;44:291–6.
- 35 Scally SW, Law SC, Ting YT *et al*. Molecular basis for increased susceptibility of Indigenous North Americans to seropositive rheumatoid arthritis. *Ann Rheum Dis* 2017;76:1915–23.
- 36 Furukawa H, Oka S, Shimada K, *et al*. Independent association of HLA-DPB1*02:01 with rheumatoid arthritis in Japanese populations. *PLoS One* 2018;13:e0204459.
- 37 Okada Y, Suzuki A, Ikari K *et al*. Contribution of a non-classical HLA gene, HLA-DOA, to the risk of rheumatoid arthritis. *Am J Hum Genet* 2016;99:366–74.
- 38 McAllister K, Yarwood A, Bowes J *et al*. Identification of BACH2 and RAD51B as rheumatoid arthritis

- susceptibility loci in a meta-analysis of genome-wide data. *Arthritis Rheum* 2013;65:3058–62.
- 39 Jiang L, Yin J, Ye L *et al.* Novel risk loci for rheumatoid arthritis in Han Chinese and congruence with risk variants in Europeans. *Arthritis Rheumatol* 2014;66:1121–32.
 - 40 Alamanos Y, Drosos AA. Epidemiology of adult rheumatoid arthritis. *Autoimmun Rev* 2005;4:130–6.
 - 41 Alarcón GS, Koopman WJ, Acton RT, Barger BO. DR antigen distribution in Blacks with rheumatoid arthritis. *J Rheumatol* 1983;10:579–83.
 - 42 Schiff B, Mizrahi Y, Orgad S, Yaron M, Gazit E. Association of HLA-Aw31 and HLA-DR1 with adult rheumatoid arthritis. *Ann Rheum Dis* 1982;41:403–4.
 - 43 Okada Y, Kim K, Han B *et al.* Risk for ACPA-positive rheumatoid arthritis is driven by shared HLA amino acid polymorphisms in Asian and European populations. *Hum Mol Genet* 2014;23:6916–26.
 - 44 van der Woude D, Lie BA, Lundström E *et al.* Protection against anti-citrullinated protein antibody-positive rheumatoid arthritis is predominantly associated with HLA-DRB1*1301: a meta-analysis of HLA-DRB1 associations with anti-citrullinated protein antibody-positive and anti-citrullinated protein antibody-negative rheumatoid arthritis in four European populations. *Arthritis Rheum* 2010;62:1236–45.
 - 45 Oka S, Furukawa H, Kawasaki A *et al.* Protective effect of the HLA-DRB1*13:02 allele in Japanese rheumatoid arthritis patients. *PLoS One* 2014;9:e99453.
 - 46 Peschken CA, Esdaile JM. Rheumatic diseases in North America's indigenous peoples. *Semin Arthritis Rheum* 1999;28:368–91.
 - 47 Williams RC, Jacobsson LT, Knowler WC *et al.* Meta-analysis reveals association between most common class II haplotype in full-heritage Native Americans and rheumatoid arthritis. *Hum Immunol* 1995;42:90–4.
 - 48 Diaz-Gallo LM, Ramsköld D, Shchetynsky K *et al.* Systematic approach demonstrates enrichment of multiple interactions between non-*HLA* risk variants and *HLA-DRB1* risk alleles in rheumatoid arthritis. *Ann Rheum Dis* 2018;77:1454–62.
 - 49 Höglund J, Rafati N, Rask-Andersen M *et al.* Improved power and precision with whole genome sequencing data in genome-wide association studies of inflammatory biomarkers. *Sci Rep* 2019;9:16844.
 - 50 Tam V, Patel N, Turcotte M *et al.* Benefits and limitations of genome-wide association studies. *Nat Rev Genet* 2019;20:467–84.
 - 51 Schaid DJ, Chen W, Larson NB. From genome-wide associations to candidate causal variants by statistical fine-mapping. *Nat Rev Genet* 2018;19:491–504.
 - 52 Westra HJ, Martínez-Bonet M, Onengut-Gumuscu S *et al.* Fine-mapping and functional studies highlight potential causal variants for rheumatoid arthritis and type 1 diabetes. *Nat Genet* 2018;50:1366–74.
 - 53 Farh KK, Marson A, Zhu J *et al.* Genetic and epigenetic fine mapping of causal autoimmune disease variants. *Nature* 2015;518:337–43.
 - 54 Suzuki A, Terao C, Yamamoto K. Linking of genetic risk variants to disease-specific gene expression via multi-omics studies in rheumatoid arthritis. *Semin Arthritis Rheum* 2019;49:S49–S53.
 - 55 Ding J, Orozco G. Identification of rheumatoid arthritis causal genes using functional genomics. *Scand J Immunol* 2019;89:e12753.
 - 56 Strand V, Cohen S, Schiff M *et al.* Treatment of active rheumatoid arthritis with leflunomide compared with placebo and methotrexate. Leflunomide Rheumatoid Arthritis Investigators Group. *Arch Intern Med* 1999;159:2542–50.
 - 57 Bathon JM, Martin RW, Fleischmann RM *et al.* A comparison of etanercept and methotrexate in patients with early rheumatoid arthritis. *N Engl J Med* 2000;343:1586–93.
 - 58 Ranganathan P. Pharmacogenomics in rheumatoid arthritis. *Methods Mol Biol.* 2008;448:413–35.
 - 59 Lee YC, Cui J, Costenbader KH *et al.* Investigation of candidate polymorphisms and disease activity in rheumatoid arthritis patients on methotrexate. *Rheumatology* 2009;48:613–7.
 - 60 Iannaccone CK, Lee YC, Cui J *et al.* Using genetic and clinical data to understand response to disease-modifying anti-rheumatic drug therapy: data from the Brigham and Women's Hospital Rheumatoid Arthritis Sequential Study. *Rheumatology* 2011;50:40–6.
 - 61 Chen Y, Zou K, Sun J, Yang Y, Liu G. Are gene polymorphisms related to treatment outcomes of methotrexate in patients with rheumatoid arthritis? A systematic review and meta-analysis. *Pharmacogenomics* 2017;18:175–95.
 - 62 Lima A, Bernardes M, Sousa H *et al.* SLC19A1 80G allele as a biomarker of methotrexate-related gastrointestinal toxicity in Portuguese rheumatoid arthritis patients. *Pharmacogenomics* 2014;15:807–20.
 - 63 Owen SA, Lunt M, Bowes J *et al.* MTHFR gene polymorphisms and outcome of methotrexate treatment in patients with rheumatoid arthritis: analysis of key polymorphisms and meta-analysis of C677T and A1298C polymorphisms. *Pharmacogenomics J* 2013;13:137–47.
 - 64 Plaza-Plaza JC, Aguilera M, Cañadas-Garre M *et al.* Pharmacogenetic polymorphisms contributing to toxicity induced by methotrexate in the southern Spanish population with rheumatoid arthritis. *OMICS* 2012;16:589–95.
 - 65 Qiu Q, Huang J, Lin Y *et al.* Polymorphisms and pharmacogenomics for the toxicity of methotrexate monotherapy in patients with rheumatoid arthritis: a systematic review and meta-analysis. *Medicine* 2017;96:e6337.
 - 66 Taylor JC, Bongartz T, Massey J *et al.* Genome-wide association study of response to methotrexate in early rheumatoid arthritis patients. *Pharmacogenomics J* 2018;18:528–38.
 - 67 Tanaka E, Taniguchi A, Urano W *et al.* Adverse effects of sulfasalazine in patients with rheumatoid arthritis are associated with diplotype configuration at the N-acetyltransferase 2 gene. *J Rheumatol* 2002;29:2492–9.
 - 68 Taniguchi A, Urano W, Tanaka E *et al.* Validation of the associations between single nucleotide polymorphisms or haplotypes and responses to disease-modifying

- antirheumatic drugs in patients with rheumatoid arthritis: a proposal for prospective pharmacogenomic study in clinical practice. *Pharmacogenet Genomics* 2007;17:383–90.
- 69 Wiese MD, Alotaibi N, O'Doherty C *et al.* Pharmacogenomics of NAT2 and ABCG2 influence the toxicity and efficacy of sulphasalazine containing DMARD regimens in early rheumatoid arthritis. *Pharmacogenomics J* 2014;14:350–5.
- 70 Yee J, Kim SM, Han JM *et al.* The association between NAT2 acetylator status and adverse drug reactions of sulfasalazine: a systematic review and meta-analysis. *Sci Rep* 2020;10:3658.
- 71 Kumagai S, Komada F, Kita T *et al.* N-acetyltransferase 2 genotype-related efficacy of sulfasalazine in patients with rheumatoid arthritis. *Pharm Res* 2004;21:324–9.
- 72 Rossi AM, Bianchi M, Guarnieri C, Barale R, Pacifici GM. Genotype-phenotype correlation for thiopurine S-methyltransferase in healthy Italian subjects. *Eur J Clin Pharmacol* 2001;57:51–4.
- 73 Yang SK, Hong M, Baek J *et al.* A common missense variant in NUDT15 confers susceptibility to thiopurine-induced leukopenia. *Nat Genet* 2014;46:1017–20.
- 74 O'Rielly DD, Roslin NM, Beyene J, Pope A, Rahman P. TNF-alpha-308 G/A polymorphism and responsiveness to TNF-alpha blockade therapy in moderate to severe rheumatoid arthritis: a systematic review and meta-analysis. *Pharmacogenomics J* 2009;9:161–7.
- 75 Zeng Z, Duan Z, Zhang T *et al.* Association between tumor necrosis factor- α (TNF- α) promoter -308 G/A and response to TNF- α blockers in rheumatoid arthritis: a meta-analysis. *Mod Rheumatol* 2013;23:489–95.
- 76 Maxwell JR, Potter C, Hyrich KL *et al.* Association of the tumour necrosis factor-308 variant with differential response to anti-TNF agents in the treatment of rheumatoid arthritis. *Hum Mol Genet* 2008;17:3532–8.
- 77 Quartuccio L, Fabris M, Pontarini E *et al.* The 158VV Fc γ receptor 3A genotype is associated with response to rituximab in rheumatoid arthritis: results of an Italian multicentre study. *Ann Rheum Dis* 2014;73:716–21.
- 78 Ruysen-Witrand A, Rouanet S, Combe B *et al.* Fc γ receptor type IIIA polymorphism influences treatment outcomes in patients with rheumatoid arthritis treated with rituximab. *Ann Rheum Dis* 2012;71:875–7.
- 79 Drozdziak M, Rudas T, Pawlik A *et al.* Reduced folate carrier-1 80G>A polymorphism affects methotrexate treatment outcome in rheumatoid arthritis. *Pharmacogenomics J* 2007;7:404–7.
- 80 Hayashi H, Fujimaki C, Daimon T *et al.* Genetic polymorphisms in folate pathway enzymes as a possible marker for predicting the outcome of methotrexate therapy in Japanese patients with rheumatoid arthritis. *J Clin Pharm Ther* 2009;34:355–61.
- 81 Liu SG, Gao C, Zhang RD *et al.* Polymorphisms in methotrexate transporters and their relationship to plasma methotrexate levels, toxicity of high-dose methotrexate, and outcome of pediatric acute lymphoblastic leukemia. *Oncotarget* 2017;8:37761–72.
- 82 Morisaki T, Gross M, Morisaki H *et al.* Molecular basis of AMP deaminase deficiency in skeletal muscle. *Proc Natl Acad Sci USA* 1992;89:6457–61.
- 83 Lennard L. Clinical implications of thiopurine methyltransferase—optimization of drug dosage and potential drug interactions. *Ther Drug Monit* 1998;20:527–31.
- 84 Black AJ, McLeod HL, Capell HA *et al.* Thiopurine methyltransferase genotype predicts therapy-limiting severe toxicity from azathioprine. *Ann Intern Med* 1998;129:716–8.
- 85 Corominas H, Domènech M, Laiz A *et al.* Is thiopurine methyltransferase genetic polymorphism a major factor for withdrawal of azathioprine in rheumatoid arthritis patients? *Rheumatology* 2003;42:40–5.
- 86 Weinshilboum RM, Sladek SL. Mercaptopurine pharmacogenetics: monogenic inheritance of erythrocyte thiopurine methyltransferase activity. *Am J Hum Genet* 1980;32:651–62.
- 87 National Institute for Health and Care Excellence. (2019). DMARDs Scenario: Azathioprine. <https://cks.nice.org.uk/dmards> (7 March 2020, date last accessed).
- 88 Schaeffeler E, Jaeger SU, Klumpp V *et al.* Impact of NUDT15 genetics on severe thiopurine-related hematotoxicity in patients with European ancestry. *Genet Med* 2019;21:2145–50.
- 89 Nam JL, Winthrop KL, van Vollenhoven RF *et al.* Current evidence for the management of rheumatoid arthritis with biological disease-modifying antirheumatic drugs: a systematic literature review informing the EULAR recommendations for the management of RA. *Ann Rheum Dis* 2010;69:976–86.
- 90 Plant D, Bowes J, Potter C *et al.* Genome-wide association study of genetic predictors of anti-tumor necrosis factor treatment efficacy in rheumatoid arthritis identifies associations with polymorphisms at seven loci. *Arthritis Rheum* 2011;63:645–53.
- 91 Massey J, Plant D, Hyrich K *et al.* Genome-wide association study of response to tumour necrosis factor inhibitor therapy in rheumatoid arthritis. *Pharmacogenomics J* 2018;18:657–64.
- 92 Rönnblom L, Eloranta ML. The interferon signature in autoimmune diseases. *Curr Opin Rheumatol* 2013;25:248–53.
- 93 National Institute for Health and Clinical Excellence. (2015) Final Appraisal Determination Adalimumab, etanercept and infliximab for the treatment of rheumatoid arthritis. <https://www.nice.org.uk/guidance/ta130/documents/adalimumab-etanercept-and-infliximab-for-the-treatment-of-rheumatoid-arthritis-final-appraisal-determination3> (7 March 2020, date last accessed).
- 94 Finckh A, Liang MH, van Herckenrode CM, de Pablo P. Long-term impact of early treatment on radiographic progression in rheumatoid arthritis: a meta-analysis. *Arthritis Rheum* 2006;55:864–72.
- 95 Martin AR, Kanai M, Kamatani Y *et al.* Clinical use of current polygenic risk scores may exacerbate health disparities. *Nat Genet* 2019;51:584–91.