

Integrating pollinator decline into macroeconomic projections reveals uneven vulnerabilities from climate-driven biodiversity loss

Campoleoni Mansi G. A.^{1,2,3}, Standardi G.^{1,4}, Bosello F.^{1,2,3}

¹ CMCC Foundation - Euro-Mediterranean Center on Climate Change, Via Marco Biagi 5, Lecce, 73100, Italy

² RFF-CMCC European Institute on Economics and the Environment, Via Bergognone 34, Milan, 20144, Italy

³ CaFoscari University, Dorsoduro 3246, Venice, 30123, Italy

⁴ Università degli Studi di Milano, Via G. Celoria, 20, Milan, 20133, Italy

June 05, 2026

Abstract

Ecosystems and biodiversity sustain human welfare through essential services such as pollination, collectively contributing a significant proportion to global GDP. Yet their representation in long-term socioeconomic projections remains critically limited, leading to biased outlooks, particularly under accelerating environmental change. To address this gap, we develop a spatially explicit, integrated biophysical and economic modelling framework in which ecological impacts propagate through agricultural production systems into the macro-economy.

This study delivers the first long-term evaluation of pollination impacts under climate change, using a multi-model integrated assessment that links pollinator species distribution models to macroeconomic outcomes at sub-national resolution. Coupling an ensemble of 18 scenarios covering 153 wild pollinator species with high-resolution data for 129 crops and a regionalized computable general equilibrium model, we quantify agricultural productivity losses and economy-wide effects across 122 EU regions to 2070.

Results highlight the spatial heterogeneity of pollination decline and its uneven effects on pollinator-dependent crops across regions, revealing critical vulnerabilities in the agricultural systems. We find that EU-wide average impacts, which shows GDP losses up to €57.7 billion/year (-0.18%) and agricultural sector losses of -1.5 to -6.0% in 2070, systematically conceal severe spatial concentration of damages. Under high-emission scenarios, cropland productivity losses for vegetables, fruits and nuts in 2070 reach -25.5% in the most exposed regions, with sectorial economic impacts spanning between -21.35 and +8.85%. Mediterranean and eastern regions, already structurally disadvantaged, bear disproportionate burdens, amplifying intra-EU economic disparities. Standard integrated assessment models, typically resolved at macro-regional level, may systematically underestimate the vulnerability to climate-biodiversity impacts as aggregation conceals distributional tails by factors of 2-5×. These results demonstrate that biodiversity impacts cannot be properly captured by national-scale assessments and that spatial resolution is not a technical choice but a determinant of whether risks are visible at all.

Context

Globally, animal pollination significantly contributes to global crop production [54][29][6][28], with annual global economic output depending on pollinators ranging between US\$235 and US\$577 billion [45]. In Europe, 84% of the crops grown benefit at least partly from animal pollination [21], including almost all fruits, vegetables and nuts, oil crops (notably oilseed rape), pulses (beans, peas, lentils, etc), cotton, flax, feed and forage plants (as many protein-rich and nitrogen-fixing legumes, beans, clovers, alfalfa, vetches, lupins, etc). Knowledge of how pollinators

provide pollination services to crops has advanced (e.g. [92] [79], [55]), showing how this service is strongly linked to species diversity [108][32] and pollinators abundance. While some pollinator species are particularly common crop flower visitors overall, different pollinator communities are important for the pollination of different crops [31][33]. The economic value of pollination for agriculture in the EU is estimated at EUR 5–15 billion annually[21]. In 2019 pollination contributed €4.977 billion into the value of agricultural production in the EU, corresponding to €2 810 per km², while pollinator-dependent crops are expanding at 5% per decade [22][23][61]. Over 78% of wild flowering plants in the EU rely on insect pollination [69], including many medicinal plants, with recent estimates suggesting that up to 90% of flowering plant species depend on animal pollination [48].

Without pollinators, these plants would disappear resulting in the cascading loss of biodiversity and ecosystem services, such as biological pest control provided by some hoverfly larvae, pest regulation and support of natural enemies in agriculture [42]. Recent assessments confirm an accelerating decline[106]: the number of threatened European butterfly species has increased by 76% over the last decade, with 15% of butterflies (65 out of 442 assessed species) now at extinction risk and more than 20% of bumblebee species classified as threatened; nearly 40% of hoverfly species, 20% of butterflies, and 9% of bees are threatened with extinction [21]. There is evidence of current pollination deficits in crops associated with low abundance of pollinators; JRC [102] estimated 50% of pollinator-demanding crops in the EU are in pollinator-deficient areas. In the absence of restoration, the pollination gap is expected to widen, as pollination potential gradually declines while demand for pollination likely increases with the expansion of pollination-dependent crops [61].

Habitat degradation and climate change are globally acting as pivotal drivers of wildlife collapse, with mounting evidence that this erosion of biodiversity will accelerate in the following decades [35][46]. Habitat for wild pollinators, as natural and semi-natural grasslands [56][45][77], have been drastically reduced due to agricultural intensification and leads to loss of natural and semi natural habitats rich in wildflowers (which pollinators use for foraging, nesting and hibernating) [8][76][37][37][117]. Best estimates assess that in 2030, 27% of European agro-ecosystems will require management and restoration practice due habitat degradation, while the ecosystem loss is estimated at 1.5% per year. Projections of pollinators communities, as well as common farmland bird and grassland butterfly indicators, suggest that these groups will continue to decline into the future, likely driven by agricultural practices and land use [71].

Moreover, although climate change impacts to date are not likely to be among the greatest[23], they are expected to increase [47], together with indirect drivers, such as increased irrigation and production of bioenergy. Rising temperatures from climate change are expected to increasingly impact pollinators, influencing their geographic ranges, population sizes, and seasonal activity patterns[44][85][86][116], possibly leading to extinctions of some species in the southern and northern edges of ranges and in alpine species, whilst driving the expansion of other species to colonise new areas EU (2023).

Ecological assessments project pollinator range shifts under climate change and land use change with increasing sophistication, leveraging species distribution models (SDMs) calibrated on observed biogeographic data. Ghisbain et al, 2024 [35], identifies future large-scale range contractions and species extirpations under all future climate and land use change scenarios considered, showing that temperature and precipitation have strong negative correlations with ecological suitability difference (ESD), and suggesting that warming and altered rainfall patterns are major contributors to habitat loss, especially for cold-adapted species. Around 38-76% of studied European bumblebee species currently classified as ‘Least Concern’ are projected to undergo losses of at least 30% of ecologically suitable territory by 2061-2080 compared to 2000-2014. Prestele et al., 2021 [82], assess the future species distribution under three RCP-SSP scenarios. They find that direct climate impacts, although variable across species, dominate responses for most species, especially under high-end climate change scenarios (up to 99% range loss). Land use may amplify, attenuate, or offset changes to suitable habitat extent expected from climate impact depending on species and scenario, especially in low-end climate change scenarios RCP2.6, indicating substantial potential for targeted conservation management. Rahimi et al., 2024 [85], 2025 [86], suggest that on a global scale approximately

65% of bees are likely to witness a decrease in their distribution, with climate change's impact on bees projected to be more severe in Europe (average reduction of 56%) under SSP585. Under the most pessimistic climate change scenario (SSP585), they estimate that Europe is expected to see a 4.7% decrease in area suitability for pollination-dependent crops by 2070. Climate change's anticipated effects on bee distributions could potentially disrupt existing pollinator-plant networks, posing ecological challenges that emphasize the importance of pollinator diversity, synchrony between plants and bees, and the necessity for focused conservation efforts. Kazenel et al., 2024 [53], document abundance patterns for a hyper-diverse bee assemblage including many bee species are ecologically and evolutionarily different from bumblebees, linking bee declines with heat and desiccation tolerances and using climate sensitivity models to project pollinators communities into the future. Models forecasted declines for 46% of species and predicted more homogeneous communities dominated by drought-tolerant taxa, even while total bee abundance may remain unchanged. Such community reordering could reduce pollination services, because diverse bee assemblages typically maximize pollination for plant communities.

While these studies produce detailed, climate-driven projections of habitat suitability changes for specific pollinator taxa, they stop short of translating ecological outputs into economy-wide consequences, leaving the macroeconomic implications of climate-biodiversity interactions unquantified. Accordingly, most economic models fail to account for the impact of biodiversity loss - particularly pollinator decline - on agricultural output, reflecting a broader gap in the representation of ecosystems and biodiversity that leads to an underestimation of their importance in future projections. Moreover, the overall impact of climate change on pollination-dependent crops, such as fruits and vegetables, is commonly not accounted for in these models. This is especially concerning given the high economic, cultural and dietary importance of such crops in the EU.

Economic assessments of pollinator decline to date have followed two distinct methodological paths, none of which fully captures climate-driven risk at high spatial resolution.

Stylized 'catastrophic loss' frameworks impose exogenous productivity shocks to the economy [29][6][28], abstracting from the spatial heterogeneity and the climate-driven dynamics of pollinator distributions. They rely on partial or general-equilibrium models to simulate economic impacts of an hypothetical collapse of wild pollinators. Feuerbacher et al., 2025 estimates crop yield reductions in Europe by 2030 of approximately 8% and global welfare losses of €34.4 billion. While these frameworks deliver important insights into the economic relevance of pollinators, they share three key limitations: (i) they impose binary or stylized loss scenarios rather than representing the climate-driven dynamics that progressively reshape pollinator ranges over time; (ii) they operate at relatively coarse spatial resolution (typically national or NUTS-1 level); and (iii) they abstract from the spatial heterogeneity of pollinator habitats and crop-specific dependencies within their aggregation units. A second stream combines static pollination-sufficiency proxies with land-use scenarios derived from integrated assessment models [109][50]. While these approaches incorporate spatial detail and consider future land allocation, they rely on aggregated indicators of pollination service rather than dynamic climate change response, limiting their capacity to represent the interactions between climate forcing, pollinator habitat suitability, and agricultural productivity.

As a result, no integrated framework has yet propagated multi-model, climate-driven pollinator projections through a spatially explicit macroeconomic model across multiple emission and socioeconomic scenarios at sub-national resolution. Our assessment aims to fill these gaps by quantifying the macroeconomic effects of expected productivity loss of the EU agricultural sector due to pollinators' decline under long-term projections of climate change. Accordingly, this study develops and applies such a framework for the EU27, delivering the first long-term evaluation of pollination impacts based on a multi-model, multi-scenario SDM projection.

Building on ecosystem accounting standards, we quantify changes in pollination ecosystem services across the European Union using observed long-term trends in wild pollinator populations and ecosystems degradation proxied by European biodiversity indicators, together with climate-driven projections derived from an ensemble of state-of-the-art species distribution models (SDMs) covering up to 153 wild pollinator species across major taxonomic groups. The resulting spatially explicit ecological projections are matched with high-resolution crop

cultivation areas (0.05°) and crop-specific pollination dependency ratios for 129 European crops to estimate high-resolution agricultural productivity impacts under multiple climate scenarios (RCP2.6, 4.5, 6.0, 7.0, and 8.5) and socioeconomic pathways (SSP1, 2, 3, and 5) up to 2070. These biophysical impacts are then translated into second-order macroeconomic shocks using a regionalized version of the ICES computable general equilibrium (CGE) model, enabling the assessment of economy-wide effects down to NUTS-2 regional resolution.

An overview of the research framework and methods is provided in here and in section Methods below. A detailed description of data processing and modelling framework is provided in *Appendix 1. Extended Methods, Materials and Data Processing*. *Appendix 2. Pollination current status and projected trends*, provides the background and scientific foundation of pollination loss dynamics supporting the Baseline and Impact scenario, while caveats, novelty and differences with existing literature are provided in *Appendix 3*.

Main

This study presents the first climate-driven, multi-model integrated assessment of the macroeconomic consequences of species-explicit pollinator decline assessing sub-national heterogeneity. By coupling an ensemble of 18 Species Distribution Model (SDM) scenarios covering up to 153 wild pollinator species with high-resolution crop data and a spatially explicit Computable General Equilibrium (CGE) model, we resolve climate-biodiversity-economy interactions down to NUTS-2 level across the EU27 to 2070. Pollination projections highlight the spatial heterogeneity of pollination decline and its uneven effects on pollinator-dependent crops across regions and climate scenarios. Our central finding is that EU-wide impacts that appear moderate in aggregate, with EU GDP losses of €29.5 (-0.09%) to €57.7 (-0.18%) billion-year by 2070, and agricultural-sector losses of -1.5% to -6.0%, systematically conceal a severe spatial concentration of damages, with most exposed regions experiencing impacts two to five times larger than the country mean.

Climate change drives spatially uneven decline in pollination potential

Projected changes in pollination service potential with respect to 2020 exhibit a clear and consistent pattern across pollination scenarios considered, with significant losses occurring in all climate change scenarios also in the first half of the century. By 2070, EU-wide averages range from about -22 to -26% under low-radiative-forcing trajectories (RCP2.6) to -62% to -66% under high-end trajectories (RCP7.0, RCP8.5), with intermediate scenarios spanning these bounds (figure 3). No scenario produces a positive EU-wide signal by 2070, confirming the directional robustness of the result even under large inter-model spread, while the model choice becomes decisive under high-forcing scenarios (up to 36 percentage points of spread across models under RCP8.5) (Appendix 1 for full data and models sensitivity).

The aggregate trend, however, hides growing heterogeneity as spatial detail increases. As spatial detail increases from EU averages to regional distributions, heterogeneity grows sharply. While most regions consistently show declines across all scenarios, the magnitude of results diverge depending on the specific Species Distribution Model (SDM) or the radiative forcing (RCP) underlying the pollination scenario. Particularly, some Central-Eastern and Northern European regions show localized gains of up to +24% (2050) and +16% (2070) under low climate forcing scenarios (LF), while others experience negligible impacts even under moderate (MOD) to high-end (HE) scenarios. These cases are few but noteworthy, as they demonstrate that the response is not uniform: while many Mediterranean and eastern European regions experience severe losses (average -30% or more), other areas may temporarily benefit from more favourable climatic conditions, suggesting a localized increase in pollination services. This spatial variability underscores the importance of regional analysis: the more granular the lens, the greater the cross-regional heterogeneity of biodiversity loss; a pattern that propagates into every downstream economic outcome. Changes in pollination services on agricultural land influence cropland productivity and are reflected in yield losses, revealing critical vulnerabilities in the EU agricultural systems.

Agricultural productivity losses show high regional and sectorial heterogeneity

Out of the 129 analysed crops and crop categories cultivated in the EU, 103 were found to be at least partially dependent on pollination for commercially viable yields (dependency rates between 0.05 and 0.95). Within this group, 29 items exhibit a high dependency (≥ 0.65), while 39 fall within the intermediate range (0.50 - 0.25). When assessing cropland productivity losses on fixed 2020 agricultural areas (Figure 1), results reveal a predominantly negative pattern across all the analysed agri-sectors and scenarios, with the magnitude, rather than the direction of losses, being the key differentiator. However, pronounced heterogeneity across scenarios and regions is found, with a minority of northern EU regions showing limited or positive yield changes under both moderate (MOD) and high-end (HE) emission scenarios, while southern and central EU countries bearing the highest losses, a nuance that is entirely masked when results are averaged across macro-regions.

EU-wide average yield losses for the aggregate agricultural area rise from -2.4% in both 2050 and 2070 under low climate forcing scenarios, to -5.3% under high-end climate change scenarios by 2070, with a cross-scenario median of -3.9% (range -2% to -8.5%). This value is consistent with recent literature[28] showing that wild-pollinator collapse yields a -7.8% (-5.2% to -10.3% range) European crop-yield decline by 2030, supporting the plausibility of our climate-driven trajectory as a more gradual realisation of the same risk.

Losses roughly double between the lowest and highest forcing trajectories across the agricultural sub-sectors considered, with limited temporal progression except for high end emission scenarios, and reflect the underlying pollination dependency and cultivation areas. Vegetables, Fruits and Nuts sectors are most affected, with EU averages reaching -7.5% and maximum NUTS-3 losses of -25.5% in 2070, followed by Other Crops sector and Oilseeds (maximum -14.7%). Cereals, Wheat, Rice, Sugar cane and Beets sectors show negligible impacts at all aggregation levels and scenarios.

While localized yield gains do emerge at all NUTS level (0-3), they are strongly scenario-dependent and vanish almost entirely as forcing increases. Under scenarios with low climate change trajectories, on average, 12.4% of NUTS-3 level regions experience positive impacts (between 0 and 2.88%) on agricultural land in 2050, while only 7.7% in 2070 (0 - 5.22%). Under the PREST-1 2.6 and PREST-CLIM 2.6, between 59 and 110 NUTS-3 regions show gains in 2050, spreading across a geographically diverse set of countries including Finland, Germany, Poland, Romania, Spain and Bulgaria. Maximum gains reach 2.9% for the whole Agricultural Sector, 6.2% for Oilseeds, and 11.5% for Vegetables and Fruits in 2070. These positive cases diminish sharply under moderate emission scenarios (fewer than 5 regions in isolated scenarios) and are negligible under high-end emission scenarios. This pattern indicates that localized positive yield responses are possible under low emissions, but are highly sensitive to the specific combination of socioeconomic and climate assumptions, and disappear almost entirely as forcing increases. Crucially, the severity of these losses is masked by aggregation, with standard deviation and tail losses growing systematically with spatial resolution. Productivity losses greater than -10% for the entire agricultural sector under high-impact scenarios in 2070 occur in 5 country-level cases (18.5%), and in 33 NUTS-2 regions and 155 NUTS-3 regions (13.5%). Losses between -15% and -20% are entirely absent at the NUTS-0 scale and only partially visible at NUTS-3, where 12 regions exceed this threshold; no regions experiences losses beyond -20%. When deepening the sectorial detail, for Vegetables, Fruits and Nuts, higher losses emerge even at higher aggregation level, with 12 (44.4%) countries crossing the -10% threshold against 371 (32.2%) NUTS-3 regions. Only 3 countries experience losses from -15% to -20% against 129 NUTS-3. The worst-case losses are borne by Bulgaria and deepens from -22% at the country level to -32.5% at NUTS-3 level. Critically, lower-tail percentiles deepen far more sharply than standard deviation as spatial resolution increases: the 1st-percentile loss moves from -21.3% at NUTS-0 to -25.1% at NUTS-3 (an 18% deepening of the tail, against only 1.5% growth in standard deviation), confirming that dispersion alone understates the aggregation effect. The masking is also not confined to traditionally vulnerable southern regions: Germany combines a NUTS-0 Vegetables, Fruits and Nuts loss of -4.2% with a worst NUTS-3 value of -12.1%, nearly triple the national figure, evidence that hidden regional vulnerabilities arise even in countries with modest national averages.

This aggregation effect is negligible for low impacted crop sectors, but critical for Vegetables, Fruits and Nuts,

Oilseeds and Other crops sectors, under high-impact scenarios, underscoring that high spatial resolution is indispensable for identifying local vulnerabilities from biodiversity loss that would otherwise be systematically masked by national averaging. This systematic invisibility of regional risk at the national scale is not a feature of pollination alone, it is a general property of how spatial aggregation interacts with biodiversity-driven productivity shocks. Standard integrated assessment models, typically resolved at country or macro-region level, are therefore likely to underestimate the spatial concentration of climate-biodiversity damages across other ecosystem services as well.

Full data on projected yield changes for 129 crops and agri-sectors in the EU at NUTS (0-3) level are provided in the Extended Material.




Figure 1 | Cropland productivity changes in 2070 relative to 2020 for different crop categories and for the aggregate Agricultural sector, shown at the EU27 country level and for NUTS (0-2) regions. Maps display the median impact values for different crop-sectors, alongside the highest and lowest impacts across all scenarios considered for the Agricultural sector as a whole, illustrating the range of uncertainty driven by pollination and climate scenario assumptions. The table reports country-level median (50th percentile), and extreme values (low, high) for each crop category. Productivity losses are expressed as percentage change on fixed 2020 cropland areas, with no SSP-induced land use adjustments.

Macroeconomic propagation absorbs aggregate losses but concentrates regional vulnerability

The preceding analysis demonstrates that aggregating biophysical impacts at the country or EU level systematically masks local extremes, particularly for high-value sectors such as Vegetables, Fruits, and Nuts. Embedding these productivity shocks in the CGE model reveals two opposing mechanisms: endogenous adjustment buffers economy-

wide losses, while regional and sectoral distress intensifies.

We estimate changes from the respective SSP baselines across 18 scenarios for the regional GDP, the value of aggregated Agricultural Sector output, and the value of agricultural sub-sectors output (Vegetables, Fruits and Nuts; Oilseeds; Cereals, Wheat & Rice; Sugar Cane & Beets; Other Crops). A comparison between EU-wide means and NUTS-level distributions reveals that the masking effect observed for physical yields is amplified for regional aggregates, while the degree of masking increases with higher levels of sectoral aggregation, becoming progressively stronger when moving from the aggregate macroeconomic variable GDP, to the sectoral aggregate Agricultural Sector, and ultimately to individual agri-sector outputs, where extremely pronounced regional differences emerge (Figure 2; Table 1).




Figure 2 | Economic impacts in 2070. Maps show the median impact values (50th percentile) for individual crop sectors, the aggregated agricultural sector, and regional GDP across 122 NUTS-2 regions under Moderate Forcing scenarios. The line graph shows the aggregated EU GDP impact, expressed as percentage deviation from the respective baselines, for all 18 scenarios considered, as well as the median values for Low Forcing (LF), Moderate Forcing (MOD), and High-end Emissions (HE) scenarios.

By 2050, European GDP losses range between approximately -0.15% and -0.02%, even under the most severe climate scenarios, reaching -0.18% by 2070, with an average loss across all scenarios of -0.09% in both years. Although these figures may appear small, their absolute magnitude is not negligible, corresponding to a net GDP loss up to €37.85 billion (-0.15%) in 2050 and €57.7 billion (-0.18%) in 2070 for high-end scenarios, expressed in 2017 values with respect to SSP2 baseline GDP level. Low-forcing scenarios display European aggregate losses between -0.10% to

-0.08% in 2050 and -0.09% to -0.06% in 2070, with an average loss across all scenarios of -0.09% and -0.07% , thus indicating a decreasing magnitude of impacts for low climate change scenarios. These figures corresponding to up to €26.6 billion (-0.10%) in 2050 and €29.5 billion (-0.09%) in 2070, expressed in 2017 values with respect to SSP1 baseline GDP level. This apparent resilience reflects sectoral reallocation and trade adjustment, together with the limited weight of agriculture in EU GDP (1.6% in the base year), which dampens the macroeconomic visibility of even substantial agricultural shocks. The aggregate, however, conceals sharp regional divergence: under high-end forcing, regional GDP impacts range from -1.61% to +0.17% by 2070, and one quarter of EU regions face losses exceeding -0.19%, more than double the EU average.

The masking strengthens systematically with sectoral resolution. For the aggregate Agricultural sector, EU-wide losses of -1.5% to -6.0% (cross-scenario average -3.2% to -3.5%) conceal a regional distribution spanning -12.78% to +7.96% in 2070, an amplification factor (NUTS over EU average) of 2.0× to 3.6×.

For Vegetables, Fruits and Nuts the concealment is extreme: EU-wide losses of -1.9% to -7.0% mask regional losses reaching -21.35%, an amplification of 3.4× to 5.6× and an absolute gap exceeding 14 percentage points between the EU value and the worst-affected region. Under high-end scenarios this divergence becomes a polarisation that aggregate indicators cannot show. For the Agricultural Sector, between 2050 and 2070 the EU-wide mean and median improve slightly, mostly driven by gains in northern and alpine regions where warming extends growing seasons and enhances pollinator habitat suitability, while the lower tail deepens and the share of regions losing more than 10% of agricultural output nearly triples (0.7% to 1.9%). For Vegetables, Fruits and Nuts, more than one third of regions (34.7%) record gains by 2070 even as the worst-affected deteriorate to -21.35%, with 26.3% and 7.6% of regions losing more than 5% and 10% respectively. This "internal divergence", with some regions benefiting while others collapse, represents a fundamental redistribution of climate–biodiversity risk that is statistically invisible in EU-level means.

The remaining sub-sectors mapped in Figure 2 mirror the cropland-productivity ordering of Figure 1: Oilseeds and Other Crops& fibers show intermediate to very high economic deviations, while Cereals, Wheat, Rice and Sugar Cane& Beet show low to negligible negative impacts. Interestingly, the economic gain deriving from macroeconomic effects is evident even under median forcing scenarios, with many regions experiencing both sectorial impacts higher than +5%/+10%, boosted by the relative smaller impact (sometimes opposite sign impact) with respect to other regions.

The emerging systematic masking results as a structural property of how spatial aggregation interacts with the joint distribution of pollination loss, crop dependency, and economic structure. Across all 18 scenarios, the amplification factor remains consistently in the 3.4-5.6× range for vegetables, fruits and nuts, substantially exceeding the 2.0-3.2× factor observed for the aggregate agricultural sector. The interquartile range of regional impacts significantly expands from GDP estimates to sectorial cropland productivity, indicating that macroeconomic adjustment mechanisms compress the visible spread of biophysical outcomes and the underlying vulnerability.

Discussion

This study delivers the first climate-driven, multi-model and multi-scenario assessment of the macroeconomic consequences of pollinator decline at sub-national resolution. Our findings complement, and contextualise, the two established strands of pollinator economics. Stylized "catastrophic collapse" frameworks, imposing binary productivity shocks, have established the first-order economic relevance of pollinators [29][6][28]. Our cross-scenario median agricultural loss of -3.9% (-2 to -8.5%) is consistent with this estimate as a gradual, climate-driven realisation of the same risk on the long term, providing mutual validation across the two approaches. A second stream combines static pollination-sufficiency proxies with land-use scenarios [109][50], adding better ecological detail but not dynamic climate response. Our framework is the first to integrate both dimensions: time-evolving, climate-driven pollinator ranges and sub-national heterogeneity in service supply and economic dependence. Our results show a structural insight of pollination decline: the propagation of biodiversity loss through the

economy generates a systematic spatial masking that intensifies at every level of aggregation, so that for any given EU-wide average the most vulnerable regions experience impacts two to five times greater, losses entirely concealed in national aggregates.

Three nested mechanisms produce this masking. First, the composition of each region's cropland plays a pivotal role in determining the resulting sectoral shocks on agriculture. The relative share of each crop in the composition of the total output, shapes the aggregate sectoral impact according to its specific degree of dependence on animal pollination and the pollination service change in the cultivated area. This means that those regions severely affected by absolute losses in pollination service may nonetheless face comparatively smaller agricultural impacts if their cropland composition is dominated by crops that are less reliant on animal pollination, and vice versa. Furthermore, regions that register some of the largest losses at finer spatial scales (NUTS 2-3) may exhibit more moderate impacts once these effects are aggregated at broader regional or national levels as biophysical and economic exposure decouple spatially, so that negative impacts in some areas may be partially offset by positive or negligible effects in others. Second, endogenous economic adjustment, such trade, substitution, and the reallocation of land and capital, compresses the visible spread of shocks at the aggregate level while concentrating residual vulnerability in the regions least able to adapt. The translation of severe regional biophysical impacts on cropland productivity into macroeconomic outcomes shows how endogenous economic adjustments simultaneously attenuate aggregate losses and concentrate vulnerabilities in already disadvantaged regions, rendering these risks nearly invisible in national-scale assessments. Third, statistical aggregation mechanically conceals tail losses, an effect that grows with the size of the aggregation unit. Together these effects explain why EU GDP appears only moderately affected (-0.09% on average) while a quarter of regions lose more than twice that, and the worst regions lose up to -21.35% in in Vegetables, Fruits and Nuts sector.

Geographically, the burden falls disproportionately on Mediterranean and eastern European regions, which combine the most severe biophysical impacts (average pollination losses up to -30% in some countries) with cropland-mix highly dependent to pollination, and the least economic capacity to absorb agricultural shocks. Under high-emission scenarios, parts of southern Italy, Greece, Bulgaria and the Iberian Peninsula record sectoral losses exceeding 43% for key crop categories, while northern regions remain stable or marginally positive. The economic burden of biodiversity loss therefore reinforces, rather than offsets, existing structural inequalities. The emerging aggregation bias is not specific to Europe: any country-level assessment of pollination loss risks understating the exposure of precisely those agricultural communities least able to absorb productivity shocks. This distributional pattern mirrors, at sub-national scale, the global inequality documented for natural capital loss between countries [5].

The implications for long-term socioeconomic projections are unambiguous, assessing severe regional and sector-specific economic losses, and showing how country-level resolution systematically underestimates the spatial concentration of climate-biodiversity damages. Our results demonstrate that for any given EU-wide average loss, the most vulnerable regions experience impacts 2-5 times greater, losses that remain entirely concealed in national aggregates: a pattern that may extend beyond pollination, reflecting a general property of any ecosystem service characterised by geographic heterogeneity in both service provision and economic dependence. Standard integrated assessment models, typically resolved at the country or macro-region level, are therefore likely to ignore the spatial concentration of climate impacts across all ecosystem services characterized by geographic heterogeneity in both service provision and economic dependence, underestimating the impacts of biodiversity loss. Spatial resolution is therefore not a technical refinement but a determinant of whether critical vulnerabilities are visible in policy-relevant assessments at all. For pollinator policy specifically, the concentration of damages in structurally disadvantaged regions argues for spatially targeted restoration and conservation under the Nature Restoration Regulation [23][24], directed to the areas where pollination deficits and economic exposure coincide.

Several considerations suggest that the aggregate figures, moderate as they appear, should be read as a conservative lower bound on the true welfare cost. Because climate enters our framework only through pollinator habitat suitability, the results capture the pollination channel in isolation: direct climate effects on yields are excluded, and

the spatial concentration of damages we document should be interpreted as the pollination-specific component of a larger, partly co-located climate burden. Second, our framework propagates pollination loss only through the commercial yield of pollinator-dependent crops, and therefore omits the cascading effects of pollinator decline on the broader ecosystems in which agriculture is embedded. Pollination is not only an agricultural input but a keystone regulating service: the large majority of wild flowering plant species in the EU depend on animal pollination [69][48], so that sustained pollinator loss propagates into the reproduction and persistence of wild flora, the food webs they support and the configurational landscape heterogeneity on which other taxa depend [42][45], deepening the regional damages reported here. The macroeconomic cost we estimate is thus best understood as the lower-bound of a wider, largely non-market, loss.

Taken together, our results show that the macroeconomic risk of pollinator decline under climate change is real but profoundly unequal, and that its most severe manifestations are precisely those that national-scale economics renders invisible. Capturing them requires resolving biodiversity–economy interactions at the scale at which they actually occur.

Methods

We developed a modular, climate-driven, biophysical-to-economic cascade framework that integrates a multi-model, multi-scenario ensemble of pollinator species distribution projections with a spatially explicit computable general equilibrium model. The framework links four sequential modules: (i) climate-driven pollination potential under 18 SSP-RCP scenarios; (ii) crop-specific yield response based on pollination dependency ratios for 129 EU crops at a resolution of 0.05°; (iii) regionalized agricultural shocks aggregated to NUTS 0-3 levels; and (iv) macroeconomic propagation through a 122-region CGE model resolved to NUTS-2. The framework operates as a sequential, modular cascade, in which outputs from each ecological and agricultural module are consistently propagated as inputs into subsequent economic modules, enabling an internally consistent assessment of ecological and socioeconomic impacts. While the framework does not explicitly model endogenous feedbacks from economic outcomes to ecological dynamics [4], it captures the one-way propagation of ecosystem services changes through agricultural systems into the macro-economy in a spatially explicit manner, consistent with the production-function method of the SEEA-EA ecosystem accounting standards (United Nations, 2024; Markandya et al., 2022).

Baseline pollination potential scenario

The KIP-INCA framework [101, 61] developed a set of ecosystem services accounts including crop pollination. The service is provided by the functional match between pollination potential, defined as the ecosystem’s ecological suitability to support wild insect pollinators, and pollination demand, defined as the spatial extent of pollinator-dependent crops (the use area). The spatial co-occurrence of pollination potential and pollination demand is used to quantify the actual flow of the service and to identify priority areas where restoration or conservation interventions are most needed to mitigate pollination deficits.

The KIP-INCA datasets [103] classifies EU land surface into categories of pollination potential based on a highly detailed assessment of land-cover composition and land-use patterns, aimed at identifying habitat elements suitable for pollinators. This classification enables us to project how future changes will differentially affect different regions based on projected future climate and land-use trends (see sections below). The assessment is based on an indicator of the environmental suitability to support wild insect pollinators which integrates two different models: an Expert-Based Model for solitary bees and a Species Distribution Model (SDM) for bumblebees. Both models are based on land use, land cover and on the distance to semi-natural areas, being the most important drivers, while their relative importance differs among the taxonomic groups. The presence of semi-natural areas in agricultural landscapes (such as hedgerows, wildflower strips, and fallow land) is included as an element providing food resources and nesting sites, based on Garibaldi et al. (2011) which shows that the stability of pollination

services decreases with isolation from natural areas. LULC data, CORINE data and TeleAtlas® Maps, are used to identify semi-natural areas and areas that cannot provide floral resources or nesting sites to insect pollinators. Current climate is also considered as an important driver of the large-scale occurrence of the groups of pollinators, but other pressures affecting pollinators are not considered, such as the use of pesticides, climate change and invasive alien species.

To build the pollination potential baseline (2020-2070), the initial level of Pollination Potential for the year 2020 relies on data from the [INCA \(2018\) potential](#) [101], normalized to a 0-1 scale. To ensure consistency with existing literature, we compared this metric against the 2015 Pollination Sufficiency baseline from Von Jeetse et al. (2023) [109], which incorporates similar ecological and spatial elements. The two metrics proved to be broadly comparable, with closely aligned mean values across the EU (See Appendix 1).

The INCA pollination potential represents a snapshot of current ecological conditions and does not explicitly incorporate temporal trends. Likewise, species distribution models (SDMs) generally omit several relevant stressors, including pollutant exposure. Future changes are modelled by scaling down the 2020 baseline potential up to 2050 to match observed historical trends in pollination loss (see Appendix 1, 2). In the absence of climate change effects, no further decline is projected between 2050 and 2070, implying a stabilisation of pollination potential in the long run under the baseline scenario. The assumption is deliberately conservative, intended to avoid overstating long-term losses under substantial uncertainty in ecological trajectories and policy effectiveness. In particular, it reflects the fact that existing policy interventions, such as the EU Pollinators Initiative and earlier reforms under the Common Agricultural Policy (CAP), may not yet have fully translated into observable trends. Extrapolating current declines over extended horizons would otherwise rely on a strong and potentially unjustified assumption of persistence, with a risk of systematic overestimation of long-run impacts. This standard approach in pollination modelling aligns with estimates from the EU [Science and Technology for Pollinating Insects \(STING\)](#) project (Potts et al., 2024 [80]), long-term projections of agro-ecosystem degradation reported by the European Commission [22, 23], and historical trend of the [EU Grassland Butterfly Index](#), a widely used proxy for pollinator-related biodiversity change [106, 26].

Introducing an explicit temporal degradation therefore complements the static baseline and better captures observed declines not otherwise represented, ensuring methodological coherence with the most recent EU-level scientific assessments and reinforcing its empirical plausibility within the modelling framework. In this context, historical trends are interpreted as broadly independent of climate-driven effects which are excluded from the no-climate-change baseline.

Climate change scenarios

Projected declines in pollination potential due to changing climate are derived from an ensemble of state-of-the-art multi-model, multi-scenario species distribution models (SDMs) for key pollinator taxa (wild bees, hoverflies, butterflies, beetles), based on the most recent and robust data currently available. This framework enables the assessment of multiple climate change scenarios (RCP2.6, 4.5, 6.0, 7.0, 8.5) and socioeconomic pathways (SSP1, 2, 3, 5) [89, 107], accounting for future impacts on pollinators up to 2070 (Figure 3). In the CLIM experiments, the input data used only capture climate-induced environmental changes such as temperature and precipitation patterns, while in the other experiments the datasets also embed land-use dynamics under different SSPs. Data sources were selected to ensure full consistency with the KIP-INCA and Status and Trends of European Pollinators (STEP) frameworks.

The selected data sources provide information on the future probability of presence for different species based on projected habitat suitability for up to 153 wild pollinator species across two major taxonomic groups: Hymenoptera (including bumblebees, wild bees, wasps, and ants) and Lepidoptera (including butterflies and moths). The expected probability of presence is then averaged across species within each SDM-scenario considered, under the assumption of perfect substitutability in the provision of pollination services, and that SDM outputs are a reasonable proxy

for overall pollinator presence. These are standard assumptions in ecological modelling of pollination services, reflecting current data constraints. A composite indicator combining species distribution, abundance, and functional trait diversity would provide a more comprehensive measure of pollination capacity[84], yet such data remain unavailable for large-scale, long-term projections. Nonetheless, our approach is grounded in substantial empirical evidence demonstrating that species richness, even as a first-order approximation, captures critical dimensions of pollination service provision. A full description of data sources and data elaboration is provided in Appendix 1.

Spatially referenced information are provided at a level of granularity comparable to that of the INCA datasets, and capture both increases and decreases in pollinator distributions due to habitat suitability as driven by projected regional climate conditions under several SSP-RCP combinations. Consequently, some areas are expected to experience gains in pollination potential, while others will face substantial declines. This spatial heterogeneity reflects the inherently local nature of climate impacts: current evidence suggests that many wild pollinator species may shift towards northern ranges, whereas most bumblebee species are projected to undergo substantial range contractions (Potts et al., 2015).

The projected per-period shocks are then cumulated pixel by pixel to obtain, for each scenario and five-year step τ , the cumulative percentage change in habitat suitability $\Delta\tilde{p}_{n,\tau}$ relative to the 2020 reference value. This cumulative shock is applied multiplicatively to the corresponding baseline value to obtain the climate-adjusted pollination potential:

$$P_{n,\tau} = \left(1 + \frac{\Delta\tilde{p}_{n,\tau}}{100}\right) \cdot P_{b,n,\tau}, \quad P_{n,\tau} \in [0, 1] \quad (1)$$

where $P_{b,n,\tau}$ is the baseline pollination potential in pixel n at time step τ , and $\Delta\tilde{p}_{n,\tau}$ is the cumulative percentage change in habitat suitability relative to 2020 (negative values indicate a decline). $P_{n,\tau}$ is clamped to the unit interval $[0, 1]$ after application of the shock.

By imposing the projected, spatially explicit climate-induced changes in habitat suitability onto the baseline pollination potential, we estimate future pollination potential and its percentage change in each time step at a high spatial resolution of $1 \text{ km} \times 1 \text{ km}$ across all SSP-RCP combinations considered (Figure 3).




Figure 3 | Projected pollination potential change in 2070 with respect to 2020 values under all scenarios considered. The table shows the combination of SSP-consistent land use variables and climate radiative forcing (RCPs) used by the species distribution models (SDMs) in each scenario.

Impacts on agriculture

We assess the agricultural impacts due to yield reduction for specific crop following a well established methodology based on the dependency of specific crops to pollination [29][6][28], and on the expected reduction of pollinator habitat suitability (potential) in the cultivation area [38][109][50]. The overlap between the pollination potential and demand for pollination is used to quantify the area generating the actual flow of service (i.e. the use area) [61]. This can be defined as the number of hectares of pollinator-dependent crops covered by areas with different pollination potential levels. Consequently, changes in crop areas (service demand) or declines in pollination (service supply) translate into variations in agricultural output. As we do not assess climate-change impacts arising from other sources (such as droughts or wildfires), this implies that in our framework climate change affects agricultural production through reductions in pollination potential. Accordingly, the impact of pollination decline is modelled as a decrease in cropland productivity for specific crops, based on each crop's dependency on pollination and its

cultivated area. Agricultural impacts are then represented as the percentage change in crop yield (crop-specific cropland productivity) relative to the baseline, across the various SSP-RCP scenario combinations considered.

To determine use areas at the highest possible spatial resolution, agricultural land (cultivated hectares) for 129 crops and crop categories from [FAO CROPGRIDS](#) [95] is mapped at the grid level and then overlapped with raster layers representing projected percentage changes in pollination under each SSP-RCP scenario combination for the period 2020-2070 relative to the baseline values with no pollination impacts. Only crops whose average pollination dependency exceeds 0.05 are retained in the analysis, as crops below this threshold generate yield shocks too small to be resolved within the macroeconomic model. This enables us to quantify the extent to which the 2020 cultivation areas of specific crops in the EU are affected by pollination decline, prior to evaluating potential SSP-driven land use changes.

The impact of pollination changes is modelled as decreased cropland productivity based on the dependency of specific crops on pollination, which measures the fractional reduction in crop yield associated with a loss of animal pollination [54] [10] [92](See Appendix 1). For example, a pollination dependency of 0.05, means that a 100% decrease in pollination results in a 5% decrease in crop yield. The impact on production for a specific crop in a given area is therefore determined by the projected change in pollination service in the area where the crop is cultivated and by that crop's dependency on animal pollination.

$$\text{yield}_{c,n,t_1} = \text{yield}_{c,n,t_0} \cdot (1 + \text{dependency_p}_c \cdot \Delta p_{n,t_1}) \quad (2)$$

where:

- c the specific crop;
- n the specific spatial unit of production area (grid cell);
- Δp the variation ($t_0 - t_1$) of pollination potential in n ;

For each crop c and NUTS region r , the area-weighted mean cumulative percentage change in pollination potential is computed across all pixels n belonging to that region:

$$\overline{\Delta \tilde{p}}_{c,r,\tau} = \frac{\sum_{n \in r} A_{c,n}^0 \cdot \Delta \tilde{p}_{n,\tau}}{\sum_{n \in r} A_{c,n}^0} \quad (3)$$

where $A_{c,n}^0$ is the baseline cultivated area (hectares) of crop c in pixel n from FAO CROPGRIDS (2020), and $\Delta \tilde{p}_{n,\tau}$ is the cumulative percentage change in pollination potential in pixel n at time step τ (equation 1). The yield shock for crop c in region r at time τ is then expressed as a fractional change in yield (negative values indicate a yield loss):

$$S_{c,r,\tau} = \text{dependency_p}_c \cdot \frac{\overline{\Delta \tilde{p}}_{c,r,\tau}}{100} \quad (4)$$

In this way, the resulting shocks fully embed spatial information on habitat-specific suitability shifts, crop-specific pollination dependency, and the spatial distribution of crop areas.

Assessing macroeconomic impacts on agricultural sub-sectors

The projected gridded percentage changes in yield for each crop and climate scenario are mapped and aggregated at NUTS levels (0 to 3) and then used as land-productivity shocks in the macroeconomic model. Macroeconomic impacts are then estimated using a regionalized version of the ICES Computable General Equilibrium (CGE) model based on GTAP11 database [1], which provides detail for 122 regions down to NUTS-2 level across the EU27, while the rest of the world is represented as a macro-regional aggregate.

Since the agricultural sector in ICES is relatively aggregated, we averaged the projected yield percentage changes based on the relative weight of each crops cultivated in each region, aligning them with the 5 crop categories (agricultural sub-sectors) used in the model. These are: Vegetable, Fruit, Nuts ; Oil seeds ; Cereals, Wheat, Rice ; Sugar cane & Sugar beet ; Other crops, Fibers (See Appendix 1 for full crops detail). This approach is also convenient because the GTAP crop-sectors categories generally correspond to similar levels of pollination dependence; for example, Vegetables and fruits have, on average, high dependence, Oil seeds medium, while Cereals and Rice low to negligible dependence. To do this, we compute a weighted average in which the weight of crops in each grid cell within a NUTS region is proportional to the number of hectares cultivated in that cell. These results are then further aggregated to match the crop categories used in the economic model, taking into account the relative contribution of each crop within the NUTS region of interest.

With the exception of the CLIM scenarios, which are based on a constant 2020 land use exercise, we incorporate projected crop-specific land-use changes from Goebel and Britz (2025)[36], both in the baseline runs and in the impact scenarios to account for future demand changes and crop expansion within each crop-sector under different SSPs [89, 107], at the highest resolution available. These changes reflects how population, GDP and demand dynamics evolve within each SSP scenario, and how these dynamics translate into agricultural sector expansion and land-use patterns across different cropland categories in the EU. Projections are made ensuring full consistency with our modelling framework, as they also rely on a Computable General Equilibrium platform built on the GTAP v11 Power Database [1, 13]. The database represents the global economy in 2017 and includes LULC information, with regional land-use extents adjusted using additional datasets such as the 2018 European CORINE land-cover (LC) data [14]. Cropland and yield developments are calibrated to projections from the Food and Agriculture Organization of the United Nations (FAO, 2018)[27], thereby maintaining full alignment across all components of our modelling chain.

Land-use change information is provided at the NUTS-2 level for eight land-cover types (e.g., cropland, forests, pastureland, shrubland) and for 65 land-use categories corresponding to individual crops and crop aggregates, which are matched to the FAO CROPGRIDS crop categories used in this study. Since information on future cropland expansion is available only at the NUTS-2 level, a grid-level spatial overlap is no longer feasible, as the exact spatial distribution of future crops within each polygon is unknown. The shock is therefore recalibrated proportionally to the expected cropland shares in each crop-sector, NUTS region, scenario, and year, so that the percentage change in crop area is applied uniformly across the entire polygon, assuming that future land allocation within each NUTS-2 region occurs in areas close to the original ones, with no significant deviations in the projected changes in pollination potential for each crop-NUTS-scenario-year combination. The resulting agricultural region and category-specific shocks are then imposed in each run of the recursive-dynamic macroeconomic model as a percentage change of land productivity specific to each agricultural crop-sector and NUTS, reflecting the impact on regional yields in each time step due to pollination loss.

In the SSP-based recalibration of cropland areas, future cultivated areas are therefore computed by applying the percentage changes associated with each SSP. Where $\Delta_{c,r,s}^{SSP}$ the projected percentage change in cultivated area, r NUTS regions, and s SSP scenario–year combinations. Each crop c belongs to one GTAP category g . Baseline cultivated areas $A_{c,r}^0$ are obtained from FAO CROPGRIDS (2020).

$$A_{c,r,s} = A_{c,r}^0 \cdot \left(1 + \frac{\Delta_{c,r,s}^{SSP}}{100} \right) \quad (5)$$

Total cultivated area for each GTAP sector g , region r , and scenario–year s , is defined as:

$$A_{g,r,s} = \sum_{c \in g} A_{c,r,s} \quad (6)$$

In the dynamic crop weighting, crop-specific weights are computed as the share of each crop area on total GTAP crop category area:

$$w_{c,g,r,s} = \frac{A_{c,r,s}}{A_{g,r,s}} \quad (7)$$

By construction, weights vary across scenarios and years and satisfy:

$$\sum_{c \in g} w_{c,g,r,s} = 1 \quad (8)$$

In the weighted aggregation of pollination shocks, let $S_{c,r,s}$ denote the pollination shock at crop level. The GTAP-sector shock is obtained as a weighted average:

$$S_{g,r,s} = \sum_{c \in g} w_{c,g,r,s} \cdot S_{c,r,s} \quad (9)$$

Substituting the weights yields,

$$S_{g,r,s} = \sum_{c \in g} \left(\frac{A_{c,r,s}}{\sum_{c \in g} A_{c,r,s}} \right) \cdot S_{c,r,s} \quad (10)$$

this formulation ensures that changes in cultivated areas across SSP scenarios and years directly affect both the magnitude and the composition of GTAP crop categories shocks.

Acknowledgments

The authors gratefully acknowledge Dr Ioanna Grammatikopoulou, Dr Chiara Polce, Dr Marialuisa Tamborra and the entire INCA team at the JRC for the valuable discussions and for their contribution to the development of the methodological framework and the baseline scenario adopted in this study. We also thank Dr Simon Potts, Dr Gemma Gerber, Dr Simon Dellicour, and Dr Reinhard Prestele for their assistance in the identification and analysis of the pollination-impact scenarios.

This work was supported by the European Union's *Assessing Climate Change Risk in EUrope (ACCREU)* project (Grant Agreement No. 101081358; Funding scheme: HORIZON-CL5-2022-D1-07-two-stage).

Extended results and data analysis

All data, extended material and appendices are available on GitHub:

<https://github.com/gabrielemansi/Macroeconomic-impacts-of-pollination-loss-in-the-EU>

Regional and sectorial heterogeneity analysis of macroeconomic results

Looking at European GDP losses across all 18 scenarios considered, the divergence between extreme values and central estimates remains limited, indicating low dispersion among scenario projections. In both 2050 and 2070, the mean and median values approximate -0.09, while the standard deviation exhibits a modest increase from 0.032 to 0.043 over time. This pattern suggests that at higher levels of spatial aggregation, results tend to become more homogeneous both within and across scenarios, with scenario uncertainty correspondingly becoming less pronounced.

At the EU level, therefore, the European economic system appears relatively resilient: aggregate losses remain limited and suggest a capacity of the macroeconomic system to absorb part of the productivity shocks through sectoral

reallocation and trade adjustments, also considering the limited weight of the value of agricultural production to the EU GDP (approximately 1.6% in the 2017 base year), which generally reduces the macroeconomic visibility of even substantial shocks occurring within agriculture.

Increasing the analytical granularity to NUTS 1-2 regions, the dispersion of outcomes is much wider than the EU-level aggregate suggests. Even under relatively moderate scenarios, regional impacts deviate substantially from the EU mean. Between 2050 and 2070, the distribution of regional GDP impacts shifts downward in its lower tail while becoming more dispersed.

Under low-forcing scenarios (GHSB-1 2.6, IIASA-2 2.6, IIASA-CLIM 2.6, PREST-1 2.6 and PREST-CLIM 2.6), regional GDP impacts range from -0.79% to +0.26% in 2050 and from -1.07% to +0.42% in 2070, with pronounced spatial heterogeneity and negative tails. By 2070 the distribution becomes significantly more polarized: the median is essentially unchanged (from -0.10% to -0.08%), yet regional extremes grow substantially in both directions. Dispersion increases markedly: the 5th percentile worsens from -0.31% to -0.47%, the 10th percentile from -0.23% to -0.31%, and the bottom 1% of regions experiences a deterioration from -0.48% to -0.66%. This means that 10% of EU regions face GDP losses exceeding 0.31% in 2070, while one quarter of regions records losses of at least 0.15% in both years (P25 unchanged at -0.15%). Even 75th percentile turns slightly negative (-0.01% compared to +0.02% in 2050), indicating that more regions are experiencing losses and the small share of regions with positive impacts shrinks. The standard deviation rises from 0.112 to 0.161 (+44%), and skewness deepens from -1.34 to -1.89. The widening gap between the mean (which worsens only modestly) and the lower-tail percentiles (which deteriorate substantially) highlights increasing left-skewness: while a few regions still display modest gains, most suffer progressively worse impacts. Overall, climate damages on regional GDP become more severe and more unequally distributed over time, whereby EU-level aggregates conceal the depth of increasing impacts borne by the most vulnerable NUTS regions.

High-end emission scenarios (GHSB-5 8.5, IIASA-2 7.0, IIASA-CLIM 7.0, IIASA-CLIM 8.5, PREST-3 8.5, PREST-5 8.5, PREST-CLIM 8.5) show an even clearer worsening from 2050 to 2070, with regional dispersions ranging from -0.99% to +0.20% in 2050 and from -1.61% to +0.17% in 2070. The distribution remains strongly left-skewed in both periods, with positive values becoming rarer and smaller over time. The interquartile range widens from 0.10 to 0.15, indicating a broader structural spread of impacts. The negative tail lengthens considerably: the 5th percentile drops from -0.41% to -0.61%, the 1st percentile from -0.64% to -1.00%, while the positive tail compresses (95th percentile from +0.01% to +0.004%). The 25th percentile worsens from -0.15% to -0.19%, meaning that one quarter of EU regions records losses of at least 0.15% in 2050 and 0.19% in 2070. The 75th percentiles remain narrowly negative in both years (-0.05% and -0.04%), but the share of regions with positive impacts contracts (from 6.2% to 5.0%) and the number of regions clustering around zero grows. Mean and median diverge increasingly (median stable at -0.09%, mean worsening from -0.12% to -0.16%), reflecting a small group of regions suffering disproportionately large damages while most see persistent but moderate declines. The standard deviation grows by 58% (from 0.134 to 0.212) and the variance more than doubles (from 0.018 to 0.045), while skewness deepens from -2.28 to -2.88 and kurtosis from 7.8 to 10.8, indicating that the distribution becomes both more dispersed and more heavy-tailed. Crucially, regional GDP losses exceeding 1% are entirely absent in 2050, but appear in 1.1% of regions in 2070. The comparison confirms a clear deterioration of regional economic impacts over time, while mean and median remaining stable (near -0.10%) show that central statistics and averaged data understates the severity of the change, which materializes almost entirely in the left tail.

Increasing the sectorial detail of the analysis allows to detect significant losses even at the EU aggregate level. Economic impacts on the aggregate Agricultural Sector reveal a sectorial contraction between -5.17% to -1.78% in 2050 and -5.95% to -1.54% in 2070, with average losses across all scenarios of -3.22% and -3.47%, respectively. As for the case of GDP, the distribution between scenario is overall modest with output spread generally consistent with scenario family and radiative forcing levels. As with GDP, the variation across scenarios remains relatively modest, with the distribution of impacts broadly consistent with scenario families and radiative forcing levels.

When the spatial granularity is increased to NUTS 1-2 regions, the dispersion of agricultural sector outcomes is substantially wider than for the EU-level aggregate, and considerably larger in absolute terms than that observed for GDP. Even under moderate forcing scenarios, regional impacts deviate dramatically from the EU mean, confirming that agricultural production is more spatially heterogeneous in its climate sensitivity than the broader economy. Under low-forcing scenarios, regional agricultural sector impacts range from -11.39% to +6.62% in 2050 and from -11.83% to +5.31% in 2070, with pronounced spatial heterogeneity and negative tails. In 2050, 83% of EU NUTS regions record negative impacts, with a mean of -2% and a median of -2.13%. By 2070, both indicators worsen (mean -2.2%, median -2.28%), and 70% of regions deteriorate relative to 2050. The standard deviation rises from 1.95 to 2.07 (+6%) and the IQR from 2.26 to 2.42. The negative tail deepens: the 5th percentile drops from -4.68% to -5.58%, the 10th from -4.08% to -4.39%, and the 1st percentile from -7.40% to -8.41%. Regions with losses exceeding 5% of agricultural output grow from 4.4% to 6.2% between the two years, while the share recording positive impacts contracts from 16.6% to 15.2%.

Under intermediate forcing scenarios (GHSB-3 6.0, IIASA-2 4.5, IIASA-CLIM 4.5, PREST-1 4.5, PREST-3 4.5 and PREST-CLIM 4.5), extreme values exhibit modest differences with low-impact, with regional distributions ranging from -10.93% to +5.63% in 2050 and -12.67% to +6.77% in 2070. The mean reaches -2.90% and the median -2.76%, while the 25th percentile stands at -4.37% in 2050, worsening to -4.68% by 2070. The standard deviation grows from 2.61 to 3.00, and the IQR expands from 3.10 to 3.80, indicating a broad structural divergence of regional trajectories. Critically, the 1st percentile deteriorates from -9.55% to -10.41%, and the share of NUTS with losses exceeding 5% rises from 18.7% to 21.7%, while regions with losses exceeding 10% more than double (from 0.7% to 1.6%), a damage class nearly absent under low-forcing conditions.

High-end emission scenarios (GHSB-5 8.5, IIASA-2 7.0, IIASA-CLIM 7.0, IIASA-CLIM 8.5, PREST-3 8.5, PREST-5 8.5 and PREST-CLIM 8.5) show the greatest structural divergence between the EU aggregate and the regional distribution. In 2050, the pooled range runs from -11.26% to +6.13%, with a mean of -2.63%, a median of -2.46%, and 85% of NUTS regions in negative territory. By 2070, the range widens from -12.78% to +7.96%, but the temporal evolution is qualitatively different from that observed under low-forcing and intermediate scenarios: the mean and median both improve slightly (+0.28 pp and +0.30 pp respectively), driven by a marked positive reorientation in northern and alpine NUTS under higher-warming conditions, while simultaneously the standard deviation grows from 2.64 to 3.13 and the IQR expands from 3.14 to 4.43. Strikingly, the 75th percentile, which stood at -0.96% in 2050, turns positive in 2070, indicating that more than one quarter of all NUTS regions now records agricultural gains under these scenarios, a structural shift that the negative mean still hides. The share of NUTS with positive impacts almost doubles between the two horizons (from 14.7% to 25.9%). At the same time, the lower tail deepens markedly: the 5th percentile drops from -7.39% to -8.11%, the 1st from -9.80% to -11.23%, and regions with losses exceeding 10% nearly triple (from 0.7% to 1.9%), a damage class entirely absent under low-forcing conditions. The skewness deepens from -0.33 to -0.46, confirming the growing asymmetry of the distribution. This dual dynamic – an improving central tendency alongside a strongly widening spread – is a clear statistical signature of deepening polarisation: a growing subset of northern and alpine regions actually benefits, while the most exposed southern NUTS deteriorate further.

The comparison across all three scenario groups highlights a systematic pattern. First, the mean and median values across groups are more similar than the scenario labels might suggest (ranging from -1.99% to -2.90% at the 2050 mean), whereas the dispersion indicators diverge dramatically: the IQR moves from 2.26 under low-forcing to 3.10 under intermediate to 3.25 under high-end forcing, widening further in all groups by 2070. Second, the share of NUTS with losses exceeding 5% – only 4.4% under low-forcing – reaches roughly 19% under both intermediate and high-end scenarios in 2050, demonstrating a threshold-like response in agricultural sensitivity to forcing level. Third, losses exceeding 10% emerge as a statistically relevant outcome class only under intermediate and high-end forcing (around 0.7-0.8% of NUTS in 2050, growing to 1.6-2.2% by 2070), and are entirely absent under low-forcing. Across all scenario groups, the EU-level aggregate consistently understates the magnitude of the worst regional impacts by a factor of 2.0× to 3.6×, with gaps growing in absolute terms as forcing increases. These results

confirm that EU-level data are structurally insufficient to assess agricultural losses from pollination decline, and that NUTS-level granularity is essential to capture both the severity and the heterogeneity of the impacts.

Finally, we examine the most disaggregated agri-sectors level. By analysing the highly impacted Vegetable, Fruit, and Nuts sector, the same pattern holds. At the EU aggregate, scenario differences remain modest. The 2050 range spans -5.86% to -1.89%, and it widens to -7.0% to -1.91% by 2070. The mean declines from -3.58% to -4.08%, and the median from -2.94% to -3.47%. The standard deviation rises from 1.37 to 1.58, indicating growing dispersion over time, while skewness is consistently moderately negative (-0.53), reflecting a slight left-skew. The 75th percentile worsens from -2.40% to -2.73%, confirming a clear negative trend across all scenarios toward 2070. The data divergence drastically increase when looking at NUTS level. Under low-forcing scenarios, regional impacts range from -15.33% to +8.37% in 2050 and from -17.22% to +6.97% in 2070, with very strong spatial heterogeneity and heavy negative tails. In 2050, 76% of EU NUTS regions record negative impacts, with a mean of -2.14% and a median of -1.80%. By 2070, the mean worsens to -2.38% while the median is essentially unchanged (-1.80%), a first signal that the bulk of the distribution shifts only marginally while the tails grow substantially. The standard deviation rises from 3.00 to 3.31 (+10%), and the IQR widens from 3.67 to 4.14. The negative tail deepens markedly: the 5th percentile drops from -7.81% to -8.44%, the 10th from -5.78% to -6.39%, and the 1st percentile from -10.97% to -12.28%. Regions with losses exceeding 5% grow from 14.6% to 19.0% between the two horizons, while regions with losses exceeding 10% almost triple (from 1.3% to 3.3%), a damage class that is essentially unknown for the broader Agricultural sector under the same scenarios. The skewness deepens from -0.77 to -0.96, confirming the growing asymmetry.

Under intermediate forcing scenarios, the regional distribution shifts substantially further, with impacts ranging from -18.04% to +6.12% in 2050 and from -18.56% to +7.78% in 2070. The mean reaches -3.12% in 2050 and the median -2.36%, while 80% of NUTS are in negative territory. The 25th percentile already stands at -5.04%, meaning that a quarter of all EU NUTS regions record losses of at least 5% in sectorial output. By 2070, although the mean marginally improves to -3.00% and the median to -2.09%, the standard deviation grows by 13% (from 3.89 to 4.40) and the IQR expands from 4.70 to 5.64. The 1st percentile deteriorates from -14.17% to -15.23%, and regions with losses exceeding 10% rise from 7.4% to 8.1%. Regions with losses exceeding -5% rise to 29.0% in 2070.

High-end emission scenarios display the most extreme structural divergence between the EU aggregate and the regional distribution. In 2050, the pooled range runs from -18.85% to +6.69%, with a mean of -2.88% and a median of -2.17%, while 77% of NUTS experience losses. By 2070 the distribution widens dramatically, ranging between -21.35% to +8.85%. The standard deviation grows by 19% (from 3.89 to 4.61), and the IQR expands from 4.77 to 5.92. As observed for the broader agricultural sector, the temporal evolution exhibits a marked dual dynamic: the mean improves slightly (+0.28 pp) and the median improves more visibly (+0.66 pp), driven by a substantial positive reorientation in northern and alpine NUTS, while simultaneously the lower tail deteriorates further. The share of NUTS with positive impacts grows from 22.6% to 34.7%, and the 75th percentile turns positive in 2070 (+0.68%) from negative in 2050 (-0.17%). At the same time, the 5th percentile drops from -10.79% to -11.79%, the 1st from -14.13% to -15.90%, and the absolute minimum reaches -21.35%. Regions with losses exceeding 10% remain stable at around 7–8% but their absolute severity intensifies; the skewness deepens from -0.93 to -1.00, kurtosis from 1.1 to 1.2.

The comparison across the three scenario groups shows that the amplification factor (NUTS over EU average) remains in the 3.4× to 5.6× range across all scenario groups, substantially higher than for Agricultural sector (2.0× to 3.2×), while the absolute gap between the EU value and the most affected NUTS reaches values of over 14 percentage points, more than double those observed for the aggregate agricultural sector. The structural pattern of polarisation observed for Agricultural sector is amplified here: under high-end scenarios, more than one third of NUTS shows positive impacts in 2070, while simultaneously the worst-affected NUTS deteriorates further, demonstrating that Vegetable, Fruit, and Nuts sector is structurally heterogeneous in its response to pollination loss under climate change and that EU-level data are unable to capture this dimension. Regions with losses exceeding 10% affect 1.3%

of NUTS already under low-forcing, a value absent in Agricultural sector under same scenarios, rising to 7–8% under intermediate and high-end forcing, confirming that specific agricultural sub-sectors may be systematically most exposed to severe regional climate damages compared to the regional and sectorial aggregates.

Table 1: Summary statistics | EU aggregate vs NUTS distribution at 2050 and 2070

Var	Group	Year	EU range		NUTS distribution								Losses (%)	
			min	max	min	max	mean	median	std	IQR	P5	P95	< -5	< -10
GDP	LF	2050	-0.10	-0.08	-0.79	+0.26	-0.11	-0.10	0.11	0.09	-0.31	+0.05	0	0
GDP	LF	2070	-0.09	-0.06	-1.07	+0.42	-0.12	-0.08	0.16	0.12	-0.47	+0.02	0	0
GDP	MOD	2050	-0.14	-0.02	-0.99	+0.20	-0.13	-0.09	0.15	0.12	-0.41	+0.04	0	0
GDP	MOD	2070	-0.16	-0.00	-1.68	+0.22	-0.16	-0.09	0.23	0.16	-0.58	+0.03	0	0
GDP	HE	2050	-0.15	-0.04	-0.99	+0.20	-0.12	-0.09	0.13	0.10	-0.41	+0.01	0	0
GDP	HE	2070	-0.18	-0.04	-1.61	+0.17	-0.16	-0.09	0.21	0.15	-0.61	+0.005	0	0
agri	LF	2050	-3.16	-2.29	-11.39	+6.62	-1.99	-2.13	1.95	2.26	-4.68	+0.95	4.4	0.2
agri	LF	2070	-3.19	-2.51	-11.83	+5.31	-2.17	-2.28	2.07	2.42	-5.58	+1.04	6.2	0.2
agri	MOD	2050	-5.07	-2.05	-10.93	+5.63	-2.90	-2.76	2.61	3.10	-7.86	+1.15	18.7	0.7
agri	MOD	2070	-5.61	-2.03	-12.67	+6.77	-2.86	-2.84	3.00	3.80	-8.12	+1.65	21.7	1.6
agri	HE	2050	-5.17	-1.78	-11.26	+6.13	-2.63	-2.46	2.64	3.14	-7.39	+1.33	16.2	0.7
agri	HE	2070	-5.96	-1.54	-12.78	+7.96	-2.35	-2.16	3.13	4.43	-8.11	+1.77	17.7	1.9
v_f	LF	2050	-4.33	-2.36	-15.33	+8.37	-2.14	-1.80	3.00	3.67	-7.81	+1.69	14.6	1.3
v_f	LF	2070	-4.84	-2.59	-17.22	+6.97	-2.38	-1.80	3.31	4.14	-8.44	+1.65	19.0	3.3
v_f	MOD	2050	-5.47	-2.24	-18.04	+6.12	-3.12	-2.36	3.89	4.70	-11.08	+1.91	25.5	7.4
v_f	MOD	2070	-6.25	-2.47	-18.56	+7.78	-3.00	-2.09	4.40	5.64	-11.97	+2.58	29.0	8.1
v_f	HE	2050	-5.86	-1.89	-18.85	+6.69	-2.88	-2.17	3.89	4.77	-10.79	+2.22	24.6	6.7
v_f	HE	2070	-6.99	-1.91	-21.35	+8.85	-2.60	-1.51	4.61	5.92	-11.79	+2.94	26.3	7.6

Notes | Analysis of the percentage deviations from the respective SSP baselines across the 18 scenarios considered for projected model variables: *GDP* = regional economic output; *agri* = value of aggregated Agricultural sector output; *v_f* = value of agricultural sub-sector Vegetables, Fruits and Nuts. Scenario groups are: *LF* = Low Forcing (GHSB-1 2.6, IIASA-2 2.6, IIASA-CLIM 2.6, PREST-1 2.6, PREST-CLIM 2.6); *MOD* = Moderate Forcing (GHSB-3 6.0, IIASA-2 4.5, IIASA-CLIM 4.5, PREST-1 4.5, PREST-3 4.5, PREST-CLIM 4.5); *HE* = High Forcing (GHSB-5 8.5, IIASA-2 7.0, IIASA-CLIM 7.0, IIASA-CLIM 8.5, PREST-3 8.5, PREST-5 8.5, PREST-CLIM 8.5). *EU range* reports the minimum and maximum aggregate EU impacts across all scenarios. *NUTS distribution* statistics are computed across 122 NUTS regions: *std* = standard deviation; *IQR* = interquartile range; *P5* and *P95* denote the 5th and 95th percentiles of the regional NUTS distribution. Columns < -5 and < -10 report the share of regions experiencing losses greater than -5% and -10%, respectively.

Sensitivity of pollination Scenarios to climate change pathways

Sensitivity analysis shows that SDMs models present heterogeneous sensitivity to emission scenarios and socioeconomic pathways (Figure 4).

PREST-CLIM and PREST-1 are highly sensitive to the RCP pathway (Figure 4c), with mean regional differences of nearly 34-40 percentage points between RCP2.6 and RCP8.5 at 2070, and with substantial spatial heterogeneity in this sensitivity (Interquartile Range IQR 30-40 pp across NUTS-2 regions). In contrast, IIASA-2 presents a low sensitivity to RCP choice, with a mean range of less than 6 pp between RCP2.6 and RCP7.0 even at 2070, as well as IIASA-CLIM (6 pp). PREST-3 occupies an intermediate position (11 pp at 2070). This indicates that for some models the emissions pathway is decisive, while for others structural assumptions dominate. Importantly, part of these differences can be explained by the different representation of species within the SDM frameworks: models vary in how they parametrise species distributions, ecological tolerances, and pollinator-crop interactions, which amplifies or dampens the sensitivity to climate forcing (see Appendix 1).

When comparing models under the same RCP, the picture shifts. At RCP2.6, models converge relatively closely (range 3-7 pp), but at RCP4.5 and RCP8.5 divergences become dramatic, reaching up to 36 percentage points at 2070 under RCP8.5. This means that model choice becomes critical in pessimistic scenarios, precisely where uncertainty is highest and policy relevance greatest.

The SSP dimension adds a further layer of uncertainty (Figure 4d). Under SSP1, inter-model spread remains

moderate (mean 11 pp at 2070), reflecting that low-forcing scenarios leave little room for model divergence. Under SSP2, SSP3 and SSP5 the spread grows substantially, reaching mean values of 31, 38, and 39 pp respectively at 2070 — with IQR extending beyond 45 pp in some regions. Notably, SSP3 and SSP5 are statistically indistinguishable in terms of inter-model spread, suggesting that beyond a warming threshold the land-use variables of the socioeconomic pathway matters less than the underlying RCP and SDM structure. The eight scenarios included under SSP2 produce the widest range of outcomes in absolute terms, reflecting the diversity of model families that project into this pathway.

The full distribution of projected impacts across all NUTS-2 regions and scenarios (Figure 4e) reveals a consistently negative but highly skewed picture. In 2030, the bulk of projections cluster between -10% and -25%, with limited positive outliers confined to a handful of northern and eastern European regions under low-emission scenarios. By 2070, the distribution broadens dramatically: the median impact across scenarios sits near -30%, but the lower tail extends below -70% for PREST variants under high emissions. The overlap between the 2050 and 2070 distributions is narrow, confirming that impacts continue to accelerate rather than plateau. Critically, no scenario produces a positive EU-wide signal by 2070, reinforcing the robustness of the directional finding even in the face of large inter-model spread.

These results suggest that while the direction of change is homogeneous (negative trend), the magnitude of decline is highly sensitive both to the emissions pathway and to the modelling framework. Yet, the signal-to-noise ratio varies: in some cases the decline is strong relative to model uncertainty, in others the spread between models is wide enough to weaken confidence. The convergence of GHSB and IIASA families around -25 to -35% by 2070, regardless of RCP, may represent a more conservative but robust lower bound, while PREST-family projections under high emissions should be interpreted as plausible upper-bound scenarios conditional on specific land-use change assumptions.



Figure 4 | Projected changes in pollination supply across EU scenarios relative to 2020. **(a)** EU-level average percentage change in pollination for 18 land-use and climate scenarios across three time horizons (2030, 2050, 2070). Colour intensity reflects the magnitude of decline, ranging from moderate losses (~10%) under low-emission scenarios to severe losses (up to -66%) under high-emission PREST variants; **(b)** Spatial variability in projected pollination change across NUTS regions for each scenario in 2070, shown as the full MIN-MAX range (shaded bar) of the regional distribution; the red square marks the EU-level average. Scenarios with low RCP (2.6) display wider ranges extending into positive values in some regions, while high-emission scenarios converge toward uniformly negative outcomes; **(c)** Mean RCP sensitivity per model family in 2070, expressed as the average range (in percentage points) between the lowest and highest RCP run within the same model. PREST-CLIM shows the highest sensitivity (~40 pp), followed by PREST-1 and 3 (~32 pp), while IIASA-2 and IIASA-CLIM exhibit near-zero RCP sensitivity (<10 pp); GHSB and PREST-5 are excluded (only one RCP per variant); **(d)** Mean SSP sensitivity in 2070 as range of EU average pp across scenarios under the same SSP (model spread at fixed SSP). Under SSP2, eight scenarios from four models are grouped. SSP1 comprises PREST-1 and GHSB-1 scenarios. SSP3 comprises GHSB-3 and PREST-3. SSP5 comprises GHSB-5, IIASA-CLIM 8.5, PREST-5, and PREST-CLIM 8.5; **(e)** Impact distribution across all NUTS (0-2) and scenarios for year 2030, 2050 and 2070 (2700 annual values). The violin shape represents the kernel density estimate (KDE) (probability density function of all values). The vertical black bar indicates the interquartile range (Q1-Q3). The white dot marks the median, and the grey dot marks the mean. **(f)** EU-level average trajectory for each scenario from 2030 to 2070, grouped by model family (colour) and RCP (line style: solid = low RCP, dashed = high RCP). All scenarios project monotonically increasing losses over time, with PREST family scenarios under high emissions reaching -65% to -70% by 2070, and IIASA and GHSB families stabilising between -25% and -35%.

References

References

- [1] Aguiar, A., Chepeliev, M., Corong, E., McDougall, R., & van der Mensbrugghe, D. (2022). The GTAP 11 Data Base. *Journal of Global Economic Analysis*, 7(2), 1–44. <https://doi.org/10.21642/JGEA.070201AF>
- [2] Aizen, M. A., Garibaldi, L. A., Cunningham, S. A., & Klein, A. M. (2009). How much does agriculture depend on pollinators? Lessons from long-term trends in crop production. *Annals of Botany*, 103(9), 1579–1588. <https://doi.org/10.1093/aob/mcp076>
- [3] Aviron, S., Poggi, S., Varennes, Y. D., & Lefèvre, A. (2023). Local landscape heterogeneity affects crop colonization by natural enemies of pests in protected horticultural cropping systems. *Agriculture, Ecosystems & Environment*, 343, 108275. <https://doi.org/10.1016/j.agee.2023.108410>
- [4] Azuero Pedraza, C., Mansi, G. A., and Drouet, L., (2026). The representation of biodiversity in integrated assessment models: State of the art and directions for improvement. *Environmental Research Letters*. <http://iopscience.iop.org/article/10.1088/1748-9326/ae7394>
- [5] Bastien-Olvera, B.A., Conte, M.N., Dong, X. et al. Unequal climate impacts on global values of natural capital. *Nature* 625, 722–727 (2024). <https://doi.org/10.1038/s41586-023-06769-z>
- [6] Bauer, D. M., & Wing, I. S. (2016). The macroeconomic cost of catastrophic pollinator declines. *Ecological Economics*, 126, 1–13. ISSN 0921-8009, <https://doi.org/10.1016/j.ecolecon.2016.01.011>. (<https://www.sciencedirect.com/science/article/pii/S0921800916301331>) <https://doi.org/10.1016/j.ecolecon.2016.01.011>
- [7] Beckman, J., Ivanic, M., Jelliffe, J. L., Baquedano, F. G., & Scott, S. G. (2020). Economic and Food Security Impacts of Agricultural Input Reduction Under the European Union Green Deal’s Farm to Fork and Biodiversity Strategies. *USDA Economic Research Service*. <https://doi.org/10.22004/AG.ECON.327231>
- [8] Biesmeijer JC, Roberts SP, Reemer M, Ohlemüller R, Edwards M, Peeters T, Schaffers AP, Potts SG, Kleukers R, Thomas CD, Settele J, Kunin WE. Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science*. 2006 Jul 21;313(5785):351-4. doi: 10.1126/science.1127863. PMID: 16857940 DOI: 10.1126/science.1127863 <https://www.science.org/doi/10.1126/science.1127863>
- [9] Billeter, R., Liira, J., Bailey, D., Bugter, R., Arens, P., Augenstein, I., ... & Edwards, P. J. (2008). Indicators for biodiversity in agricultural landscapes: a pan-European study. *Journal of Applied Ecology*, 45(1), 141–150. <https://doi.org/10.1111/j.1365-2664.2007.01393.x>
- [10] Bugin, G., Lenzi, L., Ranzani, G., Barisan, L., Porrini, C., Zanella, A., & Bolzonella, C. (2022). Agriculture and pollinating insects, no longer a choice but a need: EU agriculture’s dependence on pollinators in the 2007–2019 period. *Sustainability*, 14(6), 3644. <https://doi.org/10.3390/su14063644>
- [11] Campagnolo, L., Campoleoni Mansi, G. A., Bosello, F., Raitzer, D., (2025) Asian Development Bank, 6 ADB Avenue, Mandaluyong City, 1550 Metro Manila, Philippines. Some rights reserved. Published in 2025. ISSN 2313-6537 (print), 2313-6545 (PDF) Publication Stock No. WPS250073-2 DOI: <http://dx.doi.org/10.22617/WPS250073-2>
- [12] Carvalheiro, L. G., Kunin, W. E., Keil, P., et al. (2013). Species richness declines and biotic homogenisation have slowed down for NW-European pollinators and plants. *Ecology Letters*, 16(7), 870–878. 21 May 2013 <https://doi.org/10.1111/ele.12121> <https://onlinelibrary.wiley.com/doi/10.1111/ele.12121>

- [13] Chepeliev, M. (2023). GTAP-Power Data Base: Version 11. *Journal of Global Economic Analysis*, 8(2). <https://doi.org/10.21642/JGEA.080203AF> <https://jgea.org/ojs/index.php/jgea/article/view/222>
- [14] CORINE Land Cover 2018. European Environment Agency. DOI (raster 100 m): <https://doi.org/10.2909/960998c1-1870-4e82-8051-6485205ebbac> DOI (vector): <https://doi.org/10.2909/71c95a07-e296-44fc-b22b-415f42acfd0> <https://land.copernicus.eu/pan-european/corine-land-cover>
- [15] Dasgupta, P. (2024). *The Economics of Biodiversity: The Dasgupta Review*. HM Treasury. ISBN 978-1-911680-29-1 PU 3069 CCS1120604514 02/21 <https://www.gov.uk/government/publications/final-report-the-economics-of-biodiversity-the-dasgupta-review>
- [16] Devictor, V., van Swaay, C., Brereton, T., et al. (2012). Differences in the climatic debts of birds and butterflies at a continental scale. *Nature Climate Change*, 2(2), 121–124. <https://doi.org/10.1038/nclimate1347> <https://www.nature.com/articles/nclimate1347#Bib1>
- [17] Dicks LV, Baude M, Roberts SP, Phillips J, Green M, Carvell C. How much flower-rich habitat is enough for wild pollinators? Answering a key policy question with incomplete knowledge. *Ecol Entomol*. 2015 Sep;40(Insects and Ecosystem Services 28th Symposium of the Royal Entomological Society of LondonS1):22-35. doi: 10.1111/een.12226. Epub 2015 Jul 2. PMID: 26877581; PMCID: PMC4737402. <https://resjournals.onlinelibrary.wiley.com/doi/10.1111/een.12226>
- [18] Dicks LV, Breeze TD, Ngo HT, Senapathi D, An J, Aizen MA, Basu P, Buchori D, Galetto L, Garibaldi LA, Gemmill-Herren B, Howlett BG, Imperatriz-Fonseca VL, Johnson SD, Kovács-Hostyánszki A, Kwon YJ, Lattorff HMG, Lungharwo T, Seymour CL, Vanbergen AJ, Potts SG. A global-scale expert assessment of drivers and risks associated with pollinator decline. *Nat Ecol Evol*. 2021 Oct;5(10):1453-1461. doi: 10.1038/s41559-021-01534-9. Epub 2021 Aug 16. <https://pubmed.ncbi.nlm.nih.gov/34400826/>
- [19] Donald, P. F., & Evans, A. D. (2006). Habitat connectivity and matrix restoration: the wider implications of agri-environment schemes. *Journal of Applied Ecology*, 43(2), 209–218. <https://doi.org/10.1111/j.1365-2664.2006.01146.x>
- [20] European Environment Agency. (2020). *State of Nature in the EU: Results from Reporting under the Nature Directives 2013–2018*. EEA Report No 10/2020. ISBN 978-92-9480-260-6 ISSN 1725-9177 doi: 10.2800/705440 Environmental production. <https://www.eea.europa.eu/en/analysis/publications/state-of-nature-in-the-eu-2020>
- [21] European Environment Agency. (2025). *Protecting and Restoring Europe's Wild Pollinators and Their Habitats*. HTML: TH-01-25-015-EN-Q. ISBN: 978-92-9480-720-5. <https://doi.org/10.2800/6525185>
- [22] European Commission: Directorate-General for Environment, IEEP, IUCN, Trinomics & UNEP-WCMC. (2023a). *Impact Assessment Study to Support the Development of Legally Binding EU Nature Restoration Targets: Final Report*. Publications Office of the European Union. <https://doi.org/10.2779/275295>
- [23] European Commission: Directorate-General for Environment, Trinomics, IEEP, UNEP-WCMC and IUCN. (2023). *Impact Assessment Study to Support the Development of Legally Binding EU Nature Restoration Targets – Final Report. Annex 3, Ecosystem Specific Assessments*. Publications Office of the European Union. <https://doi.org/10.2779/673007>

- [24] European Commission. (2021). *Report from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions. Progress in the Implementation of the EU Pollinators Initiative*. Brussels, 27.5.2021 COM(2021) 261 final.
- [25] EU STEP (Status and Trends of European Pollinators). (2015). Final Report Summary. European Commission FP7 Project.
- [26] Eurostat. (2025). Grassland butterfly index – EU aggregate. Online data code: sdg_15_61. Last update: 26/05/2025 23:00. Source: European Environment Agency (EEA); Butterfly Conservation Europe (BCE). https://doi.org/10.2908/sdg_15_61
- [27] Food and Agriculture Organization of the United Nations. (2018). *The Future of Food and Agriculture: Alternative Pathways to 2050*. FAO, Rome. <https://www.fao.org/global-perspectives-studies/resources/detail/en/c/1157074/>
- [28] Feuerbacher, A., Kempen, M., Steidle, J.L.M. et al. The economic, agricultural, and food security repercussions of a wild pollinator collapse in Europe. *Nat Commun* 16, 9892 (2025). <https://doi.org/10.1038/s41467-025-65414-7> <https://www.nature.com/articles/s41467-025-65414-7>
- [29] Gallai, N., Salles, J. M., Settele, J., & Vaissière, B. E. (2009). Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecological Economics*, 68(3), 810–821. <https://doi.org/10.1016/j.ecolecon.2008.06.014>
- [30] Garibaldi, L. A., Aizen, M. A., Klein, A. M., Cunningham, S. A., & Harder, L. D. (2011). Global growth and stability of agricultural yield decrease with pollinator dependence. *Proceedings of the National Academy of Sciences*, 108(14), 5909–5914. <https://doi.org/10.1073/pnas.1012431108>
- [31] Garibaldi LA, Steffan-Dewenter I, Winfree R, Aizen MA, Bommarco R, Cunningham SA, Kremen C, Carvalheiro LG, Harder LD, Afik O, Bartomeus I, Benjamin F, Boreux V, Cariveau D, Chacoff NP, Dudenhöffer JH, Freitas BM, Ghazoul J, Greenleaf S, Hipólito J, Holzschuh A, Howlett B, Isaacs R, Javorek SK, Kennedy CM, Krewenka KM, Krishnan S, Mandelik Y, Mayfield MM, Motzke I, Munyuli T, Nault BA, Otieno M, Petersen J, Pisanty G, Potts SG, Rader R, Ricketts TH, Rundlöf M, Seymour CL, Schüepp C, Szentgyörgyi H, Taki H, Tscharrntke T, Vergara CH, Viana BF, Wanger TC, Westphal C, Williams N, Klein AM. *Wild pollinators enhance fruit set of crops regardless of honey bee abundance*. *Science*. 2013 Mar 29;339(6127):1608–11. doi: 10.1126/science.1230200. Epub 2013 Feb 28. Erratum in: *Science*. 2014 May 23;344(6186):816. PMID: 23449997. <https://pubmed.ncbi.nlm.nih.gov/23449997/>
- [32] Garibaldi, L. A., Carvalheiro, L. G., Leonhardt, S. D., et al. (2016). From research to action: enhancing crop yield through wild pollinators. *Frontiers in Ecology and the Environment*, 14(4), 223–223. <https://esajournals.onlinelibrary.wiley.com/doi/10.1890/130330>
- [33] M.P.D. Garratt, T.D. Breeze, N. Jenner, C. Polce, J.C. Biesmeijer, S.G. Potts, Avoiding a bad apple: Insect pollination enhances fruit quality and economic value, *Agriculture, Ecosystems & Environment*, Volume 184, 2014, Pages 34–40, ISSN 0167-8809, <https://doi.org/10.1016/j.agee.2013.10.032>. <https://www.sciencedirect.com/science/article/pii/S0167880913003848>
- [34] Gerber, G., Palazzo, A., et al. (2025). *ACCREU (EU), Deliverable D2.4 (Impacts on Ecosystems & Biodiversity)*. GA number: 101081358. https://www.accreu.eu/wp-content/uploads/2025/09/ACCREU_Deliverable-D2.4.pdf

- [35] Ghisbain, G., Thiery, W., Massonnet, F. et al. Projected decline in European bumblebee populations in the twenty-first century. *Nature* 628, 337–341 (2024). <https://doi.org/10.1038/s41586-023-06471-0> <https://doi.org/10.1038/s41586-023-06471-0>
- [36] Goebel T., Britz, W., Global land use and land cover projections under the shared socio-economic pathways: An integrated computable general equilibrium analysis with sub-national resolution for Europe, *Global Environmental Change Advances*, Volume 5, 2025, 100027, ISSN 2950-1385, <https://doi.org/10.1016/j.gecadv.2025.100027> <https://www.sciencedirect.com/science/article/pii/S2950138525000154>
- [37] Goulson, D., Nicholls, E., Botías, C., & Rotheray, E. L. (2015). Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science*, 347(6229), 1255957. DOI:10.1126/science.1255957 <https://www.science.org/doi/10.1126/science.1255957>
- [38] Grammatikopoulou, I., et al. (2024). *The Contribution of Ecosystem Services in Agricultural Production – An Application of the Production Function Approach*. Publications Office of the European Union. ISSN 1831-9424, ISBN 978-92-68-20818-2, DOI 10.2760/1432445, Catalogue number KJ-01-24-056-EN-N. <https://doi.org/10.2760/1432445> <https://data.europa.eu/doi/10.2760/1432445>
- [39] GTAP. (2023). GTAP 11 Data Base Documentation. Center for Global Trade Analysis, Purdue University.
- [40] Harrison, P. A., Dunford, R. W., Holman, I. P., Cojocaru, G., Madsen, M. S., Chen, P. Y., ... & Sandars, D. (2019). Differences between low-end and high-end climate change impacts in Europe across multiple sectors. *Regional Environmental Change*, 19(3), 695–709. <https://doi.org/10.1007/s10113-018-1352-4>
- [41] Hart, K., Baldock, D., Tucker, G., Allen, B., Calatrava, J., Black, H., ... & Gantioler, S. (2011). *Costing the Environmental Needs Related to Rural Land Management*. Report Prepared for DG Environment. Contract No ENV. F, 1.
- [42] Hatt, S., Lopes, T., Boeraeve, F., Chen, J., & Francis, F. (2017). Pest regulation and support of natural enemies in agriculture: experimental evidence of within field wildflower strips. *Ecological Engineering*, 98, 240–245. <https://www.sciencedirect.com/science/article/pii/S0925857416306176>
- [43] Hodgson, J. A., Thomas, C. D., Cinderby, S., Cambridge, H., Evans, P., & Hill, J. K. (2011). Habitat re-creation strategies for promoting adaptation of species to climate change. *Conservation Letters*, 4(4), 289–297. <https://doi.org/10.1111/j.1755-263X.2011.00177.x>
- [44] Hoegh-Guldberg O., et al. ,The human imperative of stabilizing global climate change at 1.5°C. *Science* 365,eaaw6974(2019).DOI:10.1126/science.aaw6974 <https://www.science.org/doi/10.1126/science.aaw6974>
- [45] IPBES (2016). *The assessment report of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services on pollinators, pollination and food production*. S.G. Potts, V. L. Imperatriz-Fonseca, and H. T. Ngo (eds). Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services, Bonn, Germany. 552 pages. <https://zenodo.org/records/3402857>
- [46] IPBES. (2019). *Global Assessment Report on Biodiversity and Ecosystem Services*. IPBES Secretariat, Bonn, Germany. <https://doi.org/10.5281/zenodo.3831673>
- [47] IPCC. (2020). *Special Report on Global Warming of 1.5°C (SR15)*. Cambridge University Press. <https://www.ipcc.ch/sr15/chapter/chapter-3/>

- [48] IUCN. (2026). European Commission: Directorate-General for Environment, Michez, D., Boustani, M., Sentil, A., Benrezkallah, J. et al., *European Red List of Bees – Measuring the Pulse of European Biodiversity*. Publications Office of the European Union. <https://doi.org/10.2779/521877>
- [49] Jacobs, C., Berglund, M., Kurnik, B., Dworak, T., Marras, S., Mereu, V., & Michetti, M. (2019). *Climate Change Adaptation in the Agriculture Sector in Europe*. European Environment Agency. <https://www.eea.europa.eu/en/analysis/publications/cc-adaptation-agriculture>
- [50] Johnson J.A., Baldos U.L., Corong,E., Hertel T., Polasky S., Cervigni R.,Roxburgh T., Ruta G., Salemi C., Thakrar S. Investing in nature can improve equity and economic returns, *PNAS. Proc. Natl. Acad. Sci. U.S.A.* 120 (27) e2220401120, (2023). <https://doi.org/10.1073/pnas.2220401120>
- [51] Jung, M. (2023a). An integrated species distribution modelling framework for heterogeneous biodiversity data. *Ecological Informatics*, 76, 102127. <https://doi.org/10.1016/j.ecoinf.2023.102127>
- [52] Jung, M. (2023b). *ibis.insights: An R Implementation of the InSiGHTS Framework*.
- [53] Kazenel, M.R., Wright, K.W., Griswold, T. et al. Heat and desiccation tolerances predict bee abundance under climate change. *Nature* 628, 342–348 (2024). <https://doi.org/10.1038/s41586-024-07241-2> <https://www.nature.com/articles/s41586-024-07241-2#citeas>
- [54] Klein, A. M., Vaissière, B. E., Cane, J. H., Steffan-Dewenter, I., Cunningham, S. A., Kremen, C., & Tscharntke, T. (2007). Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B: Biological Sciences*, 274(1608), 303–313. <https://doi.org/10.1098/rspb.2006.3721>
- [55] Kleijn, D., Winfree, R., Bartomeus, I., et al. (2015). Delivery of crop pollination services is an insufficient argument for wild pollinator conservation. *Nature Communications*, 6, 7414. <https://www.nature.com/articles/ncomms8414#Sec2>
- [56] Kudrnovsky, H., Ellmauer, T., Gotzl, M., Paternoster, D., Sonderegger, G., Schwaiger, E., EAA (AT) (2020). ETC/BD Technical paper 1/2020: Report for a list of Annex I habitat types important for Pollinators *European Environment Agency Technical Report*. <https://www.eionet.europa.eu/etcs/etc-bd/products/etc-bd-reports/etc-bd-technical-paper-1-2020-report-for-a-list-of-annex-i-habitat-types-important-for-pol>
- [57] Layek, U., Baghira, N. K., Das, A., Kundu, A., & Karmakar, P. (2023). Dependency of crops on pollinators and pollination deficits: An approach to measurement considering the influence of various reproductive traits. *Agriculture*, 13(8), 1563. <https://doi.org/10.3390/agriculture13081563>
- [58] Lechenet, M., Dessaint, F., Py, G., Makowski, D., & Munier-Jolain, N. (2017). Reducing pesticide use while preserving crop productivity and profitability on arable farms. *Nature Plants*, 3(3), 17008. <https://doi.org/10.1038/nplants.2017.8>
- [59] Lichtenberg, E. M., Kennedy, C. M., Kremen, C., Batary, P., Berendse, F., Bommarco, R., ... & Crowder, D. W. (2017). A global synthesis of the effects of diversified farming systems on arthropod diversity within fields and across agricultural landscapes. *Global Change Biology*, 23(11), 4946–4957. <https://doi.org/10.1111/gcb.13714>
- [60] Lykogianni, M., Bempelou, E., Karamaouna, F., & Aliferis, K. A. (2021). Do pesticides promote or hinder sustainability in agriculture? The challenge of sustainable use of pesticides in modern agriculture. *Science of the Total Environment*, 795, 148625. <https://doi.org/10.1016/j.scitotenv.2021.148625>

- [61] Maes, J., Teller, A., Erhard, M., et al. (2020). *Mapping and Assessment of Ecosystems and Their Services: An EU Ecosystem Assessment*. Publications Office of the European Union, Luxembourg. <https://doi.org/10.2760/757183>
- [62] Markandya, A., Barton, D. N., Caparrós, A., Edens, B., Grammatikopoulou, I., Harris, R., & Obst, C. (2022). *Monetary Valuation for Ecosystem Services and Assets for Ecosystem Accounting*. System of Environmental Economic Accounting. https://seea.un.org/sites/seea.un.org/files/techreportvaluationv15_final_21072022.pdf
- [63] Martin, E. A., Dainese, M., Clough, Y., et al. (2019). The interplay of landscape composition and configuration: new pathways to manage functional biodiversity and agroecosystem services across Europe. *Ecology Letters*, 22(7), 1083–1094. <https://doi.org/10.1111/ele.13265>
- [64] Morandin, L. A., & Kremen, C. (2013). Hedgerow restoration promotes pollinator populations and exports native bees to adjacent fields. *Ecological Applications*, 23(4), 829–839. <https://doi.org/10.1890/12-1051.1>
- [65] Munir, M., Al-Hajhoj, M. R., Ghazzawy, H. S., Sallam, A. K. M., Al-Bahigan, A. M., & Al-Muiweed, M. A. (2020). A comparative study of pollination methods effect on the changes in fruit yield and quality of date palm cultivar Khalas. *Saudi Journal of Biological Sciences*, 27(9), 2340–2349.
- [66] Nesbit, M., Whiteoak, K., et al. (2022). *Biodiversity Financing and Tracking: Final Report*. Institute for European Environmental Policy and Trinomics.
- [67] Nicholson, C.C., Knapp, J., Kiljanek, T. et al. Pesticide use negatively affects bumble bees across European landscapes. *Nature* 628, 355–358 (2024). <https://doi.org/10.1038/s41586-023-06773-3>
- [68] Nieto, A., Roberts, S.P.M., Kemp, J., Rasmont, P., Kuhlmann, M., García Criado, M., Biesmeijer, J.C., Bogusch, P., Dathe, H.H., De la Rúa, P., De Meulemeester, T., Dehon, M., Dewulf, A., Ortiz-Sánchez, F.J., Lhomme, P., Pauly, A., Potts, S.G., Praz, C., Quaranta, M., Radchenko, V.G., Scheuchl, E., Smit, J., Straka, J., Terzo, M., Tomozii, B., Window, J. and Michez, D. 2014. European Red List of bees. Luxembourg: Publication Office of the European Union. Luxembourg: Publications Office of the European Union, <http://bookshop.europa.eu> IUCN Publications Services, www.iucn.org/publications A catalogue of IUCN publications is also available. <chrome-extension://efaidnbmninnibpcajpcglclefindmkaj/https://portals.iucn.org/library/sites/library/files/documents/RL-4-019.pdf>
- [69] Ollerton, J., Winfree, R., & Tarrant, S. (2011). How many flowering plants are pollinated by animals? *Oikos*, 120(3), 321–326. <https://doi.org/10.1111/j.1600-0706.2010.18644.x>
- [70] Ollerton, J., Erenler, H., Edwards, M., & Crockett, R. (2014). Extinctions of aculeate pollinators in Britain and the role of large-scale agricultural changes. *Science*, 346(6215), 1360–1362. <https://www.science.org/doi/10.1126/science.1257259>
- [71] Olmeda, C., Keenleyside, C., Underwood, E., et al. (2019). Evaluation of the effectiveness of European Union agricultural policies on pollinators. *Ecological Indicators*, 107, 105619. <https://www.ecologic.eu/17536>
- [72] Palmer, C., Groom, B., Sileci, L., & Langton, S. (2026). Biodiversity–food trade-offs when agricultural land is spared from production. *American Journal of Agricultural Economics*, 108(1), 254–284.
- [73] Perissi, I. (2025). Assessing the EU27 potential to meet the Nature Restoration Law targets. *Environmental Management*, 75(4), 711–729. <https://doi.org/10.1007/s00267-024-02010-9>

- [74] Perrot, T., Gaba, S., Roncoroni, M., Gautier, J. L., Saintilan, A., & Bretagnolle, V. (2019). Experimental quantification of insect pollination on sunflower yield, reconciling plant and field scale estimates. *Basic and Applied Ecology*, 34, 75–84. <https://doi.org/10.1016/j.baae.2018.09.005>
- [75] Polce, C., Maes, J., Rotllan-Puig, X., Michez, D., Castro, L., Cederberg, B., Dvorak, L., Fitzpatrick, Ú., Francis, F., Neumayer, J., & Manino, A. (2018). Distribution of bumblebees across Europe. *One Ecosystem*, 3, e28143. <https://doi.org/10.3897/oneeco.3.e28143>
- [76] Potts, S. G., Biesmeijer, J. C., Kremen, C., et al. (2010). Global pollinator declines: trends, impacts and drivers. *Trends in Ecology & Evolution*, Volume 25, Issue 6, 2010, Pages 345–353, ISSN 0169-5347, <https://doi.org/10.1016/j.tree.2010.01.007> <https://www.sciencedirect.com/science/article/pii/S0169534710000364?via%3Dihub>
- [77] Potts S., Biesmeijer K., Bommarco R., Breeze T., Carvalheiro L., Franzén M., González-Varo J.P., Holzschuh A., Kleijn D., Klein A.-M., Kunin, B., Lecocq T., Lundin O., Michez D., Neumann P., Nieto A., Penev L., Rasmont P., Ratamáki O., Riedinger V., Roberts S.P.M., Rundlöf M., Scheper J., Sørensen P., Steffan-Dewenter I., Stoev P., Vilà M., Schweiger O. (2015) Status and trends of European pollinators. Key findings of the STEP project. Pensoft Publishers, Sofia, 72 pp. <chrome-extension://efaidnbmnnnibpcajpcglclefindmkaj/http://step-project.net/img/uplf/STEP%20brochure%20online-1.pdf>
- [78] Potts, S., Imperatriz-Fonseca, V., Ngo, H. et al. Safeguarding pollinators and their values to human well-being. *Nature* 540, 220–229 (2016). <https://doi.org/10.1038/nature20588> <https://www.nature.com/articles/nature20588#citeas>
- [79] Potts, S., Dauber, J., Hochkirch, A., Oteman, B., Roy, D., Ahnre, K., Biesmeijer, K., Breeze, T., Carvell, C., Ferreira, C., Fitzpatrick, Ú., Isaac, N., Kuussaari, M., Ljubomirov, T., Maes, J., Ngo, H., Pardo, A., Polce, C., Quaranta, M., Settele, J., Sorg, M., Stefanescu, C. and Vujic, A., Proposal for an EU Pollinator Monitoring Scheme, EUR 30416 EN, *Publications Office of the European Union*, Luxembourg, 2020, ISBN 978-92-76-23859-1, doi:10.2760/881843, JRC122225. <https://publications.jrc.ec.europa.eu/repository/handle/JRC122225>
- [80] Potts, S. G., Bartomeus, I., Biesmeijer, K., et al. (2024). European Commission, Joint Research Centre, *Refined Proposal for an EU Pollinator Monitoring Scheme*. Publications Office of the European Union, Luxembourg. <https://doi.org/10.2760/2005545>
- [81] Powney, G.D., Carvell, C., Edwards, M. et al. Widespread losses of pollinating insects in Britain. *Nat Commun* 10, 1018 (2019). <https://doi.org/10.1038/s41467-019-08974-9> <https://www.nature.com/articles/s41467-019-08974-9#citeas>
- [82] Prestele, R., Brown, C., Polce, C., Maes, J., & Whitehorn, P. (2021). Large variability in response to projected climate and land-use changes among European bumblebee species. *Global Change Biology*, 27(19), 4530–4545. <https://doi.org/10.1111/gcb.15780>
- [83] Qiu, T., Shi, Y., Peñuelas, J., Liu, J., Cui, Q., Sardans, J., ... & Fang, L. (2024). Optimizing cover crop practices as a sustainable solution for global agroecosystem services. *Nature Communications*, 15(1), 10617. <https://doi.org/10.1038/s41467-024-55068-8>
- [84] Rader R, Bartomeus I, Garibaldi LA, Garratt MP, Howlett BG, Winfree R, Cunningham SA, Mayfield MM, Arthur AD, Andersson GK, Bommarco R, Brittain C, Carvalheiro LG, Chacoff NP, Entling MH, Foully B, Freitas BM, Gemmill-Herren B, Ghazoul J, Griffin SR, Gross CL, Herbertsson L, Herzog F, Hipólito J, Jaggard S, Jauker F, Klein AM, Kleijn D, Krishnan S, Lemos CQ, Lindström SA, Mandelik Y, Monteiro VM, Nelson W,

- Nilsson L, Pattemore DE, Pereira Nde O, Pisanty G, Potts SG, Reemer M, Rundlöf M, Sheffield CS, Scheper J, Schüepp C, Smith HG, Stanley DA, Stout JC, Szentgyörgyi H, Taki H, Vergara CH, Viana BF, Woyciechowski M. Non-bee insects are important contributors to global crop pollination. *Proc Natl Acad Sci U S A*. 2016 Jan 5;113(1):146–51. doi: 10.1073/pnas.1517092112. Epub 2015 Nov 30. PMID: 26621730; PMCID: PMC4711867. <https://pubmed.ncbi.nlm.nih.gov/26621730/>
- [85] Rahimi, E.; Jung, C. Global Trends in Climate Suitability of Bees: Ups and Downs in a Warming World. *Insects* 2024, 15, 127. <https://doi.org/10.3390/insects15020127> <https://www.mdpi.com/2075-4450/15/2/127>
- [86] Rahimi, E., Jung, C. A Global Estimation of Potential Climate Change Effects on Pollinator-Dependent Crops. *Agric Res* 14, 812–822 (2025). <https://doi.org/10.1007/s40003-024-00802-x>. <https://link.springer.com/article/10.1007/s40003-024-00802-x#citeas>
- [87] Rasmont, P., Franzen, M., Lecocq, T., et al. (2015). Climatic risk and distribution atlas of European bumblebees. *BioRisk*, 10, 1–246. <https://doi.org/10.3897/biorisk.10.4749> <https://biorisk.pensoft.net/articles.php?id=4749>
- [88] Reif, J., & Vermouzek, Z. (2019). Collapse of farmland bird populations in an Eastern European country following its EU accession. *Conservation Letters*, 12(1), e12585. <https://doi.org/10.1111/conl.12585>
- [89] Riahi, K., van Vuuren, D. P., Kriegler, E., et al. (2017). The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview. *Global Environmental Change*, 42, 153–168. <https://doi.org/10.1016/j.gloenvcha.2016.05.009>
- [90] Ricketts, T. H., Regetz, J., Steffan-Dewenter, I., et al. (2008). Landscape effects on crop pollination services: are there general patterns? *Ecology Letters*, 11(5), 499–515. <https://doi.org/10.1111/j.1461-0248.2008.01157.x>
- [91] Settele, J., Kudrna, O., Harpke, A., et al. (2008). *Climatic Risk Atlas of European Butterflies*. Pensoft, Sofia-Moscow.
- [92] Siopa, C., Carvalheiro, L. G., Castro, H., Loureiro, J., Castro, S. (2024). Animal-pollinated crops and cultivars. A quantitative assessment of pollinator dependence values and evaluation of methodological approaches. *Journal of Applied Ecology*, 61, 1279–1288. <https://doi.org/10.1111/1365-2664.14634>
- [93] Standardi G, Dasgupta S., Parrado R, De Cian E., Bosello F. (2023). Assessing Macro-economic Effects of Climate Impacts on Energy Demand in EU Sub-national Regions, *Environmental and Resource Economics*, 86, 173–201, <https://doi.org/10.1007/s10640-023-00792-4>
- [94] Tamburini, G., Bommarco, R., Wanger, T. C., Kremen, C., Van Der Heijden, M. G., Liebman, M., & Hallin, S. (2020). Agricultural diversification promotes multiple ecosystem services without compromising yield. *Science Advances*, 6(45), eaba1715. <https://doi.org/10.1126/sciadv.aba1715>
- [95] Tang, F.H.M., Nguyen, T.H., Conchedda, G. et al. CROPGRIDS: a global geo-referenced dataset of 173 crops. *Sci Data* 11, 413 (2024). <https://doi.org/10.1038/s41597-024-03247-7> <https://www.nature.com/articles/s41597-024-03247-7#citeas>
- [96] Taylor, K. E., Stouffer, R. J., & Meehl, G. A. (2012). An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society*, 93(4), 485–498. <https://doi.org/10.1175/BAMS-D-11-00094.1>
- [97] Tsiafouli, M. A., Thébault, E., Sgardelis, S. P., et al. (2015). Intensive agriculture reduces soil biodiversity across Europe. *Global Change Biology*, 21(2), 973–985. <https://doi.org/10.1111/gcb.12752>

- [98] Tucker, G., Underwood, E., Farmer, A., Scalera, R., Dickie, I., McConville, A., & van Vliet, W. (2013). *Estimation of the Financing Needs to Implement Target 2 of the EU Biodiversity Strategy*. Report to the European Commission. Institute for European Environmental Policy, London.
- [99] Underwood, E., Mottershead, D., Hart, K., Meredith, S., Keenleyside, C., Tucker, G., & Gerritsen, E. (2020). Alliance Environnement (2019) Evaluation of the impact of the CAP on habitats, landscapes, biodiversity. <https://doi.org/10.2762/818843> https://www.researchgate.net/profile/Stephen-Meredith/publication/341578944_Alliance_Environnement_2019_Evaluation_of_the_impact_of_the_CAP_on_habitats_landscapes_biodiversity/links/60cb103fa6fdcc01d47b4f4b/Alliance-Environnement-2019-Evaluation-of-the-impact-of-the-CAP-on-habitats-landscapes-biodiversity.pdf
- [100] United Nations. (2024). *System of Environmental-Economic Accounting. Ecosystem Accounting*. ST/ESA/STAT/SER.F/124. ISBN: 978-92-1-259183-4. <https://seea.un.org/ecosystem-accounting>
- [101] Vallecillo, S., La Notte, A., Polce, C., Zulian, G., Alexandris, N., Ferrini, S., & Maes, J. (2018). *Ecosystem Services Accounting: Part I – Outdoor Recreation and Crop Pollination*. EUR 29024 EN; Publications Office of the European Union, Luxembourg. <https://publications.jrc.ec.europa.eu/repository/handle/JRC110321>
- [102] Vallecillo, S., La Notte, A., Polce, C., Garcia Bendito, E., & Maes, J. (2020). INCA – Crop Pollination. European Commission, Joint Research Centre (JRC) [Dataset]. <http://data.europa.eu/89h/650331f3-e7ce-427b-8011-bd2c8f40599c>
- [103] Vallecillo, S., Van Loo, M., La Notte, A., et al. (2022). Ecosystem services accounts: maps for 7 ecosystem services produced in KIP INCA 2018 updates. European Commission, Joint Research Centre (JRC) [Dataset]. <http://data.europa.eu/89h/4cbd7c1e-6512-4ebe-8ca5-e08209cc3efb>
- [104] van der Wijst K.I., Bosello F., Dasgupta S., Parrado R., Standardi G., et al. (2023). New damage curves and multimodel analysis suggest lower optimal temperature. *Nature Climate Change*, 13, 434–441, <https://doi.org/10.1038/s41558-023-01636-1>
- [105] Van Swaay, C. A. M., et al. (2022). EU Grassland Butterfly Index 1991–2022 Technical report. Butterfly Conservation Europe & EMBRACE/eBMS. (2022). European Grassland Butterfly Indicator 1990–2020 Technical report. Butterfly Conservation Europe & SPRING/eBMS (www.butterfly-monitoring.net) & Vlinderstichting report VS2022.039. <chrome-extension://efaidnbmnnnibpcajpgclclefindmkaj/https://assets.vlinderstichting.nl/docs/290cb16a-e90f-4c5b-a7df-9b954d511cfa.pdf> <https://zenodo.org/records/16367397>
- [106] Van Swaay, C. A. M., Schmucki, R., Roy, D. B., et al. (2025). (2025). EU Grassland Butterfly Index 1991–2023 Technical report. Butterfly Conservation Europe & EMBRACE/eBMS (www.butterfly-monitoring.net) & Vlinderstichting report VS2025.014. <https://doi.org/10.5281/zenodo.16281872> <https://zenodo.org/records/16367397>
- [107] van Vuuren, D. P., Kriegler, E., O'Neill, B. C., et al. (2014). A new scenario framework for Climate Change Research: scenario matrix architecture. *Climatic Change*, 122(3), 373–386. <https://doi.org/10.1007/s10584-013-0906-1>
- [108] Vergara, C. H., & Badano, E. I. (2009). Pollinator diversity increases fruit production in Mexican coffee plantations: The importance of rustic management systems. *Agriculture, Ecosystems & Environment*, 129(1–3), 117–123.

- [109] von Jeetze, P.J., Weindl, I., Johnson, J.A. et al. Projected landscape-scale repercussions of global action for climate and biodiversity protection. *Nat Commun* 14, 2515 (2023). <https://doi.org/10.1038/s41467-023-38043-1>
- [110] Voříšek, P., Jiguet, F., van Strien, A., Škorpilová, J., Klvaňová, A., & Gregory, R. D. (2010). Trends in abundance and biomass of widespread European farmland birds: how much have we lost. *BOU Proceedings – Lowland Farmland Birds III*, 24.
- [111] Wang, J., Zhang, L., Liu, K., Zhou, B., Gao, H., Han, X., ... & Zhang, W. (2025). Minimizing the potential risk of soil nitrogen loss through optimal fertilization practices in intensive agroecosystems. *Agronomy for Sustainable Development*, 45(1), 9. <https://doi.org/10.1007/s13593-024-00977-6>
- [112] Weibull, A. C., Östman, Ö., & Granqvist, Å. (2003). Species richness in agroecosystems: the effect of landscape, habitat and farm management. *Biodiversity & Conservation*, 12, 1335–1355. <https://doi.org/10.1023/A:1023617117780>
- [113] Winfree, R., Aguilar, R., Vázquez, D. P., LeBuhn, G., & Aizen, M. A. (2009). A meta-analysis of bees' responses to anthropogenic disturbance. *Ecology*, 90(8), 2068–2076. <https://doi.org/10.1890/08-1245.1>
- [114] Winfree, R., Reilly, J. R., Bartomeus, I., et al. (2023). Species turnover promotes the importance of bee diversity for crop pollination at regional scales. *Science*, 359(6377), 791–793. <https://doi.org/10.1126/science.aao2117>
- [115] You, L., Ros, G. H., Chen, Y., Zhang, F., & de Vries, W. (2024). Optimized agricultural management reduces global cropland nitrogen losses to air and water. *Nature Food*, 5(12), 995–1004. <https://doi.org/10.1038/s43016-024-01074-0>
- [116] Sebastiano Zanini, Matteo Dainese, Lisa Obwegs, Elia Guariento, Timo Kopf, Matteo Anderle, Georg Leitinger, Ulrike Tappeiner. Disentangling biodiversity and temperature effects on bees and pollination services in mountain agroecosystems, *Agriculture, Ecosystems & Environment*, Volume 395, 2026, 109945, ISSN 0167-8809, <https://doi.org/10.1016/j.agee.2025.109945>. <https://www.sciencedirect.com/science/article/pii/S0167880925004773>
- [117] Zattara, E. E., & Aizen, M. A. (2021). Worldwide occurrence records suggest a global decline in bee species richness. *One Earth*, 4(1), 114–123. <https://doi.org/10.1016/j.oneear.2020.12.005>