

Monotone multilevel for FOSLS linear elastic contact

G. Rovi \triangle , B. Kober \circ , G. Starke \circ , R. Krause \triangle

\circ : Universität Duisburg-Essen, Germany

\triangle : Università della Svizzera italiana, Switzerland

February 10, 2019

Università
della
Svizzera
italiana

Institute of
Computational
Science
ICS


UNIVERSITÄT
DUISBURG
ESSEN

Offen im Denken

Examples of contact problems

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle



- Contact problems with incompressible materials.
- Quantities of interest: the forces generated by the contact.

Signorini's problem: strong formulation

- **First Order System Linear Elasticity**

Given displacement \mathbf{u} , stress $\boldsymbol{\sigma}$, body Ω :

$$\begin{cases} \operatorname{div} \boldsymbol{\sigma} + \mathbf{f} = 0 & \Omega & \text{momentum balance equation} \\ \mathcal{A} \boldsymbol{\sigma} - \boldsymbol{\varepsilon}(\mathbf{u}) = 0 & \Omega & \text{constitutive law} \\ \mathbf{u} = \mathbf{u}_D & \Gamma_D & \text{Dirichlet BC} \\ \boldsymbol{\sigma} \mathbf{n} = \mathbf{t}_N & \Gamma_N & \text{Neumann BC} \end{cases}$$

where $\boldsymbol{\varepsilon}(\mathbf{u}) = \frac{1}{2}(\nabla \mathbf{u} + \nabla \mathbf{u}^T)$, $\mathcal{A} \boldsymbol{\sigma} = \frac{1}{2\mu} \left(\boldsymbol{\sigma} - \frac{\lambda}{d\lambda + 2\mu} \operatorname{tr} \boldsymbol{\sigma} \mathbf{I} \right)$ and μ, λ are the Lamé parameters

Signorini's problem: strong formulation

First Order System Linear Elasticity

Given displacement \mathbf{u} , stress $\boldsymbol{\sigma}$, body Ω :



$$\begin{cases} \operatorname{div} \boldsymbol{\sigma} + \mathbf{f} = 0 & \Omega & \text{momentum balance equation} \\ \mathcal{A} \boldsymbol{\sigma} - \boldsymbol{\varepsilon}(\mathbf{u}) = 0 & \Omega & \text{constitutive law} \\ \mathbf{u} = \mathbf{u}_D & \Gamma_D & \text{Dirichlet BC} \\ \boldsymbol{\sigma} \mathbf{n} = \mathbf{t}_N & \Gamma_N & \text{Neumann BC} \end{cases}$$

where $\boldsymbol{\varepsilon}(\mathbf{u}) = \frac{1}{2}(\nabla \mathbf{u} + \nabla \mathbf{u}^T)$, $\mathcal{A} \boldsymbol{\sigma} = \frac{1}{2\mu} \left(\boldsymbol{\sigma} - \frac{\lambda}{d\lambda + 2\mu} \operatorname{tr} \boldsymbol{\sigma} \mathbf{I} \right)$ and μ, λ are the Lamé parameters

Contact Constraints

Given the gap function g , the normal and tangent vectors \mathbf{n} and \mathbf{t} :

$$\begin{cases} \mathbf{u} \cdot \mathbf{n} - g \leq 0 & \Gamma_C \text{ impenetrability} \\ (\boldsymbol{\sigma} \mathbf{n}) \cdot \mathbf{n} \leq 0 & \Gamma_C \text{ direction of the surface pressure} \\ (\mathbf{u} \cdot \mathbf{n} - g) ((\boldsymbol{\sigma} \mathbf{n}) \cdot \mathbf{n}) = 0 & \Gamma_C \text{ complementarity condition} \\ \mathbf{t}_i^T (\boldsymbol{\sigma} \mathbf{n}) = 0 & \Gamma_C \text{ frictionless condition} \end{cases}$$



The Least-Squares Formulation

- **First Order System Least-Squares (FOSLS) Functional**

$$C_1, C_2, C_3 > 0$$

$$\mathcal{J}(\mathbf{u}, \boldsymbol{\sigma}) = C_1 \|\operatorname{div} \boldsymbol{\sigma} + \mathbf{f}\|_{L^2(\Omega)^d}^2 + C_2 \|\mathcal{A} \boldsymbol{\sigma} - \boldsymbol{\varepsilon}(\mathbf{u})\|_{L^2(\Omega)^d}^2 + C_3 \langle \mathbf{u} \cdot \mathbf{n} - g, (\boldsymbol{\sigma} \mathbf{n}) \cdot \mathbf{n} \rangle_{\Gamma_c}$$

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi[△],
B.
Kober[○],
G.
Starke[○],
R.
Krause[△]

- Rolf Krause, Benjamin Müller, and Gerhard Starke. An adaptive least-squares mixed finite element method for the Signorini problem. Numerical Methods for Partial Differential Equations, 33(1):276-289, 2017.

The Least-Squares Formulation

- First Order System Least-Squares (FOSLS) Functional

$$C_1, C_2, C_3 > 0$$

$$\mathcal{J}(\mathbf{u}, \boldsymbol{\sigma}) = C_1 \|\operatorname{div} \boldsymbol{\sigma} + \mathbf{f}\|_{L^2(\Omega)^d}^2 + C_2 \|\mathcal{A} \boldsymbol{\sigma} - \boldsymbol{\varepsilon}(\mathbf{u})\|_{L^2(\Omega)^d}^2 + C_3 \langle \mathbf{u} \cdot \mathbf{n} - g, (\boldsymbol{\sigma} \mathbf{n}) \cdot \mathbf{n} \rangle_{\Gamma_C}$$

- Convex Set K

$$K = \{(\mathbf{u}, \boldsymbol{\sigma}) \in [H_{\Gamma_d}^1(\Omega)]^d \times [H_{\operatorname{div}, \Gamma_N}(\Omega)]^d : \mathbf{u} \cdot \mathbf{n} - g \leq 0, (\boldsymbol{\sigma} \mathbf{n}) \cdot \mathbf{n} \leq 0, \mathbf{t}_i^T(\boldsymbol{\sigma} \mathbf{n}) = 0 \quad \Gamma_C\}$$

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi[△],
B.
Kober[○],
G.
Starke[○],
R.
Krause[△]

- Rolf Krause, Benjamin Müller, and Gerhard Starke. An adaptive least-squares mixed finite element method for the Signorini problem. Numerical Methods for Partial Differential Equations, 33(1):276-289, 2017.

The Least-Squares Formulation

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

- **First Order System Least-Squares (FOSLS) Functional**

$$C_1, C_2, C_3 > 0$$

$$\mathcal{J}(\mathbf{u}, \boldsymbol{\sigma}) = C_1 \|\operatorname{div} \boldsymbol{\sigma} + \mathbf{f}\|_{L^2(\Omega)^d}^2 + C_2 \|\mathcal{A} \boldsymbol{\sigma} - \boldsymbol{\varepsilon}(\mathbf{u})\|_{L^2(\Omega)^d}^2 + C_3 \langle \mathbf{u} \cdot \mathbf{n} - g, (\boldsymbol{\sigma} \mathbf{n}) \cdot \mathbf{n} \rangle_{\Gamma_C}$$

- **Convex Set K**

$$K = \{(\mathbf{u}, \boldsymbol{\sigma}) \in [H_{\Gamma_d}^1(\Omega)]^d \times [H_{\operatorname{div}, \Gamma_N}(\Omega)]^d : \mathbf{u} \cdot \mathbf{n} - g \leq 0, (\boldsymbol{\sigma} \mathbf{n}) \cdot \mathbf{n} \leq 0, \mathbf{t}_i^T(\boldsymbol{\sigma} \mathbf{n}) = 0 \quad \Gamma_C\}$$

- Find $(\mathbf{u}, \boldsymbol{\sigma}) \in K$, such that:

- **Minimization problem:** $\mathcal{J}(\mathbf{u}, \boldsymbol{\sigma}) \leq \mathcal{J}(\mathbf{v}, \boldsymbol{\tau}) \quad \forall (\mathbf{v}, \boldsymbol{\tau}) \in K$

$$\iff$$

- **Variational Inequality:**
$$\left\{ \begin{array}{l} \left\langle \frac{\partial \mathcal{J}(\mathbf{u}, \boldsymbol{\sigma}; \mathbf{f}, g)}{\partial \mathbf{u}}, \mathbf{v} - \mathbf{u} \right\rangle \geq 0 \\ \left\langle \frac{\partial \mathcal{J}(\mathbf{u}, \boldsymbol{\sigma}; \mathbf{f}, g)}{\partial \boldsymbol{\sigma}}, \boldsymbol{\tau} - \boldsymbol{\sigma} \right\rangle \geq 0 \end{array} \right. \quad \forall (\mathbf{v}, \boldsymbol{\tau}) \in K$$

● Rolf Krause, Benjamin Müller, and Gerhard Starke. An adaptive least-squares mixed finite element method for the Signorini problem. Numerical Methods for Partial Differential Equations, 33(1):276-289, 2017.

Discretization

- Discretized domain Ω_L

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi ^{\triangle} ,
B.
Kober ^{\circ} ,
G.
Starke ^{\circ} ,
R.
Krause ^{\triangle}

Discretization

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Ravi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

- Discretized domain Ω_L
- FE space $X_L = P_{\Gamma_D}^1(\Omega_L) \times \mathcal{RT}_{0,\Gamma_N}(\Omega_L)$ with $\mathbf{x}_L = (\mathbf{u}_L, \boldsymbol{\sigma}_L) \in X_L$

Discretization

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Ravi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

- **Discretized domain** Ω_L
- **FE space** $X_L = P_{\Gamma_D}^1(\Omega_L) \times \mathcal{RT}_{0,\Gamma_N}(\Omega_L)$ with $\mathbf{x}_L = (\mathbf{u}_L, \boldsymbol{\sigma}_L) \in X_L$
- $\mathbf{f}_L, \mathbf{u}_{D,L}, \mathbf{t}_{N,L}, g_L$ FE representations of $\mathbf{f}, \mathbf{u}_D, \mathbf{t}_N, g$

Discretization

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi[△],
B.
Kober[○],
G.
Starke[○],
R.
Krause[△]

- **Discretized domain** Ω_L
- **FE space** $X_L = P_{\Gamma_D}^1(\Omega_L) \times \mathcal{RT}_{0,\Gamma_N}(\Omega_L)$ with $\mathbf{x}_L = (\mathbf{u}_L, \boldsymbol{\sigma}_L) \in X_L$
- $\mathbf{f}_L, \mathbf{u}_{D,L}, \mathbf{t}_{N,L}, g_L$ FE representations of $\mathbf{f}, \mathbf{u}_D, \mathbf{t}_N, g$
- **Discrete FOSLS Functional**

$$\mathcal{J}(\mathbf{x}_L; \mathbf{f}_L) = \frac{1}{2} \mathbf{x}_L^T \mathbf{A}_L \mathbf{x}_L - \mathbf{x}_L^T \mathbf{f}_L$$

Discretization

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Ravi \triangle ,
B.
Kober o ,
G.
Starke o ,
R.
Krause \triangle

- **Discretized domain** Ω_L
- **FE space** $X_L = P_{\Gamma_D}^1(\Omega_L) \times \mathcal{RT}_{0,\Gamma_N}(\Omega_L)$ with $\mathbf{x}_L = (\mathbf{u}_L, \boldsymbol{\sigma}_L) \in X_L$
- $\mathbf{f}_L, \mathbf{u}_{D,L}, \mathbf{t}_{N,L}, g_L$ FE representations of $\mathbf{f}, \mathbf{u}_D, \mathbf{t}_N, g$
- **Discrete FOSLS Functional**

$$\mathcal{J}(\mathbf{x}_L; \mathbf{f}_L) = \frac{1}{2} \mathbf{x}_L^T \mathbf{A}_L \mathbf{x}_L - \mathbf{x}_L^T \mathbf{f}_L$$

- **Convex Set** K_L (in general $K_L \not\subseteq K$)

$$\mathbf{x}_L \in K_L \quad \Longleftrightarrow \quad \mathbf{B}_L \mathbf{x}_L \leq \mathbf{g}_L$$

Discretization

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi[△],
B.
Kober[○],
G.
Starke[○],
R.
Krause[△]

- Discretized domain Ω_L
- FE space $X_L = P_{\Gamma_D}^1(\Omega_L) \times \mathcal{RT}_{0,\Gamma_N}(\Omega_L)$ with $\mathbf{x}_L = (\mathbf{u}_L, \boldsymbol{\sigma}_L) \in X_L$
- $\mathbf{f}_L, \mathbf{u}_{D,L}, \mathbf{t}_{N,L}, g_L$ FE representations of $\mathbf{f}, \mathbf{u}_D, \mathbf{t}_N, g$
- Discrete FOSLS Functional

$$\mathcal{J}(\mathbf{x}_L; \mathbf{f}_L) = \frac{1}{2} \mathbf{x}_L^T \mathbf{A}_L \mathbf{x}_L - \mathbf{x}_L^T \mathbf{f}_L$$

- Convex Set K_L (in general $K_L \not\subseteq K$)

$$\mathbf{x}_L \in K_L \iff \mathbf{B}_L \mathbf{x}_L \leq \mathbf{g}_L$$

- Minimization problem:

Find $\mathbf{x}_L \in K_L$

$$\operatorname{argmin} \mathcal{J}(\mathbf{x}_L; \mathbf{f}_L) = \frac{1}{2} \mathbf{x}_L^T \mathbf{A}_L \mathbf{x}_L - \mathbf{x}_L^T \mathbf{f}_L$$

$$\mathbf{B}_L \mathbf{x}_L \leq \mathbf{g}_L$$

Disadvantages and Advantages of the FOSLS

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Ravi
B.
Kober^O,
G.
Starke^O,
R.
Krause[△]

Pros

- Direct access to stress σ (friction, plasticity...)
- Dealing with incompressible materials ($\lambda \rightarrow \infty$)
- FOSLS functional as an a posteriori error estimator
- Flexible choice of finite element spaces (**low order**: $\mathbf{u}_L \in P^1$, $\sigma_L \in \mathcal{RT}_0$)
- Symmetric positive definite system

- Attia, Frank S., Zhiqiang Cai, and Gerhard Starke. "First-order system least squares for the Signorini contact problem in linear elasticity". SIAM Journal on Numerical Analysis 47.4 (2009): 3027-3043.

Disadvantages and Advantages of the FOSLS

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

Pros

- Direct access to stress σ (friction, plasticity...)
- Dealing with incompressible materials ($\lambda \rightarrow \infty$)
- FOSLS functional as an a posteriori error estimator
- Flexible choice of finite element spaces (**low order**: $\mathbf{u}_L \in P^1$, $\sigma_L \in \mathcal{RT}_0$)
- Symmetric positive definite system

Cons

- The functional is fictitious, not physical
- The asymmetry of the stress tensor
- Find proper weights C_1 , C_2 , C_3
- Large condition number: **need for a preconditioner**

• Attia, Frank S., Zhiqiang Cai, and Gerhard Starke. "First-order system least squares for the Signorini contact problem in linear elasticity". SIAM Journal on Numerical Analysis 47.4 (2009): 3027-3043.

Disadvantages and Advantages of the FOSLS

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Ravi \triangle ,
B.
Kober o ,
G.
Starke o ,
R.
Krause \triangle

Pros

- Direct access to stress σ (friction, plasticity...)
- Dealing with incompressible materials ($\lambda \rightarrow \infty$)
- FOSLS functional as an a posteriori error estimator
- Flexible choice of finite element spaces (**low order**: $\mathbf{u}_L \in P^1$, $\sigma_L \in \mathcal{RT}_0$)
- Symmetric positive definite system

Cons

- The functional is fictitious, not physical
- The asymmetry of the stress tensor
- Find proper weights C_1 , C_2 , C_3
- Large condition number: **need for a preconditioner**

$\left\{ \begin{array}{l} \text{Functional to be minimized} \\ \text{Local constraints} \\ \text{Need for a preconditioner} \end{array} \right. \Rightarrow \text{Monotone Multilevel}$

• Attia, Frank S., Zhiqiang Cai, and Gerhard Starke. "First-order system least squares for the Signorini contact problem in linear elasticity". SIAM Journal on Numerical Analysis 47.4 (2009): 3027-3043.

Monotone Multilevel strategy

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

- Successive energy minimization by means of local corrections
- No correction can increase energy
- Fine space corrections on fine grid (non-linear Gauß-Seidel) \Rightarrow global convergence
- Coarse space corrections \Rightarrow accelerating convergence

Monotone Multilevel by energy minimization

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Ravi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle




Monotone Multilevel by energy minimization

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

R_i^{i-1} restriction operator ($i = L, \dots, 2$)




Monotone Multilevel by energy minimization

Monotone
multilevel
for FOSLS
linear
elastic
contact

G. Rovi[△],
B. Kober[○],
G. Starke[○],
R. Krause[△]

R_i^{i-1} restriction operator ($i = L, \dots, 2$)




Monotone Multilevel by energy minimization

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

R_i^{i-1} restriction operator, I_{i-1}^i interpolation operator ($i = L, \dots, 2$)



Smoother

- Standard non-linear Gauß-Seidel smooths H^1 , **but not** H_{div}
- The kernel $\text{Ker}(\text{div}) = \{\boldsymbol{\tau} \in H_{\text{div}}, \text{div } \boldsymbol{\tau} = 0\}$ is too large
- Patch-smoother for divergence-free components of the error

Interpolations and restrictions

- Standard P^1 and RT_0 interpolations and restrictions for primal and dual variables
- Non-linear projections for constraint representation on coarser levels

Multilevel ingredients

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Ravi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

Smoother

- Standard non-linear Gauß-Seidel smooths H^1 , **but not** H_{div}
- The kernel $\text{Ker}(\text{div}) = \{\boldsymbol{\tau} \in H_{\text{div}}, \text{div } \boldsymbol{\tau} = 0\}$ is too large
- **Patch-smoother** for divergence-free components of the error

Interpolations and restrictions


- Standard P^1 and RT_0 interpolations and restrictions for primal and dual variables
- Non-linear projections for constraint representation on coarser levels

LS Patch smoother

Monotone
multilevel
for FOSLS
linear
elastic
contact

G. \triangle ,
Rovi \triangle ,
B. \triangle ,
Kober \circ ,
G. \circ ,
Starke \circ ,
R. \triangle ,
Krause \triangle

- Mesh level $j = 1, \dots, L$, vertex $\nu = 1, \dots, N_j$
- $\text{Patch}_{j,\nu}$ = dofs of node ν and surrounding edges/faces (2D/3D)
- $K_{j,\nu}$ = local closed convex set spanned by basis functions in $\text{Patch}_{j,\nu}$
- Minimization of \mathcal{J} on $K_{j,\nu}$
- Error smoothed in H^1 and H_{div} simultaneously



- Ralf Hiptmair. Multigrid method for $H(\text{div})$ in three dimensions. Electron. Trans. Numer. Anal, 6(1):133-152, 1997.
- Douglas N Arnold, Richard S Falk, and Ragnar Winther. Multigrid in $H(\text{div})$ and $H(\text{curl})$. Numerische Mathematik, 85(2):197-217, 2000.
- Gerhard Starke. Gauss-Newton multilevel methods for least-squares finite element computations of variably saturated subsurface flow. Computing, 64(4):323-338, 2000.

Smoother

- Standard non-linear Gauß-Seidel smooths H^1 , **but not** H_{div}
- The kernel $\text{Ker}(\text{div}) = \{\boldsymbol{\tau} \in H_{\text{div}}, \text{div } \boldsymbol{\tau} = 0\}$ is too large
- Patch-smoother for divergence-free components of the error

Interpolations and restrictions

- Standard P^1 and RT_0 interpolations and restrictions for primal and dual variables
- Non-linear projections for **constraint representation on coarser levels**

Non-Linear Projections for Coarse Constraints

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle


- **Exact monotone multilevel**
Comparison of coarse function \mathbf{c}_j with **fine constraint** \Rightarrow **suboptimal complexity**
- **Approximate monotone multilevel**
Comparison of coarse function \mathbf{c}_j with **coarse constraint** \Rightarrow **optimal complexity**

Non-Linear Projections for Coarse Constraints

Monotone
multilevel
for FOSLS
linear
elastic
contact


G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

- **Exact monotone multilevel**
Comparison of coarse function c_j with **fine constraint** \Rightarrow **suboptimal complexity**
- **Approximate monotone multilevel**
Comparison of coarse function c_j with **coarse constraint** \Rightarrow **optimal complexity**



Non-Linear Projections for Coarse Constraints

- **Exact monotone multilevel**
Comparison of coarse function c_j with **fine constraint** \Rightarrow **suboptimal complexity**
- **Approximate monotone multilevel**
Comparison of coarse function c_j with **coarse constraint** \Rightarrow **optimal complexity**




Non-Linear Projections for Coarse Constraints

Monotone
multilevel
for FOSLS
linear
elastic
contact


G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

- **Exact monotone multilevel**
Comparison of coarse function c_j with **fine constraint** \Rightarrow **suboptimal complexity**
- **Approximate monotone multilevel**
Comparison of coarse function c_j with **coarse constraint** \Rightarrow **optimal complexity**



Non-Linear Projections for Coarse Constraints

- **Exact monotone multilevel**
Comparison of coarse function c_j with **fine constraint** \Rightarrow **suboptimal complexity**
- **Approximate monotone multilevel**
Comparison of coarse function c_j with **coarse constraint** \Rightarrow **optimal complexity**



Monotone
multilevel
for FOSLS
linear
elastic
contact


G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

Non-Linear Projections for Coarse Constraints



Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi[△],
B.
Kober[○],
G.
Starke[○],
R.
Krause[△]

- Exact monotone multilevel
Comparison of coarse function c_j with **fine constraint** \Rightarrow **suboptimal complexity**
- Approximate monotone multilevel
Comparison of coarse function c_j with **coarse constraint** \Rightarrow **optimal complexity**




- Different consistent coarse constraints



Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle




$\mu = 1$, $\lambda = 1, \infty$ (compressible and **incompressible**)

Hertzian Contact, two-level problem

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle




- **First phase:** non-linear, capturing high frequencies
- **Second phase:** linear, known active set (blue, green), and not already known active set (red)
- **Similar behaviour** of compressible and incompressible cases

Hertzian Contact - Setting

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi[△],
B.
Kober[○],
G.
Starke[○],
R.
Krause[△]



$\mu = 1$, $\lambda = 1, \infty$ (compressible and **incompressible**)

Signorini's problem, square mesh

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle





Figure: Square mesh. Compressible material.





Figure: Square mesh. Incompressible material.

- Purely linear problem: h – and L – independency

Conclusions

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi[△],
B.
Kober[○],
G.
Starke[○],
R.
Krause[△]

- Monotone multilevel for FOSLS linear elastic contact
- Limite case: h - and L - independency
- Similar behaviour of compressible and incompressible cases

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Ravi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

Thank you for your attention!

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Ravi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi[△],
B.
Kober[○],
G.
Starke[○],
R.
Krause[△]

Exact Monotone Multilevel

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

Define:

- $\mathbf{x}_j^k = (\mathbf{u}_j^k, \boldsymbol{\sigma}_j^k) \in K_J$ k -th iterate
- $\mathbf{x}_{J,0} = \mathbf{x}_J^k$
- $\mathbf{x}_{j,0} = \mathbf{x}_{j+1,N_{j+1}}$, for $j = J-1, \dots, 1$

Compute a sequence of intermediate iterates $\mathbf{x}_{j,\nu} = \mathbf{x}_{j,\nu-1} + \mathbf{c}_{j,\nu}$:

$$\mathcal{J} \leq \mathcal{J}(\mathbf{x}_{j,\nu} + \mathbf{y}) \quad \forall \mathbf{y} \in K_{j,\nu}^* \quad j = J, \dots, 2, \quad \nu = 1, \dots, N_j$$

$$\mathcal{J}(\mathbf{x}_{2,N_2} + \mathbf{c}_1) \leq \mathcal{J}(\mathbf{x}_{2,N_2} + \mathbf{y}) \quad \forall \mathbf{y} \in K_1^* \quad j = 1$$

with the **exact** local closed convex sets $K_{j,\nu}^*$ and K_1^* :

$$K_{j,\nu}^*(\mathbf{x}_{j,\nu}) = \{\mathbf{y} \in \text{span}\{\boldsymbol{\lambda}_{j,\nu}\} : \mathbf{y} + \mathbf{x}_{j,\nu} \in K_J\}$$

$$K_1^*(\mathbf{x}_{2,N_2}) = \{\mathbf{y} \in \text{span}\{\boldsymbol{\lambda}_1\} : \mathbf{y} + \mathbf{x}_{2,N_2} \in K_J\}$$

Ralf Kornhuber. Monotone multigrid methods for elliptic variational inequalities I. Numerische Mathematik, 69(2):167-184, 1994.

Ralf Kornhuber and Rolf Krause. Adaptive multigrid methods for Signorini's problem in linear elasticity. Computing and Visualization in Science, 4(1):9-20, 2001.

Approximate Monotone Multilevel

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

Define:

- $\mathbf{c}_{j,\nu} = (\tilde{\mathbf{u}}_{j,\nu}, \tilde{\boldsymbol{\sigma}}_{j,\nu})$ correction at level j , patch ν
- $\mathbf{c}_{J,0} = \mathbf{x}_J^k$, $\mathbf{c}_{j,0} = \mathbf{0}$ for $j = J - 1, \dots, 1$
- $\mathbf{w}_{j,\nu} = \sum_{\mu=0}^{\nu} \mathbf{c}_{j,\mu}$

Compute a sequence of intermediate corrections $\mathbf{c}_{j,\nu} \in K_{j,\nu}(\mathbf{w}_{j,\nu-1})$ and $\mathbf{c}_1 \in K_1$:

$$\begin{aligned} \mathcal{J}(\mathbf{w}_{j,\nu-1} + \mathbf{c}_{j,\nu}) &\leq \mathcal{J}(\mathbf{w}_{j,\nu-1} + \mathbf{y}) \quad \forall \mathbf{y} \in K_{j,\nu} & j = J, \dots, 2, \quad \nu = 1, \dots, N_j \\ \mathcal{J}(\mathbf{c}_1) &\leq \mathcal{J}(\mathbf{y}) & \forall \mathbf{y} \in K_1 & j = 1 \end{aligned}$$

with the **coarse convex sets** K_j and the **approximate** local closed convex sets $K_{j,\nu}$:

$$\begin{aligned} K_{j,\nu}(\mathbf{w}_{j,\nu-1}) &= \{\mathbf{y} \in \text{span}\{\lambda_{j,\nu}\} : \mathbf{y} + \mathbf{w}_{j,\nu-1} \in K_j\} \\ K_1 &\subset K_2 \subset \dots \subset K_{J-1} \subset K_J \end{aligned}$$

Coarse Convex Sets and Constraints

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

Coarse Convex Sets:

$$\begin{aligned}
 K_j &= \left\{ \mathbf{x}_j = (\mathbf{u}_j, \boldsymbol{\sigma}_j) \in X_j : \mathbf{u}_j|_{\Gamma_D} = \mathbf{u}_D, \boldsymbol{\sigma}_j|_{\Gamma_N} = \mathbf{t}_N, \right. \\
 &\quad \left. \mathbf{u}_j \cdot \mathbf{n}_j|_{\Gamma_C} \leq g_{j,u_n}, \mathbf{n}^T(\boldsymbol{\sigma}_j \mathbf{n}) \leq g_{j,\sigma_n}, \mathbf{t}_j^T(\boldsymbol{\sigma}_j \mathbf{n}_j) = 0 \right\} \quad j = J \\
 K_j &= \left\{ \mathbf{x}_j = (\mathbf{u}_j, \boldsymbol{\sigma}_j) \in X_j : \mathbf{u}_j|_{\Gamma_D} = \mathbf{0}, \boldsymbol{\sigma}_j|_{\Gamma_N} = \mathbf{0}, \right. \\
 &\quad \left. \mathbf{u}_j \cdot \mathbf{n}_j|_{\Gamma_C} \leq g_{j,u_n}, \mathbf{n}^T(\boldsymbol{\sigma}_j \mathbf{n}) \leq g_{j,\sigma_n}, \mathbf{t}_j^T(\boldsymbol{\sigma}_j \mathbf{n}_j) = 0 \right\} \quad j = J-1, \dots, 1
 \end{aligned}$$

Coarse Constraints:

- $\tilde{\mathbf{u}}_{j,\nu}$ and $\tilde{\boldsymbol{\sigma}}_{j,\nu}$ are the components of the correction $\mathbf{c}_{j,\nu}$.

$$\begin{aligned}
 g_{j,u_n} &= \begin{cases} g & j = J \\ l_{j+1,u_n}^j \left(g_{j+1,u_n} - \sum_{\nu=1}^{N_{j+1}} [\tilde{\mathbf{u}}_{j+1,\nu}|_{\Gamma_C}]_n \right) & j = J-1, \dots, 1 \end{cases} \\
 g_{j,\sigma_n} &= \begin{cases} 0 & j = J \\ l_{j+1,\sigma_n}^j \left(g_{j+1,\sigma_n} - \sum_{\nu=1}^{N_{j+1}} [\tilde{\boldsymbol{\sigma}}_{j+1,\nu}|_{\Gamma_C}]_n \right) & j = J-1, \dots, 1 \end{cases}
 \end{aligned}$$

Non-Linear Projection Operators:

$l_{j+1,u_n}^j, l_{j+1,\sigma_n}^j$ chosen so that $K_1 \subset K_2 \subset \dots \subset K_{J-1} \subset K_J$.

Normal displacement Non-Linear Projection

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

$$v_H(\nu_{H,1}) \leq v_h(\nu_{H,1})$$

$$v_H(\nu_{H,2}) \leq v_h(\nu_{H,2})$$

$$\frac{1}{2}(v_H(\nu_{H,1}) + v_H(\nu_{H,2})) \leq v_h(\nu_h)$$

$$\forall \varepsilon_H \in \mathcal{E}_H \cap \Gamma_{C,H}$$

It is easy to see that, on e_H , the following values satisfy the three conditions above:

- a) $\begin{cases} \tilde{v}_H(\nu_{H,1}) = \min(v_h(\nu_{H,1}), \max(v_h(\nu_h), 2v_h(\nu_h) - v_h(\nu_{H,2}))) \\ \tilde{v}_H(\nu_{H,2}) = \min(v_h(\nu_{H,2}), \max(v_h(\nu_h), 2v_h(\nu_h) - v_h(\nu_{H,1}))) \end{cases} \quad \forall \varepsilon_H \in \mathcal{E}_H \cap \Gamma_C$
- b) $\begin{cases} \tilde{v}_H(\nu_{H,1}) = \min(v_h(\nu_{H,1}), v_h(\nu_h)) \\ \tilde{v}_H(\nu_{H,2}) = \min(v_h(\nu_{H,2}), v_h(\nu_h)) \end{cases} \quad \forall \varepsilon_H \in \mathcal{E}_H \cap \Gamma_C$
- c) $\begin{cases} \tilde{v}_H(\nu_{H,1}) = \min(v_h(\nu_{H,1}), v_h(\nu_h), v_h(\nu_{H,2})) \\ \tilde{v}_H(\nu_{H,2}) = \min(v_h(\nu_{H,1}), v_h(\nu_h), v_h(\nu_{H,2})) \end{cases} \quad \forall \varepsilon_H \in \mathcal{E}_H \cap \Gamma_C$

Pressure Non-Linear Projection

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

$$s_H(\phi_H) \leq s_h(\phi_h) \quad \forall \phi_h \in P_{\phi_H}^{\phi_h}$$

Thus:

$$s_H = I_{h,\sigma_n}^H s_h = \sum_{\phi_{H_i} \in T_H} [\lambda_{\Sigma_H, H_i}]_n s_H(\phi_{H_i}) \quad \text{with} \quad s_H(\phi_{H_i}) = \min_{\phi_h \in P_{\phi_{H_i}}^{\phi_h}} s_h(\phi_h)$$

Truncated Basis

Monotone
multilevel
for FOSLS
linear
elastic
contact

G.
Rovi \triangle ,
B.
Kober \circ ,
G.
Starke \circ ,
R.
Krause \triangle

$$[\tilde{\lambda}_{U_j, \nu}]_i = \begin{cases} [\lambda_{U_j, \nu}]_i & \nu \in \mathcal{N}_j \setminus \mathcal{N}_j^\bullet, \quad i = n, t \\ 0 & \nu \in \mathcal{N}_j^\bullet, \quad i = n \\ [\lambda_{U_j, \nu}]_i & \nu \in \mathcal{N}_j^\bullet, \quad i = t \end{cases}$$

$$[\tilde{\lambda}_{\Sigma_j, \phi}]_i = \begin{cases} [\lambda_{\Sigma_j, \phi}]_i & \phi \in \mathcal{F}_j \setminus \mathcal{F}_j^\bullet, \quad i = n, t \\ 0 & \phi \in \mathcal{F}_j^\bullet, \quad i = n \\ [\lambda_{\Sigma_j, \phi}]_i & \phi \in \mathcal{F}_j^\bullet, \quad i = t \end{cases}$$