

Discrete Mathematics

topic

week 1

Sets

week 2

Functions

week 3

Counting

week 4

Relations

week 5

Sequences

week 6

Modular Arithmetic

week 7

Asymptotic Notation

week 8

Orders

Jonathan Barrett

jonathan.barrett@cs.ox.ac.uk

Material by Andrew Ker

University of Oxford

Department of Computer Science



Discrete Mathematics



Jonathan Barrett

jonathan.barrett@cs.ox.ac.uk

Material by Andrew Ker

University of Oxford

Department of Computer Science

Chapter 3: Counting

Laws of Sum & Subtract

Law of Sum: Let P1 and P2 be properties of objects which are **exclusive**. The number of objects with **either** property is the number with property P1 **plus** the number with property P2.

This is true because, if A and B are disjoint, $|A \cup B| = |A| + |B|$

Example How many positive integers, less than a million, have an odd number of digits?

Laws of Sum & Subtract

Law of Sum: Let P1 and P2 be properties of objects which are **exclusive**. The number of objects with **either** property is the number with property P1 **plus** the number with property P2.

This is true because, if A and B are disjoint, $|A \cup B| = |A| + |B|$

Example How many positive integers, less than a million, have an odd number of digits?

Answer 90909

Laws of Sum & Subtract

Law of Sum: Let P1 and P2 be properties of objects which are **exclusive**.
The number of objects with **either** property is the number
with property P1 **plus** the number with property P2.

This is true because, if A and B are disjoint, $|A \cup B| = |A| + |B|$

Example How many positive integers, less than a million, have an odd
number of digits?

Answer 90909

Law of Subtract: Let P1 and P2 be properties, such that P1 is true at least
whenever P2 is. Then the number of objects with property
P1 **but not** P2 is the number with property P1 **subtract** the
number with property P2.

This is true because, if $B \subseteq A$ $|A \setminus B| = |A| - |B|$

Law of Product

Law of Product: If counting the number of ways of making a sequence of choices, and the choices are **independent** (the number available at each stage does not depend on the choices made previously), then the total number of ways of making the sequence of choices is the **product** of the number of choices at each stage.

This is true because $|A \times B| = |A| \cdot |B|$

Example How many 6-digit positive integers contain at least one digit 7?

Law of Product

Law of Product: If counting the number of ways of making a sequence of choices, and the choices are **independent** (the number available at each stage does not depend on the choices made previously), then the total number of ways of making the sequence of choices is the **product** of the number of choices at each stage.

This is true because $|A \times B| = |A| \cdot |B|$

Example How many 6-digit positive integers contain at least one digit 7?

Answer 427608

Factorial Function

Example In how many different ways can we arrange n distinguishable objects?

Factorial Function

Example In how many different ways can we arrange n distinguishable objects?

Answer $n! = 1 \cdot 2 \cdots (n - 1) \cdot n$

Arrangements of distinguishable objects are called **permutations**.

Factorial Function

Example In how many different ways can we arrange n distinguishable objects?

Answer $n! = 1 \cdot 2 \cdots (n - 1) \cdot n$

Arrangements of distinguishable objects are called **permutations**.

Note: The factorial function is extended to \mathbb{N} by setting $0! = 1$

In fact it can be generalized to a function on $[0, \infty)$ using

$$\Gamma(x) = \int_0^{\infty} t^{x-1} e^{-t} dt.$$

Double Counting

Sometimes it is easier to count a set twice over then divide by 2, or count m times over and divide by m .

Example Each of n teams in a league plays every other team once.
How many games must be played in total?

Double Counting

Sometimes it is easier to count a set twice over then divide by 2, or count m times over and divide by m .

Example Each of n teams in a league plays every other team once.
How many games must be played in total?

Answer
$$\frac{n(n-1)}{2}$$

Double Counting

Sometimes it is easier to count a set twice over then divide by 2, or count m times over and divide by m .

Example Each of n teams in a league plays every other team once. How many games must be played in total?

Answer
$$\frac{n(n-1)}{2}$$

Example In how many different ways can we select k out of n distinguishable objects, disregarding the order of selection?

Double Counting

Sometimes it is easier to count a set twice over then divide by 2, or count m times over and divide by m .

Example Each of n teams in a league plays every other team once. How many games must be played in total?

Answer
$$\frac{n(n-1)}{2}$$

Example In how many different ways can we select k out of n distinguishable objects, disregarding the order of selection?

Answer
$$\frac{n!}{k!(n-k)!}$$

Combinatorial Coefficients

Definition
$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

(the number of ways to select k out of n distinguishable objects).

Think of $\binom{\cdot}{\cdot}$ as a partial function $\mathbb{N}^2 \rightarrow \mathbb{N}_+$

Example How many ways to choose an unordered pair out of n objects?

Combinatorial Coefficients

Definition
$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

(the number of ways to select k out of n distinguishable objects).

Think of $\binom{\cdot}{\cdot}$ as a partial function $\mathbb{N}^2 \rightarrow \mathbb{N}_+$

Example How many ways to choose an unordered pair out of n objects?

Answer
$$\binom{n}{2} = \frac{n(n-1)}{2}$$

Combinatorial Coefficients

Definition

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

(the number of ways to select k out of n distinguishable objects).

Think of $\binom{\cdot}{\cdot}$ as a partial function $\mathbb{N}^2 \rightarrow \mathbb{N}_+$

Example How many ways to choose an unordered pair out of n objects?

Answer

$$\binom{n}{2} = \frac{n(n-1)}{2}$$

By counting the same set in different ways, we can often prove facts about combinatorial formulae, e.g.

Claim

$$2^n = \binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \cdots + \binom{n}{n}$$

Applications

Example How many ways can n (indistinguishable) objects be placed in k boxes?

Applications

Example How many ways can n (indistinguishable) objects be placed in k boxes?

Answer $\binom{n + k - 1}{k - 1}$

Applications

Example How many ways can n (indistinguishable) objects be placed in k boxes?

Answer $\binom{n + k - 1}{k - 1}$

Example How many ways can n (indistinguishable) objects be placed in k boxes so that each box receives at least one object?

Applications

Example How many ways can n (indistinguishable) objects be placed in k boxes?

Answer
$$\binom{n + k - 1}{k - 1}$$

Example How many ways can n (indistinguishable) objects be placed in k boxes so that each box receives at least one object?

Answer
$$\binom{n - 1}{k - 1}$$

Applications

Example How many ways can n (indistinguishable) objects be placed in k boxes?

Answer $\binom{n + k - 1}{k - 1}$

Example How many ways can n (indistinguishable) objects be placed in k boxes so that each box receives at least one object?

Answer $\binom{n - 1}{k - 1}$

Example How many positive integers less than a million have their digits sum to 9?

Applications

Example How many ways can n (indistinguishable) objects be placed in k boxes?

Answer $\binom{n + k - 1}{k - 1}$

Example How many ways can n (indistinguishable) objects be placed in k boxes so that each box receives at least one object?

Answer $\binom{n - 1}{k - 1}$

Example How many positive integers less than a million have their digits sum to 9?

Answer 2002

Inclusion-Exclusion Principle

Binary case: $|A \cup B| = |A| + |B| - |A \cap B|$

Example How many 6-digit positive integers contain at least one digit 7 and at least one digit 9?

Inclusion-Exclusion Principle

Binary case: $|A \cup B| = |A| + |B| - |A \cap B|$

Example How many 6-digit positive integers contain at least one digit 7
and at least one digit 9?

Answer 184592

Inclusion-Exclusion Principle

Binary case: $|A \cup B| = |A| + |B| - |A \cap B|$

General case:

$$\begin{aligned} |A_1 \cup A_2 \cup \cdots \cup A_n| = & (|A_1| + |A_2| + \cdots + |A_n|) \\ & - (|A_1 \cap A_2| + |A_1 \cap A_3| + \cdots + |A_{n-1} \cap A_n|) \\ & + (|A_1 \cap A_2 \cap A_3| + \cdots + |A_{n-2} \cap A_{n-1} \cap A_n|) \\ & - \cdots \\ & + (-1)^{n-1} |A_1 \cap A_2 \cap \cdots \cap A_n| \end{aligned}$$

Inclusion-Exclusion Principle

Binary case: $|A \cup B| = |A| + |B| - |A \cap B|$

General case: *intersections of all pairs $A_i \cap A_j, i \neq j$*

$$\begin{aligned}
 |A_1 \cup A_2 \cup \cdots \cup A_n| = & (|A_1| + |A_2| + \cdots + |A_n|) \\
 & - (|A_1 \cap A_2| + |A_1 \cap A_3| + \cdots + |A_{n-1} \cap A_n|) \\
 & + (|A_1 \cap A_2 \cap A_3| + \cdots + |A_{n-2} \cap A_{n-1} \cap A_n|) \\
 & \dots \\
 & - \dots \\
 & + (-1)^{n-1} |A_1 \cap A_2 \cap \cdots \cap A_n|
 \end{aligned}$$

intersections of all triples

...

intersection of all sets

Inclusion-Exclusion Principle

Binary case: $|A \cup B| = |A| + |B| - |A \cap B|$

General case: *intersections of all pairs $A_i \cap A_j, i \neq j$*

$$\begin{aligned}
 |A_1 \cup A_2 \cup \dots \cup A_n| = & (|A_1| + |A_2| + \dots + |A_n|) \\
 & - (|A_1 \cap A_2| + |A_1 \cap A_3| + \dots + |A_{n-1} \cap A_n|) \\
 & + (|A_1 \cap A_2 \cap A_3| + \dots + |A_{n-2} \cap A_{n-1} \cap A_n|) \\
 & \dots \\
 & - \dots \\
 & + (-1)^{n-1} |A_1 \cap A_2 \cap \dots \cap A_n|
 \end{aligned}$$

intersections of all triples

...

intersection of all sets

Example How many **derangements** are there of n objects? (A derangement is a permutation which leaves no object in its original place.)

Inclusion-Exclusion Principle

Binary case: $|A \cup B| = |A| + |B| - |A \cap B|$

General case: *intersections of all pairs $A_i \cap A_j, i \neq j$*

$$\begin{aligned}
 |A_1 \cup A_2 \cup \dots \cup A_n| = & (|A_1| + |A_2| + \dots + |A_n|) \\
 & - (|A_1 \cap A_2| + |A_1 \cap A_3| + \dots + |A_{n-1} \cap A_n|) \\
 & + (|A_1 \cap A_2 \cap A_3| + \dots + |A_{n-2} \cap A_{n-1} \cap A_n|) \\
 & \dots \\
 & - \dots \\
 & + (-1)^{n-1} |A_1 \cap A_2 \cap \dots \cap A_n|
 \end{aligned}$$

intersections of all triples → (points to the triple intersection term)

intersection of all sets → (points to the final term)

Example How many **derangements** are there of n objects? (A derangement is a permutation which leaves no object in its original place.)

Answer
$$d_n = \frac{n!}{2!} - \frac{n!}{3!} + \frac{n!}{4!} - \dots + (-1)^n \frac{n!}{n!}$$

Ceiling and Floor

The floor function “rounds down” real numbers to integers:

$$\lfloor - \rfloor : \mathbb{R} \rightarrow \mathbb{Z} \qquad \lfloor x \rfloor = \max\{n \in \mathbb{Z} \mid n \leq x\}$$

The ceiling function “rounds up” real numbers to integers:

$$\lceil - \rceil : \mathbb{R} \rightarrow \mathbb{Z} \qquad \lceil x \rceil = \min\{n \in \mathbb{Z} \mid n \geq x\}$$

Note that negative numbers are NOT rounded the other way!

Ceiling and Floor

The floor function “rounds down” real numbers to integers:

$$\lfloor - \rfloor : \mathbb{R} \rightarrow \mathbb{Z} \qquad \lfloor x \rfloor = \max\{n \in \mathbb{Z} \mid n \leq x\}$$

The ceiling function “rounds up” real numbers to integers:

$$\lceil - \rceil : \mathbb{R} \rightarrow \mathbb{Z} \qquad \lceil x \rceil = \min\{n \in \mathbb{Z} \mid n \geq x\}$$

Note that negative numbers are NOT rounded the other way!

Example How many positive integers, less than or equal to n , are divisible by k ?

Ceiling and Floor

The floor function “rounds down” real numbers to integers:

$$\lfloor - \rfloor : \mathbb{R} \rightarrow \mathbb{Z} \qquad \lfloor x \rfloor = \max\{n \in \mathbb{Z} \mid n \leq x\}$$

The ceiling function “rounds up” real numbers to integers:

$$\lceil - \rceil : \mathbb{R} \rightarrow \mathbb{Z} \qquad \lceil x \rceil = \min\{n \in \mathbb{Z} \mid n \geq x\}$$

Note that negative numbers are NOT rounded the other way!

Example How many positive integers, less than or equal to n , are divisible by k ?

Answer $\left\lfloor \frac{n}{k} \right\rfloor$

Ceiling and Floor

The floor function “rounds down” real numbers to integers:

$$\lfloor - \rfloor : \mathbb{R} \rightarrow \mathbb{Z} \qquad \lfloor x \rfloor = \max\{n \in \mathbb{Z} \mid n \leq x\}$$

The ceiling function “rounds up” real numbers to integers:

$$\lceil - \rceil : \mathbb{R} \rightarrow \mathbb{Z} \qquad \lceil x \rceil = \min\{n \in \mathbb{Z} \mid n \geq x\}$$

Note that negative numbers are NOT rounded the other way!

Example How many positive integers, less than or equal to n , are divisible by k ?

Answer $\left\lfloor \frac{n}{k} \right\rfloor$

Example How many positive integers, less than 1000, are divisible by one or more of 2, 3 or 5?

Ceiling and Floor

The floor function “rounds down” real numbers to integers:

$$\lfloor - \rfloor : \mathbb{R} \rightarrow \mathbb{Z} \qquad \lfloor x \rfloor = \max\{n \in \mathbb{Z} \mid n \leq x\}$$

The ceiling function “rounds up” real numbers to integers:

$$\lceil - \rceil : \mathbb{R} \rightarrow \mathbb{Z} \qquad \lceil x \rceil = \min\{n \in \mathbb{Z} \mid n \geq x\}$$

Note that negative numbers are NOT rounded the other way!

Example How many positive integers, less than or equal to n , are divisible by k ?

Answer $\left\lfloor \frac{n}{k} \right\rfloor$

Example How many positive integers, less than 1000, are divisible by one or more of 2, 3 or 5?

Answer 733

More Examples

Example How many permutations of $(1,2,3,4,5,6,7)$ do not include $(1,2)$, $(3,4,5)$, or $(6,7)$ as a (consecutive) subsequence?

More Examples

Example How many permutations of $(1,2,3,4,5,6,7)$ do not include $(1,2)$, $(3,4,5)$, or $(6,7)$ as a (consecutive) subsequence?

Answer 3642

More Examples

Example How many permutations of $(1,2,3,4,5,6,7)$ do not include $(1,2)$, $(3,4,5)$, or $(6,7)$ as a (consecutive) subsequence?

Answer 3642

Example In how many different ways can 5 red, 5 green, and 5 blue balls be distributed into 3 boxes? (Order in boxes is irrelevant).

More Examples

Example How many permutations of $(1,2,3,4,5,6,7)$ do not include $(1,2)$, $(3,4,5)$, or $(6,7)$ as a (consecutive) subsequence?

Answer 3642

Example In how many different ways can 5 red, 5 green, and 5 blue balls be distributed into 3 boxes? (Order in boxes is irrelevant).

Answer 9261

More Examples

Example How many permutations of $(1,2,3,4,5,6,7)$ do not include $(1,2)$, $(3,4,5)$, or $(6,7)$ as a (consecutive) subsequence?

Answer 3642

Example In how many different ways can 5 red, 5 green, and 5 blue balls be distributed into 3 boxes? (Order in boxes is irrelevant).

Answer 9261

Claim
$$n! = \binom{n}{0}d_0 + \binom{n}{1}d_1 + \binom{n}{2}d_2 + \cdots + \binom{n}{n}d_n$$

Diversion: Multinomial Coefficients

Example How many ways can n (distinguishable) objects be placed in g groups, with n_1 in group 1, n_2 in group 2, ..., n_g in group g ?

Diversion: Multinomial Coefficients

Example How many ways can n (distinguishable) objects be placed in g groups, with n_1 in group 1, n_2 in group 2, ..., n_g in group g ?

Answer As long as $n_1 + n_2 + \cdots + n_g = n$,

$$\frac{n!}{n_1!n_2!\cdots n_g!}$$

These are called the multinomial coefficients, and the symbol used is

$$\binom{n}{n_1 n_2 \cdots n_g} = \frac{n!}{n_1!n_2!\cdots n_g!}$$

Diversion: Multinomial Coefficients

Example How many ways can n (distinguishable) objects be placed in g groups, with n_1 in group 1, n_2 in group 2, ..., n_g in group g ?

Answer As long as $n_1 + n_2 + \cdots + n_g = n$,
$$\frac{n!}{n_1!n_2!\cdots n_g!}$$

These are called the multinomial coefficients, and the symbol used is

$$\binom{n}{n_1 n_2 \cdots n_g} = \frac{n!}{n_1!n_2!\cdots n_g!}$$

Example How many different bridge deals are there (allocation of 52 playing cards to four players, 13 each)?

Answer
$$\binom{52}{13 \ 13 \ 13 \ 13} = \frac{52!}{13!^4} = 53,644,737,765,488,792,839,237,440,000$$

Discrete Mathematics



Jonathan Barrett

jonathan.barrett@cs.ox.ac.uk

Material by Andrew Ker

University of Oxford

Department of Computer Science

End of Chapter 3