Time: 10 minutes

- 1. If $10 \le n \le 99$, find the integer n such that n divided by the sum of its digits is a minimum.
- Quadrilateral ABCD is cyclic (that is, it has a circumscribed circle). If AB = AD = 5, and CB = CD = 10, compute BD^2 .

Time: 10 minutes

- 3. In Priscilla's Lottery, you and Priscilla each pick 5 distinct positive integers between 1 and 50, inclusive. You win if your five numbers match the ones that are picked by Priscilla. In Quincy's Lottery, you and Quincy each pick 5 distinct positive integers between 1 and 50, inclusive and then randomly designate one as special. You win if your special integer is the same as Quincy's and your other four integers are the same as Quincy's other four. Let p be the probability that you win Priscilla's Lottery and let q be the probability that you win Quincy's Lottery. Compute p/q.
- 4. When ice melts to water, the volume of the water is 90% of the volume of the ice. A glass containing ice and water is filled to the top. When all the ice melts, the glass is 96% full. What per cent of the original volume was ice?

Time: 10 minutes

- 5. In equilateral triangle ABC with AB = 8, points P and Q are chosen on side \overline{AB} so that AP = BQ = 2. Similarly, points R and S are chosen on side \overline{BC} so that BR = CS = 2, and points T and U are chosen on side \overline{CA} so that CT = AU = 2. If the area of hexagon PQRSTU = H, find H^2 .
- 6. Given that $\sum_{k=1}^{89} \sin^2 k^\circ = m/n$, where *m* and *n* are relatively prime positive integers, find m+n.


Solutions for Contest #4

1. Represent the tens' digit and units' digit of a two-digit number by a and b, respectively, and denote the desired quotient by a. Then a0 = a1 = a2 = a3 Thus, a4 is minimum when a5 = a5. When a5 = a6, a7 = a8 = a9 = a9

Alternate solution: $\frac{10a+b}{a+b} = \frac{10a+10b}{a+b} - \frac{9b}{a+b} = 10 - \frac{9b}{a+b}$ is a minimum when a = 1 and $b = 9 \rightarrow n = 19$.

- 2. Notice that \overrightarrow{AC} is a symmetry line for the quadrilateral and its circumscribed circle. Therefore, \overrightarrow{AC} is a diameter of the circle. Use the Pythagorean Theorem in $\triangle ABC$ to find that $AC = 5\sqrt{5}$. The symmetry implies that $BD = 2 \cdot BE$, where point E is the foot of an altitude of $\triangle ABC$ from vertex B. Because \overline{BE} is a height of $\triangle ABC$, conclude that $5 \cdot 10 = 2|ABC| = 5\sqrt{5} \cdot BE$. Thus, $BE = 2\sqrt{5}$, and so $BD = 4\sqrt{5} = \sqrt{80}$. Thus, $BD^2 = 80$. Alternate Solution 1: Use the Law of Cosines in $\triangle BAD$ and in $\triangle BCD$. Note that $\cos \angle BCD = -\cos \angle BAD$. So, $BD^2 = 50 50 \cos \angle BAD$ and $BD^2 = 200 + 200 \cos \angle BAD \rightarrow BD^2 = 80$. Alternate Solution 2: Use Ptolemy's Theorem: $AC \cdot BD = AB \cdot CD + AD \cdot BC$ together with the fact that $AC = 5\sqrt{5}$. from the first solution.
- 3. Since $p = \frac{1}{\binom{50}{5}}$ and $= \frac{1}{5} \cdot \frac{1}{\binom{50}{5}}$, $\frac{p}{q} = 5$.
- 4. Let I and W be the fractions of the volume of the glass that were ice and water, respectively, originally. Then I + W = 1. Let I' and W' be the fractions of the volume of the glass that are ice and water, respectively, after the ice melts. Then I' + W' = 96. But I' = 0, and W' = W + .9I. Thus, W + .9I = .96. Subtract the last equation from the first to find that .1I = .04, so I = .4. Hence, 40% of the original volume was ice.
- 5. Triangles APU, BQR, and CST are equilateral, and the area of each of them is 1/16 that of $\triangle ABC$. Thus, H is 13/16 the area of $\triangle ABC$. Use the formula $K = (s^2 \sqrt{3})/4$ to find that the area of $\triangle ABC$ is $16\sqrt{3}$. Then $H = 13\sqrt{3}$, so $H^2 = 507$.

 Alternate solution: Draw \overline{RU} . Using $30^\circ 60^\circ 90^\circ$ triangles, we can calculate the height of trapezoid URQP to be $\sqrt{3}$ and the height of trapezoid TSRU to be $2\sqrt{3}$. The required area is the sum of the areas of the trapezoids: $\frac{1}{2} \cdot (4+6) \cdot \sqrt{3} + \frac{1}{2} \cdot (2+6) \cdot 2\sqrt{3} = 13\sqrt{3}$.



6. Let $S = ((\sin 1^{\circ})^{2} + (\sin 2^{\circ})^{2} + \dots + (\sin 43^{\circ})^{2} + (\sin 44^{\circ})^{2}) + (\sin 45^{\circ})^{2} + ((\sin 46^{\circ})^{2} + (\sin 47^{\circ})^{2} + \dots + (\sin 89^{\circ})^{2}). \rightarrow$ $S = ((\sin 1^{\circ})^{2} + (\sin 2^{\circ})^{2} + \dots + (\sin 43^{\circ})^{2} + (\sin 44^{\circ})^{2}) + (\sin 45^{\circ})^{2} + ((\cos 44^{\circ})^{2} + (\cos 43^{\circ})^{2} + \dots + (\cos 1^{\circ})^{2}). \quad \text{Re-arrange to get}$ $S = ((\sin 1^{\circ})^{2} + (\cos 1^{\circ})^{2}) + ((\sin 2^{\circ})^{2} + (\cos 2^{\circ})^{2}) + \dots + ((\sin 44^{\circ})^{2} + (\cos 44^{\circ})^{2}) + (\sin 45^{\circ})^{2} = 44 + \frac{1}{2} = \frac{89}{2}. \quad \text{So, } m + n = 91.$