

# Quantum Generalized Weyl Algebras

Phuong Ho

Miami University

# Motivation

To study subatomic particles using Quantum Mechanics, Heisenberg developed the Uncertainty Principle believing one cannot observe the position and momentum of a particle at the same time:

# Motivation

To study subatomic particles using Quantum Mechanics, Heisenberg developed the Uncertainty Principle believing one cannot observe the position and momentum of a particle at the same time:

$$PQ - QP = \frac{i\hbar}{2\pi} I$$

( $P$  and  $Q$  - square matrices of observable position and momentum,  $\hbar$  - Planck constant,  $I$ -identity matrix of appropriate size)

# Motivation

To study subatomic particles using Quantum Mechanics, Heisenberg developed the Uncertainty Principle believing one cannot observe the position and momentum of a particle at the same time:

$$PQ - QP = \frac{i\hbar}{2\pi} I$$

( $P$  and  $Q$  - square matrices of observable position and momentum,  $\hbar$  - Planck constant,  $I$ -identity matrix of appropriate size)

Rescaling gives the normalized version of the Fundamental Equation of Quantum Mechanics:

$$PQ - QP = I$$

Algebraically, the Fundamental Equation of QM may be thought by the operators of multiplication and differentiation by  $x$  on  $\mathbb{C}[x]$ .

Algebraically, the Fundamental Equation of QM may be thought by the operators of multiplication and differentiation by  $x$  on  $\mathbb{C}[x]$ . So,

$$\left(\frac{d}{dx}x\right)(p) - \left(x\frac{d}{dx}\right)(p) = p$$

for any  $p \in \mathbb{C}[x]$ .

Algebraically, the Fundamental Equation of QM may be thought by the operators of multiplication and differentiation by  $x$  on  $\mathbb{C}[x]$ . So,

$$\left(\frac{d}{dx}x\right)(p) - \left(x\frac{d}{dx}\right)(p) = p$$

for any  $p \in \mathbb{C}[x]$ .

We are interested in studying this equation and its generalizations from the viewpoint of algebra.

A  $\mathbb{C}$ -**algebra** is a  $\mathbb{C}$ -vector space with an associative multiplication operation that distributes over addition.



A  **$\mathbb{C}$ -algebra** is a  $\mathbb{C}$ -vector space with an associative multiplication operation that distributes over addition.

Examples:

A  **$\mathbb{C}$ -algebra** is a  $\mathbb{C}$ -vector space with an associative multiplication operation that distributes over addition.

Examples:

- Polynomial rings:  $\mathbb{C}[x_1, \dots, x_n]$

A  **$\mathbb{C}$ -algebra** is a  $\mathbb{C}$ -vector space with an associative multiplication operation that distributes over addition.

Examples:

- Polynomial rings:  $\mathbb{C}[x_1, \dots, x_n]$
- Matrix rings:  $M_n(\mathbb{C})$

A  **$\mathbb{C}$ -algebra** is a  $\mathbb{C}$ -vector space with an associative multiplication operation that distributes over addition.

Examples:

- Polynomial rings:  $\mathbb{C}[x_1, \dots, x_n]$
- Matrix rings:  $M_n(\mathbb{C})$
- The Weyl algebra  $A_1(\mathbb{C})$  is the  $\mathbb{C}$ -algebra generated by  $x$  and  $y$  with the relation

$$yx - xy = 1.$$

A  **$\mathbb{C}$ -algebra** is a  $\mathbb{C}$ -vector space with an associative multiplication operation that distributes over addition.

Examples:

- Polynomial rings:  $\mathbb{C}[x_1, \dots, x_n]$
- Matrix rings:  $M_n(\mathbb{C})$
- The Weyl algebra  $A_1(\mathbb{C})$  is the  $\mathbb{C}$ -algebra generated by  $x$  and  $y$  with the relation

$$yx - xy = 1.$$

An **automorphism** of a  $\mathbb{C}$ -algebra  $A$  is a bijective, structure-preserving function from  $A$  to itself. We denote the group of automorphisms of  $A$  by  $\text{Aut}(A)$ .

GWAs were defined by Bavula in highest generality.

# Generalized Weyl Algebra

GWAs were defined by Bavula in highest generality.

Let  $D$  be a commutative algebra,  $\sigma \in \text{Aut}(D)$ , and  $a \in D$ ,  $a \neq 0$ . The **generalized Weyl algebra** (GWA)  $D(\sigma, a)$  is generated over  $D$  by  $x$  and  $y$  subject to the relations

$$xd = \sigma(d)x, \quad yx = \sigma^{-1}(d)y, \quad yx = a, \quad xy = \sigma(a).$$

# Generalized Weyl Algebra

GWA's were defined by Bavula in highest generality.

Let  $D$  be a commutative algebra,  $\sigma \in \text{Aut}(D)$ , and  $a \in D$ ,  $a \neq 0$ . The **generalized Weyl algebra** (GWA)  $D(\sigma, a)$  is generated over  $D$  by  $x$  and  $y$  subject to the relations

$$xd = \sigma(d)x, \quad yx = \sigma^{-1}(d)y, \quad yx = a, \quad xy = \sigma(a).$$

This type of algebras has been studied by many authors before Bavula. Notably in the work of Hodges, Jordan, Joseph, Smith, and Stafford.



# Classical Weyl Algebra

Let  $R = D(\sigma, a)$  be a GWA. We say  $R$  is *classical* if  $D = \mathbb{C}[h]$  and  $\sigma(h) = h - 1$ .

# Classical Weyl Algebra

Let  $R = D(\sigma, a)$  be a GWA. We say  $R$  is *classical* if  $D = \mathbb{C}[h]$  and  $\sigma(h) = h - 1$ .

This type of algebra appears frequently in Lie Theory.

# Classical Weyl Algebra

Let  $R = D(\sigma, a)$  be a GWA. We say  $R$  is *classical* if  $D = \mathbb{C}[h]$  and  $\sigma(h) = h - 1$ .

This type of algebra appears frequently in Lie Theory.

For example, let  $D = \mathbb{C}[h]$ ,  $a = h$ ,  $\sigma(h) = h - 1$ .

$$xh = \sigma(h)x = (h - 1)x$$

$$yh = \sigma(h)^{-1}y = (h + 1)y$$

$$yx = a = h$$

$$xy = \sigma(a) = \sigma(h) = h - 1 = yx - 1$$

# Classical Weyl Algebra

Let  $R = D(\sigma, a)$  be a GWA. We say  $R$  is *classical* if  $D = \mathbb{C}[h]$  and  $\sigma(h) = h - 1$ .

This type of algebra appears frequently in Lie Theory.

For example, let  $D = \mathbb{C}[h]$ ,  $a = h$ ,  $\sigma(h) = h - 1$ .

$$xh = \sigma(h)x = (h - 1)x$$

$$yh = \sigma(h)^{-1}y = (h + 1)y$$

$$yx = a = h$$

$$xy = \sigma(a) = \sigma(h) = h - 1 = yx - 1$$

This is just the first Weyl algebra  $A_1(\mathbb{C})$

# Quantum Generalized Weyl Algebra

Let  $R = D(\sigma, a)$  be a GWA. We say  $R$  is *quantum* if  $D = \mathbb{C}[h]$  or  $\mathbb{C}[h^{\pm 1}]$  and  $\sigma(h) = qh$ ,  $q \in \mathbb{C} \setminus \{0, 1\}$ .

$$xh = qhx, yh = q^{-1}hy, yx = a, xy = a(qh)$$

# Quantum Generalized Weyl Algebra

Let  $R = D(\sigma, a)$  be a GWA. We say  $R$  is *quantum* if  $D = \mathbb{C}[h]$  or  $\mathbb{C}[h^{\pm 1}]$  and  $\sigma(h) = qh$ ,  $q \in \mathbb{C} \setminus \{0, 1\}$ .

$$xh = qhx, yh = q^{-1}hy, yx = a, xy = a(qh)$$

Example: let  $D = \mathbb{C}[h]$ ,  $a = h + 1$ .

$$xh = qhx, yh = q^{-1}hy, yx = a = h + 1, xy = qh + 1$$

# Quantum Generalized Weyl Algebra

Let  $R = D(\sigma, a)$  be a GWA. We say  $R$  is *quantum* if  $D = \mathbb{C}[h]$  or  $\mathbb{C}[h^{\pm 1}]$  and  $\sigma(h) = qh$ ,  $q \in \mathbb{C} \setminus \{0, 1\}$ .

$$xh = qhx, yh = q^{-1}hy, yx = a, xy = a(qh)$$

Example: let  $D = \mathbb{C}[h]$ ,  $a = h + 1$ .

$$xh = qhx, yh = q^{-1}hy, yx = a = h + 1, xy = qh + 1$$

There are many important families of algebras that may be GWAs:

# Quantum Generalized Weyl Algebra

Let  $R = D(\sigma, a)$  be a GWA. We say  $R$  is *quantum* if  $D = \mathbb{C}[h]$  or  $\mathbb{C}[h^{\pm 1}]$  and  $\sigma(h) = qh$ ,  $q \in \mathbb{C} \setminus \{0, 1\}$ .

$$xh = qhx, yh = q^{-1}hy, yx = a, xy = a(qh)$$

Example: let  $D = \mathbb{C}[h]$ ,  $a = h + 1$ .

$$xh = qhx, yh = q^{-1}hy, yx = a = h + 1, xy = qh + 1$$

There are many important families of algebras that may be GWAs: quantum Weyl algebras and primitive quotients of  $U_q(\mathfrak{sl}_2)$ .



# Shephard, Todd, Chevalley Theorem

If  $A$  is a  $\mathbb{C}$ -algebra and  $G$  a subgroup of  $\text{Aut}(A)$ , then  $A^G = \{a \in A : g(a) = a \text{ for all } g \in G\}$ .

# Shephard, Todd, Chevalley Theorem

If  $A$  is a  $\mathbb{C}$ -algebra and  $G$  a subgroup of  $\text{Aut}(A)$ , then  $A^G = \{a \in A : g(a) = a \text{ for all } g \in G\}$ .

## Theorem (Shephard, Todd, Chevalley)

*Let  $A = \mathbb{C}[x_1, \dots, x_n]$  and let  $G$  be a finite group of linear automorphisms of  $A$ . Then the fixed ring  $A^G$  is again a polynomial ring if and only if  $G$  is generated by reflections.*

Example:  $A = \mathbb{C}[x, y]$  and  $g \in \text{Aut } A$  given by  $g(x) = y$  and  $g(y) = x$ . Then  $A^G = \mathbb{C}[x + y, xy]$ .

# Shephard, Todd, Chevalley Theorem

If  $A$  is a  $\mathbb{C}$ -algebra and  $G$  a subgroup of  $\text{Aut}(A)$ , then  $A^G = \{a \in A : g(a) = a \text{ for all } g \in G\}$ .

## Theorem (Shephard, Todd, Chevalley)

*Let  $A = \mathbb{C}[x_1, \dots, x_n]$  and let  $G$  be a finite group of linear automorphisms of  $A$ . Then the fixed ring  $A^G$  is again a polynomial ring if and only if  $G$  is generated by reflections.*

Example:  $A = \mathbb{C}[x, y]$  and  $g \in \text{Aut } A$  given by  $g(x) = y$  and  $g(y) = x$ . Then  $A^G = \mathbb{C}[x + y, xy]$ .

We want to work towards developing a STC-like theorem for (quantum) GWAs. Others have contributed to this previously, including Jordan-Wells, Kirkman-Kuzmanovich, and Gaddis-Won.

# A More Generalized Version of Jordan-Wells Theorem

Jordan and Wells studied fixed rings of GWAs by automorphisms that fix the base ring. The following result generalizes their theorem.

# A More Generalized Version of Jordan-Wells Theorem

Jordan and Wells studied fixed rings of GWAs by automorphisms that fix the base ring. The following result generalizes their theorem.

## Theorem (-, Gaddis)

*Let  $D$  be an integral domain, let  $R = D(\sigma, a)$  be a GWA, and let  $\phi \in \text{Aut}(R)$  with  $|\phi| < \infty$ . Suppose  $\phi|_D$  restricts to an automorphism of  $D$ ,  $\phi(x) = \mu^{-1}x$ , and  $\phi(y) = \mu y$  for  $\mu \in \mathbb{C}^\times$ . Set  $n = |\phi|_D|$  and  $m = |\mu|$ . If  $\gcd(n, m) = 1$ , then  $R^{\langle \phi \rangle} = D^{\langle \phi \rangle}(\sigma^m, A)$  with  $A = \prod_{i=0}^{m-1} \sigma^{-i}(a)$ .*

# Example

Let  $R = \mathbb{C}[h^{\pm 1}](\sigma, a)$  be a quantum GWA with  $a = (h^2 - 1)(h^2 - 4)$  and  $q = 1/2$ . Let  $\eta \in \text{Aut}(R)$  be given by

$$\eta(h) = \gamma h, \quad \eta(x) = \mu^{-1}x, \quad \eta(y) = \mu y$$

with  $n = 2$  and  $m = 3$ .

# Example

Let  $R = \mathbb{C}[h^{\pm 1}](\sigma, a)$  be a quantum GWA with  $a = (h^2 - 1)(h^2 - 4)$  and  $q = 1/2$ . Let  $\eta \in \text{Aut}(R)$  be given by

$$\eta(h) = \gamma h, \quad \eta(x) = \mu^{-1}x, \quad \eta(y) = \mu y$$

with  $n = 2$  and  $m = 3$ .

Then  $R^{(\eta)}$  is generated by  $X = x^3$ ,  $Y = y^3$ , and  $H = h^2$ . The defining polynomial is

$$\begin{aligned} A(H) &= (H - 1)(H - 4)(4H - 1)(4H - 4)(16H - 4)(16H - 1) \\ &= 4(H - 1)^2(H - 4)(4H - 1)(16H - 1)(16H - 4) \\ &= 4^6(H - 1)^2(H - 4) \left(H - \frac{1}{4}\right)^2 \left(H - \frac{1}{16}\right). \end{aligned}$$

Thank You for Your Listening!