Ph.D. IN MATHEMATICAL SCIENCES

[Field of Study Code : MATH (897)]

Time Allowed : 3 hours

Maximum Marks : 100

INSTRUCTIONS FOR CANDIDATES

- (i) All questions are compulsory.
- (ii) There are 10 questions in Section—A and 8 questions in each of Sections B and C.
- (iii) Answers must be written in the space provided after each question. Answers that are written elsewhere will not be evaluated.
- (iv) A question in Section—A may have more than one true option. Your answer will be considered correct provided it consists only of all the true options. A correct answer will be awarded +2 marks and an incorrect/incomplete answer or an unattempted question will get 0 mark.
- (v) Each question in Section—B carries 4 marks; each question in Section—C carries 6 marks. Answers to all the questions in Sections B and C must be justified with mathematical reasoning.
- (vi) In the following, N, Z, Q, R, C denote, respectively, the set of natural numbers, the set of integers, the set of rational numbers, the set of real numbers and the set of complex numbers.
- (vii) Cⁿ, Rⁿ and its subsets are assumed to have the usual topology arising from the usual Euclidean norm unless mentioned otherwise.
- (viii) For sets A and B, let $A \setminus B := \{x \in A \mid x \notin B\}$. For $a, b \in \mathbb{R}$, let $[a, b] := \{x \in \mathbb{R} \mid a \le x \le b\}$, $(a, b] := [a, b] \setminus \{a\}$ and $(a, b) := [a, b] \setminus \{a, b\}$.
- (ix) Extra pages are attached at the end of the question paper for Rough Work.
- (x) Use of calculator, smart phone, mobile phone or any other digital device is strictly not permitted.

ENTRANCE EXAMINATION, 2018 Ph.D. IN MATHEMATICAL SCIENCES

SUBJECT
FIELD OF STUDY CODE
NAME OF THE CANDIDATE
REGISTRATION NO.
CENTRE OF EXAMINATION

DATE
(Signature of Candidate)
(Signature of Invigilator)
(Signature and Seal of Presiding Officer)

Vor Official Use Only Not to te third in Vertilationalisate Orabita Table for Backlor-AV

Not to be filled in	al Use Only n by the candidate dding Table	egil potteen Amarded
Total of Section—A	61	8.1
Total of Section—B	62	EA.
Total of Section—C	d	āA
Grand Total	a	YA
GEA .	200	PA
Section of Section 1989		
	Takes Named at	

For Official Use Only

Not to be filled in by the candidate

Grading Table for Section-A

Question Nos.	Marks Awarded	Question Nos.	Marks Awarded
A1		A A2	e islat
А3		E A4	о [авуТ
A5		O.A6	o istol
A7		A8	п0
A9		A10	
		Total (Out of 20)	

Not to be filled in by the candidate

Grading Table for Sections B and C

Question Nos.	Marks Awarded	Question Nos.	Marks Awarded
B1		C1	Name of
B2	and display	C2	
B3	TO STATE OF	СЗ	
B4	100 Pro	C4	Swar Service
B5	- 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1	C5	yette. Does is
В6	- Maria Maria	C6	11
В7	- Ti = 100	C7	spm Certail
	- cula	C8	
Total (Out of 32)	1	Total (Out of 48)	

Each question carries 2 marks.

What is the number of zeroes at the end of (2017)!? A1.

Answer: The same of the stray range at or aldeportement of the (a)

Five friends went to A to Z ice-cream shop that was selling 26 flavours of ice cream to A2. have one scoop of ice cream each. One friend asked everyone which flavour of ice-cream they wanted and went to the counter to place the order. How many distinct orders of ice creams can be placed?

Answer : net E a R be a Lebesque mogamente set and let m dennie the Lebesque : rawanA

- Let A, B, C and D be finite sets and $\varphi: A \to B$, $\psi: B \to D$, $\theta: A \to C$ and $\eta: C \to D$ be maps A3. such that $\psi \circ \varphi = \eta \circ \theta$. Consider the following statements:
 - φ is surjective
 - B and D have the same number of elements O and model to a (Alm II (o)
 - θ is injective
 - (d) A has at least as many elements as in Image (ψ)

Which of the above statements are correct?

Answer:

Is Z/899Z an integral domain? Provide your answer in Yes or No below.

Answer:

Suppose G is a group in which every proper subgroup is cyclic. Does this imply that G is A5. cyclic? Provide your answer in Yes or No below. DIA &

Lowen

Answer:

- Let $S \subset \mathbb{R}^2$ be defined by $S := \{(t\cos(t), t\sin(t)) \mid 0 < t \le 2\pi\}$. Consider the following A6. statements:
 - (a) S is bounded
 - (b) S contains all its limit points
 - (c) S is connected
 - (d) S is not path-connected = 2 is iquoxa J = |x| this s lie to anguards (h)

Which of the above statements are correct?

Answer:

A7.	Let X be a normed linear	space.	Which of the following	statements are always true?
-----	----------------------------	--------	------------------------	-----------------------------

- (a) X is locally compact with respect to the norm topology
- (b) Every linear map $\varphi: X \to \mathbb{C}$ is continuous
- (c) X is homeomorphic to its open unit ball $B := \{x \in X \mid ||x|| < 1\}$
- (d) X is not homeomorphic to its closed unit ball $\overline{B} := \{x \in X \mid ||x|| \le 1\}$

A8. Let $E \subset \mathbb{R}$ be a Lebesgue measurable set and let m denote the Lebesgue measure. Which of the following statements are true?

- (a) If $m(E) = \infty$, then E is dense in \mathbb{R}
- (b) If m(E) = 0, then E has empty interior
- (c) If m(E) = 0, then $E \subset \mathbb{Q}$ extremals in reduction such soft evant \mathbb{Q} broken
- (d) If $m(E) = \infty$, then $E \cap \mathbb{Q}^c \neq \emptyset$

Answer:

A9. What is the value of
$$\lim_{x\to 0+} \left(1+\frac{1}{x}\right)^x$$
?

Answer:

A10. The series

$$\sum_{n=1}^{\infty} \frac{z^n}{n}$$

$$|n| \le 1 > 0 | |n| = 1$$

$$|n| \ge 1 > 0 | |n| = 1$$

$$|n| \ge 1 > 0 | |n| = 1$$

- (a) converges for all z with |z| = 1, except at z = 1
- (b) converges for all z with |z| = 1, except at z = 1, -1
- (c) diverges for all z with |z| = 1
- (d) diverges for all z with |z| = 1, except at z = -1

Pick the correct options.

Answer:

Stress in Gas a group in which every proper subgroup is cyclic. Dues this unply that G is a

weight Provide your answer in Yes or its below

struog timil ati Us amateov 2

Which of the above statements are correct?

TO BE TO SELLING THE PROPERTY OF THE PARTY O

Each question carries 4 marks.

B1. Show that $\sqrt[3]{5\sqrt{2}+7} + \sqrt[3]{5\sqrt{2}-7} = 2\sqrt{2}$. (*Hint*: First show that the given number is a root of the polynomial $x^3 - 3x - 10\sqrt{2}$.)

B2. For $x \in (0, \infty)$, define $f(x) := x^{x+1}$ and $g(x) := (x+1)^x$. Determine the values of x for which f(x) > g(x).

.

B3. Let \mathcal{T} be a collection of subsets of \mathbb{Z} containing \emptyset , \mathbb{Z} and non-empty subsets U of \mathbb{Z} which are unions of sets of the form

$$S(a,b):=\{an+b\mid n\in\mathbb{Z}\}$$

for $a, b \in \mathbb{Z}$ and $a \neq 0$. Show that T defines a topology on \mathbb{Z} . Prove or disprove that T is Hausdorff.

/283

B4. Let X and Y be sets, $\varphi: X \to \mathbb{N}$ and $\psi: \mathbb{N} \to Y$ be maps such that their composite $\psi \circ \varphi: X \to Y$ is surjective. Prove or disprove the following assertion: X and Y are both countable.

1253

B5. Let $f: \mathbb{R}^3 \to \mathbb{R}^3$ be a map such that ||f(x) - f(y)|| = ||x - y|| for all $x, y \in \mathbb{R}^3$. Prove or disprove the following assertion:

f maps every closed ball homeomorphically onto its image.

B6. Let X and Y be Banach spaces and suppose that X is finite dimensional. Let $T: X \to Y$ be a linear map. Prove or disprove the following assertion:

If B is a bounded set in X, then $\overline{T(B)}$ is compact in Y.

B7. Let $E \subseteq [0, 1]$ be Lebesgue measurable with m(E) = 1 and $f: [0, 1] \to \mathbb{R}$ be a continuous function. Prove or disprove the following assertion:

If f(E) is a singleton, then so is f([0, 1]).

B8

- B8. Let $f: \mathbb{C} \to \mathbb{C}$ be an entire function such that $|f(z)| < |e^z|$ for all z and $f(0) = \frac{1}{2}$. Find the value of $f(\pi)$.
- C1. Let S be a finite subset of R² of the form A × E, where A, R : R Suppose W is his open as in R² that contains S. Show that there exist open as a land V in R show that S C V × V C W.

1253

so he will discharge the soft of the SECTION—C is not stone we set the to have the

Each question carries 6 marks.

C1. Let S be a finite subset of \mathbb{R}^2 of the form $A \times B$, where A, $B \subset \mathbb{R}$. Suppose W is an open set in \mathbb{R}^2 that contains S. Show that there exist open sets U and V in \mathbb{R} such that $S \subset U \times V \subseteq W$.

- C2. Let $f: \mathbb{R} \to [0, \infty)$ be a measurable function such that $0 < \int f \, dm < \infty$. Consider $\mu: \mathcal{L} \to [0, \infty]$ given by $\mu(E): = \int_E f \, dm$, where m is the Lebesgue measure on the σ -algebra \mathcal{L} of Lebesgue measurable sets in \mathbb{R} .
 - (a) Show that μ is a finite measure on \mathcal{L} , i.e., $\mu(\mathbb{R}) < \infty$ and for any countable mutually disjoint collection $\{E_n\}$ in \mathcal{L} , $\mu(\cup_n E_n) = \Sigma_n \mu(E_n)$.
 - (b) If $E \in \mathcal{L}$ has m(E) = 0, then show that $\mu(E) = 0$.

C3. Let X = C([0, 1]) be the complex Banach space consisting of complex-valued continuous functions on [0, 1] with the sup norm and Y be the subspace consisting of polynomials of degree less than or equal to 5. Show that for each coset $x + Y \in X/Y$, there exists an element $z \in x + Y$ such that $||z|| = \inf\{||x + y|| : y \in Y\}$.

- C4. Consider the polynomial $f(x) = x^4 + 1$ over the field $\mathbb{Z}/5\mathbb{Z}$.
 - (a) Is it irreducible? Justify.
 - (b) Find the splitting field of the polynomial of f over $\mathbb{Z}/5\mathbb{Z}$.

C5. Consider the curve E given by the equation $y^2 = x^3 + Ax + B$, where A and B are integers. Let P = (a, b) and Q = (c, d) be points on E such that $a, b, c, d \in Q$, with $a \ne c$. Let L be the line joining P and Q. Show that E and L intersect in some point R = (e, f), where $e, f \in Q$.

- C6. (a) Exhibit a 4×4 matrix over C whose eigenvalues over R are ± 1 and whose eigenvalues over C are ± 1 and $\pm i$.
 - (b) Exhibit three non-conjugate matrices over $\mathbb{Z}/7\mathbb{Z}$ of size 3×3 for which -2 is the only eigenvalue.

- C6. (a) Exhibit a 4×4 matrix over C whose eigenvalues over R are ± 1 and whose eigenvalues over C are ± 1 and $\pm i$.
 - (b) Exhibit three non-conjugate matrices over $\mathbb{Z}/7\mathbb{Z}$ of size 3×3 for which -2 is the only eigenvalue.

C7. Find and classify the extreme values, if any, of the function $f: \mathbb{R}^2 \to \mathbb{R}$ given by $f(x, y) = y^2 - x^3$.

C8. Find all the poles of the function $f(z) = \frac{z}{\sin z - 1}$ and compute $\int_C f(z) dz$, where $C: [0, 2\pi] \to \mathbb{C}$ is the curve defined as $C(\theta) = 2e^{-i\theta}$.

32