

## General Aptitude (GA)

**Q.1 – Q.5 Multiple Choice Question (MCQ), carry ONE mark each (for each wrong answer: – 1/3).**

- Q.1** (a) Arun and Aparna are here.  
(b) Arun and Aparna is here.  
(c) Arun's families is here.  
(d) Arun's family is here.

Which of the above sentences are grammatically CORRECT?

- (a) (i) and (ii)   (b) (i) and (iv)   (c) (ii) and (iv)   (d) (iii) and (iv)



Figure 1: text in graph

**Q.2** The mirror image of the above text about the X-axis is

- (a) 
- (b) 
- (c) 


 BHARGAVA'S

(d)

(GATE PH 2021)

**Q.3** Two identical cube shaped dice each with faces numbered 1 to 6 are rolled simultaneously. The probability that an even number is rolled out on each dice is:

- (a)  $\frac{1}{36}$                       (b)  $\frac{1}{12}$                       (c)  $\frac{1}{8}$                       (d)  $\frac{1}{4}$

(GATE PH 2021)

**Q.4**  $\oplus$  and  $\odot$  are two operators on numbers  $p$  and  $q$  such that  $p \odot q = p - q$ , and  $p \oplus q = p \times q$ . Then,  $(9 \odot (6 \oplus 7)) \odot (7 \oplus (6 \odot 5)) =$

- (a) 40                      (b) -26                      (c) -33                      (d) -40

(GATE PH 2021)

**Q.5** Four persons P, Q, R and S are to be seated in a row. R should not be seated at the second position from the left end of the row. The number of distinct seating arrangements possible is:

- (a) 6                      (b) 9                      (c) 18                      (d) 24

(GATE PH 2021)

**Q. 6 – Q. 10 Multiple Choice Question (MCQ), carry TWO marks each (for each wrong answer: – 2/3).**

**Q.6** On a planar field, you travelled 3 units East from a point O. Next you travelled 4 units South to arrive at point P. Then you travelled from P in the North-East direction such that you arrive at a point that is 6 units East of point O. Next, you travelled in the North-West direction, so that you arrive at point Q that is 8 units North of point P. The distance of point Q to point O, in the same units, should be \_\_\_\_\_.

- (a) 3                      (b) 4                      (c) 5                      (d) 6

(GATE PH 2021)

- Q.7** The author said, “Musicians rehearse before their concerts. Actors rehearse their roles before the opening of a new play. On the other hand, I find it strange that many public speakers think they can just walk on to the stage and start speaking. In my opinion, it is no less important for public speakers to rehearse their talks.”

Based on the above passage, which one of the following is TRUE?

- (a) The author is of the opinion that rehearsing is important for musicians, actors and public speakers.
- (b) The author is of the opinion that rehearsing is less important for public speakers than for musicians and actors.
- (c) The author is of the opinion that rehearsing is more important only for musicians than public speakers.
- (d) The author is of the opinion that rehearsing is more important for actors than musicians.

(GATE PH 2021)

- Q.8** (a) Some football players play cricket.  
(b) All cricket players play hockey.

Among the options given below, the statement that logically follows from the two statements 1 and 2 above, is:

- (a) No football players play hockey.
- (b) Some football players play hockey.
- (c) All football players play hockey.
- (d) All hockey players play football.

(GATE PH 2021)




Figure 2: square PQRS

**Q.9** In the figure shown above, PQRS is a square. The shaded portion is formed by the intersection of sectors of circles with radius equal to the side of the square and centers at S and Q. The probability that any point picked randomly within the square falls in the shaded area is \_\_\_\_\_.

- (a)  $4 - \frac{\pi}{2}$       (b)  $\frac{1}{2}$       (c)  $\frac{\pi}{2} - 1$       (d)  $\frac{\pi}{4}$

(GATE PH 2021)

**Q.10** In an equilateral triangle PQR, side PQ is divided into four equal parts, side QR is divided into six equal parts and side PR is divided into eight equal parts. The length of each subdivided part in cm is an integer. The minimum area of the triangle PQR possible, in  $\text{cm}^2$ , is


- (a) 18      (b) 24      (c)  $48\sqrt{3}$       (d)  $144\sqrt{3}$


(GATE PH 2021)

## Physics (PH)

**Q.1 – Q.9 Multiple Choice Question (MCQ), carry ONE mark each (for each wrong answer: – 1/3).**

**Q.1** Choose the graph that best describes the variation of dielectric constant ( $\epsilon_r$ ) with temperature ( $T$ ) in a ferroelectric material. ( $T_C$  is the Curie temperature)





(GATE PH 2021)

**Q.2** A matter wave is represented by the wave function

$$\Psi(x, y, z, t) = Ae^{i(4x+3y+5z-10\pi t)}$$

where A is a constant. The unit vector representing the direction of propagation of this matter wave is

- i.  $\frac{4}{5\sqrt{2}}\hat{x} + \frac{3}{5\sqrt{2}}\hat{y} + \frac{1}{\sqrt{2}}\hat{z}$
- ii.  $\frac{3}{5\sqrt{2}}\hat{x} + \frac{4}{5\sqrt{2}}\hat{y} + \frac{1}{5\sqrt{2}}\hat{z}$
- iii.  $\frac{1}{5\sqrt{2}}\hat{x} + \frac{3}{5\sqrt{2}}\hat{y} + \frac{1}{\sqrt{2}}\hat{z}$
- iv.  $\frac{1}{\sqrt{2}}\hat{x} + \frac{4}{5\sqrt{2}}\hat{y} + \frac{3}{5\sqrt{2}}\hat{z}$

(GATE PH 2021)

**Q.3** As shown in the figure, X-ray diffraction pattern is obtained from a diatomic chain of atoms P and Q. The diffraction condition is

given by  $a \cos \theta = n\lambda$ , where  $n$  is the order of the diffraction peak. Here,  $a$  is the lattice constant and  $\lambda$  is the wavelength of the X-rays. Assume that atomic form factors and resolution of the instrument do not depend on  $\theta$ . Then, the intensity of the diffraction peaks is




Figure 3: X-ray diffraction pattern

- i. lower for even values of  $n$ , when compared to odd values of  $n$
- ii. lower for odd values of  $n$ , when compared to even values of  $n$
- iii. zero for odd values of  $n$
- iv. zero for even values of  $n$

(GATE PH 2021)

**Q.4** As shown in the figure, two metal-semiconductor junctions are formed between an n-type semiconductor  $S$  and metal  $M$ . The work functions of  $S$  and  $M$  are  $\phi_S$  and  $\phi_M$ , respectively with  $\phi_M > \phi_S$ .






Figure 4: two metal-semiconductor junctions formed between an n-type semiconductor  $S$  and metal  $M$ .

The  $I - V$  characteristics (on linear scale) of the junctions is best represented by



i.



(GATE PH 2021)

**Q.5** Consider a tiny current loop driven by a sinusoidal alternating current. If the surface integral of its time-averaged Poynting vector is constant, then the magnitude of the time-averaged magnetic field intensity, at any arbitrary position,  $\vec{r}$ , is proportional to

- i.  $\frac{1}{r^3}$       ii.  $\frac{1}{r^2}$       iii.  $\frac{1}{r}$       iv.  $r$

(GATE PH 2021)

**Q.6** Consider a solenoid of length  $L$  and radius  $R$ , where  $R \ll L$ . A steady-current flows through the solenoid. The magnetic field is uniform inside the solenoid and zero outside.





Figure 5: a solenoid of length  $L$  and radius  $R$



Among the given options, choose the one that best represents the variation in the magnitude of the vector potential,  $(0, A_\phi, 0)$  at  $z = L/2$ , as a function of the radial distance ( $r$ ) in cylindrical coordinates.

**Useful information:** The curl of a vector  $\mathbf{F}$ , in cylindrical coordinates is

$$\nabla \times \mathbf{F}(r, \phi, z) = \hat{r} \left[ \frac{1}{r} \frac{\partial F_z}{\partial \phi} - \frac{\partial F_\phi}{\partial z} \right] + \hat{\phi} \left[ \frac{\partial F_r}{\partial z} - \frac{\partial F_z}{\partial r} \right] + \hat{z} \frac{1}{r} \left[ \frac{\partial(r F_\phi)}{\partial r} - \frac{\partial F_r}{\partial \phi} \right]$$







(GATE PH 2021)

**Q.7** Assume that  $^{13}\text{N}$  ( $Z = 7$ ) undergoes first forbidden  $\beta^+$  decay from its ground state with spin-parity  $J_i^\pi$ , to a final state  $J_f^\pi$ . The possible values for  $J_i^\pi$  and  $J_f^\pi$ , respectively, are

- i.  $\frac{1}{2}^-, \frac{5}{2}^+$       ii.  $\frac{1}{2}^+, \frac{5}{2}^+$       iii.  $\frac{1}{2}^-, \frac{1}{2}^-$       iv.  $\frac{1}{2}^+, \frac{1}{2}^-$

(GATE PH 2021)

**Q.8** In an experiment, it is seen that an electric-dipole (E1) transition can connect an initial nuclear state of spin-parity  $J_i^\pi = 2^+$  to a final state  $J_f^\pi$ . All possible values of  $J_f^\pi$  are

- i.  $1^+, 2^+$       ii.  $1^+, 2^+, 3^+$       iii.  $1^-, 2^-$       iv.  $1^-, 2^-, 3^-$

(GATE PH 2021)

**Q.9** Choose the correct statement from the following.

- Silicon is a direct band gap semiconductor.
- Conductivity of metals decreases with increase in temperature.
- Conductivity of semiconductors decreases with increase in temperature.

iv. Gallium Arsenide is an indirect band gap semiconductor.

(GATE PH 2021)

**Q.10 – Q.16 Multiple Select Question (MSQ), carry ONE mark each (no negative marks).**

**Q.10** A two-dimensional square lattice has lattice constant  $a$ .  $k$  represents the wavevector in reciprocal space. The coordinates  $(k_x, k_y)$  of reciprocal space where band gap(s) can occur, are

- i.  $(0, 0)$       ii.  $(\pm \frac{\pi}{a}, \pm \frac{\pi}{a})$       iii.  $(\pm \frac{\pi}{a}, \pm \frac{\pi}{1.3a})$       iv.  $(\pm \frac{\pi}{3a}, \pm \frac{\pi}{a})$

(GATE PH 2021)

**Q.11** As shown in the figure, an electromagnetic wave with intensity  $I_I$  is incident at the interface of two media having refractive indices  $n_1 = 1$  and  $n_2 = \sqrt{3}$ . The wave is reflected with intensity  $I_R$  and transmitted with intensity  $I_T$ . Permeability of each medium is the same. (Reflection coefficient  $R = I_R/I_I$  and Transmission coefficient  $T = I_T/I_I$ ).




Figure 6: electromagnetic wave with intensity  $I_I$  is incident at the interface of two media

Choose the correct statement(s).

- $R = 0$  if  $\theta_I = 0^\circ$  and polarization of incident light is parallel to the plane of incidence.
- $T = 1$  if  $\theta_I = 60^\circ$  and polarization of incident light is parallel to the plane of incidence.
- $R = 0$  if  $\theta_I = 60^\circ$  and polarization of incident light is perpendicular to the plane of incidence.
- $T = 1$  if  $\theta_I = 60^\circ$  and polarization of incident light is perpendicular to the plane of incidence.

(GATE PH 2021)

**Q.12** A material is placed in a magnetic field intensity  $H$ . As a result, bound current density  $J_b$  is induced and magnetization of the material is  $M$ . The magnetic flux density is  $B$ . Choose the correct option(s) valid at the *surface* of the material.

- i.  $\nabla \cdot M = 0$    ii.  $\nabla \cdot B = 0$    iii.  $\nabla \cdot H = 0$    iv.  $\nabla \cdot J_b = 0$

(GATE PH 2021)

**Q.13** For a finite system of Fermions where the density of states increases with energy, the chemical potential

- i. decreases with temperature  
 ii. increases with temperature  
 iii. does not vary with temperature  
 iv. corresponds to the energy where the occupation probability is 0.5

(GATE PH 2021)

**Q.14** Among the term symbols  $^4S_1$ ,  $^2D_{7/2}$ ,  $^3S_1$  and  $^2D_{5/2}$  choose the option(s) possible in the LS coupling notation.

- i.  $^4S_1$    ii.  $^2D_{7/2}$    iii.  $^3S_1$    iv.  $^2D_{5/2}$

(GATE PH 2021)

**Q.15** To sustain lasing action in a three-level laser as shown in the figure, necessary condition(s) is(are)








Figure 7: three-level laser

- i. lifetime of the energy level 1 should be greater than that of energy level 2  
 ii. population of the particles in level 1 should be greater than that of level 0  
 iii. lifetime of the energy level 2 should be greater than that of energy level 0  
 iv. population of the particles in level 2 should be greater than that of level 1

(GATE PH 2021)

**Q.16** If  $y_n(x)$  is a solution of the differential equation  $y'' - 2xy' + 2ny = 0$  where  $n$  is an integer and the prime (') denotes differentiation with respect to  $x$ , then acceptable plot(s) of  $\psi_n(x) = e^{-x^2/2}y_n(x)$ , is(are)



(GATE PH 2021)

**Q.17 – Q.25 Numerical Answer Type (NAT), carry ONE mark each (no negative marks).**

**Q.17** The donor concentration in a sample of n-type silicon is increased by a factor of 100. Assuming the sample to be non-degenerate, the shift in the Fermi level (in meV) at 300 K (rounded off to the

nearest integer) is \_\_\_\_\_. (Given:  $k_B T = 25$  meV at 300 K) (GATE PH 2021)

- Q.18** Two observers  $O$  and  $O'$  observe two events  $P$  and  $Q$ . The observers have a constant relative speed of  $0.5c$ . In the units, where the speed of light,  $c$ , is taken as unity, the observer  $O$  obtained the following coordinates:

Event P:  $x=5, y=3, z=5, t=3$

Event Q:  $x=5, y=1, z=3, t=5$

The length of the space-time interval between these two events, as measured by  $O'$ , is  $L$ . The value of  $|L|$  (in integer) is \_\_\_\_\_. (GATE PH 2021)

- Q.19** A light source having its intensity peak at the wavelength 289.8 nm is calibrated as 10,000 K which is the temperature of an equivalent black body radiation. Considering the same calibration, the temperature of light source (in K) having its intensity peak at the wavelength 579.6 nm (rounded off to the nearest integer) is \_\_\_\_\_. (GATE PH 2021)

- Q.20** A hoop of mass  $M$  and radius  $R$  rolls without slipping along a straight line on a horizontal surface as shown in the figure. A point mass  $m$  slides without friction along the inner surface of the hoop, performing small oscillations about the mean position. The number of degrees of freedom of the system (in integer) is \_\_\_\_\_.




Figure 8: hoop of mass  $M$  and radius  $R$

(GATE PH 2021)

- Q.21** Three non-interacting bosonic particles of mass  $m$  each, are in a one-dimensional infinite potential well of width  $a$ . The energy of the third excited state of the system is  $x \times \frac{h^2 \pi^2}{ma^2}$ . The value of  $x$  (in integer) is \_\_\_\_\_. (GATE PH 2021)

**Q.22** The spacing between two consecutive S-branch lines of the rotational Raman spectra of hydrogen gas is  $243.2 \text{ cm}^{-1}$ . After excitation with a laser of wavelength  $514.5 \text{ nm}$ , the Stoke's line appeared at  $17611.4 \text{ cm}^{-1}$  for a particular energy level. The wavenumber (rounded off to the nearest integer), in  $\text{cm}^{-1}$ , at which Stoke's line will appear for the next higher energy level is \_\_\_\_\_.  
(GATE PH 2021)

**Q.23** The transition line, as shown in the figure, arises between  ${}^2D_{3/2}$  and  ${}^2P_{1/2}$  states without any external magnetic field. The number of lines that will appear in the presence of a weak magnetic field (in integer) is \_\_\_\_\_.




Figure 9: transition line

(GATE PH 2021)

**Q.24** Consider the atomic system as shown in the figure, where the Einstein A coefficients for spontaneous emission for the levels are  $A_{2 \rightarrow 1} = 2 \times 10^7 \text{ s}^{-1}$  and  $A_{1 \rightarrow 0} = 10^8 \text{ s}^{-1}$ . If  $10^{14} \text{ atoms/cm}^3$  are excited from level 0 to level 2 and a steady state population in level 2 is achieved, then the steady state population at level 1 will be  $x \times 10^{13} \text{ cm}^{-3}$ . The value of  $x$  (in integer) is \_\_\_\_\_.




Figure 10: atomic system figure

(GATE PH 2021)

**Q.25** If  $\vec{a}$  and  $\vec{b}$  are constant vectors,  $\vec{r}$  and  $\vec{p}$  are generalized positions and conjugate momenta, respectively, then for the transformation  $Q = \vec{a} \cdot \vec{p}$  and  $P = \vec{b} \cdot \vec{r}$  to be canonical, the value of  $\vec{a} \cdot \vec{b}$  (in integer) is \_\_\_\_\_.  
(GATE PH 2021)

**Q.26 – Q.41 Multiple Choice Question (MCQ), carry TWO mark each (for each wrong answer: – 2/3).**




Figure 11: Logic gate combination.

**Q.26** The above combination of logic gates represents the operation

- i. OR                      ii. NAND                      iii. AND                      iv. NOR

(GATE PH 2021)

**Q.27** In a semiconductor, the ratio of the effective mass of hole to electron is 2:11 and the ratio of average relaxation time for hole to electron is 1:2. The ratio of the mobility of the hole to electron is

- i. 4:9                      ii. 4:11                      iii. 9:4                      iv. 11:4

(GATE PH 2021)

**Q.28** Consider a spin  $S = \hbar/2$  particle in the state  $|\phi\rangle = \frac{1}{3} \begin{bmatrix} 2+i \\ 2 \end{bmatrix}$ . The probability that a measurement finds the state with  $S_x = +\hbar/2$  is

- i. 5/18                      ii. 11/18                      iii. 15/18                      iv. 17/18

(GATE PH 2021)

**Q.29** An electromagnetic wave having electric field  $E = 8 \cos(kz - \omega t) \hat{y}$  V cm<sup>-1</sup> is incident at 90° (normal incidence) on a square slab from vacuum (with refractive index  $n_0 = 1.0$ ) as shown in the figure. The slab is composed of two different materials with refractive indices  $n_1 = 2.2$  and  $n_2 = 1.1$ . Assume that the permeability of each medium is the same. After passing through the slab for the first time, the electric field amplitude, in V cm<sup>-1</sup>, of the electromagnetic wave, which emerges from the slab in region 2, is closest to




Figure 12: EM wave incident on a composite slab.

- i.  $\frac{11}{1.6}$       ii.  $\frac{11}{3.2}$       iii.  $\frac{11}{13.8}$       iv.  $\frac{11}{25.6}$

(GATE PH 2021)

**Q.30** Consider a point charge  $+Q$  of mass  $m$  suspended by a massless, inextensible string of length  $l$  in free space (permittivity  $\epsilon_0$ ) as shown in the figure. It is placed at a height  $d$  ( $d > l$ ) over an infinitely large, grounded conducting plane. The gravitational potential energy is assumed to be zero at the position of the conducting plane and is positive above the plane.




Figure 13: Charged pendulum above a conducting plane.

If  $\theta$  represents the angular position and  $p_\theta$  its corresponding canonical momentum, then the correct Hamiltonian of the system is

- i.  $\frac{p_\theta^2}{2ml^2} - \frac{Q^2}{16\pi\epsilon_0(d-l\cos\theta)} - mg(d-l\cos\theta)$   
 ii.  $\frac{p_\theta^2}{2ml^2} - \frac{Q^2}{8\pi\epsilon_0(d-l\cos\theta)} + mg(d-l\cos\theta)$   
 iii.  $\frac{p_\theta^2}{2ml^2} - \frac{Q^2}{8\pi\epsilon_0(d-l\cos\theta)} - mg(d-l\cos\theta)$   
 iv.  $\frac{p_\theta^2}{2ml^2} - \frac{Q^2}{16\pi\epsilon_0(d-l\cos\theta)} + mg(d-l\cos\theta)$

(GATE PH 2021)

**Q.31** Consider two concentric conducting spherical shells as shown in the figure. The inner shell has a radius  $a$  and carries a charge  $+Q$ . The outer shell has a radius  $b$  and carries a charge  $-Q$ . The empty space



between them is half-filled by a hemispherical shell of a dielectric having permittivity  $\epsilon_1$ . The remaining space between the shells is filled with air having the permittivity  $\epsilon_0$ .




Figure 14: Concentric spherical shells with dielectric and air.

The electric field at a radial distance  $r$  from the center and between the shells ( $a < r < b$ ) is

- i.  $\frac{Q}{2\pi(\epsilon_0 + \epsilon_1)r^2} \hat{r}$  everywhere
- ii.  $\frac{Q}{4\pi\epsilon_0 r^2} \hat{r}$  on the air side and  $\frac{Q}{4\pi\epsilon_1 r^2} \hat{r}$  on the dielectric side
- iii.  $\frac{Q}{2\pi\epsilon_0 r^2} \hat{r}$  on the air side and  $\frac{Q}{2\pi\epsilon_1 r^2} \hat{r}$  on the dielectric side
- iv.  $\frac{Q}{4\pi(\epsilon_0 + \epsilon_1)r^2} \hat{r}$  everywhere

(GATE PH 2021)




Figure 15: Nuclear energy level schemes for two nuclei.

**Q.32** For the given sets of energy levels of nuclei X and Y whose mass numbers are odd and even, respectively, choose the best suited interpretation.

- i. Set I: Rotational band of X  
Set II: Vibrational band of Y
- ii. Set I: Rotational band of Y  
Set II: Vibrational band of X
- iii. Set I: Vibrational band of X  
Set II: Rotational band of Y
- iv. Set I: Vibrational band of Y  
Set II: Rotational band of X

(GATE PH 2021)

**Q.33** Consider a system of three distinguishable particles, each having spin  $S = 1/2$  such that  $S_z = \pm 1/2$  with corresponding magnetic moments  $\mu_z = \pm\mu$ . When the system is placed in an external magnetic field  $H$  pointing along the  $z$ -axis, the total energy of the system is  $\mu H$ . Let  $x$  be the state where the first spin has  $S_z = 1/2$ . The probability of having the state  $x$  and the mean magnetic moment (in the  $+z$  direction) of the system in state  $x$  are

- i.  $\frac{1}{3}, -\frac{1}{3}\mu$
- ii.  $\frac{1}{3}, \frac{2}{3}\mu$
- iii.  $\frac{2}{3}, -\frac{2}{3}\mu$
- iv.  $\frac{2}{3}, \frac{1}{3}\mu$

(GATE PH 2021)

**Q.34** Consider a particle in a one-dimensional infinite potential well with its walls at  $x = 0$  and  $x = L$ . The system is perturbed as shown in the figure.




Figure 16: Perturbation potential in an infinite well.

The first order correction to the energy eigenvalue is

- i.  $\frac{V_0}{4}$
- ii.  $\frac{V_0}{3}$
- iii.  $\frac{V_0}{2}$
- iv.  $\frac{V_0}{5}$

(GATE PH 2021)

**Q.35** Consider a state described by  $\psi(x, t) = \psi_2(x, t) + \psi_4(x, t)$ , where  $\psi_2(x, t)$  and  $\psi_4(x, t)$  are respectively the second and fourth normalized harmonic oscillator wave functions and  $\omega$  is the angular frequency of the harmonic oscillator. The wave function  $\psi(x, t = 0)$  will be orthogonal to  $\psi(x, t)$  at time  $t$  equal to

- i.  $\frac{\pi}{2\omega}$       ii.  $\frac{\pi}{\omega}$       iii.  $\frac{\pi}{4\omega}$       iv.  $\frac{\pi}{6\omega}$

(GATE PH 2021)

**Q.36** Consider a single one-dimensional harmonic oscillator of angular frequency  $\omega$ , in equilibrium at temperature  $T = (k_B\beta)^{-1}$ . The states of the harmonic oscillator are all non-degenerate having energy  $E_n = \left(n + \frac{1}{2}\right)\hbar\omega$  with equal probability, where  $n$  is the quantum number. The Helmholtz free energy of the oscillator is

- i.  $\frac{\hbar\omega}{2} + \beta^{-1} \ln[1 - \exp(\beta\hbar\omega)]$   
 ii.  $\frac{\hbar\omega}{2} + \beta^{-1} \ln[1 - \exp(-\beta\hbar\omega)]$   
 iii.  $\frac{\hbar\omega}{2} + \beta^{-1} \ln[1 + \exp(-\beta\hbar\omega)]$   
 iv.  $\beta^{-1} \ln[1 - \exp(-\beta\hbar\omega)]$

(GATE PH 2021)

**Q.37** A system of two atoms can be in three quantum states having energies  $0, \epsilon$  and  $2\epsilon$ . The system is in equilibrium at temperature  $T = (k_B\beta)^{-1}$ . Match the following Statistics with the Partition function.





| Statistics                                  | Partition function   |
|---|--|
| CD: Classical (distinguishable particles)   | Z1: $e^{-\beta\epsilon} + e^{-2\beta\epsilon} + e^{-3\beta\epsilon}$   |
| CI: Classical (indistinguishable particles) | Z2: $1 + e^{-\beta\epsilon} + e^{-2\beta\epsilon} + e^{-3\beta\epsilon} + e^{-4\beta\epsilon}$                                 |
| FD: Fermi-Dirac                             | Z3: $1 + 2e^{-\beta\epsilon} + 3e^{-2\beta\epsilon} + 2e^{-3\beta\epsilon} + e^{-4\beta\epsilon}$                              |
| BE: Bose-Einstein                           | Z4: $\frac{1}{2} + e^{-\beta\epsilon} + \frac{3}{2}e^{-2\beta\epsilon} + e^{-3\beta\epsilon} + \frac{1}{2}e^{-4\beta\epsilon}$ |

- i. CD:Z1, CI:Z2, FD:Z3, BE:Z4  
 ii. CD:Z2, CI:Z3, FD:Z4, BE:Z1  
 iii. CD:Z3, CI:Z4, FD:Z1, BE:Z2  
 iv. CD:Z4, CI:Z1, FD:Z2, BE:Z3

(GATE PH 2021)

**Q.38** The free energy of a ferromagnet is given by  $F = F_0 + a_0(T - T_C)M^2 + bM^4$ , where  $F_0$ ,  $a_0$ , and  $b$  are positive constants,  $M$  is the





magnetization,  $T$  is the temperature, and  $T_C$  is the Curie temperature. The relation between  $M^2$  and  $T$  is best depicted by



(GATE PH 2021)

**Q.39** Consider a spherical galaxy of total mass  $M$  and radius  $R$ , having a uniform matter distribution. In this idealized situation, the orbital

speed  $v$  of a star of mass  $m$  ( $m \ll M$ ) as a function of the distance  $r$  from the galactic centre is best described by ( $G$  is the universal gravitational constant)



(GATE PH 2021)

**Q.40** Consider the potential  $U(r)$  defined as

$$U(r) = -U_0 \frac{e^{-\alpha r}}{r}$$

where  $\alpha$  and  $U_0$  are real constants of appropriate dimensions. According to the first Born approximation, the elastic scattering amplitude calculated with  $U(r)$  for a (wave-vector) momentum transfer  $q$  and  $\alpha \rightarrow 0$ , is proportional to

(Useful integral:  $\int_0^\infty \sin(qr)e^{-\alpha r} dr = \frac{q}{\alpha^2 + q^2}$ )

- i.  $q^{-2}$                       ii.  $q^{-1}$                       iii.  $q$                       iv.  $q^2$

(GATE PH 2021)

**Q.41** As shown in the figure, inverse magnetic susceptibility ( $1/\chi$ ) is plotted as a function of temperature ( $T$ ) for three different materials in paramagnetic states.




Figure 17: Inverse magnetic susceptibility versus temperature.

(Curie temperature of ferromagnetic material =  $T_C$ )

Néel temperature of antiferromagnetic material =  $T_N$ )

Choose the correct statement from the following

- Material 1 is antiferromagnetic ( $T < T_N$ ), 2 is paramagnetic, and 3 is ferromagnetic ( $T < T_C$ ).
- Material 1 is paramagnetic, 2 is antiferromagnetic ( $T < T_N$ ), and 3 is ferromagnetic ( $T < T_C$ ).
- Material 1 ferromagnetic ( $T < T_C$ ), 2 is antiferromagnetic ( $T < T_N$ ), and 3 is paramagnetic.
- Material 1 is ferromagnetic ( $T < T_C$ ), 2 is paramagnetic, and 3 is antiferromagnetic ( $T < T_N$ ).

(GATE PH 2021)

**Q.42 – Q.46 Multiple Select Question (MSQ), carry TWO mark each (no negative marks).**

**Q.42** A function  $f(t)$  is defined only for  $t \geq 0$ . The Laplace transform of  $f(t)$  is

$$\mathcal{L}\{f; s\} = \int_0^{\infty} e^{-st} f(t) dt$$

whereas the Fourier transform of  $f(t)$  is

$$\mathcal{F}(\omega) = \int_0^{\infty} f(t) e^{-i\omega t} dt.$$

The correct statement(s) is(are)

- i. The variable  $s$  is always real.
- ii. The variable  $s$  can be complex.
- iii.  $\mathcal{L}\{f; s\}$  and  $\mathcal{F}(\omega)$  can never be made connected.
- iv.  $\mathcal{L}\{f; s\}$  and  $\mathcal{F}(\omega)$  can be made connected.

(GATE PH 2021)

**Q.43**  $P$  and  $Q$  are two Hermitian matrices and there exists a matrix  $R$ , which diagonalizes both of them, such that  $RPR^{-1} = S_1$  and  $RQR^{-1} = S_2$ , where  $S_1$  and  $S_2$  are diagonal matrices. The correct statement(s) is(are)

- i. All the elements of both matrices  $S_1$  and  $S_2$  are real.
- ii. The matrix  $PQ$  can have complex eigenvalues.
- iii. The matrix  $QP$  can have complex eigenvalues.
- iv. The matrices  $P$  and  $Q$  commute.

(GATE PH 2021)

**Q.44** A uniform block of mass  $M$  slides on a smooth horizontal bar. Another mass  $m$  is connected to it by an inextensible string of length  $l$  of negligible mass, and is constrained to oscillate in the X-Y plane only. Neglect the sizes of the masses. The number of degrees of freedom of the system is two and the generalized coordinates are chosen as  $x$  and  $\theta$ , as shown in the figure.




Figure 18: A block with a pendulum attached.

If  $p_x$  and  $p_\theta$  are the generalized momenta corresponding to  $x$  and  $\theta$ , respectively, then the correct option(s) is(are)

- i.  $p_x = (m + M)\dot{x} + ml \cos \theta \dot{\theta}$
- ii.  $p_\theta = ml^2 \dot{\theta} - ml \cos \theta \dot{x}$
- iii.  $p_x$  is conserved
- iv.  $p_\theta$  is conserved

(GATE PH 2021)

**Q.45** The Gell-Mann – Okubo mass formula defines the mass of baryons as  $M = M_0 + aY + b \left[ I(I + 1) - \frac{1}{4}Y^2 \right]$ , where  $M_0$ ,  $a$  and  $b$  are constants,  $I$  represents the isospin and  $Y$  represents the hypercharge. If the mass of  $\Sigma$  hyperons is same as that of  $\Lambda$  hyperons, then the correct option(s) is(are)

- i.  $M \propto I(I + 1)$
- ii.  $M \propto Y$
- iii.  $M$  does not depend on  $I$
- iv.  $M$  does not depend on  $Y$

(GATE PH 2021)

**Q.46** The time derivative of a differentiable function  $g(q_i, t)$  is added to a Lagrangian  $L(q_i, \dot{q}_i, t)$  such that

$$L' = L(q_i, \dot{q}_i, t) + \frac{d}{dt}g(q_i, t)$$

where  $q_i, \dot{q}_i, t$  are the generalized coordinates, generalized velocities and time, respectively. Let  $p_i$  be the generalized momentum and  $H$  the Hamiltonian associated with  $L(q_i, \dot{q}_i, t)$ . If  $p'_i$  and  $H'$  are those associated with  $L'$ , then the correct option(s) is(are)

- i. Both  $L$  and  $L'$  satisfy the Euler-Lagrange's equations of motion



- ii.  $p'_i = p_i + \frac{\partial}{\partial q_i} g(q_i, t)$
- iii. If  $p_i$  is conserved, then  $p'_i$  is necessarily conserved
- iv.  $H' = H + \frac{d}{dt} g(q_i, t)$

(GATE PH 2021)

**Q.47 – Q.55 Numerical Answer Type (NAT), carry TWO mark each (no negative marks).**

- Q.47** A linear charged particle accelerator is driven by an alternating voltage source operating at 10 MHz. Assume that it is used to accelerate electrons. After a few drift-tubes, the electrons attain a velocity  $2.9 \times 10^8 \text{ m s}^{-1}$ . The minimum length of each drift-tube, in m, to accelerate the electrons further (rounded off to one decimal place) is \_\_\_\_\_. (GATE PH 2021)
- Q.48** The Coulomb energy component in the binding energy of a nucleus is 18.432 MeV. If the radius of the uniform and spherical charge distribution in the nucleus is 3 fm, the corresponding atomic number (rounded off to the nearest integer) is \_\_\_\_\_. (Given:  $\frac{e^2}{4\pi\epsilon_0} = 1.44 \text{ MeV fm}$ ) (GATE PH 2021)
- Q.49** For a two-nucleon system in spin singlet state, the spin is represented through the Pauli matrices  $\sigma_1, \sigma_2$  for particles 1 and 2, respectively. The value of  $(\sigma_1 \cdot \sigma_2)$  (in integer) is \_\_\_\_\_. (GATE PH 2021)
- Q.50** A contour integral is defined as

$$I_n = \oint_C \frac{dz}{(z - n)^2 + \pi^2}$$

where  $n$  is a positive integer and  $C$  is the closed contour, as shown in the figure, consisting of the line from  $-100$  to  $100$  and the semicircle traversed in the counter-clockwise sense.




Figure 19: Closed contour  $C$  in the complex plane.

The value of  $\sum_{n=1}^5 I_n$  (in integer) is \_\_\_\_\_.

- Q.51** The normalized radial wave function of the second excited state of hydrogen atom is

$$R(r) = \frac{1}{\sqrt{24}} a^{-3/2} \frac{r}{a} e^{-r/2a}$$

where  $a$  is the Bohr radius and  $r$  is the distance from the center of the atom. The distance at which the electron is most likely to be found is  $y \times a$ . The value of  $y$  (in integer) is \_\_\_\_\_. (**GATE PH 2021**)

- Q.52** Consider an atomic gas with number density  $n = 10^{20} \text{ m}^{-3}$ , in the ground state at 300 K. The valence electronic configuration of atoms is  $f^7$ . The paramagnetic susceptibility of the gas  $\chi = m \times 10^{-11}$ . The value of  $m$  (rounded off to two decimal places) is \_\_\_\_\_.  
(Given: Magnetic permeability of free space  $\mu_0 = 4\pi \times 10^{-7} \text{ H m}^{-1}$   
Bohr magneton  $\mu_B = 9.274 \times 10^{-24} \text{ A m}^2$   
Boltzmann constant  $k_B = 1.3807 \times 10^{-23} \text{ J K}^{-1}$ ) (**GATE PH 2021**)

- Q.53** Consider a cross-section of an electromagnet having an air-gap of 5 cm as shown in the figure. It consists of a magnetic material ( $\mu = 20000\mu_0$ ) and is driven by a coil having  $NI = 10^4 \text{ A}$ , where  $N$  is the number of turns and  $I$  is the current in Ampere.




Figure 20: Cross-section of an electromagnet.

Ignoring the fringe fields, the magnitude of the magnetic field  $\vec{B}$  (in Tesla, rounded off to two decimal places) in the air-gap between the magnetic poles is \_\_\_\_\_.

- Q.54** The spin  $\vec{S}$  and orbital angular momentum  $\vec{L}$  of an atom precess about  $\vec{J}$ , the total angular momentum.  $\vec{J}$  precesses about an axis fixed by a magnetic field  $\vec{B}_1 = 2B_0\hat{z}$ , where  $B_0$  is a constant. Now

the magnetic field is changed to  $\vec{B}_2 = B_0(\hat{x} + \sqrt{2}\hat{y} + 2\hat{z})$ . Given the orbital angular momentum quantum number  $l = 2$  and spin quantum number  $s = 1/2$ ,  $\theta$  is the angle between  $\vec{B}_1$  and  $\vec{J}$  for the largest possible values of total angular quantum number  $j$  and its z-component  $j_z$ . The value of  $\theta$  (in degree, rounded off to the nearest integer) is \_\_\_\_\_.  
(GATE PH 2021)

- Q.55** The spin-orbit effect splits the  $^2P \rightarrow ^2S$  transition (wavelength,  $\lambda = 6521 \text{ \AA}$ ) in Lithium into two lines with separation of  $\Delta\lambda = 0.14 \text{ \AA}$ . The corresponding positive value of energy difference between the above two lines, in eV, is  $m \times 10^{-5}$ . The value of  $m$  (rounded off to the nearest integer) is \_\_\_\_\_. (Given: Planck's constant,  $h = 4.125 \times 10^{-15} \text{ eV s}$   
Speed of light,  $c = 3 \times 10^8 \text{ m s}^{-1}$ ) (GATE PH 2021)

**END OF THE QUESTION PAPER**