

2.7.23

EE25BTECH11011 - Navya Banavathu

September 2,2025

Question

For what value of k ($k > 0$) is the area of the triangle with vertices $(-2, 6)$, $(2k, 4)$ and $(2k + 1, 10)$ equal to 52 sq. units?

Solution

Vertices of triangle are

$$A \begin{pmatrix} x_1 \\ y_1 \end{pmatrix} = \begin{pmatrix} -2 \\ 6 \end{pmatrix} \quad (1)$$

$$B \begin{pmatrix} x_2 \\ y_2 \end{pmatrix} = \begin{pmatrix} 2k \\ 4 \end{pmatrix} \quad (2)$$

$$C \begin{pmatrix} x_3 \\ y_3 \end{pmatrix} = \begin{pmatrix} 2k + 1 \\ 10 \end{pmatrix} \quad (3)$$

$$\text{Area} = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} \quad (4)$$

$$= \frac{1}{2} \begin{vmatrix} -2 & 6 & 1 \\ 2k & 4 & 1 \\ 2k+1 & 10 & 1 \end{vmatrix} \quad (5)$$

$$= \frac{1}{2} \left[-2 \begin{vmatrix} 4 & 1 \\ 10 & 1 \end{vmatrix} - 6 \begin{vmatrix} 2k & 1 \\ 2k+1 & 1 \end{vmatrix} + \begin{vmatrix} 2k & 4 \\ 2k+1 & 10 \end{vmatrix} \right] \quad (6)$$

$$= \frac{1}{2} \left[(-2)(4 - 10) - 6(2k - (2k + 1)) + (20k - 8k - 4) \right] \quad (7)$$

$$= \frac{1}{2} |12k + 14| \quad (8)$$

Solution

Given area of triangle = 52 sq.units,

$$\frac{1}{2}|12k + 14| = 52 \quad (9)$$

$$|12k + 14| = 104. \quad (10)$$

Now consider the two cases from the absolute value,

Case 1:

$$12k + 14 = 104 \quad (11)$$

$$12k = 90 \quad (12)$$

$$k = \frac{90}{12} = \frac{15}{2}. \quad (13)$$

Case 2:

$$12k + 14 = -104 \quad (14)$$

$$12k = -118 \quad (15)$$

$$k = -\frac{118}{12} = -\frac{59}{6}. \quad (16)$$

Because the problem requires $k > 0$, we discard the negative solution.

$$\boxed{k = \frac{15}{2}} \quad (17)$$

Code (C)

Part 1

```
#include <stdio.h>

#ifdef _WIN32
#define EXPORT __declspec(dllexport)
#else
#define EXPORT
#endif

// Function to compute the positive value of k
EXPORT double solve_k_positive() {
    double k1 = (104.0 - 14.0) / 12.0; // 7.5
    double k2 = (-104.0 - 14.0) / 12.0; // -9.8333

    return (k1 > 0) ? k1 : k2;
}
```

Part 2

```
int main() {  
    double k = solve_k_positive();  
    printf("Positive value of k: %.2lf\n", k);  
    return 0;  
}
```


Python Code:Import and Define points

Part 1

```
import ctypes
import os
import matplotlib.pyplot as plt

# Load compiled shared library from code4.c
if os.name == "nt": # Windows
    lib = ctypes.CDLL("./code4.dll")
else: # Linux/macOS
    lib = ctypes.CDLL("./code4.so")

# Define C function signature
lib.solve_k.argtypes = [ctypes.POINTER(ctypes.c_double)]
lib.solve_k.restype = None
```

Part 2

```
# Prepare array for results
results = (ctypes.c_double * 2)()
lib.solve_k(results)

# Extract values of k
k1, k2 = results[0], results[1]
print(f"Possible values of k: {k1:.2f}, {k2:.2f}")

# Pick valid solution (k > 0)
k = k1 if k1 > 0 else k2
print(f"Chosen k = {k:.2f}")
```

Part 4

```
# Define triangle vertices
A = (-2, 6)
B = (2 * k, 4)
C = (2 * k + 1, 10)

# Plot triangle
plt.figure(figsize=(7,7))
plt.plot([A[0], B[0], C[0], A[0]],
         [A[1], B[1], C[1], A[1]],
         'b-', linewidth=2)

# Plot vertices as red points
plt.scatter([A[0], B[0], C[0]], [A[1], B[1], C[1]],
           color='red', s=80, zorder=5)
```

Python Code: Plotting

Part 5

```
# Annotate vertices with coordinates
```

```
plt.text(A[0]+0.2, A[1]+0.2, f"A{A}", fontsize=12)
```

```
plt.text(B[0]+0.2, B[1]-0.4, f"B{B}", fontsize=12)
```

```
plt.text(C[0]+0.2, C[1]+0.2, f"C{C}", fontsize=12)
```

```
plt.title(f"Triangle with k = {k:.2f}")
```

```
plt.grid(True)
```

```
plt.axis("equal")
```

```
# Save as fig4.png
```

```
filename = "fig4.png"
```

```
plt.savefig(filename, dpi=200)
```

```
print(f"Figure saved as {filename}")
```

```
# Show the figure
```

```
plt.show()
```

Graphical Representation




Figure: Area of triangle