

12.255

EE25BTECH11033 - Kavin

Question:

The least squares generalized inverse of an overdetermined problem is expressed as

$$1) (\mathbf{G}^\top \mathbf{G})^{-1} \mathbf{G}^\top$$

$$2) (\mathbf{G}^\top \mathbf{G})^{-1}$$

$$3) \mathbf{G}^\top (\mathbf{G} \mathbf{G}^\top)^{-1}$$

$$4) (\mathbf{G} \mathbf{G}^\top)^{-1}$$

Solution:

Let us consider the overdetermined system of equations,

$$\mathbf{G}\mathbf{x} = \mathbf{b} \quad (1)$$

Define the residual vector as,

$$\mathbf{r} = \mathbf{b} - \mathbf{G}\mathbf{x} \quad (2)$$

The least-squares problem is to minimize the squared error:

$$\mathbf{r}^\top \mathbf{r} = (\mathbf{b} - \mathbf{G}\mathbf{x})^\top (\mathbf{b} - \mathbf{G}\mathbf{x}) \quad (3)$$

$$\mathbf{r}^\top \mathbf{r} = (\mathbf{b}^\top - \mathbf{x}^\top \mathbf{G}^\top)(\mathbf{b} - \mathbf{G}\mathbf{x}) \quad (4)$$

$$\mathbf{r}^\top \mathbf{r} = \mathbf{x}^\top \mathbf{G}^\top \mathbf{G}\mathbf{x} - \mathbf{x}^\top \mathbf{G}^\top \mathbf{b} - \mathbf{b}^\top \mathbf{G}\mathbf{x} + \mathbf{b}^\top \mathbf{b} \quad (5)$$

Now, to minimize $\mathbf{r}^\top \mathbf{r}$, define

$$f(x) = \mathbf{x}^\top \mathbf{G}^\top \mathbf{G}\mathbf{x} - 2\mathbf{x}^\top \mathbf{G}^\top \mathbf{b} + \mathbf{b}^\top \mathbf{b} \quad (6)$$

Taking gradient with respect to $\mathbf{x}$:

$$\nabla_x f(x) = 2\mathbf{G}^\top \mathbf{G}\mathbf{x} - 2\mathbf{G}^\top \mathbf{b} \quad (7)$$

Setting $\nabla_x f(x) = 0$:

$$\mathbf{G}^\top \mathbf{G}\mathbf{x} = \mathbf{G}^\top \mathbf{b} \quad (8)$$

Therefore, the least squares solution is

$$\mathbf{x} = \boxed{(\mathbf{G}^\top \mathbf{G})^{-1} \mathbf{G}^\top \mathbf{b}} \quad (9)$$

Therefore, $(\mathbf{G}^\top \mathbf{G})^{-1} \mathbf{G}^\top$ is the least squares generalized inverse of an overdetermined problem.