

Solutions to Analytical Plane Geometry by Hem Raj and Hukum Chand

G V V Sharma*

CONTENTS

1 Coordinates

1

Abstract—This book provides a vector approach to analytical geometry. The content and exercises are based on Hem Raj and Hukum Chand's book on Analytical Plane Geometry.

1 COORDINATES

1.0.1. Find the distance between the following pair of points (11,16) and (23,21).

Let

$$\mathbf{A} = \begin{pmatrix} 11 \\ 16 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 23 \\ 21 \end{pmatrix} \quad (1.0.1.1)$$

Then,

$$\mathbf{B} - \mathbf{A} = \begin{pmatrix} 12 \\ 5 \end{pmatrix} \quad (1.0.1.2)$$

and

$$\|\mathbf{B} - \mathbf{A}\|^2 = (\mathbf{B} - \mathbf{A})^T (\mathbf{B} - \mathbf{A}) = \begin{pmatrix} 12 & 5 \end{pmatrix} \begin{pmatrix} 12 \\ 5 \end{pmatrix} \quad (1.0.1.3)$$

$$\Rightarrow \|\mathbf{B} - \mathbf{A}\| = \sqrt{(12^2 + 5^2)} \quad (1.0.1.4)$$

$$= 13$$

1.0.2. Find the distance between the following pair of points (66,25) and (99,69).

Let

$$\mathbf{A} = \begin{pmatrix} 66 \\ 25 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 99 \\ 69 \end{pmatrix} \quad (1.0.2.1)$$

Then

$$\|\mathbf{B} - \mathbf{A}\| = \|\mathbf{C}\| \quad (1.0.2.2)$$

$$\|\mathbf{C}\|^2 = \mathbf{C}^T \mathbf{C} \quad (1.0.2.3)$$

$$= \begin{pmatrix} 33 & 44 \end{pmatrix} \begin{pmatrix} 33 \\ 44 \end{pmatrix} \quad (1.0.2.4)$$

$$\Rightarrow \|\mathbf{C}\| = \sqrt{(33^2 + 44^2)} \quad (1.0.2.5)$$

$$= 55 \quad (1.0.2.6)$$

1.0.3. Find the area of the quadrilateral formed by the points

$$\mathbf{A} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 3 \\ 5 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} -2 \\ 4 \end{pmatrix}, \mathbf{D} = \begin{pmatrix} -1 \\ -5 \end{pmatrix}. \quad (1.0.3.1)$$

Solution: Let

$$\mathbf{A} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 3 \\ 5 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} -2 \\ 4 \end{pmatrix}, \mathbf{D} = \begin{pmatrix} -1 \\ -5 \end{pmatrix}. \quad (1.0.3.2)$$

Then,

$$(\mathbf{B} - \mathbf{A}) = \begin{pmatrix} 3 \\ 5 \end{pmatrix} - \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 4 \end{pmatrix} \quad (1.0.3.3)$$

$$(\mathbf{D} - \mathbf{B}) = \begin{pmatrix} -1 \\ -5 \end{pmatrix} - \begin{pmatrix} 3 \\ 5 \end{pmatrix} = \begin{pmatrix} -4 \\ -10 \end{pmatrix} \quad (1.0.3.4)$$

$$(\mathbf{A} - \mathbf{D}) = \begin{pmatrix} 1 \\ 1 \end{pmatrix} - \begin{pmatrix} -1 \\ -5 \end{pmatrix} = \begin{pmatrix} 2 \\ 6 \end{pmatrix} \quad (1.0.3.5)$$

Row reducing the matrix formed by the vec-

*The author is with the Department of Electrical Engineering, Indian Institute of Technology, Hyderabad 502285 India e-mail: gadepall@iith.ac.in. All content in this manual is released under GNU GPL. Free and open source.

tors,

$$\begin{pmatrix} 2 & 4 & 1 \\ -4 & -10 & 1 \\ 2 & 6 & 1 \end{pmatrix}$$

(1.0.3.6)

$$\Rightarrow \begin{pmatrix} 2 & 4 & 1 \\ -4 & -10 & 1 \\ 2 & 6 & 1 \end{pmatrix} \xleftrightarrow{R_2 \leftrightarrow R_3} \begin{pmatrix} 2 & 4 & 1 \\ 2 & 6 & 1 \\ -4 & -10 & 1 \end{pmatrix} \xleftrightarrow{R_3 \leftrightarrow R_1 + R_2 + R_3} \begin{pmatrix} 2 & 4 & 1 \\ 2 & 6 & 1 \\ 0 & 0 & 3 \end{pmatrix} \xleftrightarrow{R_2 \leftrightarrow R_2 - R_1} \begin{pmatrix} 2 & 4 & 1 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix}$$

(1.0.3.7)

The number of non-zero rows in the matrix = 3.
Hence the matrix is full rank and AB, BD, DA are not collinear. See Fig. 1.0.3.1, which shows that the points given form a quadrilateral. Area




Fig. 1.0.3.1: Quadrilateral ABCD

of a $\triangle ABC$ is given by

$$\Delta ABC = \frac{1}{2} \begin{vmatrix} 1 & 1 & 1 \\ 1 & 3 & -2 \\ 1 & 5 & 4 \end{vmatrix} = 9 \quad (1.0.3.8)$$

Area of $\triangle ACD$ is given by

$$\Delta ACD = \frac{1}{2} \begin{vmatrix} 1 & 1 & 1 \\ 1 & -2 & -1 \\ 1 & 4 & -5 \end{vmatrix} = 11.5 \quad (1.0.3.9)$$

Thus, the area of quadrilateral ABCD = 9 + 11.5
= 20.5.