

CHAPTER

21

Probability

Section-A

JEE Advanced/ IIT-JEE

A Fill in the Blanks

1. For a biased die the probabilities for the different faces to turn up are given below :

Face	1	2	3	4	5	6
Prob.	0.1	0.32	0.21	0.15	0.05	0.17

This die is tossed and you are told that either face 1 or face 2 has turned up. Then the probability that it is face 1 is (1981 - 2 Marks)

2. $P(A \cup B) = P(A \cap B)$ if and only if the relation between $P(A)$ and $P(B)$ is (1985 - 2 Marks)

3. A box contains 100 tickets numbered 1, 2, ..., 100. Two tickets are chosen at random. It is given that the maximum number on the two chosen tickets is not more than 10. The minimum number on them is 5 with probability (1985 - 2 Marks)

4. If $\frac{1+3p}{3}$, $\frac{1-p}{4}$ and $\frac{1-2p}{2}$ are the probabilities of three mutually exclusive events, then the set of all values of p is (1986 - 2 Marks)

5. Urn A contains 6 red and 4 black balls and urn B contains 4 red and 6 black balls. One ball is drawn at random from urn A and placed in urn B . Then one ball is drawn at random from urn B and placed in urn A . If one ball is now drawn at random from urn A , the probability that it is found to be red is (1988 - 2 Marks)

6. A pair of fair dice is rolled together till a sum of either 5 or 7 is obtained. Then the probability that 5 comes before 7 is (1989 - 2 Marks)

7. Let A and B be two events such that $P(A) = 0.3$ and $P(A \cup B) = 0.8$. If A and B are independent events then $P(B) =$ (1990 - 2 Marks)

8. If the mean and the variance of a binomial variate X are 2 and 1 respectively, then the probability that X takes a value greater than one is equal to (1991 - 2 Marks)

9. Three faces of a fair die are yellow, two faces red and one blue. The die is tossed three times. The probability that the colours, yellow, red and blue, appear in the first, second and the third tosses respectively is (1992 - 2 Marks)

10. If two events A and B are such that $P(A^c) = 0.3$, $P(B) = 0.4$ and $P(A \cap B^c) = 0.5$, then $P(B/(A \cup B^c)) =$ (1994 - 2 Marks)

B True / False

1. If the letters of the word "Assassin" are written down at random in a row, the probability that no two S's occur together is $1/35$ (1983 - 1 Mark)
2. If the probability for A to fail in an examination is 0.2 and that for B is 0.3, then the probability that either A or B fails is 0.5. (1989 - 1 Mark)

C MCQs with One Correct Answer

1. Two fair dice are tossed. Let x be the event that the first die shows an even number and y be the event that the second die shows an odd number. The two events x and y are :
 (a) Mutually exclusive
 (b) Independent and mutually exclusive
 (c) Dependent
 (d) None of these (1979)
2. Two events A and B have probabilities 0.25 and 0.50 respectively. The probability that both A and B occur simultaneously is 0.14. Then the probability that neither A nor B occurs is (1980)
 (a) 0.39 (b) 0.25 (c) 0.11 (d) none of these
3. The probability that an event A happens in one trial of an experiment is 0.4. Three independent trials of the experiment are performed. The probability that the event A happens at least once is (1980)
 (a) 0.936 (b) 0.784 (c) 0.904 (d) none of these
4. If A and B are two events such that $P(A) > 0$, and $P(B) \neq 1$,

then $P\left(\frac{\bar{A}}{B}\right)$ is equal to (1982 - 2 Marks)

- (a) $1 - P\left(\frac{A}{B}\right)$ (b) $1 - P\left(\frac{\bar{A}}{\bar{B}}\right)$
 (c) $\frac{1 - P(A \cup B)}{P(\bar{B})}$ (d) $\frac{P(\bar{A})}{P(\bar{B})}$

(Here \bar{A} and \bar{B} are complements of A and B respectively). Fifteen coupons are numbered 1, 2, ..., 15, respectively. Seven coupons are selected at random one at a time with replacement. The probability that the largest number appearing on a selected coupon is 9, is (1983 - 1 Mark)

- (a) $\left(\frac{9}{16}\right)^6$ (b) $\left(\frac{8}{15}\right)^7$ (c) $\left(\frac{3}{5}\right)^7$ (d) none of these

6. Three identical dice are rolled. The probability that the same number will appear on each of them is
 (1984 - 2 Marks)
 (a) $\frac{1}{6}$ (b) $\frac{1}{36}$ (c) $\frac{1}{18}$ (d) $\frac{3}{28}$
7. A box contains 24 identical balls of which 12 are white and 12 are black. The balls are drawn at random from the box one at a time with replacement. The probability that a white ball is drawn for the 4th time on the 7th draw is
 (1984 - 2 Marks)
 (a) $\frac{5}{64}$ (b) $\frac{27}{32}$ (c) $\frac{5}{32}$ (d) $\frac{1}{2}$
8. One hundred identical coins, each with probability, p , of showing up heads are tossed once. If $0 < p < 1$ and the probability of heads showing on 50 coins is equal to that of heads showing on 51 coins, then the value of p is
 (1988 - 2 Marks)
 (a) $\frac{1}{2}$ (b) $\frac{49}{101}$ (c) $\frac{50}{101}$ (d) $\frac{51}{101}$
9. India plays two matches each with West Indies and Australia. In any match the probabilities of India getting, points 0, 1 and 2 are 0.45, 0.05 and 0.50 respectively. Assuming that the outcomes are independent, the probability of India getting at least 7 points is
 (1992 - 2 Marks)
 (a) 0.8750 (b) 0.0875 (c) 0.0625 (d) 0.0250
10. An unbiased die with faces marked 1, 2, 3, 4, 5 and 6 is rolled four times. Out of four face values obtained, the probability that the minimum face value is not less than 2 and the maximum face value is not greater than 5, is then:
 (1993 - 1 Mark)
 (a) $\frac{16}{81}$ (b) $\frac{1}{81}$ (c) $\frac{80}{81}$ (d) $\frac{65}{81}$
11. Let A , B , C be three mutually independent events. Consider the two statements S_1 and S_2
 S_1 : A and $B \cup C$ are independent
 S_2 : A and $B \cap C$ are independent
 Then,
 (a) Both S_1 and S_2 are true
 (b) Only S_1 is true
 (c) Only S_2 is true
 (d) Neither S_1 nor S_2 is true
 (1994)
12. The probability of India winning a test match against West Indies is $\frac{1}{2}$. Assuming independence from match to match the probability that in a 5 match series India's second win occurs at third test is
 (1995S)
 (a) $\frac{1}{8}$ (b) $\frac{1}{4}$ (c) $\frac{1}{2}$ (d) $\frac{2}{3}$
13. Three of the six vertices of a regular hexagon are chosen at random. The probability that the triangle with three vertices is equilateral, equals
 (1995S)
 (a) $\frac{1}{2}$ (b) $\frac{1}{5}$ (c) $\frac{1}{10}$ (d) $\frac{1}{20}$
14. For the three events A , B , and C , P (exactly one of the events A or B occurs) = P (exactly one of the two events B or C occurs) = P (exactly one of the events C or A occurs) = p and P (all the three events occur simultaneously) = p^2 , where $0 < p < 1/2$. Then the probability of at least one of the three events A , B and C occurring is
 (1996 - 2 Marks)
 (a) $\frac{3p+2p^2}{2}$ (b) $\frac{p+3p^2}{4}$
 (c) $\frac{p+3p^2}{2}$ (d) $\frac{3p+2p^2}{4}$
15. If the integers m and n are chosen at random from 1 to 100, then the probability that a number of the form $7^m + 7^n$ is divisible by 5 equals
 (1999 - 2 Marks)
 (a) $\frac{1}{4}$ (b) $\frac{1}{7}$ (c) $\frac{1}{8}$ (d) $\frac{1}{49}$
16. Two numbers are selected randomly from the set $S = \{1, 2, 3, 4, 5, 6\}$ without replacement one by one. The probability that minimum of the two numbers is less than 4 is
 (2003S)
 (a) $\frac{1}{15}$ (b) $\frac{14}{15}$ (c) $\frac{1}{5}$ (d) $\frac{4}{5}$
17. If $P(B) = \frac{3}{4}$, $P(A \cap B \cap \bar{C}) = \frac{1}{3}$ and
 $P(\bar{A} \cap B \cap \bar{C}) = \frac{1}{3}$, then $P(B \cap C)$ is
 (2003S)
 (a) $\frac{1}{12}$ (b) $\frac{1}{6}$ (c) $\frac{1}{15}$ (d) $\frac{1}{9}$
18. If three distinct numbers are chosen randomly from the first 100 natural numbers, then the probability that all three of them are divisible by both 2 and 3 is
 (2004S)
 (a) $\frac{4}{25}$ (b) $\frac{4}{35}$ (c) $\frac{4}{33}$ (d) $\frac{4}{1155}$
19. A six faced fair dice is thrown until 1 comes, then the probability that 1 comes in even no. of trials is
 (2005S)
 (a) $\frac{5}{11}$ (b) $\frac{5}{6}$ (c) $\frac{6}{11}$ (d) $\frac{1}{6}$
20. One Indian and four American men and their wives are to be seated randomly around a circular table. Then the conditional probability that the Indian man is seated adjacent to his wife given that each American man is seated adjacent to his wife is
 (2007 - 3 marks)
 (a) $\frac{1}{2}$ (b) $\frac{1}{3}$ (c) $\frac{2}{5}$ (d) $\frac{1}{5}$
21. Let E^c denote the complement of an event E . Let E, F, G be pairwise independent events with $P(G) > 0$ and $P(E \cap F \cap G) = 0$. Then $P(E^c \cap F^c | G)$ equals
 (2007 - 3 marks)
 (a) $P(E^c) + P(F^c)$ (b) $P(E^c) - P(F^c)$
 (c) $P(E^c) - P(F)$ (d) $P(E) - P(F^c)$
22. An experiment has 10 equally likely outcomes. Let A and B be non-empty events of the experiment. If A consists of 4 outcomes, the number of outcomes that B must have so that A and B are independent, is
 (2008)
 (a) 2, 4 or 8 (b) 3, 6 or 9 (c) 4 or 8 (d) 5 or 10
23. Let ω be a complex cube root of unity with $\omega \neq 1$. A fair die is thrown three times. If r_1, r_2 and r_3 are the numbers obtained on the die, then the probability that $\omega^{r_1} + \omega^{r_2} + \omega^{r_3} = 0$ is
 (2010)
 (a) $\frac{1}{18}$ (b) $\frac{1}{9}$ (c) $\frac{2}{9}$ (d) $\frac{1}{36}$
24. A signal which can be green or red with probability $\frac{4}{5}$ and $\frac{1}{5}$ respectively, is received by station A and then transmitted to station B. The probability of each station receiving the signal correctly is $\frac{3}{4}$. If the signal received at station B is green, then the probability that the original signal was green is
 (2010)
 (a) $\frac{3}{5}$ (b) $\frac{6}{7}$ (c) $\frac{20}{23}$ (d) $\frac{9}{20}$

25. Four fair dice D_1, D_2, D_3 and D_4 ; each having six faces numbered 1, 2, 3, 4, 5 and 6 are rolled simultaneously. The probability that D_4 shows a number appearing on one of D_1, D_2 and D_3 is
(2012)

(a) $\frac{91}{216}$ (b) $\frac{108}{216}$ (c) $\frac{125}{216}$ (d) $\frac{127}{216}$

26. Three boys and two girls stand in a queue. The probability, that the number of boys ahead of every girl is at least one more than the number of girls ahead of her, is

(JEE Adv. 2014)

(a) $\frac{1}{2}$ (b) $\frac{1}{3}$ (c) $\frac{2}{3}$ (d) $\frac{3}{4}$

27. A computer producing factory has only two plants T_1 and T_2 . Plant T_1 produces 20% and plant T_2 produces 80% of the total computers produced. 7% of computers produced in the factory turn out to be defective. It is known that $P(\text{computer turns out to be defective given that it is produced in plant } T_1)$

$= 10P(\text{computer turns out to be defective given that it is produced in plant } T_2)$,

where $P(E)$ denotes the probability of an event E . A computer produced in the factory is randomly selected and it does not turn out to be defective. Then the probability that it is produced in plant T_2 is
(JEE Adv. 2016)

(a) $\frac{36}{73}$ (b) $\frac{47}{79}$ (c) $\frac{78}{93}$ (d) $\frac{75}{83}$

28. Three randomly chosen non-negative integers x, y and z are found to satisfy the equation $x + y + z = 10$. Then the probability that z is even, is

(a) $\frac{36}{55}$ (b) $\frac{6}{11}$ (c) $\frac{1}{2}$ (d) $\frac{5}{11}$

D MCQs with One or More than One Correct

1. If M and N are any two events, the probability that exactly one of them occurs is
(1984 - 3 Marks)

(a) $P(M) + P(N) - 2P(M \cap N)$
 (b) $P(M) + P(N) - P(M \cap N)$
 (c) $P(M^c) + P(N^c) - 2P(M^c \cap N^c)$
 (d) $P(M \cap N^c) + P(M^c \cap N)$

2. A student appears for tests I, II and III. The student is successful if he passes either in tests I and II or tests I and III. The probabilities of the student passing in tests I, II and

III are p, q and $\frac{1}{2}$ respectively. If the probability that the

student is successful is $\frac{1}{2}$, then
(1986 - 2 Marks)

(a) $p = q = 1$ (b) $p = q = \frac{1}{2}$
 (c) $p = 1, q = 0$ (d) $p = 1, q = \frac{1}{2}$
 (e) none of these

3. The probability that at least one of the events A and B occurs is 0.6. If A and B occur simultaneously with probability 0.2, then $P(\bar{A}) + P(\bar{B})$ is
(1987 - 2 Marks)

(a) 0.4 (b) 0.8 (c) 1.2 (d) 1.4
 (e) none

(Here \bar{A} and \bar{B} are complements of A and B , respectively).

4. For two given events A and B , $P(A \cap B)$
(1988 - 2 Marks)

(a) not less than $P(A) + P(B) - 1$
 (b) not greater than $P(A) + P(B)$
 (c) equal to $P(A) + P(B) - P(A \cup B)$
 (d) equal to $P(A) + P(B) + P(A \cup B)$

5. If E and F are independent events such that $0 < P(E) < 1$ and $0 < P(F) < 1$, then
(1989 - 2 Marks)

(a) E and F are mutually exclusive
 (b) E and F^c (the complement of the event F) are independent
 (c) E^c and F^c are independent
 (d) $P(E|F) + P(E^c|F) = 1$.

6. For any two events A and B in a sample space
(1991 - 2 Marks)

(a) $P(A|B) \geq \frac{P(A) + P(B) - 1}{P(B)}$, $P(B) \neq 0$ is always true

(b) $P(A \cap \bar{B}) = P(A) - P(A \cap B)$ does not hold

(c) $P(A \cup B) = 1 - P(\bar{A})P(\bar{B})$, if A and B are independent

(d) $P(A \cup B) = 1 - P(\bar{A})P(\bar{B})$, if A and B are disjoint.

7. E and F are two independent events. The probability that both E and F happen is $1/12$ and the probability that neither E nor F happens is $1/2$. Then,
(1993 - 2 Marks)

(a) $P(E) = 1/3, P(F) = 1/4$

(b) $P(E) = 1/2, P(F) = 1/6$

(c) $P(E) = 1/6, P(F) = 1/2$

(d) $P(E) = 1/4, P(F) = 1/3$

8. Let $0 < P(A) < 1, 0 < P(B) < 1$ and

$P(A \cup B) = P(A) + P(B) - P(A)P(B)$ then
(1995S)

(a) $P(B|A) = P(B) - P(A)$

(b) $P(A' \cap B') = P(A') - P(B')$

(c) $P(A \cup B)' = P(A')P(B')$

(d) $P(A|B) = P(A)$

9. If from each of the three boxes containing 3 white and 1 black, 2 white and 2 black, 1 white and 3 black balls, one ball is drawn at random, then the probability that 2 white and 1 black ball will be drawn is
(1998 - 2 Marks)

(a) $13/32$ (b) $1/4$ (c) $1/32$ (d) $3/16$

10. If \bar{E} and \bar{F} are the complementary events of events E and F respectively and if $0 < P(F) < 1$, then
(1998 - 2 Marks)

(a) $P(E|F) + P(\bar{E}|F) = 1$

(b) $P(E|F) + P(E|\bar{F}) = 1$

(c) $P(\bar{E}|F) + P(E|\bar{F}) = 1$

(d) $P(E|\bar{F}) + P(\bar{E}|\bar{F}) = 1$

11. There are four machines and it is known that exactly two of them are faulty. They are tested, one by one, in a random order till both the faulty machines are identified. Then the probability that only two tests are needed is (1998 - 2 Marks)
- (a) $\frac{1}{3}$ (b) $\frac{1}{6}$ (c) $\frac{1}{2}$ (d) $\frac{1}{4}$
12. If E and F are events with $P(E) \leq P(F)$ and $P(E \cap F) > 0$, then (1998 - 2 Marks)
- (a) occurrence of E \Rightarrow occurrence of F
 (b) occurrence of F \Rightarrow occurrence of E
 (c) non-occurrence of E \Rightarrow non-occurrence of F
 (d) none of the above implications holds
13. A fair coin is tossed repeatedly. If the tail appears on first four tosses, then the probability of the head appearing on the fifth toss equals (1998 - 2 Marks)
- (a) $\frac{1}{2}$ (b) $\frac{1}{32}$ (c) $\frac{31}{32}$ (d) $\frac{1}{5}$
14. Seven white balls and three black balls are randomly placed in a row. The probability that no two black balls are placed adjacently equals (1998 - 2 Marks)
- (a) $\frac{1}{2}$ (b) $\frac{7}{15}$ (c) $\frac{2}{15}$ (d) $\frac{1}{3}$
15. The probabilities that a student passes in Mathematics, Physics and Chemistry are m, p and c, respectively. Of these subjects, the student has a 75% chance of passing in at least one, a 50% chance of passing in at least two, and a 40% chance of passing in exactly two. Which of the following relations are true? (1999 - 3 Marks)
- (a) $p + m + c = 19/20$ (b) $p + m + c = 27/20$
 (c) $pmc = 1/10$ (d) $pmc = 1/4$
16. Let E and F be two independent events. The probability that exactly one of them occurs is $\frac{11}{25}$ and the probability of none of them occurring is $\frac{2}{25}$. If $P(T)$ denotes the probability of occurrence of the event T, then (2011)
- (a) $P(E) = \frac{4}{5}, P(F) = \frac{3}{5}$ (b) $P(E) = \frac{1}{5}, P(F) = \frac{2}{5}$
 (c) $P(E) = \frac{2}{5}, P(F) = \frac{1}{5}$ (d) $P(E) = \frac{3}{5}, P(F) = \frac{4}{5}$
17. A ship is fitted with three engines E_1, E_2 and E_3 . The engines function independently of each other with respective probabilities $\frac{1}{2}, \frac{1}{4}$ and $\frac{1}{4}$. For the ship to be operational at least two of its engines must function. Let X denote the event that the ship is operational and let X_1, X_2 and X_3 denote respectively the events that the engines E_1, E_2 and E_3 are functioning. Which of the following is/are true? (2012)
- (a) $P[X_1^c | X] = \frac{3}{16}$
 (b) $P[\text{Exactly two engines of the ship are functioning} | X] = \frac{7}{8}$
 (c) $P[X | X_2] = \frac{5}{16}$ (d) $P[X | X_1] = \frac{7}{16}$
18. Let X and Y be two events such that $P(X|Y) = \frac{1}{2}$, $P(Y|X) = \frac{1}{3}$ and $P(X \cap Y) = \frac{1}{6}$. Which of the following is/are correct? (2012)
- (a) $P(X \cup Y) = \frac{2}{3}$
 (b) X and Y are independent
 (c) X and Y are not independent
 (d) $P(X^c \cap Y) = \frac{1}{3}$
19. Four persons independently solve a certain problem correctly with probabilities $\frac{1}{2}, \frac{3}{4}, \frac{1}{4}, \frac{1}{8}$. Then the probability that the problem is solved correctly by at least one of them is (JEE Adv. 2013)
- (a) $\frac{235}{256}$ (b) $\frac{21}{256}$ (c) $\frac{3}{256}$ (d) $\frac{253}{256}$
20. Let X and Y be two events such that $P(X) = \frac{1}{3}$, $P(X|Y) = \frac{1}{2}$ and $P(Y|X) = \frac{2}{5}$. Then (JEE Adv. 2017)
- (a) $P(Y) = \frac{4}{15}$ (b) $P(X' | Y) = \frac{1}{2}$
 (c) $P(X \cap Y) = \frac{1}{5}$ (d) $P(X \cup Y) = \frac{2}{5}$
21. There are three bags B_1, B_2 , and B_3 . The bag B_1 contains 5 red and 5 green balls, B_2 contains 3 red and 5 green balls, and B_3 contains 5 red and 3 green balls. Bags B_1, B_2 and B_3 have probabilities $\frac{3}{10}, \frac{3}{10}$ and $\frac{4}{10}$ respectively of being chosen. A bag is selected at random and a ball is chosen at random from the bag. Then which of the following options is/are correct? (JEE Adv. 2019)
- (a) Probability that the selected bag is B_3 and the chosen ball is green equals $\frac{3}{10}$
 (b) Probability that the chosen ball is green, given that the selected bag is B_3 , equals $\frac{3}{8}$
 (c) Probability that the selected bag is B_3 , given that the chosen ball is green, equals $\frac{5}{13}$
 (d) Probability that the chosen ball is green equals $\frac{39}{80}$

E Subjective Problems

1. Balls are drawn one-by-one without replacement from a box containing 2 black, 4 white and 3 red balls till all the balls are drawn. Find the probability that the balls drawn are in the order 2 black, 4 white and 3 red. (1978)
2. Six boys and six girls sit in a row randomly. Find the probability that
 (i) the six girls sit together
 (ii) the boys and girls sit alternately. (1979)
3. An anti-aircraft gun can take a maximum of four shots at an enemy plane moving away from it. The probabilities of hitting the plane at the first, second, third and fourth shot are 0.4, 0.3, 0.2 and 0.1 respectively. What is the probability that the gun hits the plane? (1981 - 2 Marks)
4. A and B are two candidates seeking admission in IIT. The probability that A is selected is 0.5 and the probability that both A and B are selected is atmost 0.3. Is it possible that the probability of B getting selected is 0.9? (1982 - 2 Marks)
5. Cards are drawn one by one at random from a well-shuffled full pack of 52 playing cards until 2 aces are obtained for the first time. If N is the number of cards required to be drawn, then show that $P_r\{N = n\} = \frac{(n-1)(52-n)(51-n)}{50 \times 49 \times 17 \times 13}$ where $2 \leq n \leq 50$ (1983 - 3 Marks)
6. A, B, C are events such that (1983 - 2 Marks)
 $P(A) = 0.3, P(B) = 0.4, P(C) = 0.8$
 $P(AB) = 0.08, P(AC) = 0.28; P(ABC) = 0.09$
- If $P(A \cup B \cup C) \geq 0.75$, then show that $P(BC)$ lies in the interval $0.23 \leq x \leq 0.48$
7. In a certain city only two newspapers A and B are published, it is known that 25% of the city population reads A and 20% reads B while 8% reads both A and B . It is also known that 30% of those who read A but not B look into advertisements and 40% of those who read B but not A look into advertisements while 50% of those who read both A and B look into advertisements. What is the percentage of the population that reads an advertisement? (1984 - 4 Marks)
8. In a multiple-choice question there are four alternative answers, of which one or more are correct. A candidate will get marks in the question only if he ticks the correct answers. The candidate decides to tick the answers at random, if he is allowed upto three chances to answer the questions, find the probability that he will get marks in the questions. (1985 - 5 Marks)
9. A lot contains 20 articles. The probability that the lot contains exactly 2 defective articles is 0.4 and the probability that the lot contains exactly 3 defective articles is 0.6. Articles are drawn from the lot at random one by one without replacement and are tested till all defective articles are found. What is the probability that the testing procedure ends at the twelfth testing. (1986 - 5 Marks)
10. A man takes a step forward with probability 0.4 and backwards with probability 0.6. Find the probability that at the end of eleven steps he is one step away from the starting point. (1987 - 3 Marks)
11. A box contains 2 fifty paise coins, 5 twenty five paise coins and a certain fixed number N (≥ 2) of ten and five paise coins. Five coins are taken out of the box at random. Find the probability that the total value of these 5 coins is less than one rupee and fifty paise. (1988 - 3 Marks)
12. Suppose the probability for A to win a game against B is 0.4. If A has an option of playing either a "best of 3 games" or a "best of 5 games" match against B , which option should be chosen so that the probability of his winning the match is higher? (No game ends in a draw). (1989 - 5 Marks)
13. A is a set containing n elements. A subset P of A is chosen at random. The set A is reconstructed by replacing the elements of P . A subset Q of A is again chosen at random. Find the probability that P and Q have no common elements. (1990 - 5 Marks)
14. In a test an examine either guesses or copies or knows the answer to a multiple choice question with four choices. The probability that he make a guess is $1/3$ and the probability that he copies the answer is $1/6$. The probability that his answer is correct given that he copied it, is $1/8$. Find the probability that he knew the answer to the question given that he correctly answered it. (1991 - 4 Marks)
15. A lot contains 50 defective and 50 non defective bulbs. Two bulbs are drawn at random, one at a time, with replacement. The events A, B, C are defined as (1992 - 6 Marks)
 $A =$ (the first bulb is defective)
 $B =$ (the second bulb is non-defective)
 $C =$ (the two bulbs are both defective or both non defective)
 Determine whether
 (i) A, B, C are pairwise independent
 (ii) A, B, C are independent
16. Numbers are selected at random, one at a time, from the two-digit numbers 00, 01, 02, ..., 99 with replacement. An event E occurs if only if the product of the two digits of a selected number is 18. If four numbers are selected, find probability that the event E occurs at least 3 times. (1993 - 5 Marks)
17. An unbiased coin is tossed. If the result is a head, a pair of unbiased dice is rolled and the number obtained by adding the numbers on the two faces is noted. If the result is a tail, a card from a well shuffled pack of eleven cards numbered 2, 3, 4, ..., 12 is picked and the number on the card is noted. What is the probability that the noted number is either 7 or 8? (1994 - 5 Marks)
18. In how many ways three girls and nine boys can be seated in two vans, each having numbered seats, 3 in the front and 4 at the back? How many seating arrangements are possible if 3 girls should sit together in a back row on adjacent seats? Now, if all the seating arrangements are equally likely, what is the probability of 3 girls sitting together in a back row on adjacent seats? (1996 - 5 Marks)
19. If p and q are chosen randomly from the set $\{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$, with replacement, determine the probability that the roots of the equation $x^2 + px + q = 0$ are real. (1997 - 5 Marks)
20. Three players, A, B and C , toss a coin cyclically in that order (that is $A, B, C, A, B, C, A, B, \dots$) till a head shows. Let p be the probability that the coin shows a head. Let α, β and γ be, respectively, the probabilities that A, B and C gets the first head. Prove that $\beta = (1-p)\alpha$. Determine α, β and γ (in terms of p). (1998 - 8 Marks)
21. Eight players P_1, P_2, \dots, P_8 play a knock-out tournament. It is known that whenever the players P_i and P_j play, the player P_i will win if $i < j$. Assuming that the players are paired at random in each round, what is the probability that the player P_4 reaches the final? (1999 - 10 Marks)

22. A coin has probability p of showing head when tossed. It is tossed n times. Let p_n denote the probability that no two (or more) consecutive heads occur. Prove that $p_1=1$, $p_2=1-p^2$ and $p_n=(1-p) \cdot p_{n-1} + p(1-p) \cdot p_{n-2}$ for all $n \geq 3$. (2000 - 5 Marks)
23. An urn contains m white and n black balls. A ball is drawn at random and is put back into the urn along with k additional balls of the same colour as that of the ball drawn. A ball is again drawn at random. What is the probability that the ball drawn now is white? (2001 - 5 Marks)
24. An unbiased die, with faces numbered 1, 2, 3, 4, 5, 6, is thrown n times and the list of n numbers showing up is noted. What is the probability that, among the numbers 1, 2, 3, 4, 5, 6, only three numbers appear in this list? (2001 - 5 Marks)
25. A box contains N coins, m of which are fair and the rest are biased. The probability of getting a head when a fair coin is tossed is $1/2$, while it is $2/3$ when a biased coin is tossed. A coin is drawn from the box at random and is tossed twice. The first time it shows head and the second time it shows tail. What is the probability that the coin drawn is fair? (2002 - 5 Marks)
26. For a student to qualify, he must pass at least two out of three exams. The probability that he will pass the 1st exam is p . If he fails in one of the exams then the probability of his passing in the next exam is $\frac{p}{2}$ otherwise it remains the same. Find the probability that he will qualify. (2003 - 2 Marks)
27. A is targeting to B , B and C are targeting to A . Probability of hitting the target by A , B and C are $\frac{2}{3}$, $\frac{1}{2}$ and $\frac{1}{3}$ respectively. If A is hit then find the probability that B hits the target and C does not. (2003 - 2 Marks)
28. A and B are two independent events. C is event in which exactly one of A or B occurs. Prove that $P(C) \geq P(A \cup B)P(\bar{A} \cap \bar{B})$ (2004 - 2 Marks)
29. A box contains 12 red and 6 white balls. Balls are drawn from the box one at a time without replacement. If in 6 draws there are at least 4 white balls, find the probability that exactly one white is drawn in the next two draws. (binomial coefficients can be left as such) (2004 - 4 Marks)
30. A person goes to office either by car, scooter, bus or train, the probability of which being $\frac{1}{7}, \frac{3}{7}, \frac{2}{7}$ and $\frac{1}{7}$ respectively. Probability that he reaches office late, if he takes car, scooter, bus or train is $\frac{2}{9}, \frac{1}{9}, \frac{4}{9}$ and $\frac{1}{9}$ respectively. Given that he reached office in time, then what is the probability that he travelled by a car. (2005 - 2 Marks)
1. If $P(u_i) \propto i$, where $i = 1, 2, 3, \dots, n$, then $\lim_{n \rightarrow \infty} P(w) =$ (2006 - 5M, -2)
- (a) 1 (b) $2/3$ (c) $3/4$ (d) $1/4$
2. If $P(u_i) = c$, (c a constant) then $P(u_n/w) =$ (2006 - 5M, -2)
- (a) $\frac{2}{n+1}$ (b) $\frac{1}{n+1}$ (c) $\frac{n}{n+1}$ (d) $\frac{1}{2}$
3. Let $P(u_i) = \frac{1}{n}$, if n is even and E denotes the event of choosing even numbered urn, then the value of $P(w/E)$ is (2006 - 5M, -2)
- (a) $\frac{n+2}{2n+1}$ (b) $\frac{n+2}{2(n+1)}$ (c) $\frac{n}{n+1}$ (d) $\frac{1}{n+1}$

PASSAGE - 2

A fair die is tossed repeatedly until a six is obtained. Let X denote the number of tosses required. (2009)

4. The probability that $X = 3$ equals
- (a) $\frac{25}{216}$ (b) $\frac{25}{36}$ (c) $\frac{5}{36}$ (d) $\frac{125}{216}$
5. The probability that $X \geq 3$ equals
- (a) $\frac{125}{216}$ (b) $\frac{25}{36}$ (c) $\frac{5}{36}$ (d) $\frac{25}{216}$
6. The conditional probability that $X \geq 6$ given $X > 3$ equals
- (a) $\frac{125}{216}$ (b) $\frac{25}{216}$ (c) $\frac{5}{36}$ (d) $\frac{25}{36}$

PASSAGE - 3

Let U_1 and U_2 be two urns such that U_1 contains 3 white and 2 red balls, and U_2 contains only 1 white ball. A fair coin is tossed. If head appears then 1 ball is drawn at random from U_1 and put into U_2 . However, if tail appears then 2 balls are drawn at random from U_1 and put into U_2 . Now 1 ball is drawn at random from U_2 . (2011)

7. The probability of the drawn ball from U_2 being white is
- (a) $\frac{13}{30}$ (b) $\frac{23}{30}$ (c) $\frac{19}{30}$ (d) $\frac{11}{30}$
8. Given that the drawn ball from U_2 is white, the probability that head appeared on the coin is
- (a) $\frac{17}{23}$ (b) $\frac{11}{23}$ (c) $\frac{15}{23}$ (d) $\frac{12}{23}$

PASSAGE - 4

A box B_1 contains 1 white ball, 3 red balls and 2 black balls. Another box B_2 contains 2 white balls, 3 red balls and 4 black balls. A third box B_3 contains 3 white balls, 4 red balls and 5 black balls.

9. If 1 ball is drawn from each of the boxes B_1 , B_2 and B_3 , the probability that all 3 drawn balls are of the same colour is (JEE Adv. 2013)
- (a) $\frac{82}{648}$ (b) $\frac{90}{648}$ (c) $\frac{558}{648}$ (d) $\frac{566}{648}$

10. If 2 balls are drawn (without replacement) from a randomly selected box and one of the balls is white and the other ball is red, the probability that these 2 balls are drawn from box B_2 is

- (a) $\frac{116}{181}$ (b) $\frac{126}{181}$
 (c) $\frac{65}{181}$ (d) $\frac{55}{181}$

G Comprehension Based Questions**PASSAGE - 1**

There are n urns, each of these contain $n+1$ balls. The i th urn contains i white balls and $(n+1-i)$ red balls. Let u_i be the event of selecting i th urn, $i = 1, 2, 3, \dots, n$ and w the event of getting a white ball.

PASSAGE - 5

Box 1 contains three cards bearing numbers 1, 2, 3; box 2 contains five cards bearing numbers 1, 2, 3, 4, 5; and box 3 contains seven cards bearing numbers 1, 2, 3, 4, 5, 6, 7. A card is drawn from each of the boxes. Let x_i be number on the card drawn from the i^{th} box, $i=1, 2, 3$.

(JEE Adv. 2014)

11. The probability that $x_1 + x_2 + x_3$ is odd, is

- (a) $\frac{29}{105}$ (b) $\frac{53}{105}$
 (c) $\frac{57}{105}$ (d) $\frac{1}{2}$

12. The probability that x_1, x_2, x_3 are in an arithmetic progression, is

- (a) $\frac{9}{105}$ (b) $\frac{10}{105}$
 (c) $\frac{11}{105}$ (d) $\frac{7}{105}$

PASSAGE - 6

Let n_1 and n_2 be the number of red and black balls, respectively, in box I. Let n_3 and n_4 be the number of red and black balls, respectively, in box II.

(JEE Adv. 2015)

13. One of the two boxes, box I and box II, was selected at random and a ball was drawn randomly out of this box. The ball was found to be red. If the probability that this red ball

was drawn from box II is $\frac{1}{3}$, then the correct option(s) with

the possible values of n_1, n_2, n_3 and n_4 is(are)

- (a) $n_1 = 3, n_2 = 3, n_3 = 5, n_4 = 15$
 (b) $n_1 = 3, n_2 = 6, n_3 = 10, n_4 = 50$
 (c) $n_1 = 8, n_2 = 6, n_3 = 5, n_4 = 20$
 (d) $n_1 = 6, n_2 = 12, n_3 = 5, n_4 = 20$

14. A ball is drawn at random from box I and transferred to box II. If the probability of drawing a red ball from box I, after

this transfer, is $\frac{1}{3}$, then the correct option(s) with the pos-

sible values of n_1 and n_2 is(are)

- (a) $n_1 = 4$ and $n_2 = 6$
 (b) $n_1 = 2$ and $n_2 = 3$
 (c) $n_1 = 10$ and $n_2 = 20$
 (d) $n_1 = 3$ and $n_2 = 6$

PASSAGE - 7

Football teams T_1 and T_2 have to play two games against each other. It is assumed that the outcomes of the two games are

independent. The probabilities of T_1 winning, drawing and losing a game against T_2 are $\frac{1}{2}, \frac{1}{6}$ and $\frac{1}{3}$ respectively. Each team gets 3 points for a win, 1 point for a draw and 0 point for a loss in a game. Let X and Y denote the total points scored by teams T_1 and T_2 respectively after two games.

15. $P(X > Y)$ is

- (a) $\frac{1}{4}$ (b) $\frac{5}{12}$ (c) $\frac{1}{2}$ (d) $\frac{7}{12}$

16. $P(X = Y)$ is

- (a) $\frac{11}{36}$ (b) $\frac{1}{3}$ (c) $\frac{13}{36}$ (d) $\frac{1}{2}$

PASSAGE - 8

There are five students S_1, S_2, S_3, S_4 and S_5 in a music class and for them there are five seats R_1, R_2, R_3, R_4 and R_5 arranged in a row, where initially the seat R_i is allotted to the student $S_i, i = 1, 2, 3, 4, 5$. But, on the examination day, the five students are randomly allotted the five seats.

17. The probability that, on examination day, the student S_1 gets the previously allotted seat R_1 , and **NONE** of the remaining students gets the seat previously allotted to him/her is

- (a) $\frac{3}{40}$ (b) $\frac{1}{8}$ (c) $\frac{7}{40}$ (d) $\frac{1}{5}$

18. For $i = 1, 2, 3, 4$, let T_i denote the event that the students S_i and S_{i+1} do **NOT** sit adjacent to each other on the day of the examination. Then, the probability of the event $T_1 \cap T_2 \cap T_3 \cap T_4$ is

- (a) $\frac{1}{15}$ (b) $\frac{1}{10}$ (c) $\frac{7}{60}$ (d) $\frac{1}{5}$

**Assertion & Reason Type Questions**

1. Let H_1, H_2, \dots, H_n be mutually exclusive and exhaustive events with $P(H_i) > 0, i = 1, 2, \dots, n$. Let E be any other event with $0 < P(E) < 1$.

STATEMENT-1:

$P(H_i | E) > P(E | H_i) \cdot P(H_i)$ for $i = 1, 2, \dots, n$ because

STATEMENT-2: $\sum_{i=1}^n P(H_i) = 1$. (2007 - 3 marks)

- (a) Statement-1 is True, statement-2 is True; Statement-2 is a correct explanation for Statement-1.
 (b) Statement-1 is True, Statement-2 is True; Statement-2 is NOT a correct explanation for Statement-1.
 (c) Statement-1 is True, Statement-2 is False.
 (d) Statement-1 is False, Statement-2 is True.

2. Consider the system of equations $ax + by = 0$; $cx + dy = 0$, where $a, b, c, d \in \{0, 1\}$

STATEMENT - 1 : The probability that the system of equations has a unique solution is $\frac{3}{8}$.

and

STATEMENT - 2 : The probability that the system of equations has a solution is 1. (2008)

- (a) STATEMENT - 1 is True, STATEMENT - 2 is True; STATEMENT - 2 is a correct explanation for STATEMENT - 1
- (b) STATEMENT - 1 is True, STATEMENT - 2 is True; STATEMENT - 2 is NOT a correct explanation for STATEMENT - 1
- (c) STATEMENT - 1 is True, STATEMENT - 2 is False
- (d) STATEMENT - 1 is False, STATEMENT - 2 is True

I Integer Value Correct Type

1. Of the three independent events E_1, E_2 and E_3 , the probability that only E_1 occurs is α , only E_2 occurs is β and only E_3 occurs is γ . Let the probability p that none of events E_1, E_2 or E_3 occurs satisfy the equations $(\alpha - 2\beta)p = \alpha\beta$ and $(\beta - 3\gamma)p = 2\beta\gamma$. All the given probabilities are assumed to lie in the interval $(0, 1)$. (JEE Adv. 2013)

$$\text{Then } \frac{\text{Probability of occurrence of } E_1}{\text{Probability of occurrence of } E_3}$$

2. The minimum number of times a fair coin needs to be tossed, so that the probability of getting at least two heads is at least 0.96, is (JEE Adv. 2015)
3. Let S be the sample space of all 3×3 matrices with entries from the set $\{0, 1\}$. Let the events E_1 and E_2 be given by $E_1 = \{A \in S : \det A = 0\}$ and $E_2 = \{A \in S : \text{sum of entries of } A \text{ is 7}\}$. If a matrix is chosen at random from S , then the conditional probability $P(E_1|E_2)$ equals _____. (JEE Adv. 2019)

Section-B JEE Main / AIEEE

1. A problem in mathematics is given to three students A, B, C and their respective probability of solving the problem is $\frac{1}{2}, \frac{1}{3}$ and $\frac{1}{4}$. Probability that the problem is solved is [2002]
- (a) $\frac{3}{4}$ (b) $\frac{1}{2}$ (c) $\frac{2}{3}$ (d) $\frac{1}{3}$
2. A and B are events such that $P(A \cup B) = 3/4$, $P(A \cap B) = 1/4$, $P(\bar{A}) = 2/3$ then $P(\bar{A} \cap B)$ is [2002]
- (a) $5/12$ (b) $3/8$ (c) $5/8$ (d) $1/4$
3. A dice is tossed 5 times. Getting an odd number is considered a success. Then the variance of distribution of success is [2002]
- (a) $8/3$ (b) $3/8$ (c) $4/5$ (d) $5/4$
4. The mean and variance of a random variable X having binomial distribution are 4 and 2 respectively, then $P(X=1)$ is [2003]
- (a) $\frac{1}{4}$ (b) $\frac{1}{32}$ (c) $\frac{1}{16}$ (d) $\frac{1}{8}$
5. Events A, B, C are mutually exclusive events such that $P(A) = \frac{3x+1}{3}$, $P(B) = \frac{1-x}{4}$ and $P(C) = \frac{1-2x}{2}$ The set of possible values of x are in the interval. [2003]
- (a) $[0, 1]$ (b) $\left[\frac{1}{3}, \frac{1}{2}\right]$ (c) $\left[\frac{1}{3}, \frac{2}{3}\right]$ (d) $\left[\frac{1}{3}, \frac{13}{3}\right]$
6. Five horses are in a race. Mr. A selects two of the horses at random and bets on them. The probability that Mr. A selected the winning horse is [2003]
- (a) $\frac{2}{5}$ (b) $\frac{4}{5}$ (c) $\frac{3}{5}$ (d) $\frac{1}{5}$
7. The probability that A speaks truth is $\frac{4}{5}$, while the probability for B is $\frac{3}{4}$. The probability that they contradict each other when asked to speak on a fact is [2004]
- (a) $\frac{4}{5}$ (b) $\frac{1}{5}$ (c) $\frac{7}{20}$ (d) $\frac{3}{20}$
8. A random variable X has the probability distribution:
- | | | | | | | | | |
|-------|-----|-----|-----|-----|-----|-----|-----|-----|
| X: | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| p(X): | 0.2 | 0.2 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 |
- For the events $E = \{X \text{ is a prime number}\}$ and $F = \{X < 4\}$, the $P(E \cup F)$ is [2004]
- (a) 0.50 (b) 0.77 (c) 0.35 (d) 0.87
9. The mean and the variance of a binomial distribution are 4 and 2 respectively. Then the probability of 2 successes is [2004]
- (a) $\frac{28}{256}$ (b) $\frac{219}{256}$ (c) $\frac{128}{256}$ (d) $\frac{37}{256}$

10. Three houses are available in a locality. Three persons apply for the houses. Each applies for one house without consulting others. The probability that all the three apply for the same house is [2005]

(a) $\frac{2}{9}$ (b) $\frac{1}{9}$ (c) $\frac{8}{9}$ (d) $\frac{7}{9}$

11. A random variable X has Poisson distribution with mean 2. Then $P(X > 1.5)$ equals [2005]

(a) $\frac{2}{e^2}$ (b) 0 (c) $1 - \frac{3}{e^2}$ (d) $\frac{3}{e^2}$

12. Let A and B be two events such that $P(\overline{A \cup B}) = \frac{1}{6}$,

$P(A \cap B) = \frac{1}{4}$ and $P(\overline{A}) = \frac{1}{4}$, where \overline{A} stands for complement of event A . Then events A and B are

- (a) equally likely and mutually exclusive [2005]
 (b) equally likely but not independent
 (c) independent but not equally likely
 (d) mutually exclusive and independent

13. At a telephone enquiry system the number of phone calls regarding relevant enquiry follow Poisson distribution with an average of 5 phone calls during 10 minute time intervals. The probability that there is at the most one phone call during a 10-minute time period is [2006]

(a) $\frac{6}{5^5}$ (b) $\frac{5}{6}$ (c) $\frac{6}{55}$ (d) $\frac{6}{e^5}$

14. Two aeroplanes I and II bomb a target in succession. The probabilities of I and II scoring a hit correctly are 0.3 and 0.2, respectively. The second plane will bomb only if the first misses the target. The probability that the target is hit by the second plane is [2007]

(a) 0.2 (b) 0.7 (c) 0.06 (d) 0.14.

15. A pair of fair dice is thrown independently three times. The probability of getting a score of exactly 9 twice is [2007]

(a) $8/729$ (b) $8/243$ (c) $1/729$ (d) $8/9$.

16. It is given that the events A and B are such that

$P(A) = \frac{1}{4}$, $P(A|B) = \frac{1}{2}$ and $P(B|A) = \frac{2}{3}$. Then $P(B)$ is [2008]

(a) $\frac{1}{6}$ (b) $\frac{1}{3}$ (c) $\frac{2}{3}$ (d) $\frac{1}{2}$

17. A die is thrown. Let A be the event that the number obtained is greater than 3. Let B be the event that the number obtained is less than 5. Then $P(A \cup B)$ is [2008]

(a) $\frac{3}{5}$ (b) 0 (c) 1 (d) $\frac{2}{5}$

18. In a binomial distribution $B\left(n, p = \frac{1}{4}\right)$, if the probability of

at least one success is greater than or equal to $\frac{9}{10}$, then n is greater than: [2009]

(a) $\frac{1}{\log_{10} 4 + \log_{10} 3}$ (b) $\frac{9}{\log_{10} 4 - \log_{10} 3}$

(c) $\frac{4}{\log_{10} 4 - \log_{10} 3}$ (d) $\frac{1}{\log_{10} 4 - \log_{10} 3}$

19. One ticket is selected at random from 50 tickets numbered 00, 01, 02, ..., 49. Then the probability that the sum of the digits on the selected ticket is 8, given that the product of these digits is zero, equals: [2009]

(a) $\frac{1}{7}$ (b) $\frac{5}{14}$ (c) $\frac{1}{50}$ (d) $\frac{1}{14}$

20. Four numbers are chosen at random (without replacement) from the set {1, 2, 3, ..., 20}. [2010]

Statement -1: The probability that the chosen numbers when

arranged in some order will form an AP is $\frac{1}{85}$.

Statement -2 : If the four chosen numbers form an AP, then the set of all possible values of common difference is $\{\pm 1, \pm 2, \pm 3, \pm 4, \pm 5\}$.

- (a) Statement -1 is true, Statement -2 is true ; Statement -2 is not a correct explanation for Statement -1
 (b) Statement -1 is true, Statement -2 is false
 (c) Statement -1 is false, Statement -2 is true.
 (d) Statement -1 is true, Statement -2 is true ; Statement -2 is a correct explanation for Statement -1.

21. An urn contains nine balls of which three are red, four are blue and two are green. Three balls are drawn at random without replacement from the urn. The probability that the three balls have different colours is [2010]

(a) $\frac{2}{7}$ (b) $\frac{1}{21}$

(c) $\frac{2}{23}$ (d) $\frac{1}{3}$

22. Consider 5 independent Bernoulli's trials each with probability of success p . If the probability of at least one failure is greater than or equal to $\frac{31}{32}$, then p lies in the interval [2011]

(a) $\left(\frac{3}{4}, \frac{11}{12}\right]$ (b) $\left[0, \frac{1}{2}\right]$

(c) $\left(\frac{11}{12}, 1\right]$ (d) $\left(\frac{1}{2}, \frac{3}{4}\right]$

23. If C and D are two events such that $C \subset D$ and $P(D) \neq 0$, then the correct statement among the following is [2011]

(a) $P(C|D) \geq P(C)$ (b) $P(C|D) < P(C)$

(c) $P(C|D) = \frac{P(D)}{P(C)}$ (d) $P(C|D) = P(C)$

24. Three numbers are chosen at random without replacement from $\{1, 2, 3, \dots, 8\}$. The probability that their minimum is 3, given that their maximum is 6, is : [2012]

(a) $\frac{3}{8}$ (b) $\frac{1}{5}$ (c) $\frac{1}{4}$ (d) $\frac{2}{5}$

25. A multiple choice examination has 5 questions. Each question has three alternative answers of which exactly one is correct. The probability that a student will get 4 or more correct answers just by guessing is: [JEE M 2013]

(a) $\frac{17}{3^5}$ (b) $\frac{13}{3^5}$ (c) $\frac{11}{3^5}$ (d) $\frac{10}{3^5}$

26. Let A and B be two events such that $P(\overline{A \cup B}) = \frac{1}{6}$, $P(\overline{A \cap B}) = \frac{1}{4}$ and $P(\overline{A}) = \frac{1}{4}$, where \overline{A} stands for the complement of the event A . Then the events A and B are [JEE M 2014]

- (a) independent but not equally likely.
 (b) independent and equally likely.
 (c) mutually exclusive and independent.
 (d) equally likely but not independent.

27. If 12 identical balls are to be placed in 3 identical boxes, then the probability that one of the boxes contains exactly 3 balls is: [JEE M 2015]

(a) $220\left(\frac{1}{3}\right)^{12}$ (b) $22\left(\frac{1}{3}\right)^{11}$
 (c) $\frac{55}{3}\left(\frac{2}{3}\right)^{11}$ (d) $55\left(\frac{2}{3}\right)^{10}$

28. Let two fair six-faced dice A and B be thrown simultaneously. If E_1 is the event that die A shows up four, E_2 is the event that die B shows up two and E_3 is the event that the sum of numbers on both dice is odd, then which of the following statements is NOT true ? [JEE M 2016]

- (a) E_1 and E_3 are independent.
 (b) E_1 , E_2 and E_3 are independent.
 (c) E_1 and E_2 are independent.
 (d) E_2 and E_3 are independent.

29. A box contains 15 green and 10 yellow balls. If 10 balls are randomly drawn, one-by-one, with replacement, then the variance of the number of green balls drawn is : [JEE M 2017]

(a) $\frac{6}{25}$ (b) $\frac{12}{5}$ (c) 6 (d) 4

30. If two different numbers are taken from the set $\{0, 1, 2, 3, \dots, 10\}$, then the probability that their sum as well as absolute difference are both multiple of 4, is : [JEE M 2017]

(a) $\frac{7}{55}$ (b) $\frac{6}{55}$
 (c) $\frac{12}{55}$ (d) $\frac{14}{55}$

31. For three events A, B and C,
 $P(\text{Exactly one of } A \text{ or } B \text{ occurs})$
 $= P(\text{Exactly one of } B \text{ or } C \text{ occurs})$

$= P(\text{Exactly one of } C \text{ or } A \text{ occurs}) = \frac{1}{4}$ and

$P(\text{All the three events occur simultaneously}) = \frac{1}{16}$.

Then the probability that at least one of the events occurs, is :

[JEE M 2017]

(a) $\frac{3}{16}$ (b) $\frac{7}{32}$ (c) $\frac{7}{16}$ (d) $\frac{7}{64}$

32. A bag contains 4 red and 6 black balls. A ball is drawn at random from the bag, its colour is observed and this ball along with two additional balls of the same colour are returned to the bag. If now a ball is drawn at random from the bag, then the probability that this drawn ball is red, is : [JEE M 2018]

(a) $\frac{2}{5}$ (b) $\frac{1}{5}$ (c) $\frac{3}{4}$ (d) $\frac{3}{10}$

33. Two cards are drawn successively with replacement from a well-shuffled deck of 52 cards. Let X denote the random variable of number of faces obtained in the two drawn cards. Then $P(X=1) + P(X=2)$ equals:

[JEE M 2019 – 9 Jan (M)]

(a) 49/169 (b) 52/169
 (c) 24/169 (d) 25/169

34. Four persons can hit a target correctly with probabilities $\frac{1}{2}, \frac{1}{3}, \frac{1}{4}$ and $\frac{1}{8}$ respectively. If all hit at the target independently, then the probability that the target would be hit, is: [JEE M 2019 – 9 April (M)]

(a) $\frac{25}{192}$ (b) $\frac{7}{32}$
 (c) $\frac{1}{192}$ (d) $\frac{25}{32}$