

Circle Assignment

Gautam Singh

Abstract—This document contains the solution to Question 6 of Exercise 4 in Chapter 10 of the class 9 NCERT textbook.

- 1) A circular park of radius 20 m is situated in a colony. Three boys Ankur, Syed and David are sitting at equal distance on its boundary each having a toy telephone in his hands to talk each other. Find the length of the string of each phone.

Solution: Let the position vectors of the boys be

$$\mathbf{A} = \begin{pmatrix} r \\ 0 \end{pmatrix}, \mathbf{S} = \begin{pmatrix} r \cos \beta \\ r \sin \beta \end{pmatrix}, \mathbf{D} = \begin{pmatrix} r \cos \gamma \\ r \sin \gamma \end{pmatrix} \quad (1)$$

where

$$\beta, \gamma \in (0, 2\pi) \quad (2)$$

We have,

$$\|\mathbf{A} - \mathbf{S}\|^2 = \|\mathbf{A} - \mathbf{D}\|^2 \quad (3)$$

$$\Rightarrow \mathbf{A}^\top \mathbf{S} = \mathbf{A}^\top \mathbf{D} \quad (4)$$

$$\Rightarrow \cos \beta = \cos \gamma \quad (5)$$

$$\Rightarrow \beta = 2n\pi \pm \gamma \quad (6)$$

where $n \in \mathbb{Z}$. From (2), we force $n = 1$ and consider the negative sign to get

$$\beta + \gamma = 2\pi \quad (7)$$

Therefore, using (7)

$$\|\mathbf{A} - \mathbf{S}\|^2 = \|\mathbf{S} - \mathbf{D}\|^2 \quad (8)$$

$$\Rightarrow \mathbf{A}^\top \mathbf{S} = \mathbf{D}^\top \mathbf{S} \quad (9)$$

$$\Rightarrow \cos \beta = \cos(\beta - \gamma) \quad (10)$$

$$\Rightarrow 2\beta - 2\pi = 2m\pi \pm \beta \quad (11)$$

$$\Rightarrow 2\beta \pm \beta = 2k\pi \quad (12)$$

where $k \in \mathbb{Z}$. From (2), we can only consider the plus sign and $k \in \{1, 2\}$ to get

$$\beta, \gamma \in \left\{ \frac{2\pi}{3}, \frac{4\pi}{3} \right\} \quad (13)$$

Therefore, the length of the thread from (13)

is

$$\|\mathbf{S} - \mathbf{D}\| = \left\| r \begin{pmatrix} \cos \beta - \cos \gamma \\ \sin \beta - \sin \gamma \end{pmatrix} \right\| \quad (14)$$

$$= r\sqrt{3} \quad (15)$$

Here, $r = 20$ m. Thus, the length is $20\sqrt{3}$ m. The situation is demonstrated in Fig. 1, plotted by the Python code `codes/equilateral.py`. The values used for construction are shown in Table 1.

Parameter	Value
r	20
β	$\frac{2\pi}{3}$
γ	$\frac{4\pi}{3}$
\mathbf{O}	$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$

TABLE 1: Parameters used in the construction of Fig. 1.




Fig. 1: ASD is an equilateral triangle of side $20\sqrt{3}$ m.