

# Linear Forms

G V V Sharma\*

## CONTENTS

<b>1</b>	<b>Definitions</b>	<b>1</b>
1.1	Two Dimensions . . . . .	1
1.2	Three Dimensions . . . . .	1

## 2 Examples

*Abstract—This manual provides a simple introduction to linear forms like lines and planes, based on the NCERT textbooks from Class 6-12.*

1.1.6. The equation of the line perpendicular to (1.1.1.1) and passing through the point  $\mathbf{P}$  is given by

$$\mathbf{m}^\top (\mathbf{x} - \mathbf{P}) = 0 \quad (1.1.6.1)$$

1.1.7. The foot of the perpendicular from  $\mathbf{P}$  to the line in (1.1.1.1) is given by

$$\begin{pmatrix} \mathbf{m} & \mathbf{n} \end{pmatrix}^\top \mathbf{x} = \begin{pmatrix} \mathbf{m}^\top \mathbf{P} \\ c \end{pmatrix} \quad (1.1.7.1)$$

## 1 DEFINITIONS

### 1.1 Two Dimensions

1.1.1. The equation of a line is given by

$$\mathbf{n}^\top \mathbf{x} = c \quad (1.1.1.1)$$

where  $\mathbf{n}$  is the normal vector of the line.

1.1.2. The parametric equation of a line is given by

$$\mathbf{x} = \mathbf{A} + \lambda \mathbf{m} \quad (1.1.2.1)$$

where  $\mathbf{m}$  is the direction vector of the line and  $\mathbf{A}$  is any point on the line.

1.1.3. The distance from a point  $\mathbf{P}$  to the line in (1.1.1.1) is given by

$$d = \frac{|\mathbf{n}^\top \mathbf{P} - c|}{\|\mathbf{n}\|} \quad (1.1.3.1)$$

1.1.4. The distance from the origin to the line in (1.1.1.1) is given by

$$d = \frac{|c|}{\|\mathbf{n}\|} \quad (1.1.4.1)$$

1.1.5. The distance between the parallel lines

$$\mathbf{n}^\top \mathbf{x} = c_1 \quad \mathbf{n}^\top \mathbf{x} = c_2 \quad (1.1.5.1)$$

is given by

$$d = \frac{|c_1 - c_2|}{\|\mathbf{n}\|} \quad (1.1.5.2)$$

### 1.2 Three Dimensions

1.2.1. The equation of a line is given by (1.1.2.1)

1.2.2. The equation of a plane is given by (1.1.1.1)

1.2.3. The distance from the origin to the line in (1.1.1.1) is given by (1.1.4.1)

1.2.4. The equation of the line perpendicular to (1.1.1.1) and passing through the point  $\mathbf{P}$  is given by

$$\mathbf{m}^\top (\mathbf{x} - \mathbf{P}) = 0 \quad (1.2.4.1)$$

1.2.5. The foot of the perpendicular from  $\mathbf{P}$  to the line in (1.1.1.1) is given by

$$\begin{pmatrix} \mathbf{m} & \mathbf{n} \end{pmatrix}^\top \mathbf{x} = \begin{pmatrix} \mathbf{m}^\top \mathbf{P} \\ c \end{pmatrix} \quad (1.2.5.1)$$

1.2.6. The distance from a point  $\mathbf{P}$  to the line in (1.1.2.1) is given by

$$d = \left\| \mathbf{A} - \mathbf{P} - \frac{\mathbf{m}^\top (\mathbf{A} - \mathbf{P})}{\|\mathbf{m}\|^2} \mathbf{m} \right\| \quad (1.2.6.1)$$

**Solution:**

$$d(\lambda) = \|\mathbf{A} + \lambda \mathbf{m} - \mathbf{P}\| \quad (1.2.6.2)$$

$$\Rightarrow d^2(\lambda) = \|\mathbf{A} + \lambda \mathbf{m} - \mathbf{P}\|^2 \quad (1.2.6.3)$$

which can be simplified to obtain

$$d^2(\lambda) = \lambda^2 \|\mathbf{m}\|^2 + 2\lambda \mathbf{m}^\top (\mathbf{A} - \mathbf{P}) + \|\mathbf{A} - \mathbf{P}\|^2 \quad (1.2.6.4)$$

\*The author is with the Department of Electrical Engineering, Indian Institute of Technology, Hyderabad 502285 India e-mail: gadepall@iith.ac.in. All content in this manual is released under GNU GPL. Free and open source.

Now,  $d(\lambda)$  is smallest when the above quadratic equation has a single root, i.e.

$$\lambda = -\frac{\mathbf{m}^\top (\mathbf{A} - \mathbf{P})}{\|\mathbf{m}\|^2} \quad (1.2.6.5)$$

From (1.2.6.2) and (1.2.6.5),

1.2.7. The distance between the parallel planes (1.1.5.1) is given by (1.1.5.2).

## 2 EXAMPLES

2.1.